



This document is published at:

Méndez-Bermúdez, J.A., Reyes, R., Rodríguez, J.M., Sigarreta, J.M. (2018). Hyperbolicity on Graph Operators. *Symmetry*, 10 (9), 360.

DOI: <https://doi.org/10.3390/sym10090360>



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

# Hyperbolicity on Graph Operators

J. A. Méndez-Bermúdez <sup>1</sup>, Rosalío Reyes <sup>2</sup>, José M. Rodríguez <sup>2</sup> and José M. Sigarreta <sup>1,3,\*</sup>

<sup>1</sup> Instituto de Física, Benemérita Universidad Autónoma de Puebla, Apartado Postal J-48, Puebla 72570, Mexico; jmendezb@ifuap.buap.mx

<sup>2</sup> Departamento de Matemáticas, Universidad Carlos III de Madrid, Avenida de la Universidad 30, 28911 Leganés, Madrid, Spain; rreyes@math.uc3m.es (R.R.); jomaro@math.uc3m.es (J.M.R.)

<sup>3</sup> Facultad de Matemáticas, Universidad Autónoma de Guerrero, Carlos E. Adame No.54 Col. Garita, Acapulco Gro. 39650, Mexico

\* Correspondence: josemariasigarretaalmira@hotmail.com; Tel.: +52-1-7441592272

Received: 24 July 2018; Accepted: 22 August 2018; Published: 24 August 2018



**Abstract:** A graph operator is a mapping  $F : \Gamma \rightarrow \Gamma'$ , where  $\Gamma$  and  $\Gamma'$  are families of graphs. The different kinds of graph operators are an important topic in Discrete Mathematics and its applications. The symmetry of this operations allows us to prove inequalities relating the hyperbolicity constants of a graph  $G$  and its graph operators: line graph,  $\Lambda(G)$ ; subdivision graph,  $S(G)$ ; total graph,  $T(G)$ ; and the operators  $R(G)$  and  $Q(G)$ . In particular, we get relationships such as  $\delta(G) \leq \delta(R(G)) \leq \delta(G) + 1/2$ ,  $\delta(\Lambda(G)) \leq \delta(Q(G)) \leq \delta(\Lambda(G)) + 1/2$ ,  $\delta(S(G)) \leq 2\delta(R(G)) \leq \delta(S(G)) + 1$  and  $\delta(R(G)) - 1/2 \leq \delta(\Lambda(G)) \leq 5\delta(R(G)) + 5/2$  for every graph which is not a tree. Moreover, we also derive some inequalities for the Gromov product and the Gromov product restricted to vertices.

**Keywords:** graph operators; gromov hyperbolicity; geodesics

## 1. Introduction

In [1], J. Krausz introduced the concept of graph operators. These operators have applications in studies of graph dynamics (see [2,3]) and topological indices (see [4–6]). Many large graphs can be obtained by applying graph operators on smaller ones, thus some of their properties are strongly related. Motivated by the above works, we study here the hyperbolicity constant of several graph operators.

Along this paper, we denote by  $G = (V(G), E(G))$  a connected simple graph with edges of length 1 (unless edge lengths are explicitly given) and  $V \neq \emptyset$ . Given an edge  $e = uv \in E(G)$  with endpoints  $u$  and  $v$ , we write  $V(e) = \{u, v\}$ . Next, we recall the definition of some of the main graph operators.

The *line graph*,  $\Lambda(G)$ , is the graph constructed from  $G$  with vertices the set of edges of  $G$ , and two vertices are adjacent if and only if their corresponding edges are incident in  $G$ .

The *subdivision graph*,  $S(G)$ , is the graph constructed from  $G$  substituting each of its edges by a path of length 2.

The graph  $Q(G)$  is the graph constructed from  $S(G)$  by adding edges between adjacent vertices in  $\Lambda(G)$ .

The graph  $R(G)$  is constructed from  $S(G)$  by adding edges between adjacent vertices in  $G$ .

The *total graph*,  $T(G)$ , is constructed from  $S(G)$  by adding edges between adjacent vertices in  $G$  or  $\Lambda(G)$ .

We define:

$$E_E(G) := \{ \{e_1, e_2\} : e_1, e_2 \in E(G), e_1 \neq e_2, |V(e_1) \cap V(e_2)| = 1 \},$$

and

$$E_V(G) := \{ \{e, u\} : e \in E(G), u \in V(e) \}.$$

So, we have the following:

$$\Lambda(G) := (E(G), E_E(G)).$$

$$S(G) := (V(G) \cup E(G), E_V(G)).$$

$$T(G) := (V(G) \cup E(G), E(G) \cup E_E(G) \cup E_V(G)).$$

$$R(G) := (V(G) \cup E(G), E(G) \cup E_V(G)).$$

$$Q(G) := (V(G) \cup E(G), E_E(G) \cup E_V(G)).$$

The Gromov hyperbolic spaces have multiple applications both theoretical and practical (see [7–10]). A space is geodesic if any two points in it can be joined by a curve whose length is the distance between them. In this paper we will consider a graph  $G$  as a geodesic metric space and any geodesic joining  $x$  and  $y$  will be denoted by  $[xy]$ .

Let  $X$  be a geodesic metric space and  $x, y, z \in X$ . A *geodesic triangle* with vertices  $x, y, z$ , denoted by  $T = \{x, y, z\}$ , is the union of three geodesics  $[xy]$ ,  $[yz]$  and  $[zx]$ . We write also  $T = \{[xy], [yz], [zx]\}$ . If the  $\delta$ -neighborhood of the union of any two sides of  $T$  contains the other side, we say that  $T$  is  $\delta$ -thin. We define  $\delta(T) := \inf\{\delta \geq 0 : T \text{ is } \delta\text{-thin}\}$ . The space  $X$  is  $\delta$ -hyperbolic if all geodesic triangles  $T$  in  $X$  are  $\delta$ -thin. Let us denote the sharp hyperbolicity constant of  $X$ , by  $\delta(X)$ , i.e.,  $\delta(X) := \sup\{\delta(T) : T \text{ is a geodesic triangle in } X\}$ .  $X$  is *Gromov hyperbolic* if  $X$  is  $\delta$ -hyperbolic for some  $\delta \geq 0$ ; then  $X$  is Gromov hyperbolic if and only if  $\delta(X) < \infty$ .

In this paper we prove inequalities relating the hyperbolicity constants of a graph  $G$  and its graph operators  $\Lambda(G)$ ,  $S(G)$ ,  $T(G)$ ,  $R(G)$  and  $Q(G)$ , using their symmetries.

## 2. Definitions and Background

There are several equivalent definitions for Gromov hyperbolicity (see, e.g., [11–13]), in particular, the definition that we use in this work has an important geometric meaning and serves as a basis for multiple applications (see [14–19]).

Given a graph  $G$ , the *Gromov product* of  $q_1, q_2 \in G$  with base point  $q_0 \in G$  is defined as

$$(q_1, q_2)_{q_0} := \frac{1}{2} (d(q_1, q_0) + d(q_2, q_0) - d(q_1, q_2)).$$

For every Gromov hyperbolic graph  $G$ , we have

$$(q_1, q_3)_{q_0} \geq \min \{ (q_1, q_2)_{q_0}, (q_2, q_3)_{q_0} \} - \delta \quad (1)$$

for every  $q_0, q_1, q_2, q_3 \in G$  and some constant  $\delta \geq 0$  ([12,13]).

We denote by  $\delta^*(G)$  the sharp constant for the inequality (1), i.e.,

$$\delta^*(G) := \sup \{ \min \{ (q_1, q_2)_{q_0}, (q_2, q_3)_{q_0} \} - (q_1, q_3)_{q_0} : q_0, q_1, q_2, q_3 \in G \}.$$

Indeed, our definition of Gromov hyperbolicity is equivalent to (1); furthermore, we have  $\delta^*(G) \leq 4\delta(G)$  and  $\delta(G) \leq 3\delta^*(G)$  ([12,13]). In [20] (Proposition II.20) we found the following improvement of the previous inequality:  $\delta^*(G) \leq 2\delta(G)$ .

We denote by  $\delta_v^*(G)$  the constant of hyperbolicity of the Gromov product restricted to the vertices of  $G$ , i.e.,

$$\delta_v^*(G) := \sup \{ \min \{ (q_1, q_2)_{q_0}, (q_2, q_3)_{q_0} \} - (q_1, q_3)_{q_0} : q_0, q_1, q_2, q_3 \in V(G) \}.$$

## 3. Main Results

The following result is immediate from the definition of  $S(G)$ .

**Proposition 1.** Let  $G$  be a graph. Then

$$\delta(S(G)) = 2\delta(G), \quad \delta^*(S(G)) = 2\delta^*(G).$$

We remark that the equality is not true for  $\delta_v^*(G)$  (e.g.,  $S(C_5) = C_{10}$  but  $2\delta_v^*(C_5) = 1 \neq 2 = \delta_v^*(S(G))$ ), but inequalities may apply. The next result appears in [21].

**Theorem 1.** Let  $B = (V_0 \cup V_1, E)$  be a bipartite graph. We have  $\delta_B(V_i) \leq \delta_v^*(B) \leq \delta_B(V_i) + 2$ , where

$$\delta_B(V_i) = \sup\{\min\{(x, y)_w, (y, z)_w\} - (x, z)_w : x, y, z, w \in V_i\}$$

for every  $i \in \{1, 2\}$ .

**Corollary 1.** Let  $G$  be a graph. Then

$$2\delta_v^*(G) \leq \delta_v^*(S(G)) \leq 2\delta_v^*(G) + 2.$$

**Proof.** Note that  $S(G)$  can be considered as a bipartite graph, where  $V(S(G)) = V(G) \cup V(\Lambda(G))$ . Theorem 1 gives  $\delta_{S(G)}(V(G)) \leq \delta_v^*(S(G)) \leq \delta_{S(G)}(V(G)) + 2$ . Since  $\delta_{S(G)}(V(G)) = 2\delta_v^*(G)$ , the desired inequalities hold.  $\square$

**Proposition 2.** Let  $G$  be a graph. Then

$$\delta_v^*(G) \leq \delta^*(G) \leq \delta_v^*(G) + 3.$$

**Proof.** The inequality  $\delta_v^*(G) \leq \delta^*(G)$  is direct. Let us prove the other inequality.

For every  $q_0, q_1, q_2 \in G$  there are  $q'_0, q'_1, q'_2 \in V(G)$  such that  $d(q_i, q'_i) \leq 1/2$  for  $i = 0, 1, 2$ . Then

$$\begin{aligned} |(q_1, q_2)_{q_0} - (q'_1, q'_2)_{q'_0}| &= \frac{1}{2} |d(q_0, q_1) + d(q_0, q_2) - d(q_1, q_2) - d(q'_0, q'_1) - d(q'_0, q'_2) + d(q'_1, q'_2)| \\ &\leq \frac{1}{2} |d(q_0, q_1) - d(q'_0, q'_1)| + \frac{1}{2} |d(q_0, q_2) - d(q'_0, q'_2)| + \frac{1}{2} |d(q_1, q_2) - d(q'_1, q'_2)| \\ &\leq \frac{3}{2}. \end{aligned}$$

Given  $q_0, q_1, q_2, q_3 \in G$ , let  $q'_0, q'_1, q'_2, q'_3 \in V(G)$ , with  $d(q_i, q'_i) \leq 1/2$  for  $i = 0, 1, 2, 3$ . We have

$$\begin{aligned} (q_1, q_3)_{q_0} &\geq (q'_1, q'_3)_{q'_0} - \frac{3}{2} \geq \min\{(q'_1, q'_2)_{q'_0}, (q'_2, q'_3)_{q'_0}\} - \delta_v^*(G) - \frac{3}{2} \\ &\geq \min\{(q_1, q_2)_{q_0} - \frac{3}{2}, (q_2, q_3)_{q_0} - \frac{3}{2}\} - \delta_v^*(G) - \frac{3}{2} \\ &= \min\{(q_1, q_2)_{q_0}, (q_2, q_3)_{q_0}\} - \delta_v^*(G) - 3, \end{aligned}$$

and we conclude  $\delta^*(G) \leq \delta_v^*(G) + 3$ .  $\square$

Let  $H$  be a subgraph of  $G$ ,  $H$  is isometric if  $d_H(x, y) = d_G(x, y)$  for every  $x, y \in H$ . We will need the following well-known result.

**Lemma 1.** Let  $H$  be an isometric subgraph of  $G$ . Then

$$\delta(H) \leq \delta(G),$$

$$\delta^*(H) \leq \delta^*(G),$$

$$\delta_v^*(H) \leq \delta_v^*(G).$$

Since  $G$  is an isometric subgraph of  $T(G)$  and  $R(G)$ , and  $\Lambda(G)$  is an isometric subgraph of  $T(G)$  and  $Q(G)$ , we have the following consequence of Lemma 1.

**Corollary 2.** For any graph  $G$ , we have

$$\begin{aligned} \delta(G) &\leq \delta(T(G)), & \delta^*(G) &\leq \delta^*(T(G)), & \delta_v^*(G) &\leq \delta_v^*(T(G)), \\ \delta(G) &\leq \delta(R(G)), & \delta^*(G) &\leq \delta^*(R(G)), & \delta_v^*(G) &\leq \delta_v^*(R(G)), \\ \delta(\Lambda(G)) &\leq \delta(T(G)), & \delta^*(\Lambda(G)) &\leq \delta^*(T(G)), & \delta_v^*(\Lambda(G)) &\leq \delta_v^*(T(G)), \\ \delta(\Lambda(G)) &\leq \delta(Q(G)), & \delta^*(\Lambda(G)) &\leq \delta^*(Q(G)), & \delta_v^*(\Lambda(G)) &\leq \delta_v^*(Q(G)). \end{aligned}$$

The hyperbolicity of the line graph has been studied previously (see [21–23]). We have the following results.

**Theorem 2.** [22] (Corollary 3.12) Let  $G$  be a graph. Then

$$\delta(G) \leq \delta(\Lambda(G)) \leq 5\delta(G) + 5/2.$$

Furthermore, the first inequality is sharp: the equality is attained by every cycle graph.

**Theorem 3.** [21] (Theorem 6) Let  $G$  be a graph. Then

$$\delta_v^*(G) - 1 \leq \delta_v^*(\Lambda(G)) \leq \delta_v^*(G) + 1.$$

**Theorem 4.** Let  $G$  be a graph. Then

$$\delta^*(G) - 4 \leq \delta^*(\Lambda(G)) \leq \delta^*(G) + 4.$$

**Proof.** Proposition 2 and Theorem 3 give  $\delta^*(G) \leq \delta_v^*(G) + 3 \leq \delta_v^*(\Lambda(G)) + 4 \leq \delta^*(\Lambda(G)) + 4$ , and  $\delta^*(\Lambda(G)) \leq \delta_v^*(\Lambda(G)) + 3 \leq \delta_v^*(G) + 4 \leq \delta^*(G) + 4$ .  $\square$

From Proposition 1, and Theorems 2 and 4 we have:

**Corollary 3.** Let  $G$  be a graph. Then

$$\begin{aligned} \delta(S(G)) &\leq 2\delta(\Lambda(G)) \leq 5\delta(S(G)) + 5, \\ \delta^*(S(G)) - 8 &\leq 2\delta^*(\Lambda(G)) \leq \delta^*(S(G)) + 8. \end{aligned}$$

Corollary 2 and Theorems 2, 3 and 4 have the following consequence.

**Corollary 4.** Let  $G$  be a graph. Then

$$\begin{aligned} \delta(G) &\leq \delta(Q(G)), \\ \delta_v^*(G) &\leq \delta_v^*(Q(G)) + 1, \\ \delta^*(G) &\leq \delta^*(Q(G)) + 4. \end{aligned}$$

Theorem 4 improves the inequality  $\delta^*(\Lambda(G)) \leq \delta^*(G) + 6$  in [23].

Given a graph  $G$  with multiple edges, we define the graph  $B(G)$ , obtained from  $G$ , substituting each multiple edge for one of its simple edges of shorter length (see [23]).

**Remark 1.** By argument in the proof of [24](Theorem 8) we have: If in each multiple edge there is at most one edge with length greater than  $j := \inf\{d(u, v) : u, v \text{ are joined by a multiple edge of } G\}$ , then  $\delta(G) \leq \max\left\{\delta(B(G)) + \frac{J-j}{2}, \frac{J+j}{4}\right\}$ , where,  $J := \sup\{L(e) : e \text{ is an edge contained in a multiple edge of } G\}$ .

**Corollary 5.** *Let  $G$  be a graph. Then*

$$\max \left\{ \delta(G), \frac{3}{4} \right\} \leq \delta(R(G)) \leq \max \left\{ \delta(G) + \frac{1}{2}, \frac{3}{4} \right\}.$$

**Proof.** Note that  $R(G)$  can be obtained by adding an edge of length 2 to each pair of adjacent vertices in  $G$ , so the graph becomes a graph with multiple edges, with  $j = 1$  and  $J = 2$ . Then [24] (Theorem 8) and Remark 1 give the result.  $\square$

From [25] (Theorem 11), we have the following result.

**Lemma 2.** *Given the following graphs with edges of length 1, we have*

- *If  $P_n$  is a path graph, then  $\delta(P_n) = 0$  for all  $n \geq 1$ .*
- *If  $C_n$  is a cycle graph, then  $\delta(C_n) = n/4$  for all  $n \geq 3$ .*
- *If  $K_n$  is a complete graph, then  $\delta(K_1) = \delta(K_2) = 0$ ,  $\delta(K_3) = 3/4$  and  $\delta(K_n) = 1$  for all  $n \geq 4$ .*

If  $G$  is not a tree, we define its *girth*  $g(G)$  by

$$g(G) := \inf\{L(C) : C \text{ is a cycle in } G\}.$$

From [26] (Theorem 17), we have:

**Theorem 5.** *If  $G$  is not a tree, then*

$$\delta(G) \geq \frac{g(G)}{4}.$$

**Corollary 6.** *If  $G$  is not a tree, then*

$$\delta(G) \geq \frac{3}{4}.$$

**Corollary 7.** *If  $G$  is not a tree, then*

$$\delta(G) \leq \delta(R(G)) \leq \delta(G) + \frac{1}{2}.$$

**Proof.** Since  $G$  is not a tree, Corollary 6 gives  $\delta(G) \geq 3/4$ , and so

$$\max \left\{ \delta(G), \frac{3}{4} \right\} = \delta(G), \quad \max \left\{ \delta(G) + \frac{1}{2}, \frac{3}{4} \right\} = \delta(G) + \frac{1}{2},$$

and Corollary 5 gives the inequalities.  $\square$

Theorem 2 and Corollary 7 have the following consequence.

**Corollary 8.** *If  $G$  is not a tree, then*

$$\delta(R(G)) - \frac{1}{2} \leq \delta(\Lambda(G)) \leq 5\delta(R(G)) + \frac{5}{2}.$$

From Proposition 1 and Corollary 7 we have the following result.

**Corollary 9.** *If  $G$  is not a tree, then*

$$\delta(S(G)) \leq 2\delta(R(G)) \leq \delta(S(G)) + 1.$$

**Theorem 6.** Let  $G$  be a graph. Then

$$\begin{aligned} \delta^*(\Lambda(G)) &\leq \delta^*(Q(G)) \leq \delta_v^*(\Lambda(G)) + 6 \leq \delta^*(\Lambda(G)) + 6, \\ \delta_v^*(\Lambda(G)) &\leq \delta_v^*(Q(G)) \leq \delta_v^*(\Lambda(G)) + 6, \\ \delta^*(\Lambda(G)) &\leq \delta^*(T(G)) \leq \delta_v^*(\Lambda(G)) + 9 \leq \delta^*(\Lambda(G)) + 9, \\ \delta_v^*(\Lambda(G)) &\leq \delta_v^*(T(G)) \leq \delta_v^*(\Lambda(G)) + 6, \\ \delta^*(G) &\leq \delta^*(R(G)) \leq \delta_v^*(G) + 6 \leq \delta^*(G) + 6, \\ \delta_v^*(G) &\leq \delta_v^*(R(G)) \leq \delta_v^*(G) + 6, \\ \delta^*(G) &\leq \delta^*(T(G)) \leq \delta_v^*(G) + 9 \leq \delta^*(G) + 9, \\ \delta_v^*(G) &\leq \delta_v^*(T(G)) \leq \delta_v^*(G) + 6. \end{aligned}$$

**Proof.** The lower bounds follow from Corollary 2. We consider the map  $P : Q(G) \rightarrow \Lambda(G)$  such that  $P(q) = q$  if  $q \in \Lambda(G)$ ,  $P(q) = v_q$  if  $q \notin \Lambda(G)$ , where  $v_q \in V(\Lambda(G))$  and  $d_{Q(G)}(q, v_q) \leq 1$ . If  $q_0, q_1, q_2, q_3 \in Q(G)$ , then

$$|d_{Q(G)}(q_i, q_j) - d_{\Lambda(G)}(P(q_i), P(q_j))| = |d_{Q(G)}(q_i, q_j) - d_{Q(G)}(P(q_i), P(q_j))| \leq 2,$$

since  $\Lambda(G)$  is an isometric subgraph of  $Q(G)$  and

$$\begin{aligned} &|(q_i, q_j)_{q_0} - (P(q_i), P(q_j))_{P(q_0)}| \\ &= \frac{1}{2} |d_{Q(G)}(q_0, q_i) + d_{Q(G)}(q_0, q_j) - d_{Q(G)}(q_i, q_j) \\ &\quad - d_{\Lambda(G)}(P(q_0), P(q_i)) - d_{\Lambda(G)}(P(q_0), P(q_j)) + d_{\Lambda(G)}(P(q_i), P(q_j))| \leq 3, \end{aligned}$$

for  $i, j \in \{1, 2, 3\}$ . Thus,

$$\begin{aligned} (q_1, q_3)_{q_0} &\geq (P(q_1), P(q_3))_{P(q_0)} - 3 \\ &\geq \min\{(P(q_1), P(q_2))_{P(q_0)}, (P(q_2), P(q_3))_{P(q_0)}\} - \delta_v^*(\Lambda(G)) - 3 \\ &\geq \min\{(q_1, q_2)_{q_0} - 3, (q_2, q_3)_{q_0} - 3\} - \delta_v^*(\Lambda(G)) - 3 \\ &= \min\{(q_1, q_2)_{q_0}, (q_2, q_3)_{q_0}\} - \delta_v^*(\Lambda(G)) - 6. \end{aligned}$$

Therefore,

$$\delta^*(\Lambda(G)) + 6 \geq \delta_v^*(\Lambda(G)) + 6 \geq \delta^*(Q(G)) \geq \delta_v^*(Q(G)).$$

These inequalities allow us to obtain the result for upper bounds of  $\delta^*(Q(G))$  and  $\delta_v^*(Q(G))$ . The other upper bounds can be obtained similarly.  $\square$

From Theorems 3 and 6 and Corollary 4 we have:

**Corollary 10.** For all graph  $G$ , we have

$$\begin{aligned} \delta_v^*(G) - 1 &\leq \delta_v^*(Q(G)) \leq \delta_v^*(G) + 7, \\ \delta^*(G) - 4 &\leq \delta^*(Q(G)) \leq \delta_v^*(G) + 7 \leq \delta^*(G) + 7. \end{aligned}$$

From Corollaries 2, 4 and 10, Theorem 6 and the inequalities  $\delta(G) \leq 3\delta^*(G)$  and  $\delta^*(G) \leq 2\delta(G)$ , we have:

**Corollary 11.** Let  $G$  be a graph. Then

$$\begin{aligned}\delta(\Lambda(G)) &\leq \delta(Q(G)) \leq 6\delta(\Lambda(G)) + 18, \\ \delta(\Lambda(G)) &\leq \delta(T(G)) \leq 6\delta(\Lambda(G)) + 27, \\ \delta(G) &\leq \delta(T(G)) \leq 6\delta(G) + 27, \\ \delta(G) &\leq \delta(Q(G)) \leq 6\delta(G) + 21.\end{aligned}$$

**Proof.** Corollaries 2 and 4 give the lower bounds. On the other hand, Theorem 6 gives  $\delta(Q(G)) \leq 3\delta^*(Q(G)) \leq 3\delta^*(\Lambda(G)) + 18 \leq 6\delta(\Lambda(G)) + 18$ ,  $\delta(T(G)) \leq 3\delta^*(T(G)) \leq 3(\delta^*(\Lambda(G)) + 9) \leq 6\delta(\Lambda(G)) + 27$ ; we obtain the third upper bound in a similar way. Corollary 10 gives  $3\delta^*(Q(G)) \leq 3(\delta^*(G) + 7) \leq 6\delta(G) + 21$ , obtaining the last upper bound.  $\square$

Let  $G$  be a graph, a family of subgraphs  $\{G_s\}_s$  of  $G$  is a  $T$ -decomposition if  $\cup_s G_s = G$  and  $G_s \cap G_r$  is either a cut-vertex or the empty set for each  $s \neq r$  (see [25]).

The following result was proved in [24] (Theorem 3).

**Lemma 3.** Given a graph  $G$  and  $\{G_s\}_s$  any  $T$ -decomposition of  $G$ , then

$$\delta(G) = \sup_s \delta(G_s).$$

The following results improve the inequality  $\delta(Q(G)) \leq 6\delta(\Lambda(G)) + 18$  in Corollary 11.

**Theorem 7.** Let  $G$  be a path graph, then

$$0 = \delta(\Lambda(G)) \leq \delta(Q(G)) \leq 3/4.$$

**Proof.** Since  $G$  is a path graph,  $\Lambda(G)$  is also a path graph, and so  $0 = \delta(\Lambda(G)) \leq \delta(Q(G))$ .

Consider the  $T$ -decomposition  $\{G_n\}$  of  $Q(G)$ . Since each connected component  $G_n$  is either a cycle  $C_3$  or a path of length 1, we have  $\delta(Q(G)) = \sup_n \{\delta(G_n)\} \leq 3/4$ , by Lemmas 2 and 3.  $\square$

The union of the set of the midpoints of the edges of a graph  $G$  and the set of vertices,  $V(G)$ , will be denoted by  $N(G)$ . Let  $\mathbb{T}_1$  be the set of geodesic triangles  $T$  in  $G$  such that every vertex of  $T$  belong to  $N(G)$  and  $\delta_1(G) := \inf\{\lambda : \text{every triangle in } \mathbb{T}_1 \text{ is } \lambda\text{-thin}\}$ .

**Lemma 4.** [27] (Theorems 2.5 and 2.7) For every graph  $G$ , we have  $\delta_1(G) = \delta(G)$ . Furthermore, if  $G$  is hyperbolic, then there exists  $T \in \mathbb{T}_1$  with  $\delta(T) = \delta(G)$ .

The previous lemma allows to reduce the study of the hyperbolicity constant of a graph  $G$  to study only the geodesic triangles of  $G$ , whose vertices are vertices of  $G$  (i.e., belong to  $V(G)$ ) or midpoints of the edges of  $G$ .

**Theorem 8.** If  $G$  is not a path graph, then

$$\delta(\Lambda(G)) \leq \delta(Q(G)) \leq \delta(\Lambda(G)) + 1/2.$$

**Proof.** By Corollary 2 we have the first inequality. We will prove the second one. If  $\delta(Q(G)) = \infty$ , then Theorem 6 gives  $\delta(\Lambda(G)) = \infty$ , and the second inequality holds. Assume now that  $\delta(Q(G)) < \infty$  (and so,  $\delta(\Lambda(G)) < \infty$  by Theorem 6). If  $G$  is not a path, then  $\Lambda(G)$  is not a tree and Corollary 6 gives  $\delta(\Lambda(G)) \geq 3/4$ .

For each  $v \in V(G)$ , let us define  $V_v := \{u \in V(Q(G)) : uv \in E(Q(G))\} = \{u \in V(\Lambda(G)) : uv \in E(Q(G))\}$ . Denote by  $G_v$  and  $G_v^*$  the subgraphs of  $Q(G)$  induced by the sets  $V_v \cup \{v\}$  and  $V_v$ , respectively. Note that both  $G_v$  and  $G_v^*$  are complete graphs for every  $v \in V(G)$ , and if



$G^*$  is a complete graph with  $r$  vertices, then  $G_v$  is a complete graph with  $r + 1$  vertices. Also,  $Q(G) = \Lambda(G) \cup (\cup_{v \in V(G)} G_v)$ .

By Lemma 4 there exists a geodesic triangle  $T \in \mathbb{T}_1$  in  $Q(G)$  with  $\delta(T) = \delta(Q(G))$ . Denote by  $\gamma_1, \gamma_2, \gamma_3$  the sides of  $T$ . Without loss of generality we can assume that there exists  $p \in \gamma_1$  with  $d_{Q(G)}(p, \gamma_2 \cup \gamma_3) = \delta(T) = \delta(Q(G))$ . Thus,  $T$  is a cycle and each vertex of  $T$  is either the midpoint of some edge of  $Q(G)$  or a vertex of  $Q(G)$ .

If  $G_v$  contains  $T$  for some  $v \in V(G)$ , then  $\delta(Q(G)) = \delta(T) \leq \delta(G_v) \leq 1 < 3/4 + 1/2 \leq \delta(\Lambda(G)) + 1/2$  by Lemma 2, since  $G_v$  is an isometric subgraph of  $Q(G)$ .

If  $\Lambda(G)$  contains  $T$ , then  $\delta(Q(G)) = \delta(T) \leq \delta(\Lambda(G))$  by Lemma 1, since  $\Lambda(G)$  is isometric.

Suppose that  $T$  is not contained either in  $\Lambda(G)$  nor  $G_v$  with  $v \in V(G)$ .

Note that if  $T \cap (G_v \setminus G_v^*) \neq \emptyset$  for some  $v \in V(G)$ , then there exists at least one vertex of  $T$  in  $G_v \setminus \Lambda(G)$ . In order to form a triangle  $T^* \subset \Lambda(G)$  from  $T$ , we define  $\gamma_i^* := \gamma_i \cap \Lambda(G)$ . Note that, for  $i \in \{1, 2, 3\}$ ,  $\gamma_i^*$  is a geodesic, since  $\Lambda(G)$  is a isometric subgraph of  $Q(G)$ .

We denote by  $x_{i,j}$  the common vertex of  $\gamma_i$  and  $\gamma_j$  and by  $u_i$  and  $u_j$  the other vertices of  $\gamma_i$  and  $\gamma_j$  respectively.

We consider the following cases:

Case A. We assume that exactly one vertex of  $T$  belongs to  $Q(G) \setminus \Lambda(G)$ . Thus, there exists  $v \in V(G)$  such that  $T \cap (G_v \setminus G_v^*) \neq \emptyset$ . By Lemma 4, we have two possibilities: the vertex of  $T$  is a vertex of  $G$  or a midpoint of an edge in  $G_v \setminus G_v^*$ .

We can suppose that  $x_{i,j} \in T \setminus \Lambda(G)$ . Let  $v$  be a vertex of  $V(G)$  such that  $x_{i,j} \in G_v \setminus \Lambda(G)$ . Let  $x_i$  (respectively,  $x_j$ ) be the closest point of  $\gamma_i^*$  (respectively,  $\gamma_j^*$ ) to  $x_{i,j}$ . Thus,  $x_i x_j \in E(\Lambda(G))$ . Let  $v^*$  be the midpoint of the edge  $x_i x_j$ . Let  $T_1$  be the connected component of  $T \setminus \Lambda(G)$  joining  $x_i$  and  $x_j$ . Note that  $L(T_1) = 2$ . We analyze the two possibilities:

Case A1. Assume that  $x_{i,j} \in V(Q(G))$ . Let us define  $\sigma_i := \gamma_i^* \cup [x_i v^*]$  and  $\sigma_j := \gamma_j^* \cup [x_j v^*]$ . We are going to prove that  $\sigma_i$  and  $\sigma_j$  are geodesics in  $\Lambda(G)$ . In fact, we prove now that if  $\gamma_j^* = [z_j x_j]$ , then  $d_{Q(G)}(z_j, x_j) \leq d_{Q(G)}(z_j, x_i)$ . Seeking for a contradiction assume that  $d_{Q(G)}(z_j, x_j) > d_{Q(G)}(z_j, x_i)$ . Thus,

$$d_{Q(G)}(z_j, x_i) + d_{Q(G)}(x_i, x_{i,j}) = d_{Q(G)}(z_j, x_i) + 1 \leq d_{Q(G)}(z_j, x_j) + d_{Q(G)}(x_j, x_{i,j})$$

therefore  $\gamma_j$  is not a geodesic obtaining the desired contradiction and we conclude  $d_{Q(G)}(z_j, x_j) \leq d_{Q(G)}(z_j, x_i)$ . Hence,  $\sigma_i$  is a geodesic in  $\Lambda(G)$ .

Case A2. There is an edge  $e \in E(Q(G)) \setminus E(\Lambda(G))$  such that  $x_{i,j}$  is the midpoint of  $e$ , thus without loss of generality we can assume that  $e = x_i v$ , and we define  $\sigma_i := \gamma_i^*$  and  $\sigma_j := \gamma_j^* \cup x_j x_i$ . Thus,  $\sigma_i$  is a geodesic in  $\Lambda(G)$ .

Note that  $\gamma_j^* \cup x_j v \cup [v x_{i,j}]$  and  $\sigma_j \cup [x_i x_{i,j}] = \gamma_j^* \cup x_j x_i \cup [x_i x_{i,j}]$  have the same endpoints and length; therefore,  $\sigma_j$  is also a geodesic in  $\Lambda(G)$ .

Case B. Assume that there are two vertices of  $T$  in some connected component of  $T \setminus \Lambda(G)$ . Thus, there exists  $v \in V(G)$  such that  $T \cap (G_v \setminus G_v^*) \neq \emptyset$ . By Lemma 4, we have two possibilities again: both vertices of  $T$  are midpoints of edges or one vertex of  $T$  is a vertex of  $G$  and the other is a midpoint of an edge.

We can assume that  $u_i, u_j \in G_v \setminus G_v^*$  for some  $v$ . We denote by  $x'_i$  (respectively,  $x'_j$ ) the closest point in  $\gamma_i^*$  (respectively,  $\gamma_j^*$ ) to  $u_i$  (respectively,  $u_j$ ); then  $x'_i x'_j \in E(\Lambda(G))$ . Let  $v'$  be the midpoint of the edge  $x'_i x'_j$ . Let  $T_2$  be the connected component of  $T \setminus \Lambda(G)$  joining  $x'_i$  and  $x'_j$ . Note that  $L(T_2) = 2$ .

We analyze the two possibilities again:

Case B1. The vertices  $u_i, u_j$  of  $T$  are the midpoints of  $x'_i v$  and  $x'_j v$ . Thus,  $\sigma_i := \gamma_i^*$ ,  $\sigma_j := \gamma_j^*$  and  $\sigma_k := x'_i x'_j$  are geodesics in  $\Lambda(G)$ .

Case B2. Otherwise, we can assume without loss of generality that  $u_j = v$  and  $u_i$  is the midpoint of  $x_i v$ . We have  $d_{Q(G)}(u_i, x_j) = d_{Q(G)}(u_i, x_i) + 1$  and so,  $\sigma_i := \gamma_i^*$  and  $\sigma_j := \gamma_j^* \cup x'_i x'_j$  are geodesics in  $\Lambda(G)$ . In this case we define  $\sigma_k := \{x'_i\}$ .

Note that the most general possible case is the following: there are at most three vertices  $v_1, v_2, v_3 \in V(G)$  such that  $T \cap (G_{v_i} \setminus G_{v_i}^*) \neq \emptyset$ , for  $i = 1, 2, 3$ . Repeating the previous process at

most three times we obtain a geodesic triangle  $T^*$  in  $\Lambda(G)$  with sides  $\gamma'_1, \gamma'_2$  and  $\gamma'_3$  containing  $\gamma_1^*, \gamma_2^*$  and  $\gamma_3^*$ , respectively.

If  $p \in \Lambda(G)$ , then one can check that  $\delta(Q(G)) = d_{Q(G)}(p, \gamma_2 \cup \gamma_3) \leq d_{Q(G)}(p, \gamma'_2 \cup \gamma'_3) + 1/2 \leq \delta(\Lambda(G)) + 1/2$ . If  $p \notin \Lambda(G)$ , then  $\delta(Q(G)) = d_{Q(G)}(p, \gamma_2 \cup \gamma_3) \leq 5/4$ ; since  $\delta(\Lambda(G)) \geq 3/4$ , we have  $\delta(\Lambda(G)) + 1/2 \geq 5/4 \geq \delta(Q(G))$ . This finishes the proof.  $\square$

Proposition 1, Theorems 2 and 8, and Corollary 3 have the following consequence.

**Corollary 12.** *Let  $G$  be a graph. If  $G$  is not a path graph, then*

$$\delta(S(G)) \leq 2\delta(Q(G)) \leq 5\delta(S(G)) + 6.$$

#### 4. Conclusions

In this paper, we obtained several inequalities and closed formulas relating the hyperbolicity constants of a graph  $G$  and its graph operators  $\Lambda(G)$ ,  $S(G)$ ,  $T(G)$ ,  $R(G)$  and  $Q(G)$ , by the use of their symmetries. As a first step, as the basis of our research, we found relations among the Gromov hyperbolicity constant (satisfying the Rips condition), the Gromov product and the Gromov product restricted to vertices. In the same direction, we derived inequalities between Gromov products and graph operators; as examples we mention:  $\delta_v^*(G) \leq \delta^*(G) \leq \delta_v^*(G) + 3$ ,  $\delta_v^*(G) \leq \delta_v^*(Q(G)) + 1$  and  $\delta^*(G) \leq \delta^*(R(G)) \leq \delta_v^*(G) + 6 \leq \delta^*(G) + 6$ .

Then, we studied relations between the Gromov hyperbolicity constant of a graph and the application of given operators to that graph. In this context, we obtained inequalities such as:  $\delta(G) \leq \delta(R(G)) \leq \delta(G) + 1/2$ ,  $\delta(\Lambda(G)) \leq \delta(Q(G)) \leq \delta(\Lambda(G)) + 1/2$ ,  $\delta(S(G)) \leq 2\delta(R(G)) \leq \delta(S(G)) + 1$  and  $\delta(R(G)) - 1/2 \leq \delta(\Lambda(G)) \leq 5\delta(R(G)) + 5/2$ , where  $G$  not a tree.

We believe that our work may motivate the investigation of related open problems such as: (i) the computation of the hyperbolicity constant on geometric graphs; (ii) the analysis of hyperbolicity on the graph operators reported here (i.e.,  $\Lambda(G)$ ,  $S(G)$ ,  $T(G)$ ,  $R(G)$  and  $Q(G)$ ) when applied to geometric graphs; (iii) the study of the hyperbolicity constants of additional graph operators; and (iv) the identification of the properties of graph operations that break or preserve hyperbolicity.

**Author Contributions:** The authors contributed equally to this work.

**Funding:** Supported in part by two grants from Ministerio de Economía y Competitividad, Agencia Estatal de Investigación (AEI) and Fondo Europeo de Desarrollo Regional (FEDER) (MTM2016-78227-C2-1-P and MTM2015-69323-REDT), Spain.

**Acknowledgments:** The authors would like to thank the editor and the anonymous referees whose comments and suggestions greatly improved the presentation of this paper.

**Conflicts of Interest:** The authors declare no conflict of interest. The founding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

#### References

1. Krausz, J. Démonstration nouvelle d'un théorème de Whitney sur les réseaux. *Mat. Fiz. Lapok* **1943**, *50*, 75–85.
2. Harary, F.; Norman, R.Z. Some properties of line digraphs. *Rend. Circ. Mat. Palermo* **1960**, *9*, 161–168. [[CrossRef](#)]
3. Prisner, E. *Graph Dynamics*; Chapman and Hall/CRC: Boca Raton, FL, USA, 1995; Volume 338.
4. Bindusree, A.R.; Naci Cangul, I.; Lokesh, V.; Sinan Cevik, A. Zagreb polynomials of three graph operators. *Filomat* **2016**, *30*, 1979–1986. [[CrossRef](#)]
5. Ranjini, P.S.; Lokesh, V. Smarandache-Zagreb index on three graph operators. *Int. J. Math. Comb.* **2010**, *3*, 1–10.
6. Yan, W.; Yang, B.-Y.; Yeh, Y.-N. The behavior of Wiener indices and polynomials of graphs under five graph decorations. *Appl. Math. Lett.* **2007**, *20*, 290–295. [[CrossRef](#)]

7. Gromov, M. Hyperbolic groups. In *Essays in Group Theory*; Gersten, S.M., Ed.; Mathematical Sciences Research Institute Publications; Springer: Berlin, Germany, 1987; Volume 8, pp. 75–263.
8. Oshika, K. *Discrete Groups*; AMS Bookstore: Providence, RI, USA, 2002.
9. Jonckheere, E.A. Contrôle du trafic sur les réseaux à géométrie hyperbolique—Vers une théorie géométrique de la sécurité l’acheminement de l’information. *J. Eur. Syst. Autom.* **2002**, *8*, 45–60.
10. Jonckheere, E.A.; Lohsoonthorn, P. Geometry of network security. In Proceedings of the 2004 American Control Conference, Boston, MA, USA, 30 June–2 July 2004; pp. 111–151.
11. Bowditch, B.H. Notes on Gromov’s hyperbolicity criterion for path-metric spaces. In *Group Theory from a Geometrical Viewpoint*; Ghys, E., Haefliger, A., Verjovsky, A., Eds.; World Scientific: River Edge, NJ, USA, 1991; pp. 64–167.
12. Alonso, J.; Brady, T.; Cooper, D.; Delzant, T.; Ferlini, V.; Lustig, M.; Mihalik, M.; Shapiro, M.; Short, H. Notes on word hyperbolic groups. In *Group Theory from a Geometrical Viewpoint*; Ghys, E., Haefliger, A., Verjovsky, A., Eds.; World Scientific: Singapore, 1992; pp. 3–63.
13. Ghys, E.; de la Harpe, P. *Sur les Groupes Hyperboliques d’après Mikhael Gromov*; Progress in Mathematics 83; Birkhäuser Boston Inc.: Boston, MA, USA, 1990.
14. Chepoi, V.; Dragan, F.F.; Vaxès, Y. Core congestion is inherent in hyperbolic networks. In Proceedings of the Twenty-Eighth Annual ACM-SIAM Symposium on Discrete Algorithms, Barcelona, Spain, 16–19 January 2017; pp. 2264–2279.
15. Grippo, E.; Jonckheere, E.A. Effective resistance criterion for negative curvature: Application to congestion control. In Proceedings of the 2016 IEEE Conference on Control Applications (CCA), Buenos Aires, Argentina, 19–22 September 2016.
16. Li, S.; Tucci, G.H. Traffic Congestion in Expanders,  $(p, \delta)$ -Hyperbolic Spaces and Product of Trees. *Int. Math.* **2015**, *11*, 134–142.
17. Shang, Y. Lack of Gromov-hyperbolicity in colored random networks. *Pan-Am. Math. J.* **2011**, *21*, 27–36.
18. Shang, Y. Lack of Gromov-hyperbolicity in small-world networks. *Cent. Eur. J. Math.* **2012**, *10*, 1152–1158. [[CrossRef](#)]
19. Shang, Y. Non-hyperbolicity of random graphs with given expected degrees. *Stoch. Models* **2013**, *29*, 451–462. [[CrossRef](#)]
20. Soto, M. Quelques Propriétés Topologiques des Graphes et Applications a Internet et aux Réseaux. Ph.D. Thesis, Université Paris Diderot, Paris, France, 2011.
21. Coudert, D.; Ducoffe, G. On the hyperbolicity of bipartite graphs and intersection graphs. *Discret. Appl. Math.* **2016**, *214*, 187–195. [[CrossRef](#)]
22. Carballosa, W.; Rodríguez, J.M.; Sigarreta, J.M. New inequalities on the hyperbolicity constant of line graphs. *ARS Comb.* **2014**, *129*, 367–386.
23. Carballosa, W.; Rodríguez, J.M.; Sigarreta, J.M.; Villeta, M. On the hyperbolicity constant of line graphs. *Electron. J. Comb.* **2011**, *18*, 210.
24. Bermudo, S.; Rodríguez, J.M.; Sigarreta, J.M.; Vilaire, J.-M. Gromov hyperbolic graphs. *Discret. Math.* **2013**, *313*, 1575–1585. [[CrossRef](#)]
25. Rodríguez, J.M.; Sigarreta, J.M.; Vilaire, J.-M.; Villeta, M. On the hyperbolicity constant in graphs. *Discret. Math.* **2011**, *311*, 211–219. [[CrossRef](#)]
26. Michel, J.; Rodríguez, J.M.; Sigarreta, J.M.; Villeta, M. Hyperbolicity and parameters of graphs. *ARS Comb.* **2011**, *100*, 43–63.
27. Bermudo, S.; Rodríguez, J.M.; Sigarreta, J.M. Computing the hyperbolicity constant. *Comput. Math. Appl.* **2011**, *62*, 4592–4595. [[CrossRef](#)]

