

This document is published at:

Carballosa, W., de la Cruz, A., Martínez-Pérez, A., Rodríguez, J.M. (2018). Hyperbolicity of Direct Products of Graphs. *Symmetry*, 10 (7), 279.

DOI: <https://doi.org/10.3390/sym10070279>



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Article

Hyperbolicity of Direct Products of Graphs

Walter Carballosa ^{1,2}, Amauris de la Cruz ³, Alvaro Martínez-Pérez ⁴ 
and José M. Rodríguez ^{3,*} 

¹ Department of Mathematics and Statistics, Florida International University, 11200 SW 8th Street, Miami, FL 33199, USA; waltercarb@gmail.com

² Department of Mathematics, Miami Dade College, 300 NE Second Ave. Miami, FL 33132, USA

³ Departamento de Matemáticas, Universidad Carlos III de Madrid, Avenida de la Universidad 30, Leganés, 28911 Madrid, Spain; alcruz@math.uc3m.es

⁴ Facultad CC. Sociales de Talavera, Universidad de Castilla La Mancha, Avda. Real Fábrica de Seda, s.n. Talavera de la Reina, 45600 Toledo, Spain; alvaro.martinezperez@uclm.es

* Correspondence: jomaro@math.uc3m.es; Tel.: +34-91-624-9098

Received: 11 June 2018; Accepted: 9 July 2018; 12 July 2018



Abstract: It is well-known that the different products of graphs are some of the more symmetric classes of graphs. Since we are interested in hyperbolicity, it is interesting to study this property in products of graphs. Some previous works characterize the hyperbolicity of several types of product graphs (Cartesian, strong, join, corona and lexicographic products). However, the problem with the direct product is more complicated. The symmetry of this product allows us to prove that, if the direct product $G_1 \times G_2$ is hyperbolic, then one factor is bounded and the other one is hyperbolic. Besides, we prove that this necessary condition is also sufficient in many cases. In other cases, we find (not so simple) characterizations of hyperbolic direct products. Furthermore, we obtain good bounds, and even formulas in many cases, for the hyperbolicity constant of the direct product of some important graphs (as products of path, cycle and even general bipartite graphs).

Keywords: direct product of graphs; geodesics; Gromov hyperbolicity; bipartite graphs

1. Introduction

An interesting topic in graph theory is the study of the different types of products of graphs [1]. In particular, given two graphs G_1, G_2 , the *direct product* $G_1 \times G_2$ is defined as the graph with vertices the (Cartesian) product of $V(G_1)$ and $V(G_2)$, and two vertices $(u_1, v_1), (u_2, v_2) \in V(G_1 \times G_2)$ are connected by an edge if and only if $[u_1, u_2] \in E(G_1)$ and $[v_1, v_2] \in E(G_2)$. The direct product is associative and commutative. Direct product was introduced in Principia Mathematica by Russell and Whitehead.

Weichsel observed that $G_1 \times G_2$ is connected if and only if the graphs G_1 and G_2 are connected and G_1 or G_2 is not a bipartite graph [2], i.e., there exists an odd cycle. The direct product is known with different names: tensor product, conjunction, categorical product, Kronecker product and cardinal product. There are many works studying several properties of direct products. These works include structural results [3–8], hamiltonian properties [9,10], and above all the well-known Hedetniemi's conjecture (see [11,12]). Imrich has an algorithm in [13] which can recognize in polynomial time if a graph is a direct product; furthermore, the algorithm provides a factorization if the graph is a direct product. This fact facilitates the computational use of the direct product of graphs.

Hyperbolic spaces are an important tool in geometry and group theory [14–16]. Gromov hyperbolicity is a meeting point for different spaces: some of them continuous (hyperbolic plane and many Riemannian manifolds with negative curvature) and some of them discrete (trees and many graphs) [14–16].

Gromov hyperbolicity was introduced in the context of finitely generated groups [16], and it was applied, in the science of computation, to the study of automatic groups [17,18]. Gromov hyperbolicity is useful in networking, algorithms and discrete mathematics [19–24]; also, many real networks are hyperbolic [25–29]. Besides, there are several important applications of hyperbolic spaces to the Internet [30–34] and to random graphs [35–37]. It has recently been pointed out that also some aspects of biological systems require hyperbolicity for proper functioning [38]. In [39], it was proven that, for a large class of Riemannian surfaces endowed with a metric of negative curvature, there is a very simple graph related with the surface such that the surface is hyperbolic if and only if the graph is hyperbolic; therefore, it is interesting to study hyperbolic graphs to understand hyperbolic surfaces.

All these facts show the increasing interest of hyperbolic graphs (see, e.g., [19,24–27,32,33,35–37,39–47] and the references therein).

In this paper, let us denote by $G = (V, E) = (V(G), E(G))$ a connected graph with $V(G) \neq \emptyset$. We consider that the length of each edge is 1. In addition, we assume that the graph does not have either multiple edges or loops.

Trees are the graphs with hyperbolicity constant zero. Thus, we can view the hyperbolicity constant as a measure of how “tree-like” the space is. This is an important subject (see, e.g., [48,49]).

From a computational viewpoint, we can obtain $\delta(G)$ in time $O(n^{3.69})$ for graphs with n vertices [50]. In addition, there is an algorithm which decides if a Cayley graph is hyperbolic [51]. In [52], this algorithm is improved, allowing to obtain $\delta(G)$ in time $O(n^2)$, but only if the graph is given in terms of its distance-matrix. However, it is usually very difficult to decide if an infinite graph is hyperbolic. Therefore, it is useful to study hyperbolicity for particular classes of graphs. There are many works dealing with the hyperbolicity of different types of graphs: median graphs [53], line graphs [54–56], cubic graphs [57], complement graphs [58], regular graphs [59], chordal graphs [25,42,45,60], planar graphs [61,62], bipartite and intersection graphs [63], vertex-symmetric graphs [64], periodic graphs [65,66], expanders [34], bridged graphs [67], short graphs [68], graph minors [69], graphs with small hyperbolicity constant [70], Mycielskian graphs [71], geometric graphs [56,72], and some types of products of graphs: Cartesian product and sum [46,73], strong product [74], lexicographic product [75], and corona and join product [76].

Some of these works give results about the hyperbolicity of some unary operations in graphs:

A line graph is hyperbolic if and only if the original graph does [54–56].

For a large class of minor graphs, the minor graph is hyperbolic if and only if the original graph does [69].

Mycielskian graphs are always hyperbolic [71].

Now, we summarize the known results about the hyperbolicity of the main class of binary operations in graphs: products of graphs.

The Cartesian product is hyperbolic if and only if one factor graph is bounded and the other one is hyperbolic [46].

The same holds for the strong product [74].

The corona product $G_1 \diamond G_2$ is hyperbolic if and only if the first factor G_1 is hyperbolic, and the join $G_1 \uplus G_2$ is always hyperbolic [76].

The Cartesian sum $G_1 \oplus G_2$ is always hyperbolic, if the factors have at least two vertices [73].

The lexicographic product graph $G_1 \circ G_2$ is hyperbolic if and only if G_1 does, if the first factor has at least two vertices [75].

The goal of this paper is the characterization in many cases of the direct product of graphs which are hyperbolic. Here, the situation is more complicated than with other products of graphs. This is partly because the direct product of two bipartite graphs (i.e., graphs without odd cycles) is already disconnected and the formula for the distance in $G_1 \times G_2$ is more complicated than in the case of other products of graphs. The symmetry of this product allows us to show that, if $G_1 \times G_2$ is hyperbolic, then one factor is hyperbolic and the other one is bounded (see Theorem 10). Besides, we prove that this necessary condition is also sufficient in many cases. If G_1 is a hyperbolic graph and G_2 is a bounded

graph, then we prove that $G_1 \times G_2$ is hyperbolic when G_2 has some odd cycle (Theorem 3) or G_1 and G_2 do not have odd cycles (Theorem 4). One could think that otherwise (if G_1 has some odd cycle and G_2 does not have odd cycles) this necessary condition is also sufficient; however, Theorem 15 allows constructing in an easy way examples G_1, G_2 (with G_1 hyperbolic and G_2 bounded) such that $G_1 \times G_2$ is not hyperbolic. This shows that the characterization of hyperbolic direct products is a more difficult task when G_1 has some odd cycle and G_2 does not have odd cycles. Theorems 11 and 12 provide sufficient conditions for non-hyperbolicity and hyperbolicity, respectively. Besides, Theorems 15 and Corollary 5 characterize the hyperbolicity of $G_1 \times G_2$ under some additional conditions. Furthermore, we obtain good bounds, and even formulas in many cases, for the hyperbolicity constant of the direct product of some important graphs; in particular, Theorem 18 provides the hyperbolicity constant of many direct products of bipartite graphs, and Theorems 17 and 19 give the hyperbolicity constant of many direct products of path and cycle graphs.

We want to remark that, in a general context, the hypothesis on the existence (or non-existence) of odd cycles is artificial in the context of Gromov hyperbolicity. However, it is an essential hypothesis in the works on direct products (see Theorem 1). Throughout the development of this work, we have verified that the existence of odd cycles is also essential in the study of hyperbolic product graphs.

2. Definitions and Background

Let (X, d) be a metric space, and denote by L the length associated to the distance d . A *geodesic* is a curve $g : [a, b] \rightarrow X$ satisfying $L(g|_{[t,s]}) = d(g(t), g(s)) = |t - s|$ for every $s, t \in [a, b]$ (here, $g|_{[t,s]}$ is the restriction of g to $[t, s]$). We say that the metric space X is a *geodesic metric space* if for each $p, q \in X$ there is a geodesic connecting them; we denote by $[pq]$ any geodesic from p to q . Hence, a geodesic metric space is a connected space. When X is a graph and $p, q \in V(X)$, $[p, q]$ denotes the edge connecting p and q if they are adjacent.

Along this paper, we consider the graphs as geodesic metric spaces. To do that, we identify any edge $[p, q] \in E(G)$ with the real interval $[0, 1]$; therefore, the points in a graph are the vertices and also the points in the interior of the edges. Hence, we can define a natural distance on the points of a connected graph G by taking shortest paths in G , and so, we consider G as a metric graph. If p and q are points in different connected components of the graph, we define $d(p, q) = \infty$.

Some authors do not consider the internal points of edges in the study. Although this approach has some advantages, we prefer to consider the internal points since these graphs are geodesic metric spaces. We use this approach since to work with geodesic metric spaces provides an interesting geometric viewpoint (for instance, Theorem 2 holds for geodesic metric spaces).

Given a geodesic metric space X and three points $x_1, x_2, x_3 \in X$, the *geodesic triangle* $T = \{x_1, x_2, x_3\}$ is the union of three geodesics $[x_1x_2]$, $[x_2x_3]$ and $[x_3x_1]$. The points x_1, x_2, x_3 are the vertices of the triangle T . The geodesic triangle T is δ -thin if any side of T is contained in the δ -neighborhood of the union of the two other sides. We define the thin constant of the triangle T by $\delta(T) := \inf\{\delta \geq 0 : T \text{ is } \delta\text{-thin}\}$, and the *hyperbolicity constant* of the space X as $\delta(X) := \sup\{\delta(T) : T \text{ is a geodesic triangle in } X\}$. The space X is *hyperbolic* if $\delta(X) < \infty$, and it is δ -hyperbolic if X is hyperbolic and the constant δ satisfies $\delta \geq \delta(X)$. We say that a triangle with two identical vertices is a "bigon". Of course, each bigon in a space (which is δ -hyperbolic) is δ -thin. If $\{X_i\}_{i \in I}$ are the connected components of X , then we can define $\delta(X) := \sup_{i \in I} \delta(X_i)$, and X is hyperbolic if and only if $\delta(X) < \infty$.

We want to remark that in the classical references on hyperbolicity [14,15,77] appear many different definitions of Gromov hyperbolicity. However, the definitions are equivalent: if X is δ_1 -hyperbolic for a definition, then it is δ_2 -hyperbolic for every definition, where the constant δ_2 can be obtained from δ_1 .

We refer to the classical book [1] for definitions and background about direct product graphs.

We need bounds for the distance between points in the direct product. We use the definition given in [1].

Definition 1. Let $G_1 = (V(G_1), E(G_1))$ and $G_2 = (V(G_2), E(G_2))$ be two graphs. The direct product $G_1 \times G_2$ of G_1 and G_2 has $V(G_1) \times V(G_2)$ as vertex set, so that two distinct vertices (u_1, v_1) and (u_2, v_2) of $G_1 \times G_2$ are adjacent if $[u_1, u_2] \in E(G_1)$ and $[v_1, v_2] \in E(G_2)$.

If G_1 and G_2 are isomorphic, we write $G_1 \simeq G_2$. It is clear that, if $G_1 \simeq G_2$, then $\delta(G_1) = \delta(G_2)$.

It is clear that the direct product of two graphs is commutative, i.e., $G_1 \times G_2 \simeq G_2 \times G_1$. Therefore, the conclusion of every result in this paper with some “non-symmetric” hypothesis also holds if we change the roles of G_1 and G_2 (see, e.g., Theorems 3, 4, 11, 12 and 15 and Corollary 5).

Denote by π_i the projection map $\pi_i : V(G_1 \times G_2) \rightarrow V(G_i)$ for $i \in \{1, 2\}$. In fact, this projection is well defined as a map $\pi_i : G_1 \times G_2 \rightarrow G_i$ for $i \in \{1, 2\}$.

We need some previous results of [1]. If $u, u' \in V(G)$, then by a u, u' -walk in G we mean a path joining u and u' where repeating vertices is allowed.

Proposition 1. ([1], Proposition 5.7) Suppose (u, v) and (u', v') are vertices of the direct product $G_1 \times G_2$, and n is an integer for which G_1 has a u, u' -walk of length n and G_2 has a v, v' -walk of length n . Then, $G_1 \times G_2$ has a walk of length n from (u, v) to (u', v') . The smallest such n (if it exists) equals $d_{G_1 \times G_2}((u, v), (u', v'))$. If no such n exists, then $d_{G_1 \times G_2}((u, v), (u', v')) = \infty$.

Proposition 2. ([1], Proposition 5.8) Suppose x and y are vertices of $G_1 \times G_2$. Then,

$$d_{G_1 \times G_2}(x, y) = \min \{n \in \mathbb{N} \mid \text{each factor } G_i \text{ has a } \pi_i(x), \pi_i(y)\text{-walk of length } n \text{ for } i = 1, 2\},$$

where it is understood that $d_{G_1 \times G_2}(x, y) = \infty$ if no such n exists.

Definition 2. If G is a connected graph, the diameter of its vertices is

$$\text{diam } V(G) := \sup\{d_G(u, v) : u, v \in V(G)\},$$

and the diameter of G is

$$\text{diam } G := \sup\{d_G(x, y) : x, y \in G\}.$$

Corollary 1. We have for every $(u, v), (u', v') \in V(G_1 \times G_2)$

$$d_{G_1 \times G_2}((u, v), (u', v')) \geq \max \{d_{G_1}(u, u'), d_{G_2}(v, v')\}$$

and, consequently,

$$\text{diam } V(G_1 \times G_2) \geq \max \{ \text{diam } V(G_1), \text{diam } V(G_2) \}.$$

Furthermore, if $d_{G_1}(u, u')$ and $d_{G_2}(v, v')$ have the same parity, then

$$d_{G_1 \times G_2}((u, v), (u', v')) = \max \{d_{G_1}(u, u'), d_{G_2}(v, v')\}$$

and, consequently,

$$\text{diam } V(G_1 \times G_2) = \max \{ \text{diam } V(G_1), \text{diam } V(G_2) \}.$$

By *trivial graph*, we mean a graph which has only a vertex.

The following result characterizes when a direct product is connected. By *cycle*, we mean a simple closed curve, i.e., a path with different vertices, unless the last one, which is equal to the first vertex.

Theorem 1. ([1], Theorem 5.9) Suppose G_1 and G_2 are connected non-trivial graphs. If at least one of G_1 or G_2 has an odd cycle, then $G_1 \times G_2$ is connected. If both G_1 and G_2 are bipartite, then $G_1 \times G_2$ has exactly two connected components.

Corollary 2. ([1], Corollary 5.10) A direct product of connected non-trivial graphs is connected if and only if at most one of the factors is bipartite. In fact, the product has $2^{\max\{k,1\}-1}$ connected components, where k is the number of bipartite factors.

Consider the metric spaces (X, d_X) and (Y, d_Y) . Given constants $\alpha \geq 1, \beta \geq 0$, a map $f : X \rightarrow Y$ is an (α, β) -quasi-isometric embedding if

$$\alpha^{-1}d_X(x, y) - \beta \leq d_Y(f(x), f(y)) \leq \alpha d_X(x, y) + \beta,$$

for $x, y \in X$. We say that f is ε -full if for each $y \in Y$ there is $x \in X$ with $d_Y(f(x), y) \leq \varepsilon$.

We say that f is a quasi-isometry if there exist constants $\alpha, \beta, \varepsilon$, such that f is an ε -full (α, β) -quasi-isometric embedding.

Two metric spaces X and Y are quasi-isometric if there exists a quasi-isometry $f : X \rightarrow Y$. One can check that to be quasi-isometric is an equivalence relation. An (α, β) -quasi-geodesic in X is an (α, β) -quasi-isometric embedding between an interval of \mathbb{R} and X .

We need the following result ([15], p. 88).

Theorem 2 (Invariance of hyperbolicity). Let $f : X \rightarrow Y$ be an (α, β) -quasi-isometric embedding between the geodesic metric spaces X and Y . If Y is δ_Y -hyperbolic, then X is δ_X -hyperbolic, where δ_X is a constant which just depends on α, β, δ_Y .

Besides, if f is ε -full for some $\varepsilon \geq 0$ (a quasi-isometry) and X is δ_X -hyperbolic, then Y is δ_Y -hyperbolic, where δ_Y is a constant which just depends on $\alpha, \beta, \delta_X, \varepsilon$.

There are several explicit expressions for $\delta_X = \delta_X(\alpha, \beta, \delta_Y)$, some of them very complicated. In [78] appears the best possible formula for δ_X :

$$\delta_X(\alpha, \beta, \delta_Y) = 8\alpha(2\alpha^2(A_1\beta + A_2\delta_Y) + 4\delta_Y + \beta).$$

for some explicit constants A_1, A_2 .

3. Hyperbolic Direct Products

Let us start with a necessary condition for hyperbolicity.

Proposition 3. Let G_1 and G_2 be two unbounded connected graphs. Then, $G_1 \times G_2$ is not hyperbolic.

Proof. Since G_1 and G_2 are unbounded graphs, for each positive integer n there exist two geodesic paths $P_1 := [w_1, w_2] \cup [w_2, w_3] \cup \dots \cup [w_{n-1}, w_n]$ in G_1 and $P_2 := [v_1, v_2] \cup [v_2, v_3] \cup \dots \cup [v_{n-1}, v_n]$ in G_2 . If n is odd, then we can consider the geodesic triangle T in $G_1 \times G_2$ (see Figure 1) defined by the following geodesics:

$$\begin{aligned} \gamma_1 &:= [(w_1, v_2), (w_2, v_1)] \cup [(w_2, v_1), (w_3, v_2)] \cup [(w_3, v_2), (w_4, v_1)] \cup \dots \cup [(w_{n-1}, v_1), (w_n, v_2)], \\ \gamma_2 &:= [(w_1, v_2), (w_2, v_3)] \cup [(w_2, v_3), (w_1, v_4)] \cup [(w_1, v_4), (w_2, v_5)] \cup \dots \cup [(w_1, v_{n-1}), (w_2, v_n)], \\ \gamma_3 &:= [(w_2, v_n), (w_3, v_{n-1})] \cup [(w_3, v_{n-1}), (w_4, v_{n-2})] \cup [(w_4, v_{n-2}), (w_5, v_{n-3})] \cup \dots \cup [(w_{n-1}, v_3), (w_n, v_2)], \end{aligned}$$

Corollary 1 gives that $\gamma_1, \gamma_2, \gamma_3$ are geodesics.

Let $m := \frac{n+1}{2}$ and consider the vertex (w_m, v_{m+1}) in γ_3 . For every vertex (w_i, v_j) in $\gamma_1, j \in \{1, 2\}$, we have $d_{G_1 \times G_2}((w_m, v_{m+1}), (w_i, v_j)) \geq d_{G_2}(v_{m+1}, v_j) \geq m + 1 - 2 = \frac{n-1}{2}$ by Corollary 1. We have for

every vertex (w_i, v_j) in γ_2 , $i \in \{1, 2\}$, by Corollary 1, $d_{G_1 \times G_2}((w_m, v_{m+1}), (w_i, v_j)) \geq d_{G_1}(w_m, w_i) \geq m - 2 = \frac{n-3}{2}$. Hence, $d_{G_1 \times G_2}((w_m, v_{m+1}), \gamma_1 \cup \gamma_2) \geq \frac{n-3}{2}$ and $\delta(G_1 \times G_2) \geq \delta(T) \geq \frac{n-3}{2}$. Since n is arbitrarily large, $G_1 \times G_2$ is not hyperbolic. \square




Figure 1. If G_1 and G_2 are unbounded, for any odd n , there is a geodesic triangle $T \subset G_1 \times G_2$ with $\delta(T) \geq \frac{n-3}{2}$.

Lemma 1. Consider two connected graphs G_1 and G_2 . If $f : V(G_1) \rightarrow V(G_2)$ is an (α, β) -quasi-isometric embedding, then there exists an $(\alpha, \alpha + \beta)$ -quasi-isometric embedding $g : G_1 \rightarrow G_2$ with $g = f$ on $V(G_1)$. In addition, if f is ϵ -full, then g is $(\epsilon + \frac{1}{2})$ -full.

Proof. For each $x \in G_1$, let us choose a closest point $v_x \in V(G_1)$ from x , and define $g(x) := f(v_x)$. Note that $v_x = x$ if $x \in V(G_1)$ and so $g = f$ on $V(G_1)$. Given $x, y \in G_1$, we have

$$d_{G_2}(g(x), g(y)) = d_{G_2}(f(v_x), f(v_y)) \leq \alpha d_{G_1}(v_x, v_y) + \beta \leq \alpha(d_{G_1}(x, y) + 1) + \beta,$$

$$d_{G_2}(g(x), g(y)) = d_{G_2}(f(v_x), f(v_y)) \geq \alpha^{-1}d_{G_1}(v_x, v_y) - \beta \geq \alpha^{-1}(d_{G_1}(x, y) - 1) - \beta,$$

and g is an $(\alpha, \alpha + \beta)$ -quasi-isometric embedding, since $\alpha \geq 1 \geq \alpha^{-1}$.

In addition, if f is ϵ -full, then g is $(\epsilon + \frac{1}{2})$ -full since $g(G_1) = f(V(G_1))$. \square

Given a graph G , let $g_I(G)$ denote the odd girth of G , that is, the length of the shortest odd cycle in G .

Theorem 3. Let G_1 be a connected graph and G_2 be a non-trivial bounded connected graph with some odd cycle. Then, $G_1 \times G_2$ is hyperbolic if and only if G_1 is hyperbolic.

Proof. Fix $v_0 \in V(G_2)$ with v_0 contained in an odd cycle C with $L(C) = g_I(G_2)$. Consider the map $i : V(G_1) \rightarrow V(G_1 \times G_2)$ such that $i(w) := (w, v_0)$ for every $w \in V(G_1)$.

By Corollary 1, for every $w_1, w_2 \in V(G_1)$, $d_{G_1}(w_1, w_2) \leq d_{G_1 \times G_2}((w_1, v_0), (w_2, v_0))$. In addition, Proposition 2 gives the following.

If a geodesic joining w_1 and w_2 has even length, then

$$d_{G_1 \times G_2}((w_1, v_0), (w_2, v_0)) = d_{G_1}(w_1, w_2).$$

If a geodesic joining w_1 and w_2 has odd length, then C defines a v_0, v_0 -walk with odd length and

$$d_{G_1 \times G_2}((w_1, v_0), (w_2, v_0)) \leq \max\{d_{G_1}(w_1, w_2), g_I(G_2)\} \leq d_{G_1}(w_1, w_2) + g_I(G_2).$$

Thus, i is a $(1, g_I(G_2))$ quasi-isometric embedding.

Consider any $(w, v) \in V(G_1 \times G_2)$. Then, if the geodesic joining v and v_0 has even length,

$$d_{G_1 \times G_2}((w, v), (w, v_0)) = d_{G_2}(v, v_0).$$

If a geodesic joining v and v_0 has odd length, $[vv_0] \cup C$ defines a v, v_0 -walk with even length. Therefore,

$$d_{G_1 \times G_2}((w, v), (w, v_0)) \leq d_{G_2}(v, v_0) + g_I(G_2).$$

Thus, i is $(\text{diam}(V(G_2)) + g_I(G_2))$ -full.

Hence, by Lemma 1, there is a $(\text{diam}(V(G_2)) + g_I(G_2) + \frac{1}{2})$ -full $(1, g_I(G_2) + 1)$ -quasi-isometry, $j : G_1 \rightarrow G_1 \times G_2$, and $G_1 \times G_2$ is hyperbolic if and only if G_1 is hyperbolic by Theorem 2. \square

Theorem 4. *Let G_1 be a connected graph without odd cycles and G_2 be a non-trivial bounded connected graph without odd cycles. Then, $G_1 \times G_2$ is hyperbolic if and only if G_1 is hyperbolic.*

Proof. Fix some vertex $w_0 \in V(G_1)$ and some edge $[v_1, v_2] \in E(G_2)$.

By Theorem 1, there are exactly two components in $G_1 \times G_2$. Since there are no odd cycles, there is no $(w_0, v_1), (w_0, v_2)$ -walk in $G_1 \times G_2$. Thus, let us denote by $(G_1 \times G_2)^1$ the component containing the vertex (w_0, v_1) and by $(G_1 \times G_2)^2$ the component containing the vertex (w_0, v_2) .

Consider $i : V(G_1) \rightarrow V(G_1 \times G_2)^1$ defined as $i(w) := (w, v_1)$ for every $w \in V(G_1)$ such that every w_0, w -walk has even length and $i(w) := (w, v_2)$ for every $w \in V(G_1)$ such that every w_0, w -walk has odd length.

By Proposition 2, $d_{G_1 \times G_2}(i(w_1), i(w_2)) = d_{G_1}(w_1, w_2)$ for every $w_1, w_2 \in V(G_1)$ and i is a $(1, 0)$ -quasi-isometric embedding.

Let $(w, v) \in V(G_1 \times G_2)^1$. Let v_j with $j \in \{1, 2\}$ such that every v, v_j -walk has even length. Then, by Proposition 2, $d_{G_1 \times G_2}((w, v), (w, v_j)) = d_{G_2}(v, v_j) \leq \text{diam}(G_2)$. Therefore, i is $\text{diam}(G_2)$ -full.

Hence, by Lemma 1, there is a $(\text{diam}(G_2) + \frac{1}{2})$ -full $(1, 1)$ -quasi-isometry, $j : G_1 \rightarrow (G_1 \times G_2)^1$, and $(G_1 \times G_2)^1$ is hyperbolic if and only if G_1 is hyperbolic by Theorem 2.

The same argument proves that $(G_1 \times G_2)^2$ is hyperbolic. \square

Denote by P_2 the path graph with two vertices and an edge.

Lemma 2. *Let G_1 be a connected graph with some odd cycle and G_2 a non-trivial bounded graph without odd cycles. Then, $G_1 \times G_2$ and $G_1 \times P_2$ are quasi-isometric and $\delta(G_1 \times P_2) \leq \delta(G_1 \times G_2)$.*

Proof. By Theorem 1, we know that $G_1 \times G_2$ and $G_1 \times P_2$ are connected graphs.

Denote by v_1 and v_2 the vertices of P_2 and fix $[w_1, w_2] \in E(G_2)$. The map $f : V(G_1 \times P_2) \rightarrow V(G_1 \times [w_1, w_2])$ defined as $f(u, v_j) := (u, w_j)$ for every $u \in V(G_1)$ and $j = 1, 2$, is an isomorphism of graphs; hence, it suffices to prove that $G_1 \times G_2$ and $G_1 \times [w_1, w_2]$ are quasi-isometric.

Consider the inclusion map $i : V(G_1 \times [w_1, w_2]) \rightarrow V(G_1 \times G_2)$. Since $G_1 \times [w_1, w_2]$ is a subgraph of $G_1 \times G_2$, we have $d_{G_1 \times G_2}(x, y) \leq d_{G_1 \times [w_1, w_2]}(x, y)$ for every $x, y \in V(G_1 \times [w_1, w_2])$.

Since G_2 is a graph without odd cycles, every w_1, w_2 -walk has odd length and every w_j, w_j -walk has even length for $j = 1, 2$. Thus, Proposition 2 gives, for every $x = (u, w_1), y = (v, w_2) \in V(G_1 \times [w_1, w_2])$,

$$d_{G_1 \times [w_1, w_2]}(x, y) = d_{G_1 \times G_2}(x, y) = \min \{L(g) \mid g \text{ is a } u, v\text{-walk of odd length}\}.$$

Furthermore, for every $x = (u, w_j), y = (v, w_j) \in V(G_1 \times [w_1, w_2])$ and $j = 1, 2$,

$$d_{G_1 \times [w_1, w_2]}(x, y) = d_{G_1 \times G_2}(x, y) = \min \{L(g) \mid g \text{ is a } u, v\text{-walk of even length}\}.$$

Hence, $d_{G_1 \times [w_1, w_2]}(x, y) = d_{G_1 \times G_2}(x, y)$ for every $x, y \in V(G_1 \times [w_1, w_2])$, and the inclusion map i is an $(1, 0)$ -quasi-isometric embedding. Therefore, $\delta(G_1 \times P_2) = \delta(G_1 \times [w_1, w_2]) \leq \delta(G_1 \times G_2)$.

Since G_2 is a graph without odd cycles, given any $w \in V(G_2)$, we have either that every w, w_1 -walk has even length and every w, w_2 -walk has odd length or that every w, w_2 -walk has even length and every w, w_1 -walk has odd length. In addition, since G_1 is connected, for each $u \in V(G_1)$ there is some $u' \in V(G_1)$ such that $[u, u'] \in E(G_1)$. Therefore, by Proposition 2, for every $(u, w) \in V(G_1 \times G_2)$, if $\min \{d_{G_2}(w, w_1), d_{G_2}(w, w_2)\}$ is even, then

$$d_{G_1 \times G_2}((u, w), V(G_1 \times [w_1, w_2])) = d_{G_1 \times G_2}((u, w), V(u \times [w_1, w_2])) = \min \{d_{G_2}(w, w_1), d_{G_2}(w, w_2)\},$$

and if $\min \{d_{G_2}(w, w_1), d_{G_2}(w, w_2)\}$ is odd, then

$$d_{G_1 \times G_2}((u, w), V(G_1 \times [w_1, w_2])) = d_{G_1 \times G_2}((u, w), V(u' \times [w_1, w_2])) = \min \{d_{G_2}(w, w_1), d_{G_2}(w, w_2)\}.$$

In both cases,

$$d_{G_1 \times G_2}((u, w), V(G_1 \times [w_1, w_2])) \leq \text{diam } V(G_2),$$

and i is $(\text{diam } V(G_2))$ -full. By Lemma 1, there exists a $(\text{diam } V(G_2) + \frac{1}{2})$ -full $(1, 1)$ -quasi-isometry $g : G_1 \times [w_1, w_2] \rightarrow G_1 \times G_2$. \square

A subgraph Γ of G is said *isometric* if $d_\Gamma(x, y) = d_G(x, y)$ for any $x, y \in \Gamma$. One can check that Γ is isometric if and only if $d_\Gamma(u, v) = d_G(u, v)$ for any $u, v \in V(\Gamma)$.

Lemma 3. ([47], Lemma 5) *If Γ is an isometric subgraph of G , then $\delta(\Gamma) \leq \delta(G)$.*

A u, v -walk g in G is a *shortcut* of a cycle C if $g \cap C = \{u, v\}$ and $L(g) < d_C(u, v)$ where d_C denotes the length metric on C .

A cycle C' is a *reduction* of the cycle C if both have odd length and C' is the union of a subarc η of C and a shortcut of C joining the endpoints of η . Note that $L(C') \leq L(C) - 2$. We say that a cycle is *minimal* if it has odd length and it does not have a reduction.

Lemma 4. *If C is a minimal cycle of G , then $L(C) \leq 4\delta(G)$.*

Proof. We prove first that C is an isometric subgraph of G . Assume that C is not an isometric subgraph. Thus, there exists a shortcut g of C with endpoints u, v . There are two subarcs η_1, η_2 of C joining u and v ; since C has odd length, we can assume that η_1 has even length and η_2 has odd length. If g has even length, then $C' := g \cup \eta_2$ is a reduction of C . If g has odd length, then $C'' := g \cup \eta_1$ is a reduction of C . Hence, C is not minimal, a contradiction, and so C is an isometric subgraph of G .

It is easy to show that any isometric cycle C has length $4\delta(C)$. This fact and Lemma 3 give $L(C) = 4\delta(C) \leq 4\delta(G)$. \square

Given any w_0, w_k -walk $g = [w_0, w_1] \cup [w_1, w_2] \cup \dots \cup [w_{k-1}, w_k]$ in G_1 and $P_2 = [v_1, v_2]$, if $L(g)$ is either odd or even, then we define the $(w_0, v_1), (w_k, v_i)$ -walk for $i \in 1, 2$,

$$\Gamma_1 g := [(w_0, v_1), (w_1, v_2)] \cup [(w_1, v_2), (w_2, v_1)] \cup [(w_2, v_1), (w_3, v_2)] \cup \dots \cup [(w_{k-1}, v_1), (w_k, v_2)],$$

$$\Gamma_1 g := [(w_0, v_1), (w_1, v_2)] \cup [(w_1, v_2), (w_2, v_1)] \cup [(w_2, v_1), (w_3, v_2)] \cup \dots \cup [(w_{k-1}, v_2), (w_k, v_1)],$$

respectively.

Remark 1. By Proposition 2, if g is a geodesic path in G_1 , then $\Gamma_1 g$ is a geodesic path in $G_1 \times P_2$.

Let us define the map $R : V(G_1 \times P_2) \rightarrow V(G_1 \times P_2)$ as $R(w, v_1) = (w, v_2)$ and $R(w, v_2) = (w, v_1)$ for every $w \in V(G_1)$, and the path $\Gamma_2 g$ as $\Gamma_2 g = R(\Gamma_1 g)$.

Let us define the map $(\Gamma_1 g)' : g \rightarrow \Gamma_1 g$ which is an isometry on the edges and such that $(\Gamma_1 g)'(w_j) = (w_j, v_1)$ if j is even and $(\Gamma_1 g)'(w_j) = (w_j, v_2)$ if j is odd. In addition, let $(\Gamma_2 g)' : g \rightarrow \Gamma_2 g$ be the map defined by $(\Gamma_2 g)' := R \circ (\Gamma_1 g)'$.

Given a graph G , denote by \mathcal{C} the set of minimal cycles of G .

Lemma 5. Let G_1 be a connected graph with some odd cycle and $P_2 = [v_1, v_2]$. Consider a geodesic $g = [w_0 w_k] = [w_0, w_1] \cup [w_1, w_2] \cup \dots \cup [w_{k-1}, w_k]$ in G_1 . Let us define $w'_0 := (\Gamma_1 g)'(w_0) = (w_0, v_1)$ and $w'_k := (\Gamma_2 g)'(w_k)$, i.e., $w'_k := (w_k, v_1)$ or $w'_k := (w_k, v_2)$ if k is odd or even, respectively. Then, $d_{G_1 \times P_2}(w'_0, w'_k) > \sqrt{d_{G_1}(w_j, \mathcal{C}(G_1))}$ for every $0 \leq j \leq k$.

Proof. Fix $0 \leq j \leq k$. Define

$$\mathcal{P} := \{ \sigma \mid \sigma \text{ is a } w_0, w_k\text{-walk such that } L(\sigma) \text{ has a parity different from that of } k \}.$$

Proposition 2 gives

$$d_{G_1 \times P_2}(w'_0, w'_k) = \min \{ L(\sigma) \mid \sigma \in \mathcal{P} \}.$$

Choose $\sigma_0 \in \mathcal{P}$ such that $L(\sigma_0) = d_{G_1 \times P_2}(w'_0, w'_k)$. Since $L(g) + L(\sigma_0)$ is odd, we have $L(g) + L(\sigma_0) = 2t + 1$ for some positive integer t . Thus, $d_{G_1 \times P_2}(w'_0, w'_k) = L(\sigma_0) > \frac{1}{2}(2t + 1)$.

If $g \cup \sigma_0$ is a cycle, then let us define $C_0 := g \cup \sigma_0$. Thus, $L(C_0) = 2t + 1$ and $d_{G_1}(w_j, C_0) = 0$ for every $0 \leq j \leq k$. Otherwise, we may assume that $g \cap \sigma_0 = [w_0 w_{i_1}] \cup [w_{i_2} w_k]$ for some $0 \leq i_1 < i_2 \leq k$. If $\sigma_1 = \sigma_0 \setminus g$, then let us define $C_0 := [w_{i_1} w_{i_2}] \cup \sigma_1$ (where $[w_{i_1} w_{i_2}] \subset g$). Hence, C_0 is a cycle, $L(C_0) \leq 2t - 1$ and $d_{G_1}(w_j, C_0) < \frac{1}{2}(2t + 1)$.

If C_0 is not minimal, then consider a reduction C_1 of C_0 . Let us repeat the process until we obtain a minimal cycle C_s . Note that $L(C_1) \leq L(C_0) - 2$ and for every point $p_1 \in C_0$, $d_{G_1}(p_1, C_1) < \frac{1}{2}L(C_0)$. Now, repeating the argument, for every $1 < i \leq s$, $L(C_i) \leq L(C_{i-1}) - 2$ and for every point $p_i \in C_{i-1}$, $d_{G_1}(p_i, C_i) < \frac{1}{2}L(C_{i-1})$. Therefore,

$$\begin{aligned} d_{G_1}(w_j, \mathcal{C}(G_1)) &\leq d_{G_1}(w_j, C_s) \leq d_{G_1}(w_j, C_0) + \frac{1}{2}L(C_0) + \frac{1}{2}L(C_1) + \dots + \frac{1}{2}L(C_s) \\ &< \frac{1}{2}(2t + 1) + \frac{1}{2}(2t - 1) + \dots + \frac{5}{2} + \frac{3}{2}. \end{aligned}$$

Hence,

$$d_{G_1}(w_j, \mathcal{C}(G_1)) < \frac{1}{2} \sum_{i=1}^t (2i + 1) = \frac{1}{2}t^2 + t < \left(\frac{1}{2}(2t + 1) \right)^2 < \left(d_{G_1 \times P_2}(w'_0, w'_k) \right)^2.$$

□

Corollary 3. Let G_1 be a hyperbolic connected graph with some odd cycle and $P_2 = [v_1, v_2]$. Consider a geodesic $g = [w_0 w_k] = [w_0, w_1] \cup [w_1, w_2] \cup \dots \cup [w_{k-1}, w_k]$ in G_1 . Let us define $w'_0 := (\Gamma_1 g)'(w_0) = (w_0, v_1)$ and $w'_k := (\Gamma_2 g)'(w_k)$. Then, we have for every $0 \leq j \leq k$,

$$\frac{1}{2} \left(k + \sqrt{d_{G_1}(w_j, \mathcal{C}(G_1))} \right) \leq d_{G_1 \times P_2}(w'_0, w'_k) \leq k + 2d_{G_1}(w_j, \mathcal{C}(G_1)) + 4\delta(G_1).$$

Proof. Corollary 1 and Lemma 5 give $d_{G_1 \times P_2}(w'_0, w'_k) \geq k$ and $d_{G_1 \times P_2}(w'_0, w'_k) \geq \sqrt{d_{G_1}(w_j, \mathcal{C}(G_1))}$, and these inequalities provide the lower bound of $d_{G_1 \times P_2}(w'_0, w'_k)$.

Consider a geodesic γ joining w_j and $C \in \mathcal{C}(G_1)$ with $L(\gamma) = d_{G_1}(w_j, C) = d_{G_1}(w_j, \mathcal{C}(G_1))$ and the w_0, w_k -walk

$$g' := [w_0 w_j] \cup \gamma \cup C \cup \gamma \cup [w_j w_k].$$

One can check that $\Gamma_1 g'$ is a w'_0, w'_k -walk in $G_1 \times P_2$, and so Lemma 4 gives

$$d_{G_1 \times P_2}(w'_0, w'_k) \leq L(\Gamma_1 g') = L(g') = k + 2d_{G_1}(w_j, \mathcal{C}(G_1)) + L(C) \leq k + 2d_{G_1}(w_j, \mathcal{C}(G_1)) + 4\delta(G_1).$$

□

If $[v_1, v_2]$ is an edge of G , then the point $x \in [v_1, v_2]$ with $d_G(x, v_1) = d_G(x, v_2) = 1/2$ is the midpoint of the edge $[v_1, v_2]$. Denote by $J(G)$ the set of vertices and midpoints of edges in G . Consider the set $\mathbb{T}_1(G)$ of geodesic triangles T in G which are cycles and such that the vertices of T are in $J(G)$. We denote by $\delta_1(G)$ the infimum of the constants μ such that any triangle in $\mathbb{T}_1(G)$ is μ -thin.

The following three results are used throughout the paper.

Theorem 5. ([40], Theorem 2.5) For every connected graph G , we have $\delta_1(G) = \delta(G)$.

Theorem 6. ([40], Theorem 2.6) Let G be any connected graph. Then, $\delta(G)$ is always a multiple of $1/4$.

Theorem 7. ([40], Theorem 2.7) For any hyperbolic connected graph G , there exists a geodesic triangle $T \in \mathbb{T}_1(G)$ such that $\delta(T) = \delta(G)$.

Consider the set $\mathbb{T}_v(G)$ of geodesic triangles T in G that are cycles and such that the three vertices of the triangle T are also vertices of G . $\delta_v(G)$ denotes the infimum of the constants μ such that every triangle in $\mathbb{T}_v(G)$ is μ -thin.

Theorem 8. For every connected graph G , we have $\delta_v(G) \leq \delta(G) \leq 4\delta_v(G) + 1/2$. Hence, G is hyperbolic if and only if $\delta_v(G) < \infty$. Furthermore, if G is hyperbolic, then there are a geodesic triangle $T = \{a, b, c\} \in \mathbb{T}_v(G)$ and $q \in [ab] \cap J(G)$ such that $d(q, [ac] \cup [cb]) = \delta(T) = \delta_v(G)$. In addition, $\delta_v(G)$ is an integer multiple of $1/2$.

Proof. The inequality $\delta_v(G) \leq \delta(G)$ is direct.

Consider the set $\mathbb{T}'_v(G)$ of geodesic triangles T in G such that the three vertices of the triangle T belong to $V(G)$, and denote by $\delta'_v(G)$ the infimum of the constants μ such that every triangle in $\mathbb{T}'_v(G)$ is μ -thin. The argument in the proof of (ref. [79], Lemma 2.1) gives that $\delta'_v(G) = \delta_v(G)$.

Let us prove now $\delta(G) \leq 4\delta_v(G) + 1/2$. Let us assume that G is hyperbolic. If $\delta'_v(G) = \infty$, then the inequality is trivial. Thus, it suffices to consider the case $\delta'_v(G) < \infty$. By Theorem 7, there is a triangle $T = \{a, b, c\}$ that is a cycle with $a, b, c \in J(G)$ and $q \in [ab]$ such that $d(q, [ac] \cup [cb]) = \delta(T) = \delta(G)$. Assume that $a, b, c \in J(G) \setminus V(G)$ (otherwise, the argument is simpler). Let $a_1, a_2, b_1, b_2, c_1, c_2 \in T \cap V(G)$ such that $a \in [a_1, a_2], b \in [b_1, b_2], c \in [c_1, c_2]$ and $a_2, b_1 \in [ab], c_2, d_1 \in [cd], d_2, a_1 \in [ac]$. Since $H := \{a_2, b_1, b_2, c_1, c_2, a_1\}$ is a geodesic hexagon with vertices in $V(G)$, it is $4\delta'_v(G)$ -thin and every point $w \in [b_1, b_2] \cup [b_2 c_1] \cup [c_1, c_2] \cup [c_2 a_1] \cup [a_1, a_2]$ verifies $d(w, [ac] \cup [cb]) \leq 1/2$, we have

$$\begin{aligned} \delta(G) &= d(q, [ac] \cup [cb]) \leq d(q, [b_1, b_2] \cup [b_2 c_1] \cup [c_1, c_2] \cup [c_2 a_1] \cup [a_1, a_2]) + 1/2 \\ &\leq 4\delta'_v(G) + 1/2 = 4\delta_v(G) + 1/2. \end{aligned}$$

Assume that G is not hyperbolic. Therefore, for each $M > 0$ there is a triangle $T = \{a, b, c\}$ which is a cycle with $a, b, c \in J(G)$ and $q \in [ab]$ with $d(q, [ac] \cup [cb]) \geq M$. The previous argument gives $M \leq 4\delta_v(G) + 1/2$ and, since M is arbitrary, we conclude $\delta_v(G) = \infty = \delta(G)$.

Finally, consider any geodesic triangle $T = \{a, b, c\}$ in $\mathbb{T}_v(G)$. Since $d(q, [ac] \cup [cb]) = d(q, ([ac] \cup [cb]) \cap V(G))$, $d(q, [ac] \cup [cb])$ attains its maximum value when $q \in J(G)$. Hence, $\delta(T)$ is a multiple

of $1/2$ for any triangle $T \in \mathbb{T}_v(G)$. Since the set of non-negative numbers that are multiple of $1/2$ is a discrete set, $\delta(G)$ is an integer multiple of $1/2$ if G is hyperbolic, and there is a triangle $T = \{a, b, c\} \in \mathbb{T}_v(G)$ and $q \in [ab] \cap J(G)$ with $d(q, [ac] \cup [cb]) = \delta(T) = \delta_v(G)$. This finishes the proof. \square

Theorem 9. *If G_1 is a non-hyperbolic connected graph, then $G_1 \times P_2$ is not hyperbolic.*

Proof. Since G_1 is not hyperbolic, by Theorem 8, given any $R > 0$ there exists a triangle $T = \{x, y, z\}$ which is a cycle, with $x, y, z \in V(G_1)$ and such that T is not R -thin. Therefore, there exists some point $m \in T$, let us assume that $m \in [xy]$, such that $d_{G_1}(m, [yz] \cup [zx]) > R$.

Seeking for a contradiction let us assume that $G_1 \times P_2$ is δ -hyperbolic.

Suppose that for some $R > \delta$, there is a geodesic triangle $T = \{x, y, z\}$ that is an even cycle in G_1 , with $x, y, z \in V(G_1)$ and such that T is not R -thin. Consider the (closed) path $\Lambda = [xy] \cup [yz] \cup [zx]$. Then, since T has even length, the path $\Gamma_1\Lambda$ defines a cycle in $G_1 \times P_2$. Let $\gamma_1, \gamma_2, \gamma_3$ be the paths in $\Gamma_1\Lambda$ corresponding to $[xy], [yz], [zx]$, respectively. By Corollary 1, the curves γ_1, γ_2 and γ_3 are geodesics, and $d_{G_1 \times P_2}((\Gamma_1\Lambda)'(m), \gamma_2 \cup \gamma_3) > \delta$, leading to contradiction.

Suppose that, for every $R > 0$, there is a geodesic triangle $T = \{x, y, z\}$ which is an odd cycle, with $x, y, z \in V(G_1)$ and such that T is not R -thin.

Let $T_1 = \{x, y, z\}$ be a geodesic triangle as above and let us assume that $\text{diam}(T_1) = D > 8\delta$.

Let $T_2 = \{x', y', z'\}$ be another triangle as above such that T_2 is not $3(D + 8\delta)$ -thin, this is, there is a point m in one of the sides, let us call it σ , of T_2 such that $d_{G_1}(m, T_2 \setminus \sigma) > 3(D + 8\delta)$.

Let $g = [w_0w_k]$ with $w_0 \in T_1$ and $w_k \in T_2$ be a shortest geodesic in G_1 joining T_1 and T_2 (if T_1 and T_2 intersect, just assume that g is a single vertex, $w_0 = w_k$, in the intersection). See Figure 2.

Let us assume that $w_0 \in [xz]$ and $w_k \in [x'z']$. Then, let us consider the closed path C in G_1 given by the union of the geodesics in T_1, g , the geodesics in T_2 and the inverse of g from w_k to w_0 , this is,

$$C := [w_0x] \cup [xy] \cup [yz] \cup [zw_0] \cup [w_0w_k] \cup [w_kx'] \cup [x'y'] \cup [y'z'] \cup [z'w_k] \cup [w_kw_0].$$

Since T_1, T_2 are odd cycles, C is an even closed cycle. Therefore, Γ_1C defines a cycle in $G_1 \times P_2$. Moreover, by Remark 1, Γ_1C is a geodesic decagon in $G_1 \times P_2$ with sides $\gamma_1 = (\Gamma_1C)'([w_0x]), \gamma_2 = (\Gamma_1C)'([xy]), \gamma_3 = (\Gamma_1C)'([yz]), \gamma_4 = (\Gamma_1C)'([zw_0]), \gamma_5 = (\Gamma_1C)'([w_0w_k]), \gamma_6 = (\Gamma_1C)'([w_kx']), \gamma_7 = (\Gamma_1C)'([x'y']), \gamma_8 = (\Gamma_1C)'([y'z']), \gamma_9 = (\Gamma_1C)'([z'w_k])$ and $\gamma_{10} = (\Gamma_1C)'([w_kw_0])$.

Since we are assuming that $G_1 \times P_2$ is δ -hyperbolic, then for every $1 \leq i \leq 10$ and every point $p \in \gamma_i, d_{G_1 \times P_2}(p, C \setminus \gamma_i) \leq 8\delta$.

Let $p := (\Gamma_1C)'(m)$.

Case 1. Suppose that $d_{G_1}(m, T_1 \cup g) > 8\delta$. See Figure 2.

By assumption, $d_{G_1}(m, T_2 \setminus \sigma) > 8\delta$. If $\sigma = [x'y']$ (resp. $\sigma = [y'z']$), then $p \in \gamma_7$ (resp. $p \in \gamma_8$) and, by Corollary 1, $d_{G_1 \times P_2}(p, C \setminus \gamma_7) > 8\delta$ (resp. $d_{G_1 \times P_2}(p, C \setminus \gamma_8) > 8\delta$) leading to contradiction. If $\sigma = [x'z']$, since $[x'z'] = [x'w_k] \cup [w_kz']$, let us assume $m \in [x'w_k]$. Then, since $d_{G_1}(m, w_k) > 8\delta$, it follows that $d_{G_1}(m, [w_kz']) > 8\delta$. Thus, $p \in \gamma_6$ and, by Corollary 1, $d_{G_1 \times P_2}(p, C \setminus \gamma_6) > 8\delta$ leading to contradiction.

Case 2. Suppose that $d_{G_1}(m, T_1 \cup g) \leq 8\delta$ and $L(g) \leq 8\delta$. See the left side of Figure 3. Then, for every point q in $T_1 \cup g, d_{G_1}(m, q) \leq 8\delta + D + 8\delta$. In particular, $d_{G_1}(m, w_k) \leq 8\delta + D + 8\delta$. Therefore, $m \in [x'z']$ and let us assume that $m \in [x'w_k]$. Since $d_{G_1}(m, x') \geq d_{G_1}(m, [x'y'] \cup [y'z']) > 3(D + 8\delta)$, there is a point $m' \in [x'm] \subset [x'w_k]$ such that $d_{G_1}(m, m') = 2(D + 8\delta)$. Then, $d_{G_1}(m', T_1 \cup g) \geq 2(D + 8\delta) - D - 8\delta - 8\delta = D > 8\delta$. In addition, it is trivial to check that $d_{G_1}(m', [x'y'] \cup [y'z']) > 3(D + 8\delta) - 2(D + 8\delta) > 8\delta$ and since $[x'z']$ is a geodesic, $d_{G_1}(m', [z'w_k]) > 8\delta$. Thus, if $p' := (\Gamma_1C)'(m')$, then $p' \in \gamma_6$ and, by Corollary 1, $d_{G_1 \times P_2}(p', C \setminus \gamma_6) > 8\delta$ leading to contradiction.

Case 3. Suppose that $d_{G_1}(m, T_1 \cup g) \leq 8\delta$ and $L(g) > 8\delta$. See the right side of Figure 3. Since g is a shortest geodesic in G_1 joining T_1 and T_2 , this implies that $d_{G_1}(T_1, T_2) > 8\delta$ and $d_{G_1}(m, [w_0w_k]) \leq 8\delta$.

Moreover, $d_{G_1}(m, w_k) \leq 16\delta$. Otherwise, there is a point $q \in [w_0w_k]$ such that $d_{G_1}(m, q) \leq 8\delta$ and $d_{G_1}(q, w_k) > 8\delta$ which means that $d_{G_1}(q, w_0) < d_{G_1}(w_0, w_k) - 8\delta$ and $d_{G_1}(m, w_0) < d_{G_1}(w_0, w_k)$ leading to contradiction.

Since $d_{G_1}(m, w_k) \leq 16\delta$, $m \in [x'z']$. Let us assume that $m \in [x'w_k]$. Since $d_{G_1}(m, [x'y'] \cup [y'z']) > 3(D + 8\delta)$, there is a point $m' \in [x'm] \subset [x'w_k]$ such that $d_{G_1}(m, m') = 2(D + 8\delta)$. Let us see that $d_{G_1}(m', [w_0w_k]) > 8\delta$. Suppose there is some $q \in [w_0w_k]$ such that $d_{G_1}(m', q) \leq 8\delta$. Since $m' \in T_2$ and g is a shortest geodesic joining T_1 and T_2 , $d_{G_1}(q, w_k) \leq 8\delta$. However, $32\delta < 2(D + 8\delta) = d_{G_1}(m', m) \leq d_{G_1}(m', q) + d_{G_1}(q, w_k) + d_{G_1}(w_k, m) \leq 8\delta + 8\delta + 16\delta$ which is a contradiction. Hence, $d_{G_1}(m', [w_0w_k]) > 8\delta$. In addition, it is trivial to check that $d_{G_1}(m', [x'y'] \cup [y'z']) > 3(D + 8\delta) - 2(D + 8\delta) > 8\delta$ and since $[x'z']$ is a geodesic, $d_{G_1}(m', [z'w_k]) > 8\delta$. Thus, if $p' := (\Gamma_1 C)'(m')$, then $p' \in \gamma_6$ and, by Corollary 1, $d_{G_1 \times P_2}(p', C \setminus \gamma_6) > 8\delta$ leading to contradiction. \square




Figure 2. Two geodesic triangles, T_1, T_2 , which are odd cycles and a geodesic g joining them define




Figure 3. If $d_{G_1}(m, T_1 \cup g) \leq 8\delta$, then $m \in [x'z']$ and there is a point $m' \in [x'm] \subset [x'w_k]$ such that $d_{G_1}(m, m') = 2(D + 8\delta)$.

Proposition 3, Lemma 2 and Theorems 3, 4 and 9 have the following consequence.

Corollary 4. *If G_1 is a non-hyperbolic connected graph and G_2 is some non-trivial connected graph, then $G_1 \times G_2$ is not hyperbolic.*

Proposition 3 and Corollary 4 provide a necessary condition for the hyperbolicity of $G_1 \times G_2$.

Theorem 10. Let G_1, G_2 be non-trivial connected graphs. If $G_1 \times G_2$ is hyperbolic, then one factor graph is hyperbolic and the other one is bounded.

Theorems 3 and 4 show that this necessary condition is also sufficient if either G_2 has some odd cycle or G_1 and G_2 do not have odd cycles (when G_1 is a hyperbolic graph and G_2 is a bounded graph). We deal now with the other case, when G_1 has some odd cycle and G_2 does not have odd cycles.

Theorem 11. Let G_1 be a connected graph with some odd cycle and G_2 a non-trivial bounded connected graph without odd cycles. Assume that G_1 satisfies the following property: for each $M > 0$ there exist a geodesic g joining two minimal cycles of G_1 and a vertex $u \in g \cap V(G_1)$ with $d_{G_1}(u, \mathcal{C}(G_1)) \geq M$. Then, $G_1 \times G_2$ is not hyperbolic.

Proof. If G_1 is not hyperbolic, then Corollary 4 gives that $G_1 \times G_2$ is not hyperbolic. Assume now that G_1 is hyperbolic. By Theorem 2 and Lemma 2, we can assume that $G_2 = P_2$ and $V(P_2) = \{v_1, v_2\}$.

Fix $M > 0$ and choose a geodesic $g = [w_0 w_k] = [w_0, w_1] \cup [w_1, w_2] \cup \dots \cup [w_{k-1}, w_k]$ joining two minimal cycles in G_1 and $0 < r < k$ with $d_{G_1}(w_r, \mathcal{C}(G_1)) \geq M$.

Define the paths g_1 and g_2 in $G_1 \times P_2$ as $g_1 := \Gamma_1 g$ and $g_2 := \Gamma_2 g$. Since $L(g_1) = L(g_2) = L(g) = d_{G_1}(w_0, w_k)$, we have

$$d_{G_1 \times P_2}(g_1(w_0), g_1(w_k)) \leq L(g_1) = d_{G_1}(w_0, w_k), \quad d_{G_1 \times P_2}(g_2(w_0), g_2(w_k)) \leq L(g_2) = d_{G_1}(w_0, w_k).$$

Corollary 1 gives that

$$d_{G_1 \times P_2}(g_1(w_0), g_1(w_k)) \geq d_{G_1}(w_0, w_k), \quad d_{G_1 \times P_2}(g_2(w_0), g_2(w_k)) \geq d_{G_1}(w_0, w_k).$$

Hence, g_1 and g_2 are geodesics in $G_1 \times P_2$. Choose geodesics $g_3 = [g_1(w_0)g_2(w_0)]$ and $g_4 = [g_1(w_k)g_2(w_k)]$ in $G_1 \times P_2$. Since $d_{P_2}(v_1, v_2) = 1$ is odd, Proposition 2 gives

$$\begin{aligned} d_{G_1 \times P_2}(g_1(w_0), g_2(w_0)) &= \min \{L(\sigma) \mid \sigma \text{ is a } w_0, w_0\text{-walk}\} \\ &= \min \{L(\sigma) \mid \sigma \text{ cycle of odd length containing } w_0\}. \end{aligned}$$

Since w_0 belongs to a minimal cycle, $L(g_3) \leq 4\delta(G_1)$ by Lemma 4. In a similar way, we obtain $L(g_4) \leq 4\delta(G_1)$.

Consider the geodesic quadrilateral $Q := \{g_1, g_2, g_3, g_4\}$ in $G_1 \times P_2$. Thus, $d_{G_1 \times P_2}(g_1(w_r), g_2 \cup g_3 \cup g_4) \leq 2\delta(G_1 \times P_2)$. Since $\max \{L(g_3), L(g_4)\} \leq 4\delta(G_1)$, we deduce $d_{G_1 \times P_2}(g_1(w_r), g_2) \leq 2\delta(G_1 \times P_2) + 4\delta(G_1)$.

Let $0 \leq j \leq k$ with $d_{G_1 \times P_2}(g_1(w_r), g_2) = d_{G_1 \times P_2}(g_1(w_r), g_2(w_j))$. Let us define $w'_r := g_1(w_r)$ and $w'_j := g_2(w_j)$. Thus, Lemma 5 gives

$$\sqrt{M} \leq \sqrt{d_{G_1}(w_r, \mathcal{C}(G_1))} \leq d_{G_1 \times P_2}(w'_r, w'_j) = d_{G_1 \times P_2}(w'_r, g_2) \leq 2\delta(G_1 \times P_2) + 4\delta(G_1),$$

and since M is arbitrarily large, we deduce that $G_1 \times P_2$ is not hyperbolic. \square

Lemma 6. Let G_1 be a hyperbolic connected graph and suppose there is some constant $K > 0$ such that for every vertex $w \in G_1$, $d_{G_1}(w, \mathcal{C}(G_1)) \leq K$. Then, $G_1 \times P_2$ is hyperbolic.

Proof. Denote by v_1 and v_2 the vertices of P_2 . Let $i : V(G_1) \rightarrow V(G_1 \times P_2)$ defined as $i(w) := (w, v_1)$ for every $w \in G_1$.

For every $x, y \in V(G_1)$, by Corollary 1, $d_{G_1}(x, y) \leq d_{G_1 \times P_2}(i(x), i(y))$. By Corollary 3,

$$d_{G_1 \times P_2}(i(x), i(y)) \leq d_{G_1}(x, y) + 2d_{G_1}(x, \mathcal{C}(G_1)) + 4\delta(G_1) \leq d_{G_1}(x, y) + 2K + 4\delta(G_1).$$

Therefore, $i : V(G_1) \rightarrow V(G_1 \times P_2)$ is a $(1, 2K + 4\delta(G_1))$ -quasi-isometric embedding.

Notice that for every $(w, v_1) \in V(G_1 \times P_2)$, $(w, v_1) = i(w)$. In addition, for any $(w, v_2) \in V(G_1 \times P_2)$, since G_1 is connected, there is some edge $[w, w'] \in E(G_1)$ and we have $[(w, v_2), (w', v_1)] \in E(G_1 \times P_2)$. Therefore, $i : V(G_1) \rightarrow V(G_1 \times P_2)$ is 1-full.

Thus, by Lemma 1, G_1 and $G_1 \times P_2$ are quasi-isometric and, by Theorem 2, $G_1 \times P_2$ is hyperbolic. \square

Theorem 3 and Lemmas 2 and 6 give the following result.

Theorem 12. *Let G_1 be a hyperbolic connected graph and G_2 some non-trivial bounded connected graph. If there is some constant $K > 0$ such that for every vertex $w \in G_1$, $d_{G_1}(w, \mathcal{C}(G_1)) \leq K$, then $G_1 \times G_2$ is hyperbolic.*

We finish this section with a characterization of the hyperbolicity of $G_1 \times G_2$, under an additional hypothesis. We present first some lemmas.

Let J be a finite or infinite index set. Now, given a graph G_1 , we define some graphs related to G_1 which will be useful in the following results. Let $B_j := B_{G_1}(w_j, K_j)$ with $w_j \in V(G_1)$ and $K_j \in \mathbb{Z}^+$, for any $j \in J$, such that $\sup_j K_j = K < \infty$, $\bar{B}_{j_1} \cap \bar{B}_{j_2} = \emptyset$ if $j_1 \neq j_2$, and every odd cycle C in G_1 satisfies $C \cap B_j \neq \emptyset$ for some $j \in J$. Denote by G'_1 the subgraph of G_1 induced by $V(G_1) \setminus (\cup_j B_j)$. Let $N_j := \partial B_j = \{w \in V(G_1) : d_{G_1}(w, w_j) = K_j\}$. Denote by G_1^* the graph with $V(G_1^*) = V(G'_1) \cup (\cup_j \{w_j^*\})$, where w_j^* are additional vertices, and $E(G_1^*) = E(G'_1) \cup (\cup_j \{[w, w_j^*] : w \in N_j\})$. We have $G'_1 = G_1 \cap G_1^*$.

Lemma 7. *Let G_1 be a connected graph as above. Then, there is a quasi-isometry $g : G_1 \rightarrow G_1^*$ such that $g(w_j) = w_j^*$ for every $j \in J$.*

Proof. Let $f : V(G_1) \rightarrow V(G_1^*)$ defined as $f(u) = u$ for every $u \in V(G'_1)$, and $f(u) = w_i^*$ for every $u \in V(B_i)$. It is clear that $f : V(G_1) \rightarrow V(G_1^*)$ is 0-full.

Now, we focus on proving that $f : V(G_1) \rightarrow V(G_1^*)$ is a $(K, 2K)$ -quasi-isometric embedding. For every $u, v \in V(G_1)$, it is clear that $d_{G_1^*}(f(u), f(v)) \leq d_{G_1}(u, v)$.

Let us prove the other inequality. Fix $u, v \in V(G_1)$ and consider an oriented geodesic γ in G_1^* from $f(u)$ to $f(v)$.

Assume that $u, v \in V(G'_1)$. If $L(\gamma) = d_{G_1}(u, v)$, then $d_{G_1}(u, v) = d_{G_1^*}(f(u), f(v))$. If $L(\gamma) < d_{G_1}(u, v)$, then γ meets some w_j^* . Since γ is a compact set, it intersects only a finite number of w_j^* 's, which we denote by $w_{j_1}^*, \dots, w_{j_r}^*$. Since γ is an oriented curve from $f(u)$ to $f(v)$, we can assume that γ meets $w_{j_1}^*, \dots, w_{j_r}^*$ in this order.

Let us define the following vertices in γ

$$w_i^1 = [f(u)w_{j_i}^*] \cap N_{j_i}, \quad w_i^2 = [w_{j_i}^*f(v)] \cap N_{j_i},$$

for every $1 \leq i \leq r$. Note that $[w_i^2w_{i+1}^1] \subset G'_1$ for every $1 \leq i < r$ (it is possible to have $w_i^2 = w_{i+1}^1$).

Since $d_{G_1^*}(w_i^1, w_i^2) = 2$ and $d_{G_1}(w_i^1, w_i^2) \leq 2K$, we have $d_{G_1^*}(w_i^1, w_i^2) \geq \frac{1}{K} d_{G_1}(w_i^1, w_i^2)$ for every $1 \leq i \leq r$. Thus,

$$\begin{aligned} d_{G_1^*}(f(u), f(v)) &= d_{G_1^*}(f(u), w_1^1) + \sum_{i=1}^r d_{G_1^*}(w_i^1, w_i^2) + \sum_{i=1}^{r-1} d_{G_1^*}(w_i^2, w_{i+1}^1) + d_{G_1^*}(w_r^2, f(v)) \\ &\geq d_{G_1}(u, w_1^1) + \frac{1}{K} \sum_{i=1}^r d_{G_1}(w_i^1, w_i^2) + \sum_{i=1}^{r-1} d_{G_1}(w_i^2, w_{i+1}^1) + d_{G_1}(w_r^2, v) \\ &\geq \frac{1}{K} \left(d_{G_1}(u, w_1^1) + \sum_{i=1}^r d_{G_1}(w_i^1, w_i^2) + \sum_{i=1}^{r-1} d_{G_1}(w_i^2, w_{i+1}^1) + d_{G_1}(w_r^2, v) \right) \\ &\geq \frac{1}{K} d_{G_1}(u, v). \end{aligned}$$

Assume that $f(u) = f(v)$. Therefore, there exists j with $u, v \in B_j$ and

$$d_{G_1^*}(f(u), f(v)) = 0 > d_{G_1}(u, v) - 2K.$$

Assume now that u and/or v does not belong to $V(G'_1)$ and $f(u) \neq f(v)$. Let u_0, v_0 be the closest vertices in $V(G'_1) \cap \gamma$ to $f(u), f(v)$, respectively (it is possible to have $u_0 = f(u)$ or $v_0 = f(v)$). Since $u_0, v_0 \in V(G'_1)$, $u_0 = f(u_0), v_0 = f(v_0)$, we have $d_{G_1}(u, u_0) < 2K$ and $d_{G_1}(v, v_0) < 2K$. Hence,

$$\begin{aligned} d_{G_1^*}(f(u), f(v)) &= d_{G_1^*}(f(u), u_0) + d_{G_1^*}(u_0, v_0) + d_{G_1^*}(v_0, f(v)) \\ &\geq d_{G_1^*}(f(u_0), f(v_0)) \\ &\geq \frac{1}{K} d_{G_1}(u_0, v_0) \\ &\geq \frac{1}{K} (d_{G_1}(u, v) - d_{G_1}(u, u_0) - d_{G_1}(v, v_0)) \\ &> \frac{1}{K} d_{G_1}(u, v) - 4. \end{aligned}$$

If $K \geq 2$, then $d_{G_1^*}(f(u), f(v)) > \frac{1}{K} d_{G_1}(u, v) - 2K$. If $K = 1$, then $d_{G_1}(u, u_0) \leq 1, d_{G_1}(v, v_0) \leq 1$, and $d_{G_1^*}(f(u), f(v)) \geq d_{G_1}(u, v) - 2$.

Finally, we conclude that $f : V(G_1) \rightarrow V(G_1^*)$ is a $(K, 2K)$ -quasi-isometric embedding. Thus, Lemma 1 provides a quasi-isometry $g : G_1 \rightarrow G_1^*$ with the required property. \square

Definition 3. Given a connected graph G_1 and some index set J , let $\mathcal{B}_J = \{B_j\}_{j \in J}$ be a family of balls where $B_j := B_{G_1}(w_j, K_j)$ with $w_j \in V(G_1)$, $K_j \in \mathbb{Z}^+$ for any $j \in J$, $\sup_j K_j = K < \infty$ and $\overline{B}_{j_1} \cap \overline{B}_{j_2} = \emptyset$ if $j_1 \neq j_2$. Suppose that every odd cycle C in G_1 satisfies that $C \cap B_j \neq \emptyset$ for some $j \in J$. If there is some constant $M > 0$ such that for every $j \in J$, there is an odd cycle C_j such that $C_j \cap B_j \neq \emptyset$ with $L(C_j) < M$, then we say that \mathcal{B}_J is M -regular.

Remark 2. If J is finite, then there exists $M > 0$ such that $\{B_j\}_{j \in J}$ is M -regular.

Denote by G^* the graph with $V(G^*) = V(G'_1 \times P_2) \cup (\cup_j \{w_j^*\})$, where G'_1 is a graph as above and w_j^* are additional vertices, and $E(G^*) = E(G'_1 \times P_2) \cup (\cup_j \{[w, w_j^*] : \pi_1(w) \in N_j\})$.

Lemma 8. Let G_1 be a connected graph as above and P_2 with $V(P_2) = \{v_1, v_2\}$. If G_1 is hyperbolic and \mathcal{B}_J as above is M -regular, then there exists a quasi-isometry $f : G_1 \times P_2 \rightarrow G^*$ with $f(w_j, v_i) = w_j^*$ for every $j \in J$ and $i \in \{1, 2\}$.

Proof. Let $F : V(G_1 \times P_2) \rightarrow V(G^*)$ defined as $F(v, v_i) = (v, v_i)$ for every $v \in V(G'_1)$, and $F(v, v_i) = w_j^*$ for every $v \in V(B_j)$. It is clear that $F : V(G_1 \times P_2) \rightarrow V(G^*)$ is 0-full. Recall that we denote by $\pi_1 : G_1 \times P_2 \rightarrow G_1$ the projection map. Define $\pi^* : G^* \rightarrow G_1$ as $\pi^* = \pi_1$ on $G'_1 \times P_2$ and $\pi^*(x) = w_j$ for every x with $d_{G^*}(x, w_j^*) < 1$ for some $j \in J$.

Now, we focus on proving that $F : V(G_1 \times P_2) \rightarrow V(G^*)$ is a quasi-isometric embedding. For every $(w, v_i), (w', v_{i'}) \in V(G_1 \times P_2)$, one can check

$$d_{G^*}(F(w, v_i), F(w', v_{i'})) \leq d_{G_1 \times P_2}((w, v_i), (w', v_{i'})).$$

To prove the other inequality, let us fix $(w, v_i), (w', v_{i'}) \in V(G'_1 \times P_2)$ (the inequalities in other cases can be obtained from the one in this case, as in the proof of Lemma 7). Consider a geodesic $\gamma := [F(w, v_i)F(w', v_{i'})]$ in G^* . If $L(\gamma) = d_{G_1 \times P_2}((w, v_i), (w', v_{i'}))$, then

$$d_{G^*}(F(w, v_i), F(w', v_{i'})) = d_{G_1 \times P_2}((w, v_i), (w', v_{i'})).$$

If $L(\gamma) < d_{G_1 \times P_2}((w, v_i), (w', v_{i'}))$, then $\pi^*(\gamma)$ meets some B_j . Since γ is a compact set, $\pi^*(\gamma)$ intersects just a finite number of B_j 's, which we denote by B_{j_1}, \dots, B_{j_r} . We consider γ as an oriented curve from $F(w, v_i)$ to $F(w', v_{i'})$; thus we can assume that $\pi^*(\gamma)$ meets B_{j_1}, \dots, B_{j_r} in this order.

Let us define the following set of vertices in γ

$$\{w_i^1, w_i^2\} := \gamma \cap (N_{j_i} \times P_2),$$

for every $1 \leq i \leq r$, such that $d_{G_1 \times P_2}((w, v_i), w_i^1) < d_{G_1 \times P_2}((w, v_i), w_i^2)$. Note that $[w_i^2 w_{i+1}^1] \subset G_1 \times P_2$ for every $1 \leq i < r$ and $d_{G_1 \times P_2}(w_i^2, w_{i+1}^1) \geq 1$ since $\bar{B}_{j_i} \cap \bar{B}_{j_{i+1}} = \emptyset$.

If $d_{G_1}(\pi(w_i^1), \pi(w_i^2)) = d_{G_1 \times P_2}(w_i^1, w_i^2)$ for some $1 \leq i \leq r$, then $d_{G_1 \times P_2}(w_i^1, w_i^2) \leq 2K$. Since $d_{G_1 \times P_2}(w_i^2, w_{i+1}^1) \geq 1$ for $1 \leq i < r$, we have that $d_{G_1 \times P_2}(w_i^1, w_i^2) \leq 2K d_{G_1 \times P_2}(w_i^2, w_{i+1}^1)$ in this case.

If $d_{G_1}(\pi_1(w_i^1), \pi_1(w_i^2)) < d_{G_1 \times P_2}(w_i^1, w_i^2)$ for some $1 \leq i \leq r$, then $d_{G_1}(\pi_1(w_i^1), \pi_1(w_i^2)) + d_{G_1 \times P_2}(w_i^1, w_i^2)$ is odd.

Since B_j is M -regular, consider an odd cycle C with $C \cap B_{j_i} \neq \emptyset$ and $L(C) < M$, and let $b_i \in C \cap B_{j_i}$ and $[\pi_1(w_i^1) b_i], [b_i \pi_1(w_i^2)]$ geodesics in G_1 . Thus, $[\pi_1(w_i^1) b_i] \cup [b_i \pi_1(w_i^2)]$ and $[\pi_1(w_i^1) b_i] \cup C \cup [b_i \pi_1(w_i^2)]$ have different parity which means that one of them has different parity from $[\pi_1(w_i^1) \pi_1(w_i^2)]$. Then, $d_{G_1 \times P_2}(w_i^1, w_i^2) \leq L([\pi_1(w_i^1) b_i] \cup C \cup [b_i \pi_1(w_i^2)]) \leq 4K + M$. Since $d_{G_1 \times P_2}(w_i^2, w_{i+1}^1) \geq 1$ for $1 \leq i < r$, we have that $d_{G_1 \times P_2}(w_i^1, w_i^2) \leq (4K + M) d_{G_1 \times P_2}(w_i^2, w_{i+1}^1)$ in this case.

Thus, we have that $d_{G_1 \times P_2}(w_i^1, w_i^2) \leq 4K + M$ for every $1 \leq i \leq r$ and $d_{G_1 \times P_2}(w_i^1, w_i^2) \leq (4K + M) d_{G_1 \times P_2}(w_i^2, w_{i+1}^1)$ for every $1 \leq i < r$. Therefore,

$$\begin{aligned} d_{G_1 \times P_2}((w, v_i), (w', v_{i'})) &\leq d_{G_1 \times P_2}((w, v_i), w_1^1) + \sum_{i=1}^r d_{G_1 \times P_2}(w_i^1, w_i^2) + \sum_{i=1}^{r-1} d_{G_1 \times P_2}(w_i^2, w_{i+1}^1) \\ &\quad + d_{G_1 \times P_2}(w_r^2, (w', v_{i'})) \\ &\leq d_{G_1 \times P_2}((w, v_i), w_1^1) + d_{G_1 \times P_2}(w_r^2, (w', v_{i'})) + (4K + M + 1) \sum_{i=1}^{r-1} d_{G_1 \times P_2}(w_i^2, w_{i+1}^1) \\ &\quad + d_{G_1 \times P_2}(w_r^1, w_r^2) \\ &= d_{G^*}(F(w, v_i), F(w_1^1)) + d_{G^*}(F(w_r^2), F(w', v_{i'})) + (4K + M + 1) \sum_{i=1}^{r-1} d_{G^*}(F(w_i^2), F(w_{i+1}^1)) \\ &\quad + d_{G_1 \times P_2}(w_r^1, w_r^2) \\ &\leq (4K + M + 1) (d_{G^*}(F(w, v_i), F(w_1^1)) + d_{G^*}(F(w_r^2), F(w', v_{i'})) + \sum_{i=1}^{r-1} d_{G^*}(F(w_i^2), F(w_{i+1}^1))) + 4K + M \\ &\leq (4K + M + 1) d_{G^*}(F(w, v_i), F(w', v_{i'})) + 4K + M. \end{aligned}$$

We conclude that $F : V(G_1 \times P_2) \rightarrow V(G^*)$ is a quasi-isometric embedding. Thus, Lemma 1 provides a quasi-isometry $f : G_1 \times P_2 \rightarrow G^*$ with the required property. \square

Definition 4. Given a geodesic metric space X and closed connected pairwise disjoint subsets $\{\eta_j\}_{j \in J}$ of X , we consider another copy X' of X . The double DX of X is the union of X and X' obtained by identifying the corresponding points in each η_j and η'_j .

Definition 5. Let us consider $H > 0$, a metric space X , and subsets $Y, Z \subseteq X$. The set $V_H(Y) := \{x \in X : d(x, Y) \leq H\}$ is called the H -neighborhood of Y in X . The Hausdorff distance of Y to Z is defined by $\mathcal{H}(Y, Z) := \inf\{H > 0 : Y \subseteq V_H(Z), Z \subseteq V_H(Y)\}$.

The following results in [15,80] will be useful.

Theorem 13. ([80], Theorem 3.2) Let us consider a geodesic metric space X and closed connected pairwise disjoint subsets $\{\eta_j\}_{j \in J}$ of X , such that the double DX is a geodesic metric space. Then, the following conditions are equivalent:

- (1) DX is hyperbolic.
- (2) X is hyperbolic and there exists a constant c_1 such that for every $k, l \in J$ and $a \in \eta_k, b \in \eta_l$ we have $d_X(x, \cup_{j \in J} \eta_j) \leq c_1$ for every $x \in [ab] \subset X$.
- (3) X is hyperbolic and there exist constants c_2, α, β such that for every $k, l \in J$ and $a \in \eta_k, b \in \eta_l$ we have $d_X(x, \cup_{j \in J} \eta_j) \leq c_2$ for every x in some (α, β) -quasi-geodesic joining a with b in X .

Theorem 14. ([15], p. 87) For each $\delta \geq 0, a \geq 1$ and $b \geq 0$, there exists a constant $H = H(\delta, a, b)$ with the following property:

Let us consider a δ -hyperbolic geodesic metric space X and an (a, b) -quasigeodesic g starting in x and finishing in y . If γ is a geodesic joining x and y , then $\mathcal{H}(g, \gamma) \leq H$.

This property is called geodesic stability. It is well-known that hyperbolicity is, in fact, equivalent to geodesic stability [81].

Theorem 15. Let G_1 be a connected graph and $B_j := B_{G_1}(w_j, K_j)$ with $w_j \in V(G_1)$ and $K_j \in \mathbb{Z}^+$, for any $j \in J$, such that $\sup_j K_j = K < \infty, \bar{B}_{j_1} \cap \bar{B}_{j_2} = \emptyset$ if $j_1 \neq j_2$, and every odd cycle C in G_1 satisfies $C \cap B_j \neq \emptyset$ for some $j \in J$. Suppose $\{B_j\}_{j \in J}$ is M -regular for some $M > 0$. Let G_2 be a non-trivial bounded connected graph without odd cycles. Then, the following statements are equivalent:

- (1) $G_1 \times G_2$ is hyperbolic.
- (2) G_1 is hyperbolic and there exists a constant c_1 , such that for every $k, l \in J$ and $w_k \in B_k, w_l \in B_l$ there exists a geodesic $[w_k w_l]$ in G_1 with $d_{G_1}(x, \cup_{j \in J} w_j) \leq c_1$ for every $x \in [w_k w_l]$.
- (3) G_1 is hyperbolic and there exist constants c_2, α, β , such that for every $k, l \in J$ we have $d_{G_1}(x, \cup_{j \in J} w_j) \leq c_2$ for every x in some (α, β) -quasi-geodesic joining w_k with w_l in G_1 .

Proof. Items (2) and (3) are equivalent by geodesic stability in G_1 (see Theorem 14).

Assume that (2) holds. By Lemma 7, there exists an (α, β) -quasi-isometry $f : G_1 \rightarrow G_1^*$ with $f(w_j) = w_j^*$ for every $j \in J$. Given $k, l \in J, f([w_k w_l])$ is an (α, β) -quasi-geodesic with endpoints w_k^* and w_l^* in G_1^* . Given $x \in f([w_k w_l])$, we have $x = f(x_0)$ with $x_0 \in [w_k w_l]$ and $d_{G_1^*}(x, \cup_{j \in J} w_j^*) \leq \alpha d_{G_1}(x_0, \cup_{j \in J} w_j) + \beta \leq \alpha c_1 + \beta$. Taking $X = G_1^*, DX = G^*$ and $\eta_j = w_j^*$ for every $j \in J$, Theorem 13 gives that G^* is hyperbolic. Now, Lemma 8 gives that $G_1 \times P_2$ is hyperbolic and we conclude that $G_1 \times G_2$ is hyperbolic by Lemma 2.

Now, suppose (1) holds. By Lemma 2, $G_1 \times P_2$ is hyperbolic and, by Theorem 9, G_1 is hyperbolic. Then, Lemma 8 gives that G^* is hyperbolic and taking $X = G_1^*, DX = G^*$ and $\eta_j = w_j^*$ for every $j \in J$, by Theorem 13, (2) holds. \square

Theorem 15 and Remark 2 have the following consequence.

Corollary 5. Let G_1 be a connected graph and suppose that there are a positive integer K and a vertex $w \in G_1$, such that every odd cycle in G_1 intersects the open ball $B := B_{G_1}(w, K)$. Let G_2 be a non-trivial bounded connected graph without odd cycles. Then, $G_1 \times G_2$ is hyperbolic if and only if G_1 is hyperbolic.

4. Bounds for the Hyperbolicity Constant of Some Direct Products

The following well-known result will be useful (see a proof, e.g., in ([47], Theorem 8)).

Theorem 16. In any connected graph G the inequality $\delta(G) \leq (\text{diam } G)/2$ holds.

Remark 3. Note that, if G_1 is a bipartite connected graph, then $\text{diam } G_1 = \text{diam } V(G_1)$. Furthermore, if G_2 is a bipartite connected graph, then the product $G_1 \times G_2$ has exactly two connected components, which are denoted

by $(G_1 \times G_2)^1$ and $(G_1 \times G_2)^2$, where each one is a bipartite graph and, consequently, $\text{diam}(G_1 \times G_2)^i = \text{diam } V((G_1 \times G_2)^i)$ for $i \in \{1, 2\}$.

Remark 4. Let P_m, P_n be two path graphs with $m \geq n \geq 2$. The product $P_m \times P_n$ has exactly two connected components, which will be denoted by $(P_m \times P_n)^1$ and $(P_m \times P_n)^2$. If $u, v \in V((P_m \times P_n)^i)$ for $i \in \{1, 2\}$, then $d_{(P_m \times P_n)^i}(u, v) = \max \{d_{P_m}(\pi_1(u), \pi_1(v)), d_{P_n}(\pi_2(u), \pi_2(v))\}$ and $\text{diam}(P_m \times P_n)^i = \text{diam } V((P_m \times P_n)^i) = m - 1$.

Furthermore, if $m_1 \leq m$ and $n_1 \leq n$, then $\delta(P_m \times P_n) \geq \delta(P_{m_1} \times P_{n_1})$.

Lemma 9. Let P_m, P_n be two path graphs with $m \geq n \geq 3$, and let γ be a geodesic in $P_m \times P_n$ such that there are two different vertices u, v in γ , with $\pi_1(u) = \pi_1(v)$. Then, $L(\gamma) \leq n - 1$.

Remark 5. Note that the conclusion of Lemma 9 does not hold for $n = 2$, since we always have $L(\gamma) \geq 2$.

Proof. Let $\gamma := [xy]$, and let $V(P_m) = \{v_1, \dots, v_m\}, V(P_n) = \{w_1, \dots, w_n\}$ be the sets of vertices in P_m, P_n , respectively, such that $[v_j, v_{j+1}] \in E(P_m)$ and $[w_i, w_{i+1}] \in E(P_n)$ for $1 \leq j < m, 1 \leq i < n$. Seeking for a contradiction, assume that $L(\gamma) > n - 1$. Notice that if $[uv]$ denotes the geodesic contained in γ joining u and v , then π_2 restricted to $[uv]$ is injective. Consider two vertices $u', v' \in \gamma$ such that $[uv] \subseteq [u'v'] \subseteq \gamma$, π_2 is injective in $[u'v']$ and $\pi_2(u') = w_{i_1}, \pi_2(v') = w_{i_2}$ with $i_2 - i_1$ maximal under these conditions. See Figure 4.

Since $L(\gamma) > n - 1 \geq i_2 - i_1$, either there is an edge $[v', w]$ in $G_1 \times G_2$ such that $[v', w] \cap (\gamma \setminus [u'v']) \neq \emptyset$ or there is an edge $[u', w']$ in $G_1 \times G_2$ such that $[u', w'] \cap (\gamma \setminus [u'v']) \neq \emptyset$. In addition, since $L(\gamma) > n - 1$, notice that π_2 is not injective in γ . Moreover, since $i_2 - i_1$ is maximal, if $\pi_2(w) = w_{i_2+1}$, then $w \notin \gamma$, and since $L(\gamma) > n - 1, u' \notin \{x, y\}$ and $\pi_2(w') = w_{i_1+1}$. Thus, either $\pi_2(w) = w_{i_2-1}$ or $\pi_2(w') = w_{i_1+1}$.

Hence, let us assume that there is an edge $[v', w]$ in $G_1 \times G_2$ such that $[v', w] \cap (\gamma \setminus [u'v']) \neq \emptyset$ with $\pi_2(w) = w_{i_2-1}$ (otherwise, if there is an edge $[u', w']$ in $G_1 \times G_2$ such that $[u', w'] \cap (\gamma \setminus [u'v']) \neq \emptyset$ with $\pi_2(w') = w_{i_1+1}$, the proof is similar).

Suppose $\pi_1(v') = v_j$. Let v'' be the vertex in $[u'v']$ such that $\pi_2(v'') = w_{i_2-1}$. Then, by construction of $G_1 \times G_2$, since $v'' \neq w$, it follows that $\{\pi_1(v''), \pi_1(w)\} = \{v_{j-1}, v_{j+1}\}$. Therefore, in particular, $1 < j < m$.

Assume that $v'' = (v_{j-1}, w_{i_2-1})$ (if $v'' = (v_{j+1}, w_{i_2-1})$, then the argument is similar). Therefore, $w = (v_{j+1}, w_{i_2-1})$.

Consider the geodesic

$$\sigma = [(v_{j+1}, w_{i_2-1}), (v_j, w_{i_2-2})] \cup [(v_j, w_{i_2-2}), (v_{j-1}, w_{i_2-3})] \cup [(v_{j-1}, w_{i_2-3}), (v_{j-2}, w_{i_2-4})] \cup \dots$$

Since $\pi_1(u) = \pi_1(v)$, there is a vertex ξ of $V(P_m \times P_n)$ in $[u'v'] \cap \sigma$. Let $s \in [v', w] \cap \gamma$ with $s \neq v'$. Let σ_0 be the geodesic contained in σ joining ξ and w . Let γ_0 be the geodesic contained in γ joining ξ and s . Hence, $L(\sigma_0 \cup [ws]) < L(\sigma_0) + 1 < L(\gamma_0)$ leading to contradiction. \square




Figure 4. For any geodesic γ in $P_m \times P_n$ with $\pi_1(u) = \pi_1(v)$ for some different vertices u, v in γ , then $L(\gamma) \leq n - 1$.

Theorem 17. Let P_m, P_n be two path graphs with $m \geq n \geq 2$. If $n = 2$, then $\delta(P_m \times P_2) = 0$. If $n \geq 3$, then

$$\min \left\{ \frac{m}{2}, n - 1 \right\} - 1 \leq \delta(P_m \times P_n) \leq \min \left\{ \frac{m}{2}, n \right\} - \frac{1}{2}.$$

Furthermore, if $m \leq 2n - 3$ and m is odd, then $\delta(P_m \times P_n) = (m - 1)/2$.

Proof. If $m \geq 2$, then $P_m \times P_2$ has two connected components isomorphic to P_m , and $\delta(P_m \times P_2) = 0$.

Assume that $n \geq 3$. By symmetry, it suffices to prove the inequalities for $\delta((P_m \times P_n)^1)$. Hence, Theorem 16 and Remark 4 give $\delta((P_m \times P_n)^1) \leq \frac{m-1}{2}$. By Theorem 7, there exists a geodesic triangle $T = \{x, y, z\} \in \mathbb{T}_1(P_m \times P_n)$ with $p \in \gamma_1 := [xy], \gamma_2 := [xz], \gamma_3 := [yz]$, and $\delta((P_m \times P_n)^1) = \delta(T) = d_{(P_m \times P_n)^1}(p, \gamma_2 \cup \gamma_3)$. Let $u \in V(\gamma_1)$ such that $d_{(P_m \times P_n)^1}(p, u) \leq 1/2$.

To prove $\delta((P_m \times P_n)^1) \leq n - 1/2$, we consider two cases.

Assume first that there is at least a vertex $v \in V((P_m \times P_n)^1) \cap T \setminus \{u\}$ such that $\pi_1(u) = \pi_1(v)$. If $v \notin \gamma_1$, then $v \in \gamma_2 \cup \gamma_3$ and

$$\delta(T) = d_{(P_m \times P_n)^1}(p, \gamma_2 \cup \gamma_3) \leq 1/2 + d_{(P_m \times P_n)^1}(u, v) \leq n - 1/2.$$

If $v \in \gamma_1$, then $L(\gamma_1) \leq n - 1$ by Lemma 9, and

$$\delta(T) = d_{(P_m \times P_n)^1}(p, \gamma_2 \cup \gamma_3) \leq d_{(P_m \times P_n)^1}(p, \{x, y\}) \leq (n - 1)/2 < n - 1/2.$$

Assume now that there is not a vertex $v \in V((P_m \times P_n)^1) \cap T \setminus \{u\}$ such that $\pi_1(u) = \pi_1(v)$. Then, there exist two different vertices v_1, v_2 in $T \setminus \{u\}$ such that $d_{(P_m \times P_n)^1}(u, v_1) = d_{(P_m \times P_n)^1}(u, v_2) = 1$, and $\pi_1(v_1) = \pi_1(v_2)$. If v_1 or v_2 belongs to $\gamma_2 \cup \gamma_3$, then $\delta(T) = d_{(P_m \times P_n)^1}(p, \gamma_2 \cup \gamma_3) \leq 3/2 \leq n - 1/2$. Otherwise, $v_1, v_2 \in \gamma_1 \setminus \{u\}$. Lemma 9 gives $L(\gamma_1) \leq n - 1$, and we have that

$$\delta(T) = d_{(P_m \times P_n)^1}(p, \gamma_2 \cup \gamma_3) \leq d_{(P_m \times P_n)^1}(p, \{x, y\}) \leq (n - 1)/2 < n - 1/2.$$

To prove the lower bound, denote the vertices of P_m and P_n by $V(P_m) = \{w_1, w_2, w_3, \dots, w_m\}$ and $V(P_n) = \{v_1, v_2, v_3, \dots, v_n\}$, with $[w_i, w_{i+1}] \in E(P_m)$ for $1 \leq i < m$ and $[v_i, v_{i+1}] \in E(P_n)$ for $1 \leq i < n$.

Let $(P_m \times P_n)^1$ be the connected component of $P_m \times P_n$ containing (w_1, v_{n-1}) .

Assume first that $m \geq 2n - 3$. Consider the following curves in $(P_m \times P_n)^1$:

$$\begin{aligned} \gamma_1 &:= [(w_1, v_{n-1}), (w_2, v_n)] \cup [(w_2, v_n), (w_3, v_{n-1})] \cup [(w_3, v_{n-1}), (w_4, v_n)] \cup \dots \cup [(w_{2n-4}, v_n), (w_{2n-3}, v_{n-1})], \\ \gamma_2 &:= [(w_1, v_{n-1}), (w_2, v_{n-2})] \cup [(w_2, v_{n-2}), (w_3, v_{n-3})] \cup \dots \cup [(w_{n-2}, v_2), (w_{n-1}, v_1)] \cup [(w_{n-1}, v_1), (w_n, v_2)] \\ &\quad \cup \dots \cup [(w_{2n-4}, v_{n-2}), (w_{2n-3}, v_{n-1})]. \end{aligned}$$

Corollary 1 gives that γ_1, γ_2 are geodesics. If B is the geodesic bigon $B = \{\gamma_1, \gamma_2\}$, then Remark 4 gives that

$$\delta(P_m \times P_n) \geq \delta(B) \geq d_{(P_m \times P_n)^1}((w_{n-1}, v_1), \gamma_1) = n - 2.$$

If m is odd with $m \leq 2n - 3$, then $n - (m + 1)/2 \geq 1$ and we can consider the curves in $(P_m \times P_n)^1$:

$$\begin{aligned} \gamma_1 &:= [(w_1, v_{n-1}), (w_2, v_n)] \cup [(w_2, v_n), (w_3, v_{n-1})] \cup [(w_3, v_{n-1}), (w_4, v_n)] \cup \dots \cup [(w_{m-1}, v_n), (w_m, v_{n-1})], \\ \gamma_2 &:= [(w_1, v_{n-1}), (w_2, v_{n-2})] \cup [(w_2, v_{n-2}), (w_3, v_{n-3})] \cup \dots \cup [(w_{(m+1)/2-1}, v_{n-(m+1)/2+1}), (w_{(m+1)/2}, v_{n-(m+1)/2})] \\ &\quad \cup [(w_{(m+1)/2}, v_{n-(m+1)/2}), (w_{(m+1)/2+1}, v_{n-(m+1)/2+1})] \cup \dots \cup [(w_{m-1}, v_{n-2}), (w_m, v_{n-1})]. \end{aligned}$$

Corollary 1 gives that γ_1, γ_2 are geodesics. If $B = \{\gamma_1, \gamma_2\}$, then Remark 4 gives that

$$\delta(P_m \times P_n) \geq \delta(B) \geq d_{(P_m \times P_n)^1}((w_{(m+1)/2}, v_{n-(m+1)/2}), \gamma_1) = (m - 1)/2.$$

By Remark 4, if m is even with $m - 1 \leq 2n - 3$, then we have that

$$\delta(P_m \times P_n) \geq \delta(P_{m-1} \times P_n) \geq (m - 2)/2.$$

Hence,

$$\delta(P_m \times P_n) \geq \begin{cases} n - 2, & \text{if } m \geq 2n - 3 \\ (m - 2)/2, & \text{if } m \leq 2n - 2 \end{cases} = \min \left\{ n - 2, \frac{m - 2}{2} \right\} = \min \left\{ \frac{m}{2}, n - 1 \right\} - 1.$$

Furthermore, if $m \leq 2n - 3$ and m is odd, then we have proven $(m - 1)/2 \leq \delta(P_m \times P_n) \leq (m - 1)/2$. \square

Theorem 18. If G_1 and G_2 are bipartite connected graphs with $k_1 := \text{diam } V(G_1)$ and $k_2 := \text{diam } V(G_2)$ such that $k_1 \geq k_2 \geq 1$, then

$$\max \left\{ \min \left\{ \frac{k_1 - 1}{2}, k_2 - 1 \right\}, \delta(G_1), \delta(G_2) \right\} \leq \delta(G_1 \times G_2) \leq \frac{k_1}{2}.$$

Furthermore, if $k_1 \leq 2k_2 - 2$ and k_1 is even, then $\delta(G_1 \times G_2) = k_1/2$.

Proof. Corollary 1, Theorem 16 and Remark 3 give us the upper bound.

To prove the lower bound, we can see that there exist two path graphs P_{k_1+1}, P_{k_2+1} which are isometric subgraphs of G_1 and G_2 , respectively. It is easy to check that $P_{k_1+1} \times P_{k_2+1}$ is an isometric subgraph of $G_1 \times G_2$. By Lemma 3 and Theorem 17, we have

$$\min \left\{ \frac{k_1 - 1}{2}, k_2 - 1 \right\} \leq \delta(P_{k_1+1} \times P_{k_2+1}) \leq \delta(G_1 \times G_2).$$

Using a similar argument as above, we have $\delta(P_2 \times G_2) \leq \delta(G_1 \times G_2)$ and $\delta(G_1 \times P_2) \leq \delta(G_1 \times G_2)$. Thus, since $(G_1 \times P_2)^i \simeq G_1$ and $(P_2 \times G_2)^i \simeq G_2$ for $i \in \{1, 2\}$, we obtain the first statement.

Furthermore, if $k_1 + 1 \leq 2(k_2 + 1) - 3$ and $k_1 + 1$ is odd, then Theorem 17 gives $\delta(P_{k_1+1} \times P_{k_2+1}) = k_1/2$, and we conclude $\delta(G_1 \times G_2) = k_1/2$. \square

The following result deals just with odd cycles since otherwise we can apply Theorem 18.

Theorem 19. For every odd number $m \geq 3$ and every $n \geq 2$,

$$\delta(C_m \times P_n) = \begin{cases} m/2, & \text{if } n - 1 \leq m, \\ (n - 1)/2, & \text{if } m < n - 1 < 2m, \\ m - 1/2, & \text{if } n - 1 \geq 2m. \end{cases}$$

Proof. Let $V(C_m) = \{w_1, \dots, w_m\}$ and $V(P_n) = \{v_1, \dots, v_n\}$ be the sets of vertices in C_m and P_n , respectively, such that $[w_1, w_m], [w_j, w_{j+1}] \in E(C_m)$ and $[v_i, v_{i+1}] \in E(P_n)$ for $j \in \{1, \dots, m - 1\}$, $i \in \{1, \dots, n - 1\}$. Note that for $1 \leq j, r \leq m$ and $1 \leq i, s \leq n$, we have $d_{C_m \times P_n}((w_j, v_i), (w_r, v_s)) = \max\{|i - s|, |j - r|\}$, if $|i - s| \equiv |j - r| \pmod{2}$, or $d_{C_m \times P_n}((w_j, v_i), (w_r, v_s)) = \max\{|i - s|, m - |j - r|\}$, if $|i - s| \not\equiv |j - r| \pmod{2}$. Besides, we have $\text{diam}(C_m \times P_n) = \text{diam}(V(C_m \times P_n))$, i.e., $\text{diam}(C_m \times P_n) = m$ if $n - 1 \leq m$, and $\text{diam}(C_m \times P_n) = n - 1$ if $n - 1 > m$. Thus, by Theorem 16, we have

$$\delta(C_m \times P_n) \leq \begin{cases} m/2, & \text{if } n - 1 \leq m, \\ (n - 1)/2, & \text{if } n - 1 > m. \end{cases}$$

Assume first that $n - 1 \leq m$. Note that $C_m \times P_2 \simeq C_{2m}$ and $C_m \times P_{n'}$ is an isometric subgraph of $C_m \times P_n$, if $n' \leq n$. By Lemma 3, we have $\delta(C_m \times P_n) \geq \delta(C_{2m}) = m/2$, and we obtain the result in this case.

Assume now that $n - 1 > m$. Consider the geodesic triangle T in $C_m \times P_n$ defined by the following geodesics

$$\begin{aligned} \gamma_1 &:= [(w_1, v_n), (w_2, v_{n-1})] \cup [(w_2, v_{n-1}), (w_3, v_n)] \cup [(w_3, v_n), (w_4, v_{n-1})] \cup \dots \cup [(w_{m-1}, v_{n-1}), (w_m, v_n)], \\ \gamma_2 &:= [(w_{(m+1)/2}, v_1), (w_{(m-1)/2}, v_2)] \cup [(w_{(m-1)/2}, v_2), (w_{(m-3)/2}, v_3)] \cup \dots \cup [(w_2, v_{(m-1)/2}), (w_1, v_{(m+1)/2})] \cup \\ &\quad [(w_1, v_{(m+1)/2}), (w_m, v_{(m+3)/2})] \cup [(w_m, v_{(m+3)/2}), (w_1, v_{(m+5)/2})] \cup [(w_1, v_{(m+5)/2}), (w_m, v_{(m+7)/2})] \cup \dots, \\ \gamma_3 &:= [(w_{(m+1)/2}, v_1), (w_{(m+3)/2}, v_2)] \cup [(w_{(m+3)/2}, v_2), (w_{(m+5)/2}, v_3)] \cup \dots \cup [(w_{m-1}, v_{(m-1)/2}), (w_m, v_{(m+1)/2})] \cup \\ &\quad [(w_m, v_{(m+1)/2}), (w_1, v_{(m+3)/2})] \cup [(w_1, v_{(m+3)/2}), (w_m, v_{(m+5)/2})] \cup [(w_m, v_{(m+5)/2}), (w_1, v_{(m+7)/2})] \cup \dots, \end{aligned}$$

where (w_1, v_n) (respectively, (w_m, v_n)) is an endpoint of either γ_2 or γ_3 , depending of the parity of n . Since T is a geodesic triangle in $C_m \times P_n$, we have $\delta(C_m \times P_n) \geq \delta(T)$. If $n - 1 < 2m$ and M is the midpoint of the geodesic γ_3 , then $\delta(C_m \times P_n) \geq \delta(T) = d_{C_m \times P_n}(M, \gamma_1 \cup \gamma_2) = L(\gamma_3)/2 = (n - 1)/2$. Therefore, the result for $m < n - 1 < 2m$ follows.

Finally, assume that $n - 1 \geq 2m$. Let us consider $N \in \gamma_3$ such that $d_{C_m \times P_n}(N, (w_{(m+1)/2}, v_1)) = m - 1/2$. Thus, $\delta(C_m \times P_n) \geq \delta(T) \geq d_{C_m \times P_n}(N, \gamma_1 \cup \gamma_2) = d_{C_m \times P_n}(N, (w_{(m+1)/2}, v_1)) = m - 1/2$. To finish the proof, it suffices to prove that $\delta(C_m \times P_n) \leq m - 1/2$. Seeking for a contradiction, assume that $\delta(C_m \times P_n) > m - 1/2$. By Theorems 6 and 7, there is a geodesic triangle $\Delta = \{x, y, z\} \in \mathbb{T}_1(C_m \times P_n)$ and $p \in [xy]$ with $d_{C_m \times P_n}(p, [yz] \cup [zx]) = \delta(C_m \times P_n) \geq m - 1/4$. Then, $L([xy]) = d_{C_m \times P_n}(x, p) + d_{C_m \times P_n}(p, y) \geq 2m - 1/2$. Let V_x (respectively, V_y) be the closest vertex to x (respectively, y) in $[xy]$, and consider a vertex V_p in $[xy]$ such that $d_{C_m \times P_n}(p, V(C_m \times P_n)) = d_{C_m \times P_n}(p, V_p)$. Note that $d_{C_m \times P_n}(p, [yz] \cup [zx]) \geq m - 1/4$ implies that $d_{C_m \times P_n}(p, V_p) \leq 1/2$. Since $x, y, z \in J(C_m \times P_n)$, we have $d_{C_m \times P_n}(V_x, V_y) \geq 2m - 1 > m$ and, consequently, $\pi_2([xy])$ is a geodesic in P_n . Since $\pi_2([yz] \cup [zx])$ is a path in P_n joining $\pi_2(x)$ and $\pi_2(y)$, there exists a vertex $(u, v) \in [xz] \cup [zy]$ such that $\pi_2(V_p) = v$ and $u \neq \pi_1(V_p)$. Therefore, $d_{C_m \times P_n}(V_p, (u, v)) \leq m - 1$ and, consequently, $d_{C_m \times P_n}(p, [xz] \cup [zy]) \leq d_{C_m \times P_n}(p, V_p) + d_{C_m \times P_n}(V_p, [xz] \cup [zy]) \leq 1/2 + m - 1$, leading to contradiction. \square

5. Conclusions

In this paper, we characterize in many cases the hyperbolic direct product of graphs. Here, the situation is more complex than with other graph products, partly because the direct product of two bipartite graphs is already disconnected and the formula for the distance in $G_1 \times G_2$ is more complicated than in the case of other products of graphs. Although in the study of hyperbolicity in a general context the hypothesis on the existence (or non-existence) of odd cycles is artificial, in the study of hyperbolic direct products, it is an essential hypothesis. We have proven that, if $G_1 \times G_2$ is hyperbolic, then one factor is hyperbolic and the other one is bounded. Besides, we prove that this necessary condition is also sufficient in many cases. If G_1 is a hyperbolic graph and G_2 is a bounded graph, then we prove that $G_1 \times G_2$ is hyperbolic when G_2 has some odd cycle or G_1 and G_2 do not have odd cycles. Otherwise, the characterization of hyperbolic direct products is a more difficult task. If G_1 has some odd cycle and G_2 does not have odd cycles, we provide sufficient conditions for non-hyperbolicity and hyperbolicity, respectively. Besides, we characterize the hyperbolicity of $G_1 \times G_2$ under some additional conditions.

A natural open problem is the complete characterization of hyperbolic direct products.

A second open problem is to compute the precise value of the hyperbolicity constant of the graphs appearing in Theorems 17 and 18 with unknown hyperbolicity constant.

Direct product of graphs is a subject closely related to lift of graphs, which have been intensively studied (see, e.g., [82] and the references therein). Another interesting problem is to study the hyperbolicity of lift of graphs. We think that it is possible to obtain some similar results in this context, although the odd cycles may not play an important role in the study of hyperbolic lifts of graphs.

Author Contributions: The authors contributed equally to this work.

Funding: This work was supported in part by four grants from Ministerio de Economía y Competitividad (MTM2012-30719, MTM2013-46374-P, MTM2016-78227-C2-1-P and MTM2015-69323-REDT), Spain.

Acknowledgments: We thank the referees for their suggestions and helpful remarks.

Conflicts of Interest: The authors declare no conflict of interest. The founding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

References

1. Hammack, R.; Imrich, W.; Klavžar, S. *Handbook of Product Graphs*, 2nd ed.; Discrete Mathematics and Its Applications Series; CRC Press: Boca Raton, FL, USA, 2011.
2. Weichsel, P.M. The Kronecker product of graphs. *Proc. Am. Math. Soc.* **1962**, *13*, 47–52. [[CrossRef](#)]
3. Bendall, S.; Hammack, R. Centers of n -fold tensor products of graphs. *Discuss. Math. Graph Theory* **2004**, *24*, 491–501. [[CrossRef](#)]
4. Brešar, B.; Imrich, W.; Klavžar, S.; Zmazek, B. Hypercubes as direct products. *SIAM J. Discret. Math.* **2005**, *18*, 778–786. [[CrossRef](#)]
5. Hammack, R. Minimum cycle bases of direct products of bipartite graphs. *Australas. J. Combin.* **2006**, *36*, 213–222.
6. Imrich, W.; Rall, D.F. Finite and infinite hypercubes as direct products. *Australas. J. Combin.* **2006**, *36*, 83–90.
7. Imrich, W.; Stadler, P. A prime factor theorem for a generalized direct product. *Discuss. Math. Graph Theory* **2006**, *26*, 135–140. [[CrossRef](#)]
8. Jha, P.K.; Klavžar, S. Independence in direct-product graphs. *Ars Combin.* **1998**, *50*, 53–63.
9. Balakrishnan, R.; Paulraja, P. Hamilton cycles in tensor product of graphs. *Discret. Math.* **1998**, *186*, 1–13. [[CrossRef](#)]
10. Kheddouci, H.; Kouider, M. Hamiltonian cycle decomposition of Kronecker product of some cubic graphs by cycles. *J. Combin. Math. Combin. Comput.* **2000**, *32*, 3–22.

11. Imrich, W.; Klavžar, S. *Product Graphs: Structure and Recognition*; John Wiley & Sons: New York, NY, USA, 2000.
12. Zhu, X. A survey on Hedetniemi's conjecture. *Taiwanese J. Math.* **1998**, *2*, 1–24. [[CrossRef](#)]
13. Imrich, W. Factoring cardinal product graphs in polynomial time. *Discr. Math.* **1998**, *192*, 119–144. [[CrossRef](#)]
14. Alonso, J.; Brady, T.; Cooper, D.; Delzant, T.; Ferlini, V.; Lustig, M.; Mihalik, M.; Shapiro, M.; Short, H. Notes on word hyperbolic groups. In *Group Theory from a Geometrical Viewpoint*; Ghys, E., Haefliger, A., Verjovsky, A., Eds.; World Scientific: Singapore, 1992.
15. Ghys, E.; de la Harpe, P. *Sur les Groupes Hyperboliques d'après Mikhael Gromov*; Progress in Mathematics 83; Birkhäuser Boston Inc.: Boston, MA, USA, 1990.
16. Gromov, M. Hyperbolic groups. In *Essays in Group Theory*; Gersten, S.M., Ed.; Mathematical Sciences Research Institute Publications; Springer: Berlin, Germany, 1987; Volume 8, pp. 75–263.
17. Oshika, K. *Discrete Groups*; AMS Bookstore: Providence, RI, USA, 2002.
18. Charney, R. Artin groups of finite type are biautomatic. *Math. Ann.* **1992**, *292*, 671–683. [[CrossRef](#)]
19. Chepoi, V.; Estellon, B. Packing and covering δ -hyperbolic spaces by balls. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques*; Springer: Berlin/Heidelberg, Germany, 2007; pp. 59–73.
20. Eppstein, D. Squarepants in a tree: Sum of subtree clustering and hyperbolic pants decomposition. In Proceedings of the eighteenth annual ACM-SIAM symposium on Discrete algorithms (SODA'2007), Barcelona, Spain, 16–19 January 2017; pp. 29–38.
21. Gavoille, C.; Ly, O. Distance labeling in hyperbolic graphs. In *Proceedings of the International Symposium on Algorithms and Computation (ISAAC)*; Springer: Berlin/Heidelberg, Germany, 2005; pp. 171–179.
22. Krauthgamer, R.; Lee, J.R. Algorithms on negatively curved spaces. In Proceedings of the 47th Annual IEEE Symposium on Foundations of Computer Science (FOCS'06), Berkeley, CA, USA, 21–24 October 2006; pp. 1–11.
23. Shavitt, Y.; Tankel, T. On Internet embedding in hyperbolic spaces for overlay construction and distance estimation. *IEEE/ACM Trans. Netw.* **2008**, *16*, 25–36. [[CrossRef](#)]
24. Verbeek, K.; Suri, S. Metric embeddings, hyperbolic space and social networks. In Proceedings of the 30th Annual Symposium on Computational Geometry, Kyoto, Japan, 8–11 June 2014; pp. 501–510.
25. Wu, Y.; Zhang, C. Chordality and hyperbolicity of a graph. *Electr. J. Comb.* **2011**, *18*, P43.
26. Abu-Ata, M.; Dragan, F.F. Metric tree-like structures in real-life networks: An empirical study. *Networks* **2016**, *67*, 49–68. [[CrossRef](#)]
27. Adcock, A.B.; Sullivan, B.D.; Mahoney, M.W. Tree-like structure in large social and information networks. In Proceedings of the 13th Int Conference Data Mining (ICDM), Dallas, TX, USA, 7–10 December 2013; pp. 1–10.
28. Krioukov, D.; Papadopoulos, F.; Kitsak, M.; Vahdat, A.; Boguná, M. Hyperbolic geometry of complex networks. *Phys. Rev. E* **2010**, *82*, 036106. [[CrossRef](#)] [[PubMed](#)]
29. Montgolfier, F.; Soto, M.; Viennot, L. Treewidth and Hyperbolicity of the Internet. In Proceedings of the 10th IEEE International Symposium on Network Computing and Applications (NCA), Cambridge, MA, USA, 25–27 August 2011; pp. 25–32.
30. Chepoi, V.; Dragan, F.F.; Vaxès, Y. Core congestion is inherent in hyperbolic networks. In Proceedings of the Twenty-Eighth Annual ACM-SIAM Symposium on Discrete Algorithms, Barcelona, Spain, 16–19 January 2017; pp. 2264–2279.
31. Grippo, E.; Jonckheere, E.A. Effective resistance criterion for negative curvature: application to congestion control. In Proceedings of the 2016 IEEE Conference on Control Applications (CCA), Buenos Aires, Argentina, 19–22 September 2016.
32. Jonckheere, E.A. Contrôle du trafic sur les réseaux à géométrie hyperbolique—Vers une théorie géométrique de la sécurité l'acheminement de l'information. *J. Eur. Syst. Autom.* **2002**, *8*, 45–60.
33. Jonckheere, E.A.; Lohsoonthorn, P. Geometry of network security. In Proceedings of the 2004 American Control Conference, Boston, MA, USA, 30 June–2 July 2004; pp. 111–151.
34. Li, S.; Tucci, G.H. Traffic Congestion in Expanders, (p, δ) -Hyperbolic Spaces and Product of Trees. *Int. Math.* **2015**, *11*, 134–142.
35. Shang, Y. Lack of Gromov-hyperbolicity in colored random networks. *Pan-Am. Math. J.* **2011**, *21*, 27–36.

36. Shang, Y. Lack of Gromov-hyperbolicity in small-world networks. *Cent. Eur. J. Math.* **2012**, *10*, 1152–1158. [[CrossRef](#)]
37. Shang, Y. Non-hyperbolicity of random graphs with given expected degrees. *Stoch. Models* **2013**, *29*, 451–462. [[CrossRef](#)]
38. Gosak, M.; Markovič, R.; Dolensšek, J.; Slak Rupnik, M.; Marhl, M.; Stožer, A.; Perc, M. Network science of biological systems at different scales: A review. *Phys. Life Rev.* **2018**, *24*, 118–135. [[CrossRef](#)] [[PubMed](#)]
39. Tourís, E. Graphs and Gromov hyperbolicity of non-constant negatively curved surfaces. *J. Math. Anal. Appl.* **2011**, *380*, 865–881. [[CrossRef](#)]
40. Bermudo, S.; Rodríguez, J.M.; Sigarreta, J.M. Computing the hyperbolicity constant. *Comput. Math. Appl.* **2011**, *62*, 4592–4595. [[CrossRef](#)]
41. Bermudo, S.; Rodríguez, J.M.; Sigarreta, J.M.; Vilaire, J.-M. Gromov hyperbolic graphs. *Discr. Math.* **2013**, *313*, 1575–1585. [[CrossRef](#)]
42. Brinkmann, G.; Koolen J.; Moulton, V. On the hyperbolicity of chordal graphs. *Ann. Comb.* **2001**, *5*, 61–69. [[CrossRef](#)]
43. Chepoi, V.; Dragan, F.F.; Estellon, B.; Habib, M.; Vaxes, Y. Notes on diameters, centers, and approximating trees of δ -hyperbolic geodesic spaces and graphs. *Electr. Notes Discr. Math.* **2008**, *31*, 231–234. [[CrossRef](#)]
44. Dragan, F.; Mohammed, A. Slimness of graphs. *arXiv* **2017**, arXiv:1705.09797.
45. Martínez-Pérez, A. Chordality properties and hyperbolicity on graphs. *Electr. J. Comb.* **2016**, *23*, P3.51.
46. Michel, J.; Rodríguez, J.M.; Sigarreta, J.M.; Villeta, M. Gromov hyperbolicity in Cartesian product graphs. *Proc. Indian Acad. Sci. Math. Sci.* **2010**, *120*, 1–17. [[CrossRef](#)]
47. Rodríguez, J.M.; Sigarreta, J.M.; Vilaire, J.-M.; Villeta, M. On the hyperbolicity constant in graphs. *Discr. Math.* **2011**, *311*, 211–219. [[CrossRef](#)]
48. Chen, B.; Yau, S.-T.; Yeh, Y.-N. Graph homotopy and Graham homotopy. *Discret. Math.* **2001**, *241*, 153–170. [[CrossRef](#)]
49. Shang, Y. On the likelihood of forests. *Phys. A Stat. Mech. Appl.* **2016**, *456*, 157–166. [[CrossRef](#)]
50. Fournier, H.; Ismail, A.; Vigneron, A. Computing the Gromov hyperbolicity of a discrete metric space. *Inf. Process. Lett.* **2015**, *115*, 576–579. [[CrossRef](#)]
51. Papasoglu, P. An algorithm detecting hyperbolicity. In *Geometric and Computational Perspectives on Infinite Groups, DIMACS; Series in Discrete Mathematics and Theoretical Computer Science*; AMS: Hong Kong, China, 1996; Volume 25, pp.193–200.
52. Chalopin, J.; Chepoi, V.; Papasoglu, P.; Pecatte, T. Cop and robber game and hyperbolicity. *SIAM J. Discr. Math.* **2015**, *28*, 1987–2007. [[CrossRef](#)]
53. Sigarreta, J.M. Hyperbolicity in median graphs. *Proc. Indian Acad. Sci. Math. Sci.* **2013**, *123*, 455–467. [[CrossRef](#)]
54. Carballosa, W.; Rodríguez, J.M.; Sigarreta, J.M. New inequalities on the hyperbolicity constant of line graphs. *Ars Combin.* **2016**, *129*, 367–386.
55. Carballosa, W.; Rodríguez, J.M.; Sigarreta, J.M.; Villeta, M. Gromov hyperbolicity of line graphs. *Electr. J. Comb.* **2011**, *18*, P210.
56. Cohen, N.; Coudert, D.; Ducoffe, G.; Lancin, A. Applying clique-decomposition for computing Gromov hyperbolicity. *Theor. Comp. Sci.* **2017**./j.tcs.2017.06.001. [[CrossRef](#)]
57. Pestana, D.; Rodríguez, J.M.; Sigarreta, J.M.; Villeta, M. Gromov hyperbolic cubic graphs. *Central Eur. J. Math.* **2012**, *10*, 1141–1151. [[CrossRef](#)]
58. Bermudo, S.; Rodríguez, J.M.; Sigarreta, J.M.; Tourís, E. Hyperbolicity and complement of graphs. *Appl. Math. Lett.* **2011**, *24*, 1882–1887. [[CrossRef](#)]
59. Hernández, J.C.; Rodríguez, J.M.; Sigarreta, J.M.; Torres-Nuñez, Y.; Villeta, M. Gromov hyperbolicity of regular graphs. *Ars Combin.* **2017**, *130*, 395–416.
60. Bermudo, S.; Carballosa, W.; Rodríguez, J.M.; Sigarreta, J.M. On the hyperbolicity of edge-chordal and path-chordal graphs. *Filomat* **2016**, *30*, 2599–2607. [[CrossRef](#)]
61. Carballosa, W.; Portilla, A.; Rodríguez, J.M.; Sigarreta, J.M. Planarity and hyperbolicity in graphs. *Graphs Combin.* **2015**, *31*, 1311–1324. [[CrossRef](#)]
62. Portilla, A.; Rodríguez, J.M.; Sigarreta, J.M.; Vilaire, J.-M. Gromov hyperbolic tessellation graphs. *Utilitas Math.* **2015**, *97*, 193–212.

63. Coudert, D.; Ducoffe, G. On the hyperbolicity of bipartite graphs and intersection graphs. *Discr. Appl. Math.* **2016**, *214*, 187–195. [[CrossRef](#)]
64. Calegari, D.; Fujiwara, K. Counting subgraphs in hyperbolic graphs with symmetry. *J. Math. Soc. Jpn.* **2015**, *67*, 1213–1226. [[CrossRef](#)]
65. Cantón, A.; Granados, A.; Pestana, D.; Rodríguez, J.M. Gromov hyperbolicity of periodic graphs. *Bull. Malays. Math. Sci. Soc.* **2016**, *39*, S89–S116. [[CrossRef](#)]
66. Cantón, A.; Granados, A.; Pestana, D.; Rodríguez, J.M. Gromov hyperbolicity of periodic planar graphs. *Acta Math. Sin.* **2014**, *30*, 79–90. [[CrossRef](#)]
67. Koolen, J.H.; Moulton, V. Hyperbolic Bridged Graphs. *Eur. J. Comb.* **2002**, *23*, 683–699. [[CrossRef](#)]
68. Rodríguez, J.M. Characterization of Gromov hyperbolic short graphs. *Acta Math. Sin.* **2014**, *30*, 197–212. [[CrossRef](#)]
69. Carballosa, W.; Rodríguez, J.M.; Rosario, O.; Sigarreta, J.M. Gromov hyperbolicity of graph minors. *Bull. Iran. Math. Soc.* **2018**, *44*, 481–503. [[CrossRef](#)]
70. Bermudo, S.; Rodríguez, J.M.; Rosario, O.; Sigarreta, J.M. Small values of the hyperbolicity constant in graphs. *Discr. Math.* **2016**, *339*, 3073–3084. [[CrossRef](#)]
71. Granados, A.; Pestana, D.; Portilla, A.; Rodríguez, J.M. Gromov hyperbolicity in Mycielskian Graphs. *Symmetry* **2017**, *9*, 131. [[CrossRef](#)]
72. Hernández, J.C.; Reyes, R.; Rodríguez, J.M.; Sigarreta, J.M. Mathematical properties on the hyperbolicity of interval graphs. *Symmetry* **2017**, *9*, 255. [[CrossRef](#)]
73. Carballosa, W.; de la Cruz, A.; Rodríguez, J.M. Gromov hyperbolicity in the Cartesian sum of graphs. *Bull. Iran. Math. Soc.* **2017**, 1–20. [[CrossRef](#)]
74. Carballosa, W.; Casablanca, R.M.; de la Cruz, A.; Rodríguez, J.M. Gromov hyperbolicity in strong product graphs. *Electr. J. Comb.* **2013**, *20*, 2.
75. Carballosa, W.; de la Cruz, A.; Rodríguez, J.M. Gromov hyperbolicity in lexicographic product graphs. *arXiv* **2015**, arXiv 1506.06034
76. Carballosa, W.; Rodríguez, J.M.; Sigarreta, J.M. Hyperbolicity in the corona and join of graphs. *Aequ. Math.* **2015**, *89*, 1311–1327. [[CrossRef](#)]
77. Bowditch, B.H. Notes on Gromov’s hyperbolicity criterion for path-metric spaces. In *Group Theory from a Geometrical Viewpoint*; Ghys, E., Haefliger, A., Verjovsky, A., Eds.; World Scientific: River Edge, NJ, USA, 1991; pp. 64–167.
78. Shchur, V. A quantitative version of the Morse lemma and quasi-isometries fixing the ideal boundary. *J. Funct. Anal.* **2013**, *264*, 815–836. [[CrossRef](#)]
79. Rodríguez, J.M.; Tourís, E. Gromov hyperbolicity through decomposition of metric spaces. *Acta Math. Hung.* **2004**, *103*, 53–84. [[CrossRef](#)]
80. Alvarez, V.; Portilla, A.; Rodríguez, J.M.; Tourís, E. Gromov hyperbolicity of Denjoy domains. *Geom. Dedic.* **2006**, *121*, 221–245. [[CrossRef](#)]
81. Bonk, M. Quasi-geodesics segments and Gromov hyperbolic spaces. *Geom. Dedicata* **1996**, *62*, 281–298. [[CrossRef](#)]
82. Shang, Y. Random Lifts Of Graphs: Network Robustness Based On The Estrada Index. *Appl. Math. E-Notes* **2012**, *12*, 53–61.

