Derivation of a homogenized von-Kármán shell theory from 3D elasticity

Peter Hornung

Max Planck Institute for Mathematics in the Sciences Inselstr. 22 D-04103 Leipzig Germany email: hornung@mis.mpg.de

Igor Velčić

Max Planck Institute for Mathematics in the Sciences Inselstr. 22 D-04103 Leipzig Germany BCAM-Basque Center for Applied Mathematics Alameda de Mazarredo 14 48009 Bilbao Bizkaia (Basque-Country, Spain) email: velcic@mis.mpg.de

March 15, 2013

Abstract

We derive the model of homogenized von Kármán shell theory, starting from three dimensional nonlinear elasticity. The original three dimensional model contains two small parameters: the oscillations of the material ε and the thickness of the shell h. Depending on the asymptotic ratio of these two parameters, we obtain different asymptotic theories. In the case $h \ll \varepsilon$ we identify two different asymptotic theories, depending on the ratio of h and ε^2 . In the case of convex shells we obtain a complete picture in the whole regime $h \ll \varepsilon$.

Keywords: elasticity, dimension reduction, homogenization, shell theory, two-scale convergence.

Contents

1	Intr	roduction	2
2	Geo	ometric preliminaries and general framework	4
	2.1	Displacements and infinitesimal bendings	7

3	Elas	ticity framework and main result	10
	3.1	Unit thickness rescaling	16
	3.2	FJM-compactness	17
	3.3	Two-scale convergence	20
4	Two	-scale compactness and lower bound	21
5	Upp	er bound	29
6	Con	vex shell	38
A	Aux	iliary results	42
$\lfloor 1 \rfloor$			

1 Introduction

This paper is about von-Kármán theory for thin elastic shells. There is a vast literature on shell theory in elasticity. An overview about the derivation of models for linear and nonlinear shells by the method of formal asymptotic expansions can be found in [Cia00]. In the case of linearly elastic shells, the models thus obtained can also be justified by a rigorous convergence result, starting from three dimensional linearized elasticity (see also [CL96, CLM96]).

In the last two decades, rigorous justifications of nonlinear models for rods, curved rods, plates and shells were obtained by means of Γ -convergence, starting from three dimensional nonlinear elasticity. The first papers in that direction are [ABP91, LDR95, LDR96] for the string model, membrane plate and shell model respectively. The rigorous derivation of nonlinear bending theory of plate was achieved in [FJM02]; see also [Pan01] for an earlier result in this direction. Föppl-von Kármán theory for plates was derived in [FJM06]. In [MM03, MM04], bending and von Kármán theories for rods were derived. In [FJM03] the nonlinear bending theory shell model was derived, and in [LMP10] the von Kármán shell model was derived.

Here we are interested in an the ansatz-free derivation of a homogenized von Kármán shell theory by simultaneous homogenization and dimension reduction. Our starting point is the energy functional from 3d nonlinear elasticity. It attributes to a deformation u of a given shell $S^h \subset \mathbb{R}^3$ of small thickness h > 0 around a surface $S \subset \mathbb{R}^3$ the stored elastic energy

(1)
$$\frac{1}{h^4 |S^h|} \int_{S^h} W_{\varepsilon}(x, \nabla u(x)) \, dx, \qquad u \in H^1(S^h, \mathbb{R}^3).$$

Here W_{ε} is a non-degenerate stored energy function that oscillates periodically on the surface, with some period $\varepsilon \ll 1$. We are interested in the effective behavior when both the thickness h and the period ε are small. The separate limits $h \to 0$ and $\varepsilon \to 0$ are reasonably well understood: In [LMP10] it is shown that, when W_{ε} does not depend on ε , then the functionals (1) Γ -converge as $h \to 0$ to a two-dimensional von Kármán shell

theory. Regarding the limit $\varepsilon \to 0$, which is related to homogenization, the first rigorous results relevant in nonlinear elasticity were obtained by Braides [Bra85] and independently by Müller [Mül87]. They proved that, under suitable growth assumptions on W_{ε} , the energy (1) Γ -converges as $\varepsilon \to 0$ (and h fixed) to the functional obtained by replacing W_{ε} in (1) with the homogenized energy density given by an infinite-cell homogenization formula.

In this paper we study the asymptotic behavior when both the thickness h and the period ε tend to zero simultaneously. As a Γ -limit we obtain a two-dimensional von Kármán shell model with homogenized material properties. Recently, the von Kárman plate model (see [NV]), the bending plate models (see [HNV, Vela]), and bending rod models (see [Neu10, Neu12]), were analyzed in this way. Simultaneous homogenization and dimensional reduction was also done in the case of periodically wrinkled plate (see [Velb]). As explained there, in these cases one does not obtain infinite-cell homogenization formula like in the membrane case (see [BFF00, BB06]). The basic reason for that is the fact since we are in small strain regimes, the energy is essentially convex in the strain. This is the main reason why we can use two scale convergence techniques in all these cases. However, every case has its own peculiarities. In the von Kármán theory of plates, one obtains a limiting quadratic energy density which is continuous in the asymptotic ratio γ between h and ε , for all $\gamma \in [0,\infty]$. Moreover, the case $\gamma = 0$ corresponds to the situation when the dimensional reduction dominates and the obtained model is just the homogenized von Kármán plate model. The situation $\gamma = \infty$ corresponds to the case when homogenization dominates and the obtained model is the von Kármán plate model of the homogenized functional. The case of bending plate is more involving; we are able to obtain the models in the case $\gamma \in (0, \infty]$ (see [HNV]) and in the case $\gamma = 0$ under the additional assumption that $\varepsilon^2 \ll h \ll \varepsilon$ (see [Vela]). This model does not correspond to the situation of the homogenized bending plate model, but is the limiting situation of the models when $\gamma \to 0$ and $\gamma > 0$.

In case of von Kármán shell theory studied in the present paper, we encounter two different scenarios in the regime $h \ll \varepsilon$, depending whether $h \sim \varepsilon^2$ or $h \ll \varepsilon^2$. Our main result is presented in Theorem 3.6. We are not able to cover the case $h \ll \varepsilon^2$ in a generic way for arbitrary reference surfaces S. A stronger influence of the geometry of the reference surface S is expected in this case. In fact, in the case when S is a convex surface, we successfully derive the limiting model even for the regime $h \ll \varepsilon^2$, see Theorem 6.3.

Our analysis requires both techniques from dimension reduction, in particular, the quantitative rigidity estimate and approximation schemes developed in [FJM02, FJM06]; and techniques from homogenization methods, in particular, two-scale convergence [Ngu89, All92, Vis06, Vis07]. To our knowledge our result is the first rigorous result combining homogenization and dimension reduction for shells in the von Kármán regime. The homogenization for linearly elastic shells was carried out in [Lut85].

This paper is organised as follows: after introducing the setting and basic objects in Section 2 and 3 we state the main result in Section 3. In Section 4 we identify the two scale limit of the strain and prove lower bound for Γ -limit. In Section 5 we construct the recovery sequences and thus prove the upper bound. All these results are given for general surfaces and the cases $h \gg \varepsilon^2$ or $h \sim \varepsilon^2$. In the last section we analyze the case of convex shells for the situation when $h \ll \varepsilon^2$.

Notation

The notation $A \leq B$ means that $A \leq CB$ with C depending only on quantities regarded as constant in the context in question.

In this paper we frequently encounter function spaces of periodic functions. We denote by \mathcal{Y} the real line \mathbb{R} equipped with the torus topology, that is y+1 and y are identified in \mathcal{Y} . We write $C(\mathcal{Y})$ to denote the space of continuous functions $f : \mathbb{R} \to \mathbb{R}$ satisfying f(y+1) = f(y) for all $y \in \mathbb{R}$. Clearly, $C(\mathcal{Y})$ endowed with the norm $||f||_{\infty} := \sup_{y \in \mathcal{Y}} |f(y)|$ is a Banach space. Moreover, we set $C^k(\mathcal{Y}) := C^k(\mathbb{R}) \cap C(\mathcal{Y})$ and denote by $L^2(\mathcal{Y})$, $H^1(\mathcal{Y})$ and $H^1(S \times \mathcal{Y})$ the closure of $C^{\infty}(\mathcal{Y})$ and $C^{\infty}(\bar{S}; C^{\infty}(\mathcal{Y}))$ w. r. t. the norm in $L^2(\mathcal{Y})$, $H^1(\mathcal{Y})$ and $H^1(S \times \mathcal{Y})$, respectively. By $\dot{L}^2(\mathcal{Y})$, $\dot{H}^k(\mathcal{Y})$ we denote the subspace of functions $H^k(\mathcal{Y})$ whose mid-value over \mathcal{Y} is zero. Obviously, all these spaces are Banach spaces. For $A \subset \mathbb{R}^d$ measurable and X a Banach space, $L^2(A; X)$ is understood in the sense of Bochner. We tacitly identify the spaces $L^2(A; L^2(B))$ and $L^2(A \times B)$; since whenever $f \in L^2(A \times B)$, then there exists a function $\tilde{f} \in L^2(A; L^2(B))$ with $f = \tilde{f}$ almost everywhere in $A \times B$. By (e_1, e_2, e_3) we denote the standard basis on \mathbb{R}^3 .

2 Geometric preliminaries and general framework

In this subsection we do not always display the explicit regularity assumptions; the minimal requirements are obvious. We assume that $\omega \subset \mathbb{R}^2$ is a bounded domain with boundary of class C^3 . We set $I := (-\frac{1}{2}, \frac{1}{2})$ and $\Omega^h := \omega \times (hI)$, and $\Omega := \omega \times I$. The variables on ω (resp. Ω) will be denoted by ξ_1, ξ_2 (resp. ξ_1, ξ_2, t). For a function $f : \Omega \to \mathbb{R}^3$ we define $\nabla_h f := (\partial_1 f, \partial_2 f, \frac{1}{h} \partial_3 f)$.

Let S be a compact connected oriented surface with boundary which is embedded in \mathbb{R}^3 . For convenience we assume that S is parametrized by a single chart: From now on, $\psi \in C^3(\overline{\omega}; \mathbb{R}^3)$ denotes an embedding with $\psi(\omega) = S$. The inverse of ψ is denoted by $r: S \to \omega$, and we assume it to be of class C^3 . We leave it to the interested reader to verify to which extent these regularity assumptions on S can be weakened without altering our arguments.

The nearest point retraction of a tubular neighbourhood of S onto S will be denoted by π . Hence

$$\pi(x + tn(x)) = x$$
 whenever $|t|$ is small enough.

We introduce the basis vectors of the tangent bundle determined by ψ , namely the pushforwards $\tau_i = \psi_* e_i$. Explicitly, this means

$$\tau_i(x) = (\partial_i \psi)(r(x))$$
 for $i = 1, 2$ and all $x \in S$.

By our hypotheses on ψ there exist $\eta_1, \eta_2 > 0$ such that

(2)
$$\eta_1 \leq \det([\tau_1 \ \tau_2]^T[\tau_1 \ \tau_2]) \leq \eta_2, \ \|\tau_1\|_{W^{2,\infty}(S)} \leq \eta_2, \|\tau_2\|_{W^{2,\infty}(S)} \leq \eta_2.$$

We denote by $(\tau^1(x), \tau^2(x))$ the dual base to the base $(\tau_1(x), \tau_2(x))$, that is,

$$\tau^{i}(x) = (\partial^{i}\psi)(r(x)).$$

By $n: S \to \mathbb{S}^2$ we denote the unit normal, that is,

$$n(x) = \frac{\tau_1(x) \wedge \tau_2(x)}{|\tau_1(x) \wedge \tau_2(x)|} \text{ for all } x \in S.$$

By $T_x S$ = span { $\tau_1(x), \tau_2(x)$ } we denote the tangent space to S at x For each $x \in S$, the vectors $\tau_1(x), \tau_2(x)$ and n(x) form a basis of \mathbb{R}^3 . Its dual basis is ($\tau^1(x), \tau^2(x), n(x)$). We define $\tau_3(x) = \tau^3(x) = n(x)$.

For a subset $A \subset S$ we set

$$A^h = \{ x + tn(x); \ x \in S, -h/2 < t < h/2 \}.$$

In particular, the shell is given by

$$S^h = \{x + tn(x); x \in S, -h/2 < t < h/2\}.$$

We introduce the function $t: S^1 \to \mathbb{R}$ by

(3)
$$t(x) = (x - \pi(x)) \cdot n(x) \text{ for all } x \in S^1.$$

By $r_e: S^1 \to \Omega$ we denote the map (see below why we assume that π and t is well-defined on S^1)

$$r_e(x) = r(\pi(x)) + t(x)e_3.$$

Clearly,

(4)
$$r_e^{-1}(\xi_1,\xi_2,t) = \psi(\xi_1,\xi_2) + tn(\psi(\xi_1,\xi_2)),$$

and thus:

(5)
$$\nabla r_e^{-1}(\xi_1, \xi_2, t) = (I + t\mathbf{S}(x))[\tau_1(x), \tau_2(x), \tau_3(x)], \text{ where } x = \psi(\xi_1, \xi_2)$$

(6) $\nabla r_e(x) = [\tau^1(x), \tau^2(x), \tau^3(x)]^T (I + t(x)\mathbf{S}(x))^{-1}.$

We denote by

$$T_S(x) := I - n(x) \otimes n(x)$$

the orthogonal projection from \mathbb{R}^3 onto T_xS . We will frequently deal with vector fields $V: S \to \mathbb{R}^3$ on the surface. We extend all such vector fields trivially from S to S^1 , simply by defining $V(x) = V(\pi(x))$ for all $x \in S^1$. By V_{tan} we denote the projection of vector field V on the tangential space i.e. $V_{\text{tan}} = T_S V$. We will denote by \tilde{V} the corresponding vector field along ω , i.e. we set $\tilde{V}(x) = V(\psi(\xi))$ for all $\xi \in \omega$.

The space of quadratic forms on S is denoted by S. It consists of all maps B on S such that, for each $x \in S$, the map

$$B(x): T_x S \times T_x S \to \mathbb{R}$$

is symmetric and bilinear. We will frequently regard B as a map from S into $\mathbb{R}^{3\times 3}$ via the embedding ι defined by

$$\iota(B) = B(T_S, T_S).$$

On the right-hand side and elsewhere we identify bilinear maps from \mathbb{R}^3 into itself with $\mathbb{R}^{3\times 3}$. By definition, $B(T_S, T_S): S \to \mathbb{R}^{3\times 3}$ takes the vector fields $v, w: S \to \mathbb{R}^3$ into the function

$$B(T_S v, T_S w).$$

By definition, $B \in L^2(S; \mathbb{S})$ means that (using the above embedding) $B \in L^2(S; \mathbb{R}^{3 \times 3})$ and $B \in \mathbb{S}$. The spaces $H^1(S; \mathbb{S})$ etc. are defined similarly. By $\mathbb{S}(x)$ we denote the set of all quadratic forms on $T_x S$ which can be embedded in the space $\mathbb{R}^{3\times 3}$. By $\mathbb{S}(x)_{sym}$ we denote the set of symmetric quadratic forms on $T_x S$ which can be embedded in $\mathbb{R}^{3\times 3}_{sym}$.

For a function $f: S \to \mathbb{R}^3$ we regard its tangential derivative $\nabla_{tan} f(x)$ as a linear map from $T_x S$ into \mathbb{R} . For a tangent vector field τ along S we write $\partial_{\tau} f = \nabla_{tan} f \tau$. A similar notation applies to vector fields instead of functions. By $\nabla_{tan} \nabla_{tan} f$ we denote the triilinear form $\nabla_{\tan} \nabla_{\tan} f(\eta_1, \eta_2, \eta_3) = \partial_{\eta_2} \partial_{\eta_3} f \cdot \eta_1$. For scalar $f, \nabla_{\tan} \nabla_{\tan} f$ is just the bilinear form $\nabla_{\tan} \nabla_{\tan} f(\eta_1, \eta_2) = \partial_{\eta_1} \partial_{\eta_2} f$

The Weingarten map **S** on the surface S is given by $\mathbf{S} = \nabla_{tan} n$, i.e.,

$$\mathbf{S}(x)\tau = (\partial_{\tau}n)(x)$$
 for all $x \in S, \ \tau \in T_xM$

We extend **S** to a linear map on \mathbb{R}^3 by setting $\mathbf{S} = \mathbf{S} T_S$, i.e., we define $\mathbf{S}(x)n(x) = 0$. Moreover, we extend **S** trivially from S to S^1 , i.e., we have $\mathbf{S}(x) = \mathbf{S}(\pi(x))$. With a slight abuse of notation, we denote by \mathbf{S} also the (negated) second fundamental form of S defined by

$$\mathbf{S}_{ij}(x) := \mathbf{S}(x)\tau_i(x) \cdot \tau_j(x).$$

In general, for a given bilinear form B on S we denote its local coordinates by

$$B_{ij} := B\tau_j \cdot \tau_i.$$

Obviously $B = \sum_{i,j=1}^{2} B_{ij} \tau^i \otimes \tau^j$. After rescaling the ambient space, we may assume that the curvature of S is as small as we please. In particular, we may assume without loss of generality that π is well-defined on a domain containing the closure of $\{x + tn(x); x \in S, -1 < t < 1\}$, and that

$$1/2 < |Id + t\mathbf{S}(x)| < 3/2$$

for all $t \in (-1, 1)$ and all $x \in S$.

Lemma 2.1. For all $x \in S^1$ we have

$$(\nabla \pi)(x) = T_S(\pi(x)) (I + t(x)\mathbf{S}(\pi(x)))^{-1}$$

Proof. Let $x \in S$, let $\tau \in T_x S$ and let $\gamma \in C^1((-1,1), M)$ with $\gamma(0) = x$ and $\dot{\gamma}(0) = \tau$. Then

$$\pi(\gamma + sn(\gamma)) = \gamma \text{ on } (-1,1)$$

Taking the derivative with respect to the arclength of γ , this implies

$$(\nabla \pi)(\gamma + tn(\gamma))(\tau + t\mathbf{S}(\gamma)\tau) = \tau.$$

As $x \in S$ and $\tau \in T_x S$ were arbitrary, we conclude that

(7)
$$(\nabla \pi)(x + tn(x))(I + t\mathbf{S}(x)) = T_S(x)$$

on T_xS . But by definition $\mathbf{S}(x)n(x) = 0$, and clearly $(\nabla \pi)(x + tn(x))n(x) = 0$, too. Hence both sides of (7) agree on all of \mathbb{R}^3 . We will frequently extend functions $f: S \to \mathbb{R}$ defined on S only to functions defined on S^1 in the following way:

$$f(x) = f(\pi(x))$$
 for all $x \in S^1$,

with a slight abuse of notation on the left-hand side. When referring to this extension, we will say that we extend f trivially to S^1 . By Lemma 2.1 we have for all $x \in S^1$ the following formula for the full derivative of f in terms of its tangential derivative:

$$(\nabla f)(x) = (\nabla_{tan} f)(\pi(x))T_S(\pi(x))(I + t(x)\mathbf{S}(\pi(x)))^{-1}$$

Extending $\nabla_{tan} f$, T_S and **S** trivially from S to S^1 , as we will do from now on, this formula reads

(8)
$$(\nabla f)(x) = (\nabla_{tan} f)(x) T_S(x) (I + t(x) \mathbf{S}(x))^{-1}.$$

From now on we tacitly also extend r trivially to S^1 .

2.1 Displacements and infinitesimal bendings

For a given displacement $V:S\to \mathbb{R}^3$ we introduce the quadratic form $(dV)^2$ on S which is defined by

$$(dV)^2(x)(\tau,\eta) = \partial_\tau V(x) \cdot \partial_\eta V(x)$$
 for all $\tau, \eta \in T_x S$.

We also introduce the quadratic form q_V on S which is defined by its action on tangent vectors $\tau, \eta \in T_x S$ as follows:

$$q_V(x)(\tau,\eta) = \frac{1}{2} \left(\eta \cdot \partial_\tau V(x) + \tau \cdot \partial_\eta V(x) \right)$$

In the geometry literature, this form is usually denoted by $d\psi \cdot dV$. In local coordinates, it is given by the matrix field

$$\operatorname{sym}\left((\nabla\psi^T)\nabla\tilde{V}\right)$$

on ω . Obviously,

(9)
$$\operatorname{sym}\left((\nabla\psi)^T\nabla\tilde{V}_{tan}\right) = \operatorname{sym}\nabla\bar{V} - \Gamma\cdot\bar{V},$$

where $\bar{V}_{\alpha} = \tilde{V} \cdot \partial_{\alpha} \psi$ for $\alpha = 1, 2$ and where for brevity we have set $(\Gamma \cdot \bar{V})_{ij} := \sum_{k=1,2} \Gamma_{ij}^k \bar{V}_k$. Here Γ_{ij}^k denote the Christoffel symbols of the metric induced by ψ . For our purposes it will be enough to know that $\Gamma \in L^{\infty}(\omega; \mathbb{R}^{2 \times 2 \times 2})$. Using (9) we see that

(10)
$$\operatorname{sym}\left((\nabla\psi)^T\nabla\tilde{V}\right) = \operatorname{sym}\nabla\bar{V} - \Gamma\cdot\bar{V} + (\tilde{V}\cdot n)\mathbf{S},$$

were **S** denotes the pulled back (negated) second fundamental form. Equivalently, we have the following equality between quadratic forms on S:

(11)
$$q_V = q_{V_{\text{tan}}} + (V \cdot n)\mathbf{S}.$$

It is well-known that the quadratic form q_V typically arises in the context of thin elastic shells, because it is just the first variation of the metric of S under the displacement V. For example, in [GSP95] it is denoted (in coordinates) by $\gamma_{\alpha\beta}$ and in [LMP10] it is denoted by sym ∇V .

A displacement $V: S \to \mathbb{R}^3$ is called an *infinitesimal bending* of S provided that $q_V = 0$, i.e., that

$$\operatorname{sym}\left((\nabla\psi)^T\nabla\tilde{V}\right)_{ij} = \partial_j\psi\cdot\partial_i\tilde{V} + \partial_i\psi\cdot\partial_j\tilde{V} = 0 \text{ for all } i, j = 1, 2.$$

Infinitesimal bendings have been studied extensively both in the applied literature (see e.g. [Cia00], [Cho97], [GSP95]) and in the geometry literature (see e.g. the references in [Hor12]). Recently, they have been found to be relevant as well to fully nonlinear bending theories, cf. [Hor].

For any displacement V,

we define $\mu_V: S \to \mathbb{R}^3$ by setting

(12)
$$\mu_V = T_S \frac{\partial_{\tau_1} V \wedge \tau_2 + \tau_1 \wedge \partial_{\tau_2} V}{|\tau_1 \wedge \tau_2|}$$

Note that

(13)
$$\mu(x) \cdot \tau = -n(x) \cdot \partial_{\tau} V(x) \text{ for all } \tau \in T_x S.$$

In fact, we compute

$$\mu \cdot \tau = \frac{1}{|\tau_1 \wedge \tau_2|} \left(\partial_{\tau_1} V \cdot \tau_2 \wedge \tau - \partial_{\tau_2} V \cdot \tau_1 \wedge \tau \right).$$

Hence $\mu \cdot \tau_i = -n \cdot \partial_{\tau_i} V$ for i = 1, 2. This proves (13). For a given displacement $V : S \to \mathbb{R}^3$ we define $\Omega_V : S \to \mathbb{R}^{3 \times 3}$ by

(14)
$$\Omega_V = \nabla_{tan} V \ T_S + \mu_V \otimes n.$$

Lemma 2.2. If $V \in H^1(S; \mathbb{R}^3)$ then sym $\Omega_V = q_V(T_S, T_S)$ almost everywhere on S.

Proof. Clearly $n \cdot \Omega_V n = n \cdot \mu_V = 0$ and for any tangent vector field τ along S we have

$$\tau \cdot \Omega_V n + n \cdot \Omega_V \tau = \tau \cdot \mu_V + n \cdot \partial_\tau V = 0$$

by (13). For any tangent vector field σ we have

$$\tau \cdot \Omega_V \sigma + \sigma \cdot \Omega_V \tau = 2q_V(\sigma, \tau).$$

If V is an infinitesimal bending, then

(15)
$$\Omega_V^2(T_S, T_S) = -(dV)^2(T_S, T_S),$$

that is, $\Omega_V^2(\tau, \sigma) = -\partial_\tau V \cdot \partial_\sigma V$ for all tangent vector fields τ , σ along S. In fact, by skew symmetry, $\Omega_V^2(\tau, \sigma) = -\Omega_V \tau \cdot \Omega_V \sigma$, and $\partial_\tau V = \Omega_V \tau$.

For any displacement V, the *linearised Weingarten map* $b_V(x)$ is the linear map on T_xS given by

(16)
$$b_V = \nabla_{tan} V \mathbf{S} - \nabla_{tan} \mu_V.$$

An infinitesimal bending V determines a linearized second fundamental form b_V , which can be regarded as the first order change of the second fundamental form of the surface ψ under the displacement V. In coordinates, the (negated) linearized second fundamental form of V is given by

(17)
$$(b_V)_{ij} = n \cdot (\partial_i \partial_j \tilde{V} - \Gamma_{ij}^k \partial_k \tilde{V}),$$

cf. [Hor12] and the references therein. The linearized second fundamental form also occurs e.g. in the analysis in [GSP95].

The following lemma justifies our use of the symbol b_V here.

Lemma 2.3. If V is an infinitesimal bending, then $\partial_i \psi \cdot b_V \partial_j \psi = (b_V)_{ij}$.

Proof. We write μ , V, n etc. instead of $\tilde{\mu}_V$, \tilde{V} , \tilde{n} , and the coordinates of the second fundamental form are denoted h_{ij} , and we use the common convention regarding the raising and lowering of indices. By definition of the linearised Weingarten map, we have

$$b_V \partial_j \psi = -\partial_j \mu + \nabla_{\partial_j n} V = -\partial_j \mu - h_j^k \partial_k V.$$

Hence using $n \cdot \partial_i V + \mu \cdot \partial_i \psi = 0$ (which follows from $n \cdot \partial_i \psi = 0$), we see

$$\begin{aligned} \partial_i \psi \cdot b_V \partial_j \psi &= -\partial_i \psi \cdot \partial_j \mu - h_j^k \partial_i \psi \cdot \partial_k V \\ &= -\partial_j (\partial_i \psi \cdot \mu) + \partial_j \partial_i \psi \cdot \mu - h_j^k \partial_i \psi \cdot \partial_k V \\ &= \partial_j (\partial_i V \cdot n) + \Gamma_{ij}^k \partial_k \psi \cdot \mu - h_j^k \partial_i \psi \cdot \partial_k V \\ &= \partial_j \partial_i V \cdot n + \partial_i V \cdot \partial_j n - \Gamma_{ij}^k \partial_k V \cdot n - h_j^k \partial_i \psi \cdot \partial_k V \\ &= n \cdot \left(\partial_j \partial_i V - \Gamma_{ij}^k \partial_k V \right) - h_j^k \partial_i V \cdot \partial_k \psi - h_j^k \partial_i \psi \cdot \partial_k V \end{aligned}$$

This indeed agrees with $(b_V)_{ij}$ as defined in (17), because the last terms cancel by the definition of infinitesimal bendings.

We will frequently need the following diffeomorphism $\Phi^h: S^h \to S^1$:

$$\Phi^h(x) = \pi(x) + \frac{t(x)}{h}n(x).$$

The following lemma summarizes a computation that will later be used for the generic type of ansatz functions.

Lemma 2.4. Let $h \in (0, 1/2)$, let $V \in H^2(S; \mathbb{R}^3)$, and for $x \in S^h$ define

$$\rho(x) = V(x) + t(x)\mu_V(x).$$

Then the following equality holds on S^h :

(18)
$$\nabla \rho = \Omega_V - tb_V(T_S, T_S) - t^2 \nabla_{tan} \mu_V \mathbf{S} + (\nabla_{tan} V + t \nabla_{tan} \mu_V) T_S \left((I + t\mathbf{S})^{-1} - (I - t\mathbf{S}) \right),$$

where we extend V, μ_V , Ω_V , b_V , $\nabla_{tan}V$ etc. trivially from S to S^h .

Proof. For all $x \in S^h$ define

(19)
$$Q(x) = (I + t(x)\mathbf{S}(x))^{-1} - (I - t(x)\mathbf{S}(x)).$$

Since clearly $\nabla t = n$, formula (8) shows that on S^h :

$$\begin{aligned} \nabla \rho &= \left(\nabla_{tan} V + t \nabla_{tan} \mu_V \right) T_S \left(I + t \mathbf{S} \right)^{-1} + \mu_V \otimes n \\ &= \left(\nabla_{tan} V + t \nabla_{tan} \mu_V \right) T_S \left(I - t \mathbf{S} \right) + \mu_V \otimes n + \left(\nabla_{tan} V + t \nabla_{tan} \mu_V \right) T_S Q \\ &= \nabla_{tan} V + \mu_V \otimes n - t \nabla_{tan} V \mathbf{S} + t \nabla_{tan} \mu_V T_S - t^2 \nabla_{tan} \mu_V \mathbf{S} + \left(\nabla_{tan} V + t \nabla_{tan} \mu_V \right) T_S Q. \end{aligned}$$

By the definition of Ω_V and b_V this is the claim.

3 Elasticity framework and main result

Throughout this paper we assume that the limit

$$\gamma := \lim_{h \to 0} \frac{h}{\varepsilon(h)}$$

exists in $[0, \infty]$. We will frequently write ε instead of $\varepsilon(h)$, but always with the understanding that ε depends on h via γ .

Definition 3.1 (nonlinear material law). Let $0 < \alpha \leq \beta$ and $\rho > 0$. The class $\mathcal{W}(\alpha, \beta, \rho)$ consists of all measurable functions $W : \mathbb{R}^{3\times3} \to [0, +\infty]$ that satisfy the following properties:

- $\begin{array}{ll} \text{(W1)} & W \text{ is frame indifferent, i.e.} \\ & W(RF) = W(F) \quad \text{ for all } F \in \mathbb{R}^{3 \times 3}, \, R \in \mathrm{SO}(3); \\ \text{(W2)} & W \text{ is non degenerate, i.e.} \\ & W(F) \geq \alpha \operatorname{dist}^2(F, \mathrm{SO}(3)) \quad \text{ for all } F \in \mathbb{R}^{3 \times 3}; \\ & W(F) \leq \beta \operatorname{dist}^2(F, \mathrm{SO}(3)) \quad \text{ for all } F \in \mathbb{R}^{3 \times 3} \text{ with } \operatorname{dist}^2(F, \mathrm{SO}(3)) \leq \rho; \\ \text{(W2)} & W \text{ is minimal at } L \text{ is a} \end{array}$
- (W3) W is minimal at I, i.e.

$$W(I) = 0;$$

(W4) W admits a quadratic expansion at I, i.e. $W(I+G) = \mathcal{O}(G) + o(|G|^2) \quad \text{for all } G \in \mathbb{R}^{3 \times 3}$

$$W(I+G) = \mathcal{Q}(G) + O(|G|) \quad \text{for all } G \in \mathcal{Q}(G)$$

where \mathcal{Q} : $\mathbb{R}^{3 \times 3} \to \mathbb{R}$ is a quadratic form.

Definition 3.2 (admissible composite material). Let $0 < \alpha \leq \beta$ and $\rho > 0$. We say

$$W: S^1 \times \mathbb{R}^2 \times \mathbb{R}^{3 \times 3} \to \mathbb{R}^+ \cup \{+\infty\}$$

describes an admissible composite material of class $\mathcal{W}(\alpha, \beta, \rho)$ if

- (i) W is almost everywhere equal to a Borel function on $S^1 \times \mathbb{R}^2 \times \mathbb{R}^{3 \times 3}$,
- (ii) $W(\cdot, y, F)$ is continuous for almost every $y \in \mathbb{R}^2$ and $F \in \mathbb{R}^{3 \times 3}$,
- (iii) $W(x, \cdot, F)$ is Y-periodic for all $x \in \Omega$ and almost every $F \in \mathbb{R}^{3 \times 3}$,

(iv) $W(x, y, \cdot) \in \mathcal{W}(\alpha, \beta, \rho)$ for all $x \in S^1$ and almost every $y \in \mathbb{R}^2$.

Assumption 3.3. We assume that

- W describes an admissible composite material of class W(α, β, ρ) in the sense of Definition 3.2.
- Q is the quadratic energy density associated to W through expansion (W4) in Definition 3.1.
- The following uniformity is valid

$$\lim_{G \to 0} \operatorname{ess\,sup}_{(x,y) \in S^1 \times \mathcal{Y}} \frac{|W(x,y,I+G) - \mathcal{Q}(x,y,G)|}{|G|^2} = 0.$$

We collect some basic properties of admissible W and the associated quadratic forms Q; a proof can be found in [Neu12, Lemma 2.7].

Lemma 3.4. Let W and Q satisfy the assumption (3.3). Then

- (Q1) $\mathcal{Q}(\cdot, y, \cdot)$ is continuous for almost every $y \in \mathbb{R}^2$,
- (Q2) $\mathcal{Q}(x,\cdot,G)$ is Y-periodic and measurable for all $x \in S^1$ and all $F \in \mathbb{R}^{3\times 3}$,
- (Q3) for all $x \in S^1$ and almost every $y \in \mathbb{R}^2$ the map $\mathcal{Q}(x, y, \cdot)$ is quadratic and satisfies

$$||\operatorname{sym} G||^2 \leq \mathcal{Q}(x, y, G) = \mathcal{Q}(x, y, \operatorname{sym} G) \leq \beta |\operatorname{sym} G|^2 \quad \text{for all } G \in \mathbb{R}^{3 \times 3}.$$

Furthermore, there exists a monotone function $m : \mathbb{R}^+ \to \mathbb{R}^+ \cup \{+\infty\}$ such that $m(\delta) \to 0$ as $\delta \to 0$ and

(20)
$$\forall G \in \mathbb{R}^{3 \times 3} : |W(x, y, I + G) - \mathcal{Q}(x, y, G)| \le |G|^2 m(|G|)$$

for all $x \in S^1$ and almost every $y \in \mathbb{R}^2$.

Let W be an energy density satisfying Assumption 3.3. The elastic energy per unit thickness of a deformation $u^h \in H^1(S^h; \mathbb{R}^3)$ of the shell S^h is given by

$$E^{h}(u^{h}) = \frac{1}{h} \int_{S^{h}} W\left(\Phi^{h}(x), r(x)/\varepsilon, \nabla u^{h}(x)\right) dx.$$

We denote by \mathcal{B} the L^2 -closure of the set

$$\{q_w: w \in H^1(S; \mathbb{R}^3)\}.$$

As this is a linear space, its strong and its weak L^2 -closure coincide. The set \mathcal{B} is a closed linear subspace of $L^2(S; \mathbb{S})$. The space \mathcal{B} is also encountered in the context of shell models derived from linearized elasticity; see [SP89a, SP89b, GSP95] for details.

Before we give the main statement we have to define the limit functionals. To do that we need the definition of the relaxation fields and the cell formulas.

Definition 3.5. We define the following operators:

$$\begin{split} \mathcal{U}_{0} &: \dot{H}^{1}(\mathcal{Y}; \mathbb{R}^{2}) \times L^{2}(I \times \mathcal{Y}; \mathbb{R}^{3}) \to L^{2}(I \times \mathcal{Y}; \mathbb{R}^{3 \times 3}_{\text{sym}}), \\ \mathcal{U}_{0}(\zeta, g) &= \begin{pmatrix} \text{sym} \nabla_{y} \zeta & g_{1} \\ (g_{1}, g_{2}) & g_{3} \end{pmatrix}_{ij} \tau^{i} \otimes \tau^{j}, \\ \mathcal{U}_{0}^{0} &: \dot{H}^{1}(\mathcal{Y}; \mathbb{R}^{2}) \times \dot{H}^{2}(\mathcal{Y}) \times L^{2}(I \times \mathcal{Y}; \mathbb{R}^{3}) \to L^{2}(I \times \mathcal{Y}; \mathbb{R}^{3 \times 3}), \\ \mathcal{U}_{0}^{0}(\zeta, \varphi, g) &= \begin{pmatrix} \text{sym} \nabla_{y} \zeta - t \nabla_{y}^{2} \varphi & g_{1} \\ (g_{1}, g_{2}) & g_{3} \end{pmatrix}_{ij} \tau^{i} \otimes \tau^{j}, \\ \mathcal{U}_{0,\gamma_{1}}^{1} &: \dot{H}^{1}(\mathcal{Y}; \mathbb{R}^{2}) \times \dot{H}^{2}(\mathcal{Y}) \times L^{2}(I \times \mathcal{Y}; \mathbb{R}^{3}) \to L^{2}(I \times \mathcal{Y}; \mathbb{R}^{3 \times 3}), \\ \mathcal{U}_{0,\gamma_{1}}^{1}(\zeta, \varphi, g) &= \begin{pmatrix} \text{sym} \nabla_{y} \zeta + \frac{1}{\gamma_{1}} \varphi \mathbf{S}(x) - t \hat{\nabla}_{y}^{2} \varphi & g_{1} \\ (g_{1}, g_{2}) & g_{3} \end{pmatrix}_{ij} \tau^{i} \otimes \tau^{j}, \\ \mathcal{U}_{\infty} &: L^{2}(I; \dot{H}^{1}(\mathcal{Y}; \mathbb{R}^{2})) \times L^{2}(I; \dot{H}^{1}(\mathcal{Y})) \times L^{2}(I; \mathbb{R}^{3}) \to L^{2}(I \times \mathcal{Y}; \mathbb{R}^{3 \times 3}), \\ \mathcal{U}_{\infty}(\zeta, \psi, c) &= \begin{pmatrix} \text{sym} \nabla_{y} \zeta & \partial_{y_{1}} \psi + c_{1} \\ \nabla_{y} \psi + (c_{1}, c_{2}) & c_{3} \end{pmatrix}_{ij} \tau^{i} \otimes \tau^{j}, \\ \mathcal{U}_{\gamma}(\phi) &= \text{sym}(\nabla_{y} \phi, \frac{1}{\gamma} \partial_{3} \phi)_{ij} \tau^{i} \otimes \tau^{j}. \end{split}$$

For $\gamma \in (0,\infty)$ we introduce the function spaces of *relaxation fields*

$$L_{\gamma}(I \times \mathcal{Y}) = \left\{ \mathcal{U}_{\gamma}(\phi) : \phi \in \dot{H}^{1}(I \times \mathcal{Y}; \mathbb{R}^{3}) \right\}$$

For $\gamma = \infty$ and $\gamma = 0$ we define

$$L_{\infty}(I \times \mathcal{Y}) = \left\{ U_{\infty}(\zeta, \psi, c) : (\zeta, \psi, c) \in L^{2}(I; \dot{H}^{1}(\mathcal{Y}; \mathbb{R}^{2})) \times L^{2}(I; \dot{H}^{1}(\mathcal{Y})) \times L^{2}(I; \mathbb{R}^{3}) \right\},\$$
$$L_{0}(I \times \mathcal{Y}) = \left\{ U_{0}(\zeta, g) : (\zeta, g) \in \dot{H}^{1}(\mathcal{Y}; \mathbb{R}^{2}) \times L^{2}(I \times \mathcal{Y}; \mathbb{R}^{3}) \right\}.$$

We also introduce

$$L_0^0(I \times \mathcal{Y}) = \left\{ \begin{pmatrix} \operatorname{sym} \nabla_y \zeta - t \nabla_y^2 \varphi & g_1 \\ (g_1, g_2) & g_3 \end{pmatrix}_{ij} \tau^i \otimes \tau^j : \zeta \in \dot{H}^1(\mathcal{Y}; \mathbb{R}^2), \\ \varphi \in \dot{H}^2(\mathcal{Y}), \ g \in L^2(I \times \mathcal{Y}; \mathbb{R}^3) \right\}$$

and for $x \in S, \gamma_1 \in (0, \infty)$ we define

$$L^{1}_{0,\gamma_{1}}(I \times \mathcal{Y}) = \left\{ \begin{pmatrix} \operatorname{sym} \nabla_{y} \zeta + \frac{1}{\gamma_{1}} \varphi \mathbf{S}(x) - t \nabla_{y}^{2} \varphi & g_{1} \\ (g_{1}, g_{2}) & g_{3} \end{pmatrix}_{ij} \tau^{i} \otimes \tau^{j} : \zeta \in \dot{H}^{1}(\mathcal{Y}; \mathbb{R}^{2}), \\ \varphi \in \dot{H}^{2}(\mathcal{Y}), \ g \in L^{2}(I \times \mathcal{Y}; \mathbb{R}^{3}) \right\}.$$

Remark 1. Notice that all the operators \mathcal{U} and the appropriate spaces also depend on $x \in S$. For simplicity of writing we do not write x in the notation.

For $\gamma \in (0,\infty]$ and $x \in S$ we define the functions $\mathcal{Q}_{\gamma}(x) : \mathbb{S}(x)_{\text{sym}} \times \mathbb{S}(x)_{\text{sym}} \to \mathbb{R}$ as follows:

(21)
$$\mathcal{Q}_{\gamma}(x,q^1,q^2) = \inf_{U \in L_{\gamma}(I \times \mathcal{Y})} \int_I \int_{\mathcal{Y}} \mathcal{Q}\left(x + tn(x), y, q^1 + tq^2 + U\right) dy dt.$$

For $\gamma_1 \in (0,\infty)$ we define $\mathcal{Q}_0^0(x), Q_{0,\gamma_1}^1(x) : \mathbb{S}(x)_{\text{sym}} \times \mathbb{S}(x)_{\text{sym}} \to \mathbb{R}$ as follows:

(22)
$$\mathcal{Q}_{0}^{0}(x,q^{1},q^{2}) = \inf_{U \in L_{0}^{0}(I \times \mathcal{Y})} \int_{I} \int_{\mathcal{Y}} \mathcal{Q}\left(x + tn(x), y, q^{1} + tq^{2} + U\right) dy dt$$

(23)
$$\mathcal{Q}_{0,\gamma_{1}}^{1}(x,q^{1},q^{2}) = \inf_{U \in L_{0,\gamma_{1}}^{1}(I \times \mathcal{Y})} \int_{I} \int_{\mathcal{Y}} \mathcal{Q}\left(x + tn(x), y, q^{1} + tq^{2} + U\right) dy dt$$

Remark 2. We discuss the cell formula in the limiting cases $\gamma = 0$ and $\gamma = \infty$.

(i) In the case $\gamma = \infty$ define

$$\widetilde{L}_{\infty}(I \times \mathcal{Y}) := \left\{ \sum_{i,j=1}^{3} \left(\begin{array}{cc} \operatorname{sym} \nabla_{y} \zeta & \partial_{y_{1}} \psi + c_{1} \\ \partial_{y_{2}} \psi + c_{2} \\ \nabla_{y} \psi + (c_{1}, c_{2}) & c_{3} \end{array} \right)_{ij} \tau^{i} \otimes \tau^{j} :$$
$$\zeta \in \dot{H}^{1}(\mathcal{Y}, \mathbb{R}^{2}), \psi \in \dot{H}^{1}(\mathcal{Y}), \ c \in \mathbb{R}^{3} \right\}.$$

Also define for $(x,t)\in S\times I$

(24)
$$\widetilde{\mathcal{Q}}_{\infty}(x,t,q^1,q^2) = \inf_{U \in \widetilde{L}_{\infty}(I \times \mathcal{Y};\mathbb{R}^{3\times3}_{\mathrm{sym}})} \int_{\mathcal{Y}} \mathcal{Q}\Big(x+tn(x),y,q^1+tq^2+U\Big) \,\mathrm{d}y.$$

It is easy to see that $\widetilde{\mathcal{Q}}_{\infty}$ is, for a fixed $x \in S, t \in I$, a quadratic in q^1, q^2 . We have

(25)
$$\mathcal{Q}_{\infty}(x,q^1,q^2) = \int_I \widetilde{\mathcal{Q}}_{\infty}(x+tn(x),q^1,q^2) \,\mathrm{d}t$$

(ii) Define as in [LMP10]

(26)
$$\mathcal{Q}_2(x,t,q^1,q^2) = \min_{M \in \mathbb{R}^{3 \times 3}_{\text{sym}}} \{ \mathcal{Q}(x+tn(x),M) : q^1 + tq^2 - \sum_{i,j=1,2} (M\tau_j \cdot \tau_i)\tau^i \otimes \tau^j = 0 \}.$$

Also define

$$\begin{split} \widetilde{L}_{0}^{0}(I \times \mathcal{Y}) &= \left\{ \sum_{i,j=1}^{2} \left(\operatorname{sym} \nabla_{y} \zeta - t \nabla_{y}^{2} \varphi \right)_{ij} \tau^{i} \otimes \tau^{j} : \zeta \in \dot{H}^{1}(\mathcal{Y}, \mathbb{R}^{2}), \varphi \in \dot{H}^{2}(\mathcal{Y}) \right\}, \\ \widetilde{L}_{0,\gamma_{1}}^{1}(I \times \mathcal{Y}) &= \left\{ \sum_{i,j=1}^{2} \left(\operatorname{sym} \nabla_{y} \zeta + \frac{1}{\gamma_{1}} \varphi \mathbf{S}_{ij}(x) - t \nabla_{y}^{2} \varphi \right)_{ij} \tau^{i} \otimes \tau^{j} : \\ \zeta \in \dot{H}^{1}(\mathcal{Y}, \mathbb{R}^{2}), \varphi \in \dot{H}^{2}(\mathcal{Y}) \right\} \end{split}$$

It can be easily seen that we have for the cell formula

(27)
$$\mathcal{Q}_0^0(x,q^1,q^2) = \inf_{U \in \widetilde{L}_0^0(I \times \mathcal{Y})} \iint_{I \times \mathcal{Y}} \mathcal{Q}_2\left(x + tn(x), y, q^1 + tq^2 + U\right) \mathrm{d}t \,\mathrm{d}y,$$

for i=0,2 i.e.

(28)
$$\mathcal{Q}_{0,\gamma_1}^1(x,q^1,q^2) = \inf_{\substack{U \in \widetilde{L}_{0,\gamma_1}^1(I \times \mathcal{Y})}} \iint_{I \times \mathcal{Y}} \mathcal{Q}_2\left(x + tn(x), y, q^1 + tq^2 + U\right) \mathrm{d}t \,\mathrm{d}y.$$

In the case when Q does not depend on t we have that

$$\begin{aligned} \mathcal{Q}_{0}^{0}(x,q^{1},q^{2}) &= \inf_{\zeta \in \dot{H}^{1}(\mathcal{Y},\mathbb{R}^{2})} \int_{\mathcal{Y}} \mathcal{Q}_{2}(x,y,q^{1}+\sum_{i,j=1}^{2}(\operatorname{sym}\nabla_{y}\zeta)_{ij}\tau^{i}\otimes\tau^{j}) \,\mathrm{d}y \\ &+ \frac{1}{12}\inf_{\varphi \in \dot{H}^{2}(\mathcal{Y})} \int_{\mathcal{Y}} \mathcal{Q}_{2}(x,y,q^{2}+\sum_{i,j=1}^{2}(\nabla_{y}^{2}\varphi)_{ij}\tau_{i}\otimes\tau_{j}) \,\mathrm{d}y, \\ \mathcal{Q}_{0}^{1,\gamma_{1}}(x,q^{1},q^{2}) &= \\ \inf_{\zeta \in \dot{H}^{1}(\mathcal{Y},\mathbb{R}^{2}),\varphi \in \dot{H}^{2}(\mathcal{Y})} & \left(\int_{\mathcal{Y}} \mathcal{Q}_{2}\Big(x,y,q^{1}+\sum_{i,j=1}^{2}\big((\operatorname{sym}\nabla_{y}\zeta)_{ij}+\frac{1}{\gamma_{1}}\varphi\mathbf{S}_{ij}\big)\tau^{i}\otimes\tau^{j}\Big) \,\mathrm{d}y \\ &+ \frac{1}{12}\int_{\mathcal{Y}} \mathcal{Q}_{2}(x,y,q^{2}+\sum_{i,j=1}^{2}(\nabla_{y}^{2}\varphi)_{ij}\tau_{i}\otimes\tau_{j}) \,\mathrm{d}y \Big). \end{aligned}$$

Remark 3. In the same way as in [NV] we can prove the following: For every $q^1, q^2 \in S(x)_{sym}$ and $x \in S$ we have that

$$\lim_{\gamma \to \infty} \mathcal{Q}_{\gamma}(x, q^1, q^2) = \mathcal{Q}_{\infty}(x, q^1, q^2)$$
$$\lim_{\gamma \to 0} \mathcal{Q}_{\gamma}(x, q^1, q^2) = \mathcal{Q}_0^0(x, q^1, q^2)$$

Remark 4. Notice that when $\mathbf{S} = 0$ then all spaces L_0^0 and L_{0,γ_1}^1 coincide for $\gamma_1 \in (0,\infty)$. This corresponds to the observation in the von Kármán plate theory that for $\gamma = 0$ one obtains only one relaxation space, cf. [NV] for details.

For $\gamma \in (0,\infty]$ define the functionals $I_{\gamma}: H^2(S;\mathbb{R}^3) \times L^2(S;\mathbb{S}) \to \mathbb{R}$ by setting

(29)
$$I_{\gamma}(V, B_w) = \int_S \mathcal{Q}_{\gamma}(\cdot, B_w + \frac{1}{2}(dV)^2, -b_V) d\mathcal{H}^2,$$

and define the functionals $I^0_0: H^2(S;\mathbb{R}^3) \times L^2(S;\mathbb{S}) \to \mathbb{R}$ by

(30)
$$I_0^0(V, B_w) = \int_S \mathcal{Q}_0^0(\cdot, B_w + \frac{1}{2}(dV)^2, -b_V) d\mathcal{H}^2$$

as well as, for $\gamma_1 \in (0,\infty)$, define the functionals $I^1_{0,\gamma_1}: H^2(S;\mathbb{R}^3) \times L^2(S;\mathbb{S}) \to \mathbb{R}$ by

(31)
$$I_{0,\gamma_1}^1(V,B_w) = \int_S \mathcal{Q}_{0,\gamma_1}^1(\cdot,B_w + \frac{1}{2}(dV)^2, -b_V) d\mathcal{H}^2$$

This is our main result:

Theorem 3.6. Let W satisfy Assumption 3.3 and assume that $u^h \in H^1(S^h; \mathbb{R}^3)$ satisfy

(32)
$$\limsup_{h \to 0} h^{-4} E^h(u^h) < \infty.$$

Then the following are true:

(i) (compactness). There exists a subsequence, still denoted by (\bar{y}^h) , and there exist $Q^h \in SO(3)$ and $c^h \in \mathbb{R}^3$ such that the sequences y^h and V^h defined by

$$y^h = (Q^h)^T \bar{y}^h - c^h$$

and

$$V^{h}(x) = \frac{1}{h} \left(\int_{I} y^{h}(x + tn(x))dt - x \right) \text{ for all } x \in S$$

satisfy the following:

(a) We have

$$y^h \to \pi$$
 strongly in $H^1(S^1; \mathbb{R}^3)$.

(b) There exists an infinitesimal bending $V \in H^2(S; \mathbb{R}^3)$ of S such that

 $V^h \to V$ strongly in $H^1(S; \mathbb{R}^3)$.

(c) There exists $B_w \in L^2(S; \mathbb{S})$ such that

$$\frac{1}{h}q_{V^h} \rightharpoonup B_w \text{ weakly in } L^2(S; \mathbb{S}).$$

(ii) (lower bound). Defining I_{γ} by (29) and I_0^0 by (30) and I_0^{1,γ_1} by (31), we have

$$\liminf_{h \to 0} h^{-4} E^h(u^h) \ge \begin{cases} I_{\gamma}(V, B_w) & \text{if } h/\varepsilon \to \gamma \in (0, \infty] \\ I_0^0(V, B_w) & \text{if } \varepsilon \gg h \gg \varepsilon^2 \\ I_{0,\gamma_1}^1(V, B_w) & \text{if } \varepsilon^2/h \to \frac{1}{\gamma_1} \in (0, \infty) \end{cases}$$

(iii) (recovery sequence) For any infinitesimal bending $V \in H^2(S, \mathbb{R}^3)$ of S and any $B_w \in \mathcal{B}$, there exist $u^h \in H^1(S^h; \mathbb{R}^3)$ satisfying (32), and such that the conclusions of part (i) are true with $Q^h = I$ and $c^h = 0$, and

$$\lim_{h \to 0} h^{-4} E^h(u^h) = \begin{cases} I_{\gamma}(V, B_w) & \text{if } h/\varepsilon \to \gamma \in (0, \infty] \\ I_0^0(V, B_w) & \text{if } \varepsilon \gg h \gg \varepsilon^2 \\ I_{0,\gamma_1}^1(V, B_w) & \text{if } \varepsilon^2/h \to \frac{1}{\gamma_1} \in (0, \infty). \end{cases}$$

From now on $u^h \in H^1(S^h; \mathbb{R}^3)$ will always denote a sequence satisfying (32).

3.1 Unit thickness rescaling

Recall that $\Phi^h: S^h \to S^1$ is given by

$$\Phi^h(x) = \pi(x) + \frac{t(x)}{h}n(x).$$

Since $\nabla t = n$, Lemma 2.1 and formula (8) show (recall that n is extended trivially to S^1):

$$\nabla \Phi^{h} = \nabla \pi + \frac{t}{h} \nabla n + \frac{1}{h} n \otimes n$$
$$= T_{S} (I + t\mathbf{S})^{-1} + \frac{t}{h} \mathbf{S} T_{S} (I + t\mathbf{S})^{-1} + \frac{1}{h} n \otimes n$$

Since T_S clearly commutes with **S**, we see that T_S commutes with $(I + t\mathbf{S})^{-1}$ as well. Hence

(33)
$$\nabla \Phi^h = (I_h + \frac{t}{h} \mathbf{S})(I + t\mathbf{S})^{-1} \text{ on } S^h,$$

where $I_h = T_S + \frac{1}{h}n \otimes n$. Following [FJM02], for given $u: S^h \to \mathbb{R}^3$ we define its rescaled version $y: S^1 \to \mathbb{R}^3$ by

$$y(\Phi^h) = u$$
 on S^h .

We define the rescaled gradient of y by the condition

(34)
$$\nabla_h y(\Phi^h) = \nabla u \text{ on } S^h.$$

To compute ∇_h more explicitly, insert the definition of y into (34) and use (33) to find

(35)
$$\nabla_h y = \nabla y \ (I_h + t\mathbf{S})(I + ht\mathbf{S})^{-1} \text{ on } S^1,$$

In order to express the elastic energy in terms of the new variables, we associate with $y:S^1\to\mathbb{R}^3$ the energy

$$I^{h}(y) = \int_{S^{1}} W(x, r(x)/\varepsilon, \nabla_{h} y(x)) \det (I + t(x) \mathbf{S}(x))^{-1} dx$$
$$= \int_{S} \int_{I} W(x + tn(x), r(x)/\varepsilon, \nabla_{h} y(x)) dt d\mathcal{H}^{2}$$

By a change of variables we have

$$E^{h}(u^{h}) = \frac{1}{h} \int_{S^{1}} W\left(\cdot, r/\varepsilon, \nabla_{h} y^{h}\right) \left| \det \nabla(\Phi^{h})^{-1} \right|.$$

Using (33) it is easy to see that

$$E^{h}(u^{h}) = I^{h}(y^{h}) + o(h^{4})$$
 as $h \to 0$.

3.2 FJM-compactness

The following lemma proves the first part of Theorem 3.6. It is a direct consequences of [FJM02, Theorem 3.1] and of arguments in [FJM06]. We refer to [LMP10] for the extension to the present setting.

Lemma 3.7. There exist a constant C > 0, independent of h, and a sequence of matrix fields $(\mathbb{R}^h) \subset H^1(S; SO(3))$ (extended trivially to S^h) and there exists a sequence of matrices $(Q^h) \subset SO(3)$ such that:

- (i) $\limsup_{h \to 0} h^{-5/2} \|\nabla u^h R^h\|_{L^2(S^h)} < \infty$
- (*ii*) $\limsup_{h\to 0} h^{-1} \|\nabla R^h\|_{L^2(S)} < \infty$
- (*iii*) $\limsup_{h\to 0} h^{-1} \| (Q^h)^T R^h I \|_{L^p(S)} < \infty$, for all $p \in [1, \infty)$.
- (iv) $(Q^h)^T R^h \to I$ strongly in H^1 .

Moreover, there exists a matrix field $A \in H^1(S, so(3))$ taking values in the space of skew symmetric matrices, such that (after passing to subsequences)

(v)
$$\frac{1}{h}((Q^h)^T R^h - I) \rightarrow A$$
, weakly in $H^1(S; \mathbb{R}^{3 \times 3})$.
(vi) $\frac{1}{h^2} \operatorname{sym}((Q^h)^T R^h - I) \rightarrow \frac{1}{2}A^2$, strongly in $L^p(S; \mathbb{R}^{3 \times 3})$, for all $p \in [1, \infty)$.

Moreover, the following are true:

- (i) $\limsup_{h\to 0} \frac{1}{h^2} \|\nabla_h \bar{y}^h R^h\|_{L^2(S^1)} < \infty.$
- (ii) $\frac{1}{h}((Q^h)^T \nabla_h y^h I) \rightharpoonup A$, weakly in H^1 up to a subsequence.

Define $y^h \in H^1(S^1; \mathbb{R}^3)$ by

$$y^h = (Q^h)^T \bar{y}^h - c^h,$$

where

$$c^{h} = \oint_{S} \int_{I} \left((Q^{h})^{T} \bar{y}^{h}(x + tn(x)) - x \right) dt \ d\mathcal{H}^{2}(x).$$

Introduce the (average) midplane displacements $V^h: S \to \mathbb{R}^3$ by setting

(36)
$$V^{h}(x) := \frac{1}{h} \left(\int_{I} \left(y^{h}(x+tn(x)) \right) dt - x \right) \text{ for all } x \in S.$$

Then $\int_{S} V^{h} = 0$ and (after passing to a subsequence)

- (iii) $y^h \to \pi$, strongly in $H^1(S^1; \mathbb{R}^3)$.
- (iv) There exists an infinitesimal bending $V \in H^2(S; \mathbb{R}^3)$ of S with $\Omega_V = A$ and such that $V^h \to V$ strongly in $H^1(S; \mathbb{R}^3)$.
- (v) $\frac{1}{h}q_{V^h}$ is bounded in $L^2(S; \mathbb{R}^{3\times 3})$.

In what follows we replace the sequence R^h by $(Q^h)^T R^h$ and the sequence y^h by \bar{y}^h , so we assume without loss of generality that $Q^h = Id$. Expressed in the unrescaled variables, we have

 $V^{h}(x) = \frac{1}{h^{2}} \left(\int_{I^{h}} u^{h}(x + tn(x)) dt - x \right),$

 $\text{i.e. } x+hV^h(x)=f_{I^h}\,u^h(x+tn(x))\ dt.$

Next we modify the displacement fields V^h into more regular fields V^h_s enjoying a similar compactness.

Lemma 3.8. There exist $V_s^h \in H^2(S; \mathbb{R}^3)$ with $f_S V_s^h = 0$ satisfying

(37)
$$\limsup_{h \to 0} h^{-1} \| V_s^h - V^h \|_{H^1(S)} < \infty$$

and

(38)
$$\left\| \left(\nabla_{tan} V_s^h - \frac{R^h - I}{h} \right) T_S \right\|_{L^2(S)} \le \left\| \left(\nabla_{tan} V^h - \frac{R^h - I}{h} \right) T_S \right\|_{L^2(S)}$$

Moreover, (V_s^h) is uniformly bounded in $H^2(S)$ and

(39)
$$V_s^h \rightharpoonup V \text{ weakly in } H^2(S; \mathbb{R}^3).$$

Proof. We follow [NV, Proposition 3.1]. For i = 1, 2, 3 denote by p_i the *i*-th row of the matrix $\frac{R^h(\psi)-I}{h}\nabla\psi$. We define $\tilde{V}_s^h \in H^2(\tilde{\omega}; \mathbb{R}^3)$ such that $(\tilde{V}_s^h)_i$ is a minimiser of the functional

$$v\mapsto \int_{\tilde{\omega}} |\nabla v - p_i|^2 \, dx$$

among all $v \in H^1(\tilde{\omega})$ satisfying $\int_{\omega} v = 0$, and we define V_s^h via $V_s^h(\psi) = \tilde{V}_s^h$. The bound (38) follows from the minimality of V_s^h . Combining the tangential components of (40) and (41) below, we obtain

$$\|\nabla_{tan}V_s^h - \nabla_{tan}V^h\|_{L^2(S)} \le Ch.$$

Hence (37) follows from Poincaré's inequality on S.

Since $\partial \tilde{\omega}$ is $C^{1,1}$, standard regularity estimates for minimisers imply that $V_s^h \in H^2(S)$ with bounds

$$\|V_s^h\|_{H^2(S)} \le C \left(\|\operatorname{div} p\|_{L^2(\tilde{\omega})} + \|p\|_{L^2(\tilde{\omega})} \right).$$

Hence Lemma 3.7 (v) ensures that (V_s^h) is uniformly bounded in $H^2(S)$. Since $V^h \to V$ in H^1 , the bound (37) therefore implies (39).

Lemma 3.9. There exist maps F_s^h , $F^h \in L^2(S; \mathbb{R}^{3 \times 3})$ with

$$\limsup_{h \to 0} h^{-2} \left(\|F_s^h\|_{L^2(S)} + \|F^h\|_{L^2(S)} \right) < \infty,$$

such that

(40) $R^h = I + h\Omega_{V^h} + F^h$

and

(41)
$$R^h = I + h\Omega_{V_s^h} + F_s^h$$

Proof. For brevity, we set $\mu_s^h = \mu_{V_s^h}$ and $\mu^h = \mu_{V^h}$.

We first verify the tangential component of (40). Let τ be a C^1 tangent vector field along S. Then by the definition of V^h and using $\int_{I_h} R^h(x) t \mathbf{S}(x) dt = 0$, we see that $(I + h \nabla_{tan} V^h T_S) \tau$ equals

$$\partial_{\tau}(id+hV^{h}) = \partial_{\tau} \left(\int_{I} y^{h}(x+tn(x)) dt \right) = \frac{1}{h} \partial_{\tau} \left(\int_{I_{h}} u^{h}(x+tn(x)) dt \right)$$
$$= \frac{1}{h} \int_{I_{h}} \nabla u^{h}(x+tn(x))(I+t\mathbf{S}(x)) dt\tau(x)$$
$$= R^{h}(x)\tau(x) - M^{h}(x)\tau(x),$$

where we have introduced

$$M^{h}(x) = -\frac{1}{h} \int_{I_{h}} \left(\nabla u^{h}(x+tn(x)) - R^{h}(x) \right) \left(I + t\mathbf{S}(x) \right) dt.$$

Clearly,

$$\int_{S} |M^{h}|^{2} d\mathcal{H}^{2} = \frac{1}{h} \int_{S \times I_{h}} |\nabla u^{h}(x + tn(x)) - R^{h}(x)|^{2} |I + t\mathbf{S}(x)|^{2} d\mathcal{H}^{2}(x) dt$$
$$\leq \frac{C}{h} \int_{S^{h}} |\nabla u^{h}(x) - R^{h}(x)|^{2} dx \leq Ch^{4}.$$

To verify the normal component of (40),

we compute using the tangential part of (40):

$$\tau \cdot R^{h}n = -n \cdot R^{h}\tau + 2n \cdot (\operatorname{sym} R^{h})\tau$$
$$= -n \cdot (R^{h} - I)\tau + 2n \cdot \operatorname{sym} \left(R^{h} - I\right)\tau$$
$$= -hn \cdot \partial_{\tau}V^{h} - n \cdot F^{h}\tau + 2n \cdot \operatorname{sym} \left(R^{h} - I\right)\tau$$
$$= h\mu^{h} \cdot \tau - n \cdot F^{h}\tau + 2n \cdot \operatorname{sym} \left(R^{h} - I\right)\tau.$$

In the last step we used (13). As

(42)
$$\left\| \operatorname{sym} \left(R^h - I \right) \right\|_{L^2(S)} \le Ch^2$$

by Lemma 3.7, we conclude that

(43)
$$\left\|T_{S}R^{h}n - h\mu^{h}\right\| \le Ch^{2}.$$

But again by (42) we have

$$||n \cdot (R^h n - n)||_{L^2(S)} = ||n \cdot \operatorname{sym}(R^h - I)n||_{L^2(S)} \le Ch^2.$$

Since $R^h n = (n \cdot R^h n) n + T_S R^h n$, we conclude that $R^h n$ agrees – up to an error term whose $L^2(S)$ -norm is dominated by h^2 – with $n + h\mu^h$. This concludes the proof of (40). The tangential component of (40) together with (38) imply that the tangential component of (41) is satisfied. But then the normal component of (41) follows from its tangential component in exactly the same way in which the normal component of (40) followed from its tangential component.

3.3 Two-scale convergence

Recall that we extend the chart r trivially from S to S^1 .

Definition 3.10 (two-scale convergence). We say that a sequence $g^h \in L^2(S^1)$, weakly two-scale converges in L^2 to the function $g \in L^2(S^1, L^2(\mathcal{Y}))$ as $h \to 0$, if the sequence g^h is bounded in $L^2(S^1)$ and

$$\lim_{h \to 0} \int_{S^1} g^h(x) \, \psi(x, r(x)/\varepsilon) \, \mathrm{d}x = \iint_{S^1 \times \mathcal{Y}} g(x, y) \, \psi(x, y) \, \mathrm{d}y \, \mathrm{d}x$$

for all $\psi \in C^{\infty}_{c}(S^{1}, C(\mathcal{Y}))$. We say that g^{h} strongly two-scale converges to g if, in addition,

$$\lim_{h \to 0} \|g^h\|_{L^2(S^1)} = \|g\|_{L^2(S^1 \times Y)}.$$

We write $g^h \xrightarrow{2,\gamma} g$ in L^2 (resp. $g^h \xrightarrow{2,\gamma} g$ in L^2) for weak (resp. strong) two-scale convergence in L^2 .

For the basic properties of two-scale convergence we refer to [Ngu89, All92, Vis06]. If $g^h \xrightarrow{2,\gamma} g$ then $g^h \xrightarrow{} \int_{\mathcal{Y}} g(\cdot, y) dy$ weakly in L^2 . If g^h is bounded in $L^2(S^1)$ then it has subsequence which weakly two scale converges to some $g \in L^2(S^1; L^2(\mathcal{Y}))$.

The following lemma summarizes standard results about two scale convergence and adapts them to a possibly curved surface. Its proof follows easily from the analogous statements for the planar case (see v) and vi) of Lemma A.1 in the Appendix).

Lemma 3.11. (i) if $(g^h)_{h>0} \subset H^1(S^1)$ is bounded, then there exist $g_0 \in H^1(S^1)$ and $g_1 \in L^2(S^1; \dot{H}^1(\mathcal{Y}))$ such that, after passing to a subsequence, $\nabla g^h \xrightarrow{2,\gamma} g$, where

$$g = \nabla g_0 + \nabla_y g_1(x, y).$$

(ii) if $(g^h) \subset H^2(S^1; \mathbb{R}^3)$ is bounded, then there exist $g_0 \in H^2(S^1)$ and $g_1 \in L^2(S^1; \dot{H}^2(\mathcal{Y}))$ such that, after passing to a subsequence,

$$\nabla^2 g^h \xrightarrow{2,\gamma} g,$$

where

$$g = \nabla^2 g_0 + \sum_{i,j \le 2} \left(\partial_{y_i y_j}^2 g_1(x, y) \right) \tau^i \otimes \tau^j.$$

(iii) if (g^h) is bounded in $H^1(S^1; \mathbb{R}^3)$ then we have $\nabla g^h \xrightarrow{2,\gamma} g$, along a subsequence, and there exist $g_0 \in H^1(S^1; \mathbb{R}^3)$ and $g_1 \in L^2(S^1; \dot{H}^1(\mathcal{Y}; \mathbb{R}^3))$ such that

$$g = \nabla g_0 + \sum_{i \le 3; j \le 2} \left(\nabla_y g_1(x, y) \right)_{ij} \tau^i \otimes \tau^j.$$

(iv) if (g^h) is bounded in $H^2(S^1; \mathbb{R}^3)$ then we have $\nabla^2 g^h \xrightarrow{2,\gamma} g$, along a subsequence, and there exist $g_0 \in H^2(S^1; \mathbb{R}^3)$ and $g_1 \in L^2(S^1; \dot{H}^2(\mathcal{Y}; \mathbb{R}^3))$ such that

$$g = \nabla^2 g_0 + \sum_{i \le 3; j,k \le 2} \left(\partial^2_{y_j y_k} g_{1,i}(x,y) \right) \tau^i \otimes \tau^j \otimes \tau^k$$

4 Two-scale compactness and lower bound

Next we will identify the space of possible two scale limits of symmetrized gradients. The following auxiliary result is standard and it can be easily derived, e.g., using Fourier transforms.

Lemma 4.1. Let $B \in L^2(\omega; L^2(\mathcal{Y}; \mathbb{R}^{2 \times 2}_{sym}))$ have the following property: for every

 $\Psi \in C_0^{\infty}(\omega; C^{\infty}(\mathcal{Y}; \mathbb{R}^{2 \times 2}_{\mathrm{sym}}))$

satisfying

(44)
$$\Psi(\xi, y) = \psi(\xi) \operatorname{cof} \nabla^2 F(y),$$

for some $\psi \in C_c^{\infty}(\omega)$, $F \in C^{\infty}(\mathcal{Y})$ such that $\int_{\mathcal{Y}} F(y) \, \mathrm{d}y = 0$,

we have

$$\iint_{\omega \times \mathcal{Y}} B(\xi, y) : \Psi(x, y) \, \mathrm{d}y \, \mathrm{d}\xi = 0.$$

Then there exist unique $B \in L^2(\omega; \mathbb{R}^{3 \times 3}_{sym})$ and $w \in L^2(\omega; \dot{H}^1(\mathcal{Y}; \mathbb{R}^2))$ such that

$$B(\xi, y) = B(\xi) + \operatorname{sym} \nabla_y w(\xi, y)$$

In what follows, we will use the notation $\frac{osc,\gamma}{\gamma}$ introduced in the appendix. We prove the next proposition in local coordinates; of course, this is equivalent to performing computations on the level of the surface.

Proposition 4.2. Let (w^h) be a bounded sequence in $H^2(S; \mathbb{R}^3)$ such that

 $\frac{1}{h}q_{w^h}$

is bounded in $L^2(S; \mathbb{S})$. Then there exist $w_0 \in H^2(S)$, $w_1 \in L^2(S; \dot{H}^2(\mathcal{Y}; \mathbb{R}^3))$ and $B \in \dot{L}^2(S \times Y; \mathbb{S})$ such that, after passing to a subsequence,

(45)
$$\nabla_{tan} \nabla_{tan} w^h \xrightarrow{2,\gamma} \nabla_{tan} \nabla_{tan} w_0 + \sum_{i \le 3, j,k \le 2} \left(\partial_{y_j y_k}^2 w_{1,i}(x,y) \right) \tau^i \otimes \tau^j \otimes \tau^k$$

and

$$\frac{1}{h}q_{w^h} \xrightarrow{2,\gamma} B$$

Set $B_w = \int_Y B(\cdot, y) \, dy$. Then the following are true:

(i) If $h \gg \varepsilon^2$ then there exists a unique $v \in L^2(S; \dot{H}^1(\mathcal{Y}; \mathbb{R}^2))$ such that

$$B = B_w + \sum_{i,j=1,2} \left(\operatorname{sym} \nabla_y v \right)_{ij} \tau^i \otimes \tau^j.$$

(ii) If $h \sim \varepsilon^2$ and if we set $\lim_{h\to 0} \frac{\varepsilon(h)^2}{h} = \frac{1}{\gamma_1}$, then there exists a unique $v \in L^2(S; \dot{H}^1(\mathcal{Y}; \mathbb{R}^2))$ such that

$$B = B_w + \sum_{i,j=1,2} \left(\operatorname{sym} \nabla_y v \right)_{ij} \tau^i \otimes \tau^j + \frac{1}{\gamma_1} w_{1,3} \mathbf{S}$$

(iii) If $h \ll \varepsilon^2$, then there exists a unique $v \in L^2(S; \dot{H}^1(\mathcal{Y}; \mathbb{R}^2))$ such that

$$\sum_{i,j=1,2} \left(\operatorname{sym} \nabla_y v \right)_{ij} \tau^i \otimes \tau^j + (w_{1,3}) \mathbf{S} = 0.$$

Proof. The existence of w_0 and w_1 follows from Lemma 3.11. In local coordinates (45) can be expressed as

$$\nabla^2 \tilde{w}^h \xrightarrow{2,\gamma} \nabla^2 \tilde{w}_0 + \nabla^2_y \check{w}_1,$$

where $\tilde{w}^h = w^h \circ \psi$, $\tilde{w}_0 = w_0 \circ \psi$, $\tilde{w}_1 = \tilde{w}_{1,i}\tilde{\tau}^i$, $\tilde{w}_1 = w_1 \circ \psi$, $\tilde{\tau}^i = \tau^i \circ \psi$ where by slight abuse of notation we write $(w_{1,i} \circ \psi)(x, y) = w_{1,i}(\psi(x), y)$, for $(x, y) \in \omega \times \mathcal{Y}$. Denote also $\bar{w}^h_\alpha = \tilde{w}^h \cdot \partial_\alpha \psi$, $\bar{w}_{0,\alpha} = w_0 \cdot \partial_\alpha \psi$, for $\alpha = 1, 2$ and $\bar{w}_1 = (\check{w}_{1,1}, \check{w}_{1,2})$. By Lemma A.4 in the appendix we have

(46)
$$\frac{1}{\varepsilon} \operatorname{sym} \nabla \bar{w}^h \xrightarrow{\operatorname{osc},\gamma} \operatorname{sym} \nabla_y \bar{w}_1, \quad \frac{1}{\varepsilon} \operatorname{sym} \left((\nabla \psi)^T \nabla \tilde{w}^h_{\operatorname{tan}} \right) \xrightarrow{\operatorname{osc},\gamma} \operatorname{sym} \nabla_y \bar{w}_1.$$

and

(47)
$$\frac{\tilde{w}^h}{\varepsilon^2} \xrightarrow{osc,\gamma} \check{w}_1, \quad \frac{\bar{w}^h}{\varepsilon^2} \xrightarrow{osc,\gamma} \bar{w}_1, \quad \frac{\tilde{w}^h \cdot n}{\varepsilon^2} \xrightarrow{osc,\gamma} \tilde{w}_{1,3}$$

Now let $F \in C^{\infty}(\mathcal{Y})$ and $\varphi \in C_0^{\infty}(\omega)$. With (9) in mind, we compute:

$$\begin{split} &\int_{\omega} \operatorname{sym} \nabla \bar{w}^{h} : \left(\operatorname{cof} \nabla^{2} F \right) \left(\frac{\cdot}{\varepsilon} \right) \varphi \\ &= \varepsilon^{2} \int_{\omega} \operatorname{sym} \nabla \bar{w}^{h} : \operatorname{cof} \left[\nabla^{2} \left(F \left(\frac{\cdot}{\varepsilon} \right) \varphi \right) - \frac{2}{\varepsilon} (\nabla F) \left(\frac{\cdot}{\varepsilon} \right) \otimes \nabla \varphi - F \left(\frac{\cdot}{\varepsilon} \right) \nabla^{2} \varphi \right] \\ &= -\varepsilon^{2} \int_{\omega} \operatorname{sym} \nabla \bar{w}^{h} : \operatorname{cof} \left[\frac{2}{\varepsilon} (\nabla F) \left(\frac{\cdot}{\varepsilon} \right) \otimes \nabla \varphi + F \left(\frac{\cdot}{\varepsilon} \right) \nabla^{2} \varphi \right]. \end{split}$$

We used that the term sym $\nabla \tilde{w}^h$ is L^2 -orthogonal to test matrix fields of the form $\operatorname{cof} \nabla^2 F$. From this and from (9) and (47) we deduce that

Recall the identity

(49)
$$q_{w^h} \equiv \operatorname{sym}\left((\nabla\psi)^T \nabla \tilde{w}^h\right) = \operatorname{sym}\left((\nabla\psi)^T \nabla \tilde{w}^h_{\operatorname{tan}}\right) + (\tilde{w}^h \cdot n)\mathbf{S}$$

Assume first that $h \gg \varepsilon^2$. From the assumption $\frac{1}{h}q_{w^h} \xrightarrow{2,\gamma} B$ and (47), (48) and Lemma 4.1 we conclude that there exists $\check{v} \in L^2(\omega; \dot{H}^1(\mathcal{Y}; \mathbb{R}^3))$ such that

$$\operatorname{sym}\left((\nabla\psi)^T\nabla\tilde{w}^h\right)\xrightarrow{2,\gamma}\tilde{B}_w+\operatorname{sym}\nabla_y\check{v}$$

Here $\tilde{B}_w = B_w \circ \psi$. This is i) in local coordinates, after defining $v_i = (\check{v} \circ \psi^{-1}) \cdot \tau_i$ for i = 1, 2, 3. Case ii) we conclude as follows: By dividing the identity (49) by ε and using

$$\frac{q^h}{\varepsilon} \to 0 \text{ strongly in } L^2, \quad \text{when } h \ll \varepsilon,$$

as well as (46) and (47) we conclude that $\bar{w}_1 = 0$ from Korn's inequality. Using (48) and the identity (49), after dividing it by ε^2 , we conclude

$$\lim_{h \to 0} \int_{\omega} \frac{\operatorname{sym}\left((\nabla \psi)^T \nabla \tilde{w}^h\right)}{\varepsilon^2} : \left(\operatorname{cof} \nabla^2 F\right) \left(\frac{\cdot}{\varepsilon}\right) \varphi = \lim_{h \to 0} \int_{\omega} \left(\frac{\tilde{w}^h}{\varepsilon^2} \cdot n\right) \mathbf{S} : \left(\operatorname{cof} \nabla^2 F\right) \left(\frac{\cdot}{\varepsilon}\right) \varphi$$

$$= \int_{\omega \times Y} \tilde{w}_{1,3}(\cdot, y) \mathbf{S} : \left(\operatorname{cof} \nabla^2 F\right)(y) \varphi.$$
(50)

Lemma 4.1 again shows that there exists v such that

$$\frac{q^h}{h} \xrightarrow{2,\gamma} B_w + \sum_{i,j=1}^2 (\operatorname{sym} \nabla_y v)_{ij} \tau^i \otimes \tau^j + \left(\lim_{h \to 0} \frac{\varepsilon^2}{h} \right) w_{1,3} \mathbf{S}.$$

For the case (iii) we argue similarly: as in the case of ii) we conclude $\bar{w}_1 = 0$. Also we know that $q^h/\varepsilon^2 \to 0$ strongly in L^2 . Hence the left-hand side of (50) converges to zero, so

$$\int_{\omega \times Y} \tilde{w}_{1,3}(\cdot, y) \mathbf{S} : (\operatorname{cof} \nabla^2 F)(y) \varphi = 0,$$

which by Lemma 4.1 implies the claim.

Lemma 4.3. Let $(w^h)_{h>0} \subset H^1(S^1; \mathbb{R}^3)$ be such that

$$\limsup_{h \to 0} \left(\|w^h\|_{L^2} + \|\nabla_h w^h\|_{L^2} \right) < \infty.$$

Then there exists a map $w_0 \in H^1(S; \mathbb{R}^3)$ and a field $H_{\gamma} \in L(S \times I \times \mathcal{Y}; \mathbb{R}^{3 \times 3})$ such that

$$(51) \quad H_{\gamma} = \begin{cases} \sum_{i,j=1}^{3} \left(\nabla_{y} w_{1}, w_{2} \right)_{ij} \tau^{i} \otimes \tau^{j} & \text{for some } \begin{cases} w_{1} \in L^{2}(S; \dot{H}^{1}(\mathcal{Y}; \mathbb{R}^{3})) \\ w_{2} \in L^{2}(S \times Y \times I; \mathbb{R}^{3})) \end{cases} \\ if \gamma = 0, \\ \sum_{i,j=1}^{3} \left(\nabla_{y} w_{1}, \frac{1}{\gamma} \partial_{3} w_{1} \right)_{ij} \tau^{i} \otimes \tau^{j} & \text{for some } w_{1} \in L^{2}(S, \dot{H}^{1}(I \times \mathcal{Y}; \mathbb{R}^{3})) \\ if \gamma \in (0, \infty), \\ \sum_{i,j=1}^{3} \left(\nabla_{y} w_{1}, w_{2} \right)_{ij} \tau^{i} \otimes \tau^{j} & \text{for some } \begin{cases} w_{1} \in L^{2}(S \times I; \dot{H}^{1}(\mathcal{Y}; \mathbb{R}^{3})) \\ w_{2} \in L^{2}(S \times I; \mathcal{H}^{1}(\mathcal{Y}; \mathbb{R}^{3})) \\ w_{2} \in L^{2}(S \times I; \mathbb{R}^{3})) \end{cases} \\ if \gamma = \infty, \end{cases}$$

such that, up to a subsequence, $w^h \to w_0$ in L^2 and

$$\nabla_h w^h \xrightarrow{2,\gamma} \nabla_{tan} w_0 T_S + H_{\gamma}$$
 weakly two-scale in L^2 .

Here, w_0 is the weak limit in $H^1(S)$ of $\int_I w^h(x + tn(x))dt$.

Proof. The lemma is an analogue of Proposition 6.3.5 in [Neu10], adapted to the manifold S and the definition of two scale convergence on the manifold. Thus we will only prove the case $\gamma \in (0, \infty)$. Since the sequence is bounded in H^1 norm there exists a weak limit $w_0 \in H^1(S^1; \mathbb{R}^3)$. Let us denote by \tilde{w}^h , \tilde{w}_0 the elements of $H^1(\Omega; \mathbb{R}^3)$, defined by:

(52)
$$\tilde{w}^h = w^h \circ r_e^{-1}, \quad \tilde{w}_0 = w_0 \circ r_e^{-1}.$$

By Proposition 6.3.5 in [Neu10], which is proved for planar domains, \tilde{w}_0 does not depend on t and there exists $\tilde{w}_1 \in L^2(\omega; \dot{H}^1(I \times \mathcal{Y}; \mathbb{R}^3))$ such that:

(53)
$$\nabla_h \tilde{w}^h \xrightarrow{2,\gamma} (\nabla_{\xi} \tilde{w}_0, 0) + (\nabla_y \check{w}_1, \frac{1}{\gamma} \partial_t \check{w}_1).$$

Then we have that $\check{w}_1 \in L^2(S; \dot{H}^1(I \times \mathcal{Y}; \mathbb{R}^3))$. Using (5) and (35) we conclude

(54)
$$\nabla_h w^h = \nabla w^h (I + t\mathbf{S})(I_h + th\mathbf{S})^{-1}$$
$$= \nabla \tilde{w}^h [\tau^1, \tau^2, \tau^3]^T (I + t\mathbf{S})^{-1} (I + t\mathbf{S})(I_h + th\mathbf{S})^{-1}$$
$$= \nabla_h \tilde{w}^h [\tau^1, \tau^2, \tau^3]^T (I + th\mathbf{S})^{-1}.$$

By using (53) we conclude that

(55)
$$\nabla_h w^h \xrightarrow{2,\gamma} ((\nabla_\xi \tilde{w}_0) \circ r, 0) [\tau^1, \tau^2, \tau^3]^T + (\nabla_y \breve{w}_1 \circ r_e, \frac{1}{\gamma} \partial_t \breve{w}_1 \circ r_e) [\tau^1, \tau^2, \tau^3]^T$$

$$= \nabla w_0 T_S + \sum_{i,j=1}^3 (\nabla_y w_1, \frac{1}{\gamma} \partial_t w_1)_{ij} \tau^i \otimes \tau^j,$$

where $(w_1)_i = (\breve{w}_1 \circ r_e) \cdot \tau_i$. The last property follows from the fact that \tilde{w}_0 does not depend on t.

The following lemma is fairly straightforward; we refer to [Neu10, Corollary 2.3.4] for a proof.

Lemma 4.4. Let $(E^h_{app}) \subset L^2(\Omega; \mathbb{R}^{3 \times 3})$ be such that

$$E^h_{app} \xrightarrow{2,\gamma} E_{app} \text{ in } L^2(\Omega \times \mathcal{Y}; \mathbb{R}^{3 \times 3}).$$

Then

$$h^{-2}\left(\sqrt{(I+h^2 E^h_{app})^T (I+h^2 E^h_{app})} - I\right) \xrightarrow{2,\gamma} \operatorname{sym} E_{app} \quad in \ L^2(\Omega \times \mathcal{Y}; \mathbb{R}^{3\times 3})$$

Proposition 4.5. Assume that there is $B \in L^2(S \times \mathcal{Y}; \mathbb{S})$ such that

$$\frac{1}{h}q_{V_s^h} \xrightarrow{2,\gamma} B \text{ in } L^2(S;\mathbb{S}),$$

and assume that there is $B_w \in L^2(S; \mathbb{S})$ such that

$$\frac{1}{h}q_{V^h} \rightharpoonup B_w$$
, weakly in $L^2(S; \mathbb{S})$.

Assume also that

$$\nabla_{tan} \nabla_{tan} (V_s^h \cdot n) \xrightarrow{2,\gamma} \nabla_{tan} \nabla_{tan} (V \cdot n) + \sum_{i,j=1,2} (\partial_{y_i y_j}^2 \varphi) \tau^i \otimes \tau^j,$$

and set

(56)
$$E^{h} = \frac{\sqrt{(\nabla_{h}y^{h})^{T}\nabla_{h}y^{h}} - I}{h^{2}}.$$

Then there exist

$$U_{\gamma} \in L^2(S; L_{\gamma}(I \times \mathcal{Y}))$$

such that (after passing to a subsequence)

$$E^h \xrightarrow{2,\gamma} E \text{ in } L^2(S^1; \mathbb{R}^{3 \times 3}),$$

where E is given by

(57)
$$E = \left(B_w + \dot{B} + \frac{1}{2}(dV)^2 - tb_V\right)(T_S, T_S) - t\sum_{i,j=1}^2 (\partial_{y_i y_j}^2 \varphi)\tau^i \otimes \tau^j + U_\gamma,$$

and where $\dot{B} = B - \int_Y B(\cdot, y) \, dy$. In particular, the following are true:

(i) If $\gamma \in (0, \infty]$ then there exists $U_{\gamma} \in L^2(S; L_{\gamma}(I \times \mathcal{Y}))$ such that (58) $E = B_w + \frac{1}{2} (dV)^2 - tb_V + U_{\gamma}.$

(ii) If $\varepsilon^2 \ll h \ll \varepsilon$, there exists $U \in L^2(S; L^0_0(I \times \mathcal{Y}))$ such that

(59)
$$E = B_w + \frac{1}{2}(dV)^2 - tb_V + U.$$

(iii) If $h \sim \varepsilon^2$, with $\lim_{h\to 0} \frac{\varepsilon^2}{h} = \frac{1}{\gamma_1} \in (0,\infty)$, there exists $U \in L^2(S; L^1_{0,\gamma_1}(I \times \mathcal{Y}))$ such that

(60)
$$E = B_w + \frac{1}{2} (dV)^2 - tb_V + U.$$

Proof. By Lemma 3.11, there exists a subsequence such that $E^h \xrightarrow{2,\gamma} E$ for some E. Denote by E_{app}^h the approximate strain

(61)
$$E_{\text{app}}^h := \frac{(R^h)^T \nabla_h y^h - I}{h^2}$$

By Lemma 4.4 it is enough to identify the two-scale limit of sym E_{app}^{h} . Let us write

(62)
$$R^{h}E^{h}_{\text{app}} = \frac{\nabla_{h}y^{h} - I}{h^{2}} - \frac{R^{h} - I}{h^{2}}.$$

We have sym $(R^h E^h_{app}) \xrightarrow{2,\gamma} E$, because $R^h \to I$ boundedly in measure. By property (vi) of Lemma 3.7 the symmetric part of the second term converges strongly in L^2 (and thus two-scale) to $\Omega_V^2/2$. So we need to identify the two scale limit of

$$\operatorname{sym}\left(\frac{\nabla_h y^h - I}{h^2}\right)$$

For brevity we set $\mu_s^h = \mu_{V_s^h}$. As usual, we extend V_s^h , n and μ_s^h trivially to S^h . In what follows we abuse notation using t also as an independent variable. We define the maps $z^h : S^h \to \mathbb{R}^3$ by setting

$$z^{h}(x) = x + h\left(V_{s}^{h}(x) + t(x)\mu_{s}^{h}(x)\right) \text{ for all } x \in S^{h}.$$

Define Q(x) as in (19) and define (compare (16))

$$b^{h}(x) = b_{V_{s}^{h}}(x) \equiv -\nabla_{tan}\mu_{s}^{h}(x)T_{S}(x) + \nabla_{tan}V_{s}^{h}(x)\mathbf{S}(x)$$

and (compare (14))

$$\Omega^h(x) = \Omega_{V_s^h}(x) \equiv \nabla_{tan} V_s^h(x) T_S(x) + \mu_s^h(x) \otimes n(x).$$

Then Lemma 2.4 shows that

(63)
$$\nabla z^{h} = I + h\Omega^{h} - ht b^{h} - ht^{2} \nabla_{tan} \mu_{s}^{h} \mathbf{S} + h \left(\nabla_{tan} V_{s}^{h} + t \nabla_{tan} \mu_{s}^{h} \right) T_{S} Q.$$

Note that $|Q| \leq Ct^2 \leq Ch^2$ on S^h , so

$$\|Q\|_{L^2(S^h)} \le Ch^{5/2}.$$

In what follows $\Theta^h \in L^2(S^h)$ denote maps which may change from expression to expression, but which always satisfy

$$\|\Theta^h\|_{L^2(S^h)} \le Ch^{5/2}$$

We see from (63) that

$$\nabla z^h = I + h\Omega^h + \Theta^h = R^h + \Theta^h,$$

by (41). On the other hand, Lemma 3.7 shows that $\nabla u^h = R^h + \Theta^h$. Hence

(64)
$$\|\nabla u^h - \nabla z^h\|_{L^2(S^h)} \le Ch^{5/2}.$$

However, by the definition of V^h and of z^h we have, for $x \in S$,

$$\frac{1}{h} \int_{I^h} z^h(x + tn(x)) - u^h(x + tn(x)) \, dt = h\left(V^h_s(x) - V^h(x)\right).$$

Hence Poincaré's inequality implies that

$$\begin{aligned} \|u^{h} - z^{h}\|_{L^{2}(S^{h})} &\leq \left\|u^{h} - z^{h} - h\left(V_{s}^{h} - V^{h}\right)\right\|_{L^{2}(S^{h})} + h\|V_{s}^{h} - V^{h}\|_{L^{2}(S^{h})} \\ &\leq \|\nabla u^{h} - \nabla z^{h}\|_{L^{2}(S^{h})} + h^{3/2}\|V_{s}^{h} - V^{h}\|_{L^{2}(S)} \leq Ch^{5/2}, \end{aligned}$$

by (64) and (37). Defining $Z^h: S^1 \to \mathbb{R}^3$ by setting $Z^h(\Phi^h) = z^h$ on S^h , we have the equivalent bounds

$$\|y^h - Z^h\|_{L^2(S^1)} + \|\nabla_h(y^h - Z^h)\|_{L^2(S^1)} \le Ch^2.$$

Thus, using Lemma 4.3, we conclude that there exists

$$H_{\gamma} \in L^2(S^1 \times \mathcal{Y}; R^{3 \times 3})$$

of the form given in Lemma 4.3 and $c \in H^1(S; \mathbb{R}^3)$ such that (after passing to a subsequence)

(65)
$$\frac{1}{h^2} \nabla_h \left(y^h - Z^h \right) \xrightarrow{2,\gamma} \nabla_{tan} c T_S + H_{\gamma}.$$

Here c is a weak limit in $H^1(S; \mathbb{R}^3)$ of $\frac{V^h - V_s^h}{h}$.

We will now identify the two scale limit on S^1 of the quantity sym $\left(\frac{\nabla_h Z^h - I}{h^2}\right)$. By (63) we have for all $x \in S^h$:

(66)
$$\frac{\nabla z^h(x) - I}{h^2} = \frac{1}{h} \Omega^h(x) - \frac{t(x)}{h} b^h(x) + M^h(x),$$

where

$$\|M^h\|_{L^2(S^h)} \le Ch^{3/2}.$$

We must therefore identify the two-scale limits on S of the first two terms. Lemma 2.2 implies

(67)
$$\frac{1}{h}\operatorname{sym}\Omega^{h} = \frac{1}{h}q_{V_{s}^{h}}(T_{S}, T_{S}) \xrightarrow{2, \gamma} B(T_{S}, T_{S})$$

weakly two-scale in $L^2(S; \mathbb{R}^{3 \times 3})$, by definition of *B*.

It remains to identify the two-scale limit of b^h on S. Its weak L^2 -limit clearly is b_V . This follows by comparing the definition of b^h with (16). Next note that, since $V_s^h \to V$ strongly in $H^1(S)$, we know that $\nabla_{tan} V_s^h \mathbf{S}$ does not contribute to the oscillating part. On the other hand, we have from (13):

$$\nabla_{tan}\mu^{h} = -\nabla_{tan}\nabla_{tan}\left(n\cdot V_{s}^{h}\right) + \nabla_{tan}\left(V_{s}^{h}\cdot\mathbf{S}\right).$$

The last term converges strongly in $L^2(S)$ to $\nabla_{tan} (V \cdot \mathbf{S})$, so it does not contribute to the oscillating part. The contribution of the term $\nabla_{tan} \nabla_{tan} (n \cdot V_s^h)$ is given by the assumption.

We conclude that

(68)
$$b^{h} \xrightarrow{2,\gamma} b_{V} + \partial^{2}_{y_{i}y_{j}} \varphi \ \tau^{i} \otimes \tau^{j}$$

on S. By (66), the above convergence results on S imply that

(69)
$$\operatorname{sym}\left(\frac{\nabla_h Z^h - I}{h^2}\right) \xrightarrow{2,\gamma} (B - tb_V) (T_S, T_S) - t(\partial_{y_i y_j}^2 \varphi) \tau^i \otimes \tau^j$$

weakly two-scale on S^1 .

We conclude from (65) and (69) that

$$E = B + \operatorname{sym}\left(\nabla_{\tan} c \, T_S\right) - \frac{1}{2}(\Omega_V^2) - t \sum_{i,j=1}^2 (\partial_{y_i y_j}^2 \varphi) \tau^i \otimes \tau^j - t b_V + \tilde{U}_{\gamma}$$

for some $\tilde{U}_{\gamma} \in L^2(S; L_{\gamma}(I \times \mathcal{Y}))$. Notice that

(70)
$$B + \operatorname{sym}\left(\nabla_{\operatorname{tan}} c \, T_S\right) = B_w + \dot{B} + \ddot{U}_{\gamma}$$

where

$$\bar{U}_{\gamma} := \frac{1}{2} \sum_{i=1}^{2} (\nabla c \tau_i, n) (n \otimes \tau^i + \tau^i \otimes n).$$

Notice that $\bar{U}_{\gamma} \in L^2(S; L_{\gamma}(I \times \mathcal{Y})).$

Define

$$U_{\gamma} = \tilde{U}_{\gamma} + \bar{U}_{\gamma} \in L^2(S; L_{\gamma}(I \times \mathcal{Y})).$$

Hence (57) is proven, using the identity (15). The remaining claims now follow from Proposition 4.2. Namely, ii) and iii) are direct and i) is the consequence of the identity

$$\begin{pmatrix} t(\partial_{y_iy_j}\varphi)_{i,j=1}^2 & 0\\ 0 & 0 \end{pmatrix} = \operatorname{sym}(\nabla_y, \frac{1}{\gamma}\partial_3) \begin{pmatrix} t\partial_{y_1}\varphi\\ t\partial_{y_2}\varphi\\ \frac{1}{\gamma}\varphi \end{pmatrix} \in L_{\gamma}(I \times \mathcal{Y}),$$

for $\gamma \in (0, \infty)$.

For lower bound we need the following lemma.

Lemma 4.6. Let $(y^h) \subset H^1(S^1; \mathbb{R}^3)$, define $E^h : S^1 \to \mathbb{R}^{3 \times 3}$ by

$$\sqrt{(\nabla_h y^h)^T (\nabla_h y^h)} = I + h^2 E^h,$$

and assume that $E^h \xrightarrow{2,\gamma}: E$. Then we have

$$\begin{split} \liminf_{h \to 0} &\int_{S} \int_{I} \mathcal{Q}(x + tn(x), r(x + tn(x)) / \varepsilon, E^{h}(x + tn(x))) \, \mathrm{d}t \, \mathrm{d}\mathcal{H}^{2} \geq \\ &\int_{S} \int_{I} \int_{\mathcal{Y}} \mathcal{Q}(x + tn(x), y, E(x + tn(x), y)) \, \mathrm{d}y \, \mathrm{d}t \, \mathrm{d}\mathcal{H}^{2} \end{split}$$

and

$$\begin{split} \liminf_{h\to 0} \frac{1}{h^4} \int_S \int_I W(x+tn(x), r(x+tn(x))/\varepsilon, I+h^2 E^h(x+tn(x))) \, \mathrm{d}t \, \mathrm{d}\mathcal{H}^2 \geq \\ \int_S \int_I \int_{\mathcal{Y}} \mathcal{Q}(x+tn(x), y, E(x+tn(x), y)) \, \mathrm{d}y \, \mathrm{d}t \, d\mathcal{H}^2. \end{split}$$

Proof. For the first claim we refer to [Vis06, Vis07]. The second claim then follows from the standard truncation argument. \Box

The lower bound parts of Theorem 3.6 and Theorem 6.3 is now a direct consequence of Proposition 4.5 and of Lemma 4.6.

5 Upper bound

We start with the following observation.

Remark 5. It is easy to see, by using Korn's inequality, that $L_{\gamma}(I \times \mathcal{Y}, \mathbb{R}^{3 \times 3}_{\text{sym}})$ as well as $L_{0}^{0}(I \times \mathcal{Y}, \mathbb{R}^{3 \times 3}_{\text{sym}}), L_{0,\gamma_{1}}^{1}(I \times \mathcal{Y}, \mathbb{R}^{3 \times 3}_{\text{sym}})$ for $\gamma \in [0, \infty]$ and $\gamma_{1} \in (0, \infty)$ are closed subspaces of $L^{2}(I \times \mathcal{Y}, \mathbb{R}^{3 \times 3}_{\text{sym}})$. Also by using Korn's inequality it is easy to see (see also [Neu12, Neu10, NV]) that the following coercivity bounds are satisfied:

$$\begin{aligned} \|\zeta\|_{H^1}^2 + \|g\|_{L^2}^2 &\lesssim \quad \|\mathcal{U}_0(\zeta,g)\|_{L^2}^2, \\ \forall \zeta \in \dot{H}^1(\mathcal{Y};\mathbb{R}^2), g \in L^2(I \times \mathcal{Y};\mathbb{R}^3), \end{aligned}$$

$$\begin{split} \|\zeta\|_{H^1}^2 + \|\varphi\|_{H^2} + \|g\|_{L^2}^2 &\lesssim \quad \|\mathcal{U}_0^0(\zeta,\varphi,g)\|_{L^2}^2, \\ \forall \zeta \in \dot{H}^1(\mathcal{Y};\mathbb{R}^2), \varphi \in \dot{H}^2(\mathcal{Y}), g \in L^2(I \times \mathcal{Y};\mathbb{R}^3) \text{ and} \end{split}$$

$$\begin{aligned} \|\zeta\|_{H^1}^2 + \|\varphi\|_{H^2} + \|g\|_{L^2}^2 &\lesssim \quad \|\mathcal{U}_{0,\gamma_1}^1(\zeta,\varphi,g)\|_{L^2}^2, \\ \forall \zeta \in \dot{H}^1(\mathcal{Y};\mathbb{R}^2), \varphi \in \dot{H}^2(\mathcal{Y}), g \in L^2(I \times \mathcal{Y};\mathbb{R}^3) \text{ and } \gamma_1 \in (0,\infty) \end{aligned}$$

$$\begin{aligned} \|\zeta\|_{H^1}^2 + \|\psi\|_{H^1} + \|c\|_{L^2}^2 &\lesssim \quad \|\mathcal{U}_{\infty}(\zeta,\psi,c)\|_{L^2}^2, \\ \forall \zeta \in \dot{H}^1(\mathcal{Y};\mathbb{R}^2), \psi \in L^2(I;\dot{H}^1(\mathcal{Y})), c \in L^2(I;\mathbb{R}^3), \end{aligned}$$

$$\|\phi\|_{H^1}^2 \leq C(\gamma) \|\mathcal{U}_{\gamma}(\phi)\|_{L^2}^2, \ \forall \phi \in \dot{H}^1(I \times \mathcal{Y}; \mathbb{R}^3).$$

Here the constant absorbed into the symbol \lesssim depends on η_1, η_2 .

The following two lemmas and remark are analogous to [NV, Lemma 2.10, 2.11].

Lemma 5.1. For $\gamma \in (0, \infty]$ there exists a bounded linear operator

$$\Pi_{\gamma}: L^2(S, \mathbb{S}) \times L^2(S, \mathbb{S}) \to L^2(S, L_{\gamma}(I \times \mathcal{Y}))$$

such that for almost every $x \in S$ we have

$$\mathcal{Q}_{\gamma}(x,q^1,q^2) = \int_I \int_{\mathcal{Y}} \mathcal{Q}\left(x + tn(x), y, q^1 + tq^2 + \Pi_{\gamma}[q^1,q^2](x,t,y)\right) \mathrm{d}y \,\mathrm{d}t.$$

Moreover, if $q^1, q^2 \in C(S, \mathbb{S})$ then

$$(x,t,y) \mapsto \Pi_{\gamma}[q^1,q^2](x,t,y)$$

is continuous as well.

Lemma 5.2. The function $Q_{\gamma}: S \times \mathbb{S} \to \mathbb{R}^+$ is continuous. Moreover for any $x \in S$ the function $Q_{\gamma}(x): \mathbb{S}(x)_{\text{sym}} \times \mathbb{S}(x)_{\text{sym}} \to \mathbb{R}$ satisfies

$$\begin{aligned} (|q^1|^2 + |q^2|^2) &\lesssim & \mathcal{Q}_{\gamma}(x, q^1, q^2) = \mathcal{Q}_{\gamma}(x, q^1, q^2) \\ &\lesssim & |q^1|^2 + |q^2|^2. \end{aligned}$$

The constant in \lesssim depends only on $\alpha, \beta, \eta_1, \eta_2$.

Lemma 5.1 and Lemma 5.2 remain true for the quadratic forms \mathcal{Q}_0^0 , $\mathcal{Q}_0^{1,\gamma_1}$, the spaces $L_0^0(I \times \mathcal{Y})$, $L_{0,\gamma_1}^1(I \times \mathcal{Y})$, and the appropriate operators

 $\Pi^0_0: L^2(S, \mathbb{S}) \times L^2(S, \mathbb{S}) \to L^2(S, L^0_0(I \times \mathcal{Y}))$

and

$$\Pi^1_{0,\gamma_1}: L^2(S,\mathbb{S}) \times L^2(S,\mathbb{S}) \to L^2(S,L^1_{0,\gamma_1}(I \times \mathcal{Y}))$$

for $\gamma_1 \in (0, \infty)$.

We need the following auxiliary result concerning the linearization of the square root of a matrix. Its proof is straighforward by Taylor expansion.

Lemma 5.3. There exists $\eta > 0$ and a nondecreasing function $m : (0, \eta) \to \mathbb{R}^+$ such that $m(\delta) \to 0$ as $\delta \to 0$ such that the following is true: Let $G^h \in L^2(S^1, \mathbb{R}^{3 \times 3})$ and $K^h \in L^4(S^1, \mathbb{R}^{3 \times 3})$ satisfy the following conditions:

(i) There exists $M \in \mathbb{R}$ such that

$$\limsup_{h \to 0} \left(\|\operatorname{sym} G^h\|_{L^2(\Omega)} + \|K^h\|_{L^4(\Omega)} \right) \le M.$$

- (ii) We have $h^{-1} \operatorname{sym} K^h \to 0$ strongly in $L^2(\Omega)$ as $h \to 0$.
- (iii) There exists $\delta < \eta$ such that

$$\limsup_{h \to 0} h \| K^h \|_{L^{\infty}} \le \delta$$

and

$$\limsup_{h \to 0} h^2 \|G^h\|_{L^{\infty}} \le \delta.$$

(iv) We have $hG^h \to 0$ strongly in $L^4(\Omega)$ as $h \to 0$.

Set

$$E^{h} = \frac{1}{h^{2}} \left(\sqrt{(I + hK^{h} + h^{2}G^{h})^{t}(I + hK^{h} + h^{2}G^{h})} - I \right)$$

and

(71)
$$E^{h}_{app} = \operatorname{sym} G^{h} - \frac{1}{2} (K^{h})^{2}.$$

Then we have

(72)
$$\limsup_{h \to 0} \|E^h - E^h_{app}\|_{L^2} \le M^2 m(\delta),$$

and

(73)

$$\lim_{h \to 0} \sup_{h \to 0} \left| \frac{1}{h^4} \int_S \int_I W(x + tn(x), r(x + tn(x))/\varepsilon, I + h^2 E^h(x + tn(x))) \, \mathrm{d}t \, \mathrm{d}\mathcal{H}^2 - \int_S \int_I \mathcal{Q}(x + tn(x), r(x + tn(x))/\varepsilon, E^h_{app}(x + tn(x)) \, \mathrm{d}t \, \mathrm{d}\mathcal{H}^2 \right| \le (M+1)^4 m(\delta)$$

If, moreover, $E^h_{app} \xrightarrow{2,\gamma} E(x,y)$ strongly two scale, then

(74)
$$\limsup_{h \to 0} \left| \frac{1}{h^4} \int_S \int_I W(x + tn(x), r(x + tn(x))/\varepsilon, I + h^2 E^h(x + tn(x))) \, \mathrm{d}t \, \mathrm{d}\mathcal{H}^2 - \int_S \int_I \int_{\mathcal{Y}} \mathcal{Q}(x + tn(x), y, E(x + tn(x), y)) \, \mathrm{d}y \, \mathrm{d}t \, \mathrm{d}\mathcal{H}^2 \, \mathrm{d}x \right| \le (M + 1)^4 m(\delta).$$

Proof. We will just give the sketch of the proof. By Taylor expansion there exists $\eta_1 > 0$ and nondecreasing function $m_1 : (0, \eta_1) \to \mathbb{R}^+$ such that $m_1(\delta) \to 0$ as $\delta \to 0$ and for every $A \in \mathbb{R}^{3\times 3}$ which satisfies $|A - I| < \eta_1$ we have

(75)
$$\left|\sqrt{(I+A^t)(I+A)} - \left(I + \operatorname{sym} A + \frac{1}{2}A^t A\right)\right| \le m_1(|A-I|)\left(\operatorname{sym} A + \frac{1}{2}A^T A\right).$$

If we plug into this identity $A = hK^h + h^2G^h$, divide by h^2 and integrate and let $h \to 0$ we obtain

$$\limsup_{h \to 0} \|E^h - \tilde{E}^h_{app}\|_{L^2} \le m_1(\delta) \|\tilde{E}^h_{app}\|_{L^2},$$

where

$$\tilde{E}^{h}_{app} = \frac{\operatorname{sym} K^{h}}{h} + \operatorname{sym} G^{h} + \frac{1}{2} (K^{h})^{t} K^{h} + h \operatorname{sym} \left((G^{h})^{t} K^{h} \right) + \frac{1}{2} h^{2} (G^{h})^{t} G^{h}.$$

Using the assumptions it is easy to prove that

$$\lim_{h \to 0} \|\tilde{E}^h_{app} - E^h_{app}\|_{L^2} \to 0.$$

Now (72) immediately follows. (73) follows from Lemma 3.4 and triangular inequality. (73) follows from the fact that

$$\begin{split} &\int_{S} \int_{I} \mathcal{Q}(x+tn(x),r(x+tn(x))/\varepsilon,E^{h}_{app}(x+tn(x))\,\mathrm{d}t\,\mathrm{d}\mathcal{H}^{2} \rightarrow \\ &\int_{S} \int_{I} \int_{\mathcal{Y}} \mathcal{Q}(x+tn(x),y,E(x+tn(x),y))\,\mathrm{d}y\,\mathrm{d}t\,\mathrm{d}\mathcal{H}^{2}\,\mathrm{d}x, \end{split}$$

as $h \to 0$ which is the consequence of the continuity of the integral functionals with respect to strong two scale convergence, see [Vis06, Vis07].

We give here a general computation that will be needed in the proof of the next proposition. Let $P \in C^1(S^1; C^1(\mathcal{Y}; \mathbb{R}^3))$ define $P^h : S^1 \to \mathbb{R}^3$ by $P^h = P(\cdot, r/\varepsilon)$, where, as usual, r is extended trivially from S to S^1 . Then by (35)

$$\begin{aligned} \nabla_h P^h &= \frac{1}{h} \partial_n P^h \otimes n + \nabla P^h T_S (I + t\mathbf{S}) (I + ht\mathbf{S})^{-1} T_S \\ &= \frac{1}{h} \partial_n P(\cdot, r/\varepsilon) \otimes n \\ &+ \left(\nabla P(\cdot, r/\varepsilon) + \frac{1}{\varepsilon} \nabla_y P(\cdot, r/\varepsilon) \nabla_{tan} r \ T_S (I + t\mathbf{S})^{-1} \right) (I + t\mathbf{S}) (I + ht\mathbf{S})^{-1} T_S, \end{aligned}$$

because having extended r trivially to S^1 , we have $\partial_n r = 0$ and (8) applies. We use the notation $\partial_n P$ to denote the *n*-derivative with respect to the first argument only. Since $(I + ht\mathbf{S})^{-1}$ agrees with I up to a term that on S^1 is uniformly bounded by h, and since $h \leq C\varepsilon$, we conclude that

(76)
$$\left\| \nabla_h P^h - \frac{1}{h} \partial_n P(\cdot, r/\varepsilon) \otimes n - \frac{1}{\varepsilon} \nabla_y P(\cdot, r/\varepsilon) \nabla_{tan} r T_S \right\|_{L^{\infty}(S^1)} \le C$$

Note that the linear operator $\nabla_{tan} rT_S$ on the tangent space can be expressed as, see (6),

$$\nabla_{tan} r T_S = e_1 \otimes \tau^1 + e_2 \otimes \tau^2,$$

which is just the pullback operator ψ^* from the tangent space to \mathbb{R}^2 .

Proposition 5.4. For every $B_w \in \mathcal{B}$ and for every infinitesimal bending $V \in H^2(S; \mathbb{R}^{3\times 3})$ of S, there exists a sequence $(y^h) \subset H^1(S^1, \mathbb{R}^3)$ satisfying the following:

(i)
$$y^h \to \pi$$
 strongly in $H^1(S^1; \mathbb{R}^3)$.

(ii) The maps $V^h(x) = h^{-1} \int_I y^h(x + tn(x)) dt - x$ satisfy

$$V^h \to V$$
 strongly in $H^1(S; \mathbb{R}^3)$

and

$$\frac{1}{h}q_{V^h} \rightarrow B_w$$
 weakly in $L^2(S; \mathbb{S})$.

(iii) We have

$$\lim_{h \to 0} h^{-4} I^h(y^h) = \begin{cases} I_{\gamma}(V, B_w) & \text{if } \lim h/\varepsilon = \gamma \in (0, \infty];\\ I_0^0(V, B_w) & \text{if } \varepsilon \gg h \gg \varepsilon^2;\\ I_{0,\gamma_1}^1(V, B_w) & \text{if } h \sim \varepsilon^2 \text{ with } \lim \varepsilon^2/h = 1/\gamma_1. \end{cases}$$

Proof. The proof is a modification of the proof of the recovery sequence in [LMP10], cf. also [FJM06].

By definition and by density, since $B_w \in \mathcal{B}$, there exist $w_n \in C_0^{\infty}(S; \mathbb{R}^3)$ such that $q_{w_n} \to B_w$ strongly in $L^2(S)$. Hence

$$\limsup_{n \to \infty} \limsup_{h \to 0} \left(\|q_{w_n} - B_w\|_{L^2(S;\mathbb{S})} + h \|w_n\|_{W^{2,\infty}(S;\mathbb{R}^3)} \right) = 0.$$

Lemma A.6 yields a sequence n_h with $n_h \to \infty$ as $h \to 0$, such that the maps

$$w^n = w_{n_h}$$

satisfy

(77)
$$\limsup_{h \to 0} h \|w^h\|_{W^{2,\infty}(S;\mathbb{R}^3)} = 0.$$

and

(78)
$$q_{w^h} \to B_w$$
 strongly in $L^2(S; \mathbb{S})$.

In order to have common proof for all cases (see the case $\gamma = 0$) we will assume that there exists a constant $M_1 > 0$ such that and double index sequence $q_{w^{\delta,h}}$ such that

(79)
$$||q_{w^{\delta,h}}||_{L^2} \le M_1,$$

and for all $\delta > 0$ we have

(80)
$$\limsup_{h \to 0} h \|\nabla w^{\delta,h}\|_{L^4} = 0, \ \limsup_{h \to 0} h \|\nabla^2 w^{\delta,h}\|_{L^2} \le M_1;$$
$$\limsup_{h \to 0} h^2 \|\nabla w^{\delta,h}\|_{L^{\infty}} = 0, \ \limsup_{h \to 0} h^3 \|\nabla^2 w^{\delta,h}\|_{L^{\infty}} = 0.$$

and

(81)
$$q_{w^{\delta,h}} \rightharpoonup B_w$$
, weakly in L^2

Now let $V \in H^2(S, \mathbb{R}^3)$ be an infinitesimal bending of S. We approximate V by a sequence $v^{\delta,h} \in W^{2,\infty}(S; \mathbb{R}^3)$ such that, for each $\delta > 0$ we have:

$$\limsup_{h \to 0} \|v^{\delta,h} - V\|_{H^2(S)} = 0, \quad \limsup_{h \to 0} h \|v^{\delta,h}\|_{W^{2,\infty}(S)} \le \delta$$

(82)

$$\lim_{h \to 0} \frac{1}{h^2} \mathcal{H}^2\left(\left\{x \in S : v^{\delta,h}(x) \neq V(x)\right\}\right) = 0.$$

The existence of such $v^{\delta,h}$ follows by Proposition A.5 in the appendix. We claim that

(83)
$$\frac{q_{v^{\delta,h}}}{h} \to 0 \text{ strongly in } L^2(S)$$

as $h \to 0$. In fact,

$$(84) \quad \frac{1}{h} \|q_{v^{\delta,h}}\|_{L^{2}(S)} \leq \frac{1}{h} \left| \{x \in S; v^{\delta,h}(x) \neq V(x)\} \right|^{1/2} \cdot \|\nabla_{tan} v^{\delta,h}\|_{L^{\infty}(\{v^{\delta,h} \neq V\})},$$

and this converges to zero by (82) and because $q_{v^{\delta,h}}$ is uniformly bounded in $L^{\infty}(S)$ (see below). From this we deduce (83).

To see that $q_{v^{\delta,h}}$ is uniformly bounded in $L^{\infty}(S)$, note that the Lipschitz constants of all $q_{v^{\delta,h}}$ are bounded by δ/h . Since $q_{v^{\delta,h}} = 0$ almost everywhere on $\{v^{\delta,h} = V\}$, we have

$$|q_{v^{\delta,h}}(x)| \le C\frac{1}{h}\operatorname{dist}\left(x, \{v^{\delta,h}=V\}\right) \le C.$$

The last estimate is true because (due to (82) and bounded curvature of S) for small h the set $\{v^{\delta,h} \neq V\}$ cannot contain a disk of radius h.

Now let
$$o^{\delta}, p^{\delta} \in C^1(S^1; C^1(\mathcal{Y}; \mathbb{R}^3))$$
 and set $p^{\delta,h} = p^{\delta}(\cdot, r/\varepsilon)$ and $o^{\delta,h} = o^{\delta}(\cdot, r/\varepsilon)$.
We define $z^{\delta,h} : S^h \to \mathbb{R}^3$ by

$$z^{\delta,h} = id + h\left(v^{\delta,h} + hw^{\delta,h} + t\mu_{v^{\delta,h} + hw^{\delta,h}}\right) + h^3 p^{\delta,h}(\Phi^h) + \varepsilon h^2 o^{\delta,h}(\Phi^h),$$

and we define $y^{\delta,h}: S^1 \to \mathbb{R}^3$ by $y^{\delta,h}(\Phi^h) = z^{\delta,h}$. Clearly, for each δ , as $h \to 0$ we have

$$y^{\delta,h} \to \pi$$
 strongly in $H^1(S^1)$

For $x \in S$ we define

$$V^{\delta,h}(x) = \frac{1}{h^2} \left(\int_{I^h} z^{\delta,h}(x+tn(x)) \, dt - x \right) = \frac{1}{h} \left(\int_{I} y^{\delta,h}(x+tn(x)) \, dt - x \right).$$

Hence

$$V^{\delta,h} = v^{\delta,h} + hw^{\delta,h} + \varepsilon h \tilde{o}^{\delta} + h^2 \tilde{p}^{\delta}.$$

Here we have introduced $\widetilde{o}^{\delta,h}:S\to \mathbb{R}^3$ by

$$\widetilde{o}^{\delta,h}(x) = \int_{I} o^{\delta,h} \left(x + tn(x) \right) dt,$$

and $\widetilde{p}^{\delta,h}$ is defined similarly.

From (81) and (83) we deduce that for each δ , as $h \to 0$ we have

(85)
$$\frac{1}{h}q_{V^{\delta,h}} = \frac{1}{h}q_{v^{\delta,h}} + q_{w^{\delta,h}} + \varepsilon q_{\tilde{o}^{\delta}} + hq_{\tilde{p}^{\delta}} \rightharpoonup B_w \text{ weakly in } L^2(S;\mathbb{S}).$$

Thus the first two parts of the claim are satisfied (for each δ). Also from (83) we deduce

(86)
$$\lim_{h \to 0} \sup_{h \to 0} \left\| \frac{1}{h} q_{V^{\delta,h}} \right\|_{L^2} \leq R \Big(\limsup_{h \to 0} \|q_{w^{\delta,h}}\|_{L^2} + \|o^{\delta}\|_{L^2(S;H^1(I \times \mathcal{Y};\mathbb{R}^3))} + \limsup_{h \to 0} h \|q_{\int_I p^{\delta} dt}\|_{L^2} \Big),$$

for some R > 0. Notice that for $w^{\delta,h}$ that satisfies (78) we have for all $\delta > 0$

(87)
$$\limsup_{h \to 0} \|q_{w^{\delta,h}}\|_{L^2} = \|B_w\|_{L^2}.$$

In order to prove the third part, we need to understand the limiting behaviour of

$$E^{\delta,h} = \frac{1}{h^2} \left(\sqrt{(\nabla_h y^{\delta,h})^T \nabla_h y^{\delta,h}} - I \right).$$

Define $Q: S^h \to \mathbb{R}^{3 \times 3}$ by (19). Lemma 2.4 shows that

$$\nabla_h y^{\delta,h}(\Phi^h) = \nabla z^{\delta,h} = I + h\Omega_{v^{\delta,h}} + h^2 G^{\delta,h}(\Phi^h),$$

where $G^{\delta,h}: S^1 \to \mathbb{R}^{3 \times 3}$ is defined by the following equation on S^h :

$$\begin{aligned} G^{\delta,h}(\Phi^{h}) &= \Omega_{w^{\delta,h}} - \frac{t}{h} b_{v^{\delta,h}}(T_{S}, T_{S}) - t b_{w^{\delta,h}}(T_{S}, T_{S}) - \frac{t^{2}}{h} \nabla_{tan} \mu_{v^{\delta,h} + hw^{\delta,h}} \mathbf{S} \\ &+ \frac{1}{h} \left(\nabla_{tan} v^{\delta,h} + h \nabla_{tan} w^{\delta,h} + t \nabla_{tan} \mu_{v^{\delta,h} + hw^{\delta,h}} \right) T_{S} Q \\ &+ h \nabla_{h} p^{\delta,h}(\Phi^{h}) + \varepsilon \nabla_{h} o^{\delta,h}(\Phi^{h}). \end{aligned}$$

In the case $\gamma = \lim h/\varepsilon$ is nonzero, we deduce from (76) (applied with $P = p^{\delta}$ and $P = o^{\delta}$) that, as $h \to 0$,

(88)
$$h\nabla_h p^{\delta,h} + \varepsilon \nabla_h o^{\delta,h} \xrightarrow{2,\gamma} \Xi^{\delta}.$$

Here we have introduced $\Xi^{\delta}: S^1 \times \mathcal{Y} \to \mathbb{R}^{3 \times 3}$ by

$$\Xi^{\delta}(x,y) = \left(\partial_n p^{\delta}(x,y) + \frac{1}{\gamma}\partial_n o^{\delta}(x,y)\right) \otimes n(x) + \left(\gamma \nabla_y p^{\delta}(x,y) + \nabla_y o^{\delta}(x,y)\right) \nabla_{tan} r(x) T_S(x)$$

Observe that (88) remains true for $\gamma = 0$ provided that $\partial_n o^{\delta} \equiv 0$, and for $\gamma = \infty$ provided that $p^{\delta,h}$ does not depend on y. This follows from (76). Lemma 2.2 and (78) shows that

(89)
$$\operatorname{sym} \Omega_{w^{\delta,h}} = q_{w^{\delta,h}}(T_S, T_S) \to B_w(T_S, T_S) \text{ strongly in } L^2(S).$$

And

$$b_{v^{\delta,h}}(T_S, T_S) \to b_V(T_S, T_S)$$
 in $L^2(S)$.

This holds for each $\delta > 0$ as $h \to 0$.

We conclude that

(90)
$$\operatorname{sym} G^{\delta,h} \xrightarrow{2,\gamma} B_w(T_S,T_S) - tb_V(T_S,T_S) + \operatorname{sym} \Xi^{\delta}$$

strongly two-scale in S^1 as $h \to 0$. Hence the map $E_{app}^{\delta,h}: S^1 \to \mathbb{R}^{3 \times 3}$ defined by

$$E_{app}^{\delta,h} = \operatorname{sym} G^{\delta,h} - \frac{1}{2} (\Omega_{v^{\delta,h}})^2$$

converges strongly two-scale on S^1 to $E^{\delta}: S^1 \times \mathcal{Y} \to \mathbb{R}^{3 \times 3}$, given by

$$E^{\delta}(x,y) = B_w(x)(T_S(x), T_S(x)) - t(x)b_V(x)(T_S(x), T_S(x)) - \frac{1}{2}\Omega_V^2(x) + \operatorname{sym}\Xi^{\delta}(x,y).$$

We now wish to apply Lemma 5.3 to $E^{\delta,h}$ and $E^{\delta,h}_{app}$ in order to conclude that

(91)
$$h^{-4} \int_{S} \int_{I} W\left(x + tn(x), r(x + tn(x))/\varepsilon, I + h^{2} E^{\delta,h}(x + tn(x))\right) dt d\mathcal{H}^{2} \rightarrow \int_{S} \int_{I} \int_{\mathcal{Y}} \mathcal{Q}(x + tn(x), E^{\delta}(x + tn(x), y)) dy dt d\mathcal{H}^{2} + \tilde{E}(\delta),$$

where $|\tilde{E}(\delta)| \to 0$ as $\delta \to 0$. In order to prove this, it remains to verify that the hypotheses of Lemma 5.3 are indeed satisfied.

But in fact, it is not difficult to see that there exists a constant M such that for all $\delta > 0$ small enough

(92)
$$\limsup_{h \to 0} \left(\| \operatorname{sym} G^{\delta,h} \|_{L^2(S^1)} + \| \Omega_{v^{\delta,h}} \|_{L^4(S)} \right) \le M$$

This follows from (86) and the construction of p^{δ} , o^{δ} below.

Moreover, hypotheses (ii) and (iii) of Lemma 5.3 are clearly satisfied by virtue of Sobolev embedding and by (80), (82), (83). Hypothesis (iv) is a direct consequence of (80) and that, as $h \to 0$,

$$h \|\nabla^2 v^{\delta,h}\|_{L^4} \le Ch \sqrt{\|\nabla^2 v^{\delta,h}\|_{L^{\infty}}} \sqrt{\|\nabla^2 v^{\delta,h}\|_{L^2}} \to 0,$$

$$h^2 \|\nabla^2 w^{\delta,h}\|_{L^4} \le h^2 \sqrt{\|\nabla^2 w^{\delta,h}\|_{L^{\infty}}} \sqrt{\|\nabla^2 w^{\delta,h}\|_{L^2}} \to 0,$$

which follow from (80) and (82). Hence the hypotheses of Lemma 5.3 are indeed satisfied, and (91) follows.

It remains to choose the oscillations p^{δ} and o^{δ} in an optimal way. We have to distinguish the three cases.

The case $\gamma \in (0, \infty)$. Let $A_{\delta} \in C^2(S; \mathrm{so}(3))$ be such that $A_{\delta} \to \Omega_V$ strongly in $H^1(S)$ as $\delta \to 0$. Define $p^{\delta} : S^1 \times \mathcal{Y} \to \mathbb{R}^3$ by

$$p^{\delta}(x,y) = t(x) \left(\frac{|A_{\delta}(x)n(x)|^2}{2} I + A_{\delta}^2(x)\right) n(x).$$

Then clearly $\nabla_y p^{\delta} \equiv 0$. We claim that

(93)
$$\operatorname{sym}\left(\partial_n p^\delta \otimes n\right) - \frac{1}{2}A_\delta^2 = -\frac{1}{2}A_\delta^2(T_S, T_S)$$

In fact,

$$\begin{aligned} A_{\delta}^2 - T_S A_{\delta}^2 T_S &= (n \otimes n) A_{\delta}^2 - (n \otimes n) A_{\delta}^2 (n \otimes n) + A_{\delta}^2 (n \otimes n) \\ &= 2 \operatorname{sym} \left(A_{\delta}^2 n \otimes n + \frac{|A_{\delta} n|^2}{2} \ n \otimes n \right) \end{aligned}$$

because A_{δ} is skew symmetric. And this equals $2 \operatorname{sym}(\partial_n p^{\delta} \otimes n)$ because clearly

$$\partial_n p^\delta = \frac{|A_\delta n|^2}{2}n + A_\delta^2 n$$

Thus (93) is proven.

From (93) and from the definition of E^{δ} and Ξ^{δ} we conclude:

$$E^{\delta} = (B_w - tb_V - \frac{1}{2}(A^{\delta})^2)(T_S, T_S) + \tilde{\mathcal{U}}_{\gamma}(o^{\delta}),$$

where

$$\tilde{\mathcal{U}}_{\gamma}(o^{\delta}) := \operatorname{sym}\left(\nabla_{y}o^{\delta}\nabla_{tan}rT_{S} + \gamma^{-1}\partial_{n}o^{\delta}\otimes n\right).$$

Notice that $\tilde{\mathcal{U}}_{\gamma}(o^{\delta}) = \mathcal{U}_{\gamma}(\tilde{o}^{\delta})$, which is just \mathcal{U}_{γ} as in Definition 3.5 and $\tilde{o}_i^{\delta} = o^{\delta} \cdot \tau_i$. Now we choose $\tilde{o}^{\delta} \in C^1(S^1; \dot{C}^1(\mathcal{Y}; \mathbb{R}^3))$ in such a way that

(94)
$$\mathcal{U}_{\gamma}(\tilde{o}^{\delta}) \to \Pi_{\gamma}\left(B_w + \frac{(dV)^2}{2}, -b_V\right) \text{ strongly in } L^2(S; \mathbb{R}^{3\times 3})$$

as $\delta \to 0$. Here, the operator Π_{γ} is as in Lemma 5.1. Then

(95)
$$E^{\delta} \to (B_w - tb_V)(T_S, T_S) + \frac{(dV)^2}{2} + \Pi_{\gamma} (B_w + \frac{(dV)^2}{2}, -b_V)$$
 strongly in $L^2(S; \mathbb{R}^{3\times 3}),$

as $\delta \to 0$ because $-A_{\delta}^2(T_S, T_S) \to (dV)^2$ by (15). By (95) and by the above results, we see that

$$g(\delta,h) = \|y^{\delta,h} - \pi\|_{H^1(S^1)} + \|V^{\delta,h} - V\|_{H^1(S)} + \tilde{d}^K(\frac{1}{h}q_{V^{\delta,h}}, B_w) + \left|\frac{1}{h^4}I^h(y^h) - I_\gamma(V, B_w)\right|$$

satisfies $\limsup_{\delta\to 0} \limsup_{h\to 0} g(\delta, h) = 0$. Here $\tilde{d}^K : L^2(S; M^3) \times L^2(S; M^3) \to \mathbb{R}$ is defined in the following way: For K > 0 we know there exists a metric d^K which defines the weak topology on the ball of radius K. We define:

$$\tilde{d}^{K}(M_{1}, M_{2}) = \begin{cases} d^{K}(M_{1}, M_{2}), \text{ if } \|M_{1}\|_{L^{2}} < K \text{ and } \|M_{2}\|_{L^{2}} < K, \\ +\infty, \text{ otherwise.} \end{cases}$$

The constant $K = K(R, ||B_w||_{L^2}, ||V||_{H^2})$ we choose in a way that the right hand side of (86) is bounded by e.g. $\frac{K}{2} - 1$.

Lemma A.6 then yields a sequence $\delta_h \to 0$ such that $g(\delta_h, h) \to 0$ as $h \to 0$.

The case $\gamma = \infty$. Let $A_{\delta} \in C^2(S; \mathrm{so}(3))$ be such that $A_{\delta} \to A$ strongly in $H^1(S)$ as $\delta \to 0$. Define $\zeta^{\delta} \in C^1(S; \dot{C}^1(I \times \mathcal{Y}; \mathbb{R}^2)), \psi^{\delta} \in C^1(S; \dot{C}^1(I \times \mathcal{Y})), c^{\delta} \in C^1(S; \dot{C}^1(I; \mathbb{R}^3))$ such that

$$\mathcal{U}_{\infty}(\zeta^{\delta},\psi^{\delta},c^{\delta}) \to \Pi_{\infty}\left(B_w + \frac{(dV)^2}{2}, -b_V\right)$$
 strongly in $L^2(S; \mathbb{R}^{3\times 3})$

as $\delta \to 0$. We will use the following fact: if $f: I \times \mathcal{Y} \to \mathbb{R}^3$ then $F(x, y) = \int_0^{t(x)} f(s, y) ds$ satisfies $\partial_n F(x, y) = f(t(x), y)$. Again we wish to have p^{δ} independent of y, in order to ensure the validity of (88). We define

$$p^{\delta}(x,y) = t(x) \left(\frac{|A_{\delta}(x)n(x)|^2}{2} I + A_{\delta}^2(x)\right) n(x) + 2\int_0^{t(x)} c_{\alpha}(s) \, ds\tau^{\alpha} + \int_0^{t(x)} c_3(s) \, ds\tau^3.$$

Then

$$\partial_n p^{\delta} = \frac{|A_{\delta}n|^2}{2}n + A_{\delta}^2 n + 2c_{\alpha}(t)\tau^{\alpha} + c_3(t)\tau^3.$$

For $x \in S^1$ and $y \in \mathcal{Y}$ set

$$o^{\delta}(x,y) = \zeta^{\delta}_{\alpha}(\pi(x),t(x),y)\tau^{\alpha}(x) + 2\psi^{\delta}(\pi(x),t(x),y)n(x).$$

Then

$$\nabla_y o^\delta = \tau^\alpha \otimes \nabla_y \zeta^\delta_\alpha + 2n \otimes \nabla_y \psi^\delta.$$

Since $\gamma = \infty$, we have

sym
$$\Xi^{\delta}$$
 = sym $\left(\partial_n p^{\delta} \otimes n + \nabla_y o^{\delta} \nabla_{tan} r T_S\right) = \mathcal{U}_{\infty}(\zeta^{\delta}, \psi^{\delta}, c^{\delta}),$

From now on the proof is analogous to the case $\gamma \in (0, \infty)$.

Construction for $\gamma = 0$ and $\lim_{h \to 0} \frac{\varepsilon(h)^2}{h} \in [0, \infty)$:

In this one has to modify, in addition, the maps $w^{\delta,h}$. Let A_{δ} be as before. In the case $\lim_{h\to 0} \frac{\varepsilon(h)^2}{h} = 0$ choose $\zeta^{\delta} \in C^1(S; \dot{C}^1(\mathcal{Y}; \mathbb{R}^2)), \varphi^{\delta} \in C^1(S; \dot{C}^2(\mathcal{Y})), g^{\delta} \in C^1(S; \dot{C}^1(I \times \mathcal{Y}; \mathbb{R}^3))$ such that

(96)
$$\mathcal{U}_0^0(\zeta^{\delta},\varphi^{\delta},g^{\delta}) \to \Pi_0^0(B_w + \frac{(dV)^2}{2}, -b_V) \text{ strongly in } L^2(S; \mathbb{R}^{3\times 3}),$$

as $\delta \to 0$. In the case $\lim_{h\to 0} \frac{\varepsilon(h)^2}{h} = \frac{1}{\gamma_1} \in (0,\infty)$ choose them such that

(97)
$$\mathcal{U}^{1}_{0,\gamma_{1}}(\zeta^{\delta},\varphi^{\delta},g^{\delta}) \to \Pi^{1,\gamma_{1}}_{0}\left(B_{w} + \frac{(dV)^{2}}{2}, -b_{V}\right) \text{ strongly in } L^{2}(S^{1}).$$

Extend ζ^{δ} trivially to S^1 and define $o^{\delta}(x,y) = \zeta^{\delta}_{\alpha}(x,y)\tau^{\alpha}(x)$. Then $\partial_n o^{\delta} \equiv 0$, so (88) remains true. We define

$$p^{\delta}(x,y) = t(x) \left(\frac{|A_{\delta}(x)n(x)|^2}{2} I + A_{\delta}^2(x) \right) n(x) + 2 \int_0^{t(x)} g_{\alpha}^{\delta}(\pi(x), s, y) \, ds \, \tau^{\alpha}(x) \\ + \int_0^{t(x)} g_3^{\delta}(\pi(x), s, y) \, ds \, \tau^3(x).$$

We define the modified fields

$$\tilde{w}^{\delta,h} = w^{\delta,h} + \frac{\varepsilon^2}{h} \varphi^{\delta}(\cdot, r/\varepsilon)n,$$

where $w^{\delta,h}$ is defined by the property (77).

Notice that $\tilde{w}^{\delta,h}$ satisfies the condition (80) with $M_1 \leq C \|\nabla_y^2 \varphi^{\delta}\|$, for some constant C > 0, independent of δ . Also using the following facts valid for every fixed $\delta > 0$:

(98)
$$\frac{h}{\varepsilon} \left(\nabla \widetilde{w}^{\delta,h} - \nabla w^{\delta,h} \right) \text{ bounded in } L^{\infty}(S)$$

(99)
$$\frac{h}{\varepsilon} \left(\Omega_{w^{\delta,h}} - \Omega_{\tilde{w}^{\delta,h}} \right) \text{ bounded in } L^{\infty}(S)$$

(100)
$$\frac{h}{\varepsilon} \left(b_{\tilde{w}^{\delta,h}} - \left(b_{w}^{\delta,h} + \frac{1}{h} \sum_{i,j=1}^{2} \partial_{y_{i}y_{j}} \varphi^{\delta} \tau^{i} \otimes \tau^{j} \right) \right) \text{ bounded in } L^{\infty}(S)$$

(101)
$$q_{\tilde{w}^{\delta,h}} = q_{w^{\delta,h}} + \frac{\varepsilon^2}{h} \varphi^{\delta}(\cdot, r/\varepsilon) \mathbf{S} \xrightarrow{2,\gamma} B_w + \frac{1}{\gamma_1} \varphi^{\delta}(x,y) \mathbf{S},$$

we can repeat the same argument as in the case $\gamma \in (0, \infty)$. Namely, notice that (79) is valid and thus the right hand side of (86) can be bounded, by a bound independently of δ . It can be easily seen that (80), (81) are valid. Instead of (89) we have (101).

6 Convex shell

In this chapter we shall identify the Γ -limit for convex shells in the remaining case, i.e. $h \ll \varepsilon^2$. We want to demonstrate the stronger influence of the geometry in this case. We work under the assumption that there exists C > 0 such that

(102)
$$\mathbf{S}(x)\tau \cdot \tau \ge C\tau \cdot \tau, \forall x \in S, \tau \in T_x S.$$

Definition 6.1. For $x \in S$ we define the following operator

$$\mathcal{U}_0^{2,c}: \dot{L}^2(\mathcal{Y}; \mathbb{R}^{2\times 2}_{\text{sym}}) \times L^2(I \times \mathcal{Y}; \mathbb{R}^3) \to L^2(I \times \mathcal{Y}; \mathbb{R}^{3\times 3}_{\text{sym}})$$
$$\mathcal{U}_0^{2,c}(B,g) = \sum_{i,j=1}^3 \begin{pmatrix} B & g_1 \\ g_2 \\ (g_1, g_2) & g_3 \end{pmatrix}_{ij} \tau^i \otimes \tau^j$$

and function space of relaxation fields

(103)
$$L_0^{2,c}(I \times \mathcal{Y}; \mathbb{R}^{3 \times 3}_{\text{sym}}) := \left\{ \mathcal{U}_0^{2,c}(B,g) : B \in \dot{L}^2(\mathcal{Y}; \mathbb{R}^{2 \times 2}_{\text{sym}}), g \in L^2(I \times \mathcal{Y}; \mathbb{R}^3) \right\}.$$

Again as before it can be seen that $L_0^{2,c}(I \times \mathcal{Y}; \mathbb{R}^{3 \times 3}_{sym})$ is a closed subspace of $L^2(I \times \mathcal{Y}; \mathbb{R}^{3 \times 3}_{sym})$. We also define the functional $I_0^{2,c}: H^2(S; \mathbb{R}^3) \times L^2(S; \mathbb{S})) \to \mathbb{R}$

(104)
$$I_0^{2,c}(V, B_w) = \int_S \mathcal{Q}_0^{2,c}(x, B_w + \frac{1}{2}(dV)^2, -b_V) \,\mathrm{d}x,$$

with the quadratic form $\mathcal{Q}_0^{2,c}(x) : \mathbb{S}(x)_{\text{sym}} \times \mathbb{S}(x)_{\text{sym}} \to \mathbb{R}$:

(105)
$$\mathcal{Q}_{0}^{2,c}(x,q^{1},q^{2}) = \inf_{U \in L_{0}^{2,c}(I \times \mathcal{Y};\mathbb{R}^{3\times3}_{\text{sym}})} \iint_{I \times \mathcal{Y}} \mathcal{Q}\left(x + tn(x), y, q^{1} + tq^{2} + U\right) dt dy.$$

As before, it is easy to see that the definition is equivalent to the following one:

(106)
$$\mathcal{Q}_0^{2,c}(x,q^1,q^2) = \inf_{\substack{\dot{B}\in\dot{L}^2(\mathcal{Y};\mathbb{M}^2_{\mathrm{sym}})}} \iint_{I\times\mathcal{Y}} \mathcal{Q}_2\Big(x+tn(x),y,q^1+tq^2+\sum_{i,j=1}^2\dot{B}_{ij}\tau_i\otimes\tau_j)\,\mathrm{d}t\,\mathrm{d}y.$$

In the case when Q does not depend on t we have that

$$\begin{split} \tilde{\mathcal{Q}}_0^{2,c}(x,q^1,q^2) &= \inf_{\dot{B}\in\dot{L}^2(\mathcal{Y};\mathbb{M}^2_{\mathrm{sym}})} \int_{\mathcal{Y}} \mathcal{Q}_2(x,y,q^1+\sum_{i,j=1}^2 (\dot{B})_{ij}\tau^i\otimes\tau^j) \,\mathrm{d}y \\ &+ \frac{1}{12} \int_{\mathcal{Y}} \mathcal{Q}_2(x,y,q^2) \,\mathrm{d}y. \end{split}$$

Under the assumption (102) it is well-known that $\mathcal{B} = L^2(S, \mathbb{S})$, cf. e.g. [Cia00], [LMP10]. Thus if one wants to additionally to relax the functional $I_0^{2,c}$ with respect to B_w , one in this case obtains the functional $\tilde{I}_0^{2,c} : H^2(S; \mathbb{R}^3) \to \mathbb{R}$

(107)
$$\tilde{I}_0^{2,c}(V) = \frac{1}{12} \int_S \int_{\mathcal{Y}} \mathcal{Q}_2(x, y, b_V) \, \mathrm{d}y \, \mathrm{d}x.$$

This functional is the same as in the ordinary von Kármán model. For the form $\mathcal{Q}_0^{2,c}$ one can make assertions analogous to Lemma 5.1 with the appropriate operator $\Pi_0^{2,c}$ and Lemma 5.2. We introduce the space

$$FL(S; \dot{C}^{\infty}(\mathcal{Y})) = \Big\{ (x, y) \mapsto \sum_{k \in \mathbb{Z}^2, \ |k| \le n, \ k \ne 0} c^k(x) e^{2\pi i k \cdot y} :$$
$$n \in \mathbf{N} \text{ and } c^k \in C_0^1(S; \mathbb{C}) \text{ with } \overline{c}^k = c^{-k} \Big\}.$$

By Fourier transform it can be easily seen that $FL(S; C^{\infty}(\mathcal{Y}))$ is dense in $L^{2}(S; \dot{H}^{m}(\mathcal{Y}))$, for any $m \in \mathbb{N}_{0}$. The following lemma resembles Lemma 3.3 in [Sch07]:

Lemma 6.2. Assume (102) and let $\dot{B} \in L^2(S; \dot{L}^2(\mathcal{Y}; \mathbb{S}))$. Then there exist unique $w \in L^2(S; \dot{H}^1(\mathcal{Y}; \mathbb{R}^2))$ and $\varphi \in L^2(S; \dot{L}^2(Y))$ such that

(108)
$$\sum_{i,j=1,2} \left(\operatorname{sym} \nabla_y w \right)_{ij} \tau^i \otimes \tau^j + \varphi \mathbf{S} = \dot{B}.$$

Moreover, if $\dot{B}_{ij} \in FL(S; \dot{C}^{\infty}(\mathcal{Y}))$ for every i, j = 1, 2 then $w_i \in FL(S; \dot{H}^1(\mathcal{Y}))$, for i = 1, 2 and $\varphi \in FL(S; \dot{H}^1(\mathcal{Y}))$

Proof. One possible proof is to apply the operator $\operatorname{curl}_y \operatorname{curl}_y$ to both sides of (108), which leads to the PDE

$$\operatorname{cof} \mathbf{S} : \nabla_y^2 \varphi = \operatorname{curl}_y \operatorname{curl}_y \dot{B},$$

which by virtue of (102) is an elliptic PDE with constant coefficients (for each x). We prefer to give a direct proof.

There exist b_{ij}^k such that for all i, j = 1, 2:

$$\dot{B}(x,y)_{ij} = \sum_{k \in \mathbb{Z}^2} b_{ij}^k(x) e^{2\pi i k \cdot y}, \text{ where } \sum_{k \in \mathbb{Z}^2} \|b_{ij}^k\|_{L^2} < \infty, \overline{b}_{ij}^k = b_{ij}^{-k}, \ b_{ij}^0 = 0.$$

We assume that for i = 1, 2:

$$w_j = \sum_{k \in \mathbb{Z}^2} c_j^k(x) e^{2\pi i k \cdot y}, \ \overline{c}_j^k = c_j^{-k}, \ c_j^0 = 0$$

and

$$\varphi = \sum_{k \in \mathbb{Z}^2} d^k(x) e^{2\pi i k \cdot y}, \ \overline{d}^k = d^{-k}, \ d^0 = 0.$$

The equation (108) is equivalent to the following problem for every $(k_1, k_2) \in \mathbb{Z}^2$ find complex coefficients c_i^k, b_{ij}^k, d^k such that

$$k_1 c_1^k + d^k \mathbf{S}_{11} = b_{11}^k$$
$$\frac{1}{2} (k_2 c_1^k + k_1 c_2^k) + d^k \mathbf{S}_{12} = b_{12}^k$$
$$k_2 c_2^k + d^k \mathbf{S}_{22} = b_{22}^k.$$

By (2) and (102) it is easy to see that there exists C > 0 such that the determinant of the system is bounded from below by $C(k_1^2 + k_2^2)$. Using this it follows that there exists C > 0 such that

$$|d^{k}(x)|^{2} + \sum_{i=1}^{2} |k|^{2} |c_{i}^{k}(x)|^{2} \leq C(\sum_{i,j=1}^{2} |b_{ij}^{k}(x)|^{2}), \forall x \in S.$$

Now all claims follow easily.

Theorem 6.3. Assume (102) and that $h \ll \varepsilon^2$ and that W satisfies Assumption 3.3. Let $(u^h) \subset H^1(S^h; \mathbb{R}^3)$ satisfy

(109)
$$\limsup_{h \to 0} \frac{1}{h^4} E^h(u^h) < \infty,$$

where

$$E^{h}(u^{h}) = \frac{1}{h} \int_{S^{h}} \bar{W}(\Phi^{h}(x), r(x)/\varepsilon, \nabla u^{h}) dx.$$

Define $y^h \in H^1(S^1; \mathbb{R}^3)$ by the equation $\bar{y}^h(\Phi^h(x)) = u^h(x)$. Then the following are true

(i) (compactness). There exists a subsequence of (\bar{y}^h) , still denoted by (\bar{y}^h) and there exist $Q^h \in SO(3)$ and $c^h \in \mathbb{R}^3$ such that the sequences $y^h := (Q^h)^T \bar{y}^h - c^h$ and $V^h := \frac{1}{h} \left(\int_I y^h(x + tn(x)) \, \mathrm{d}t - x \right)$ satisfy the following

- (a) $y^h \to \pi$ strongly in $H^1(S^1; \mathbb{R}^3)$.
- (b) There exists an infinitesimal bending $V \in H^2(S; \mathbb{R}^3)$ of S such that $V^h \to V$ strongly in $H^1(S; \mathbb{R}^3)$.
- (c) There exists $B_w \in L^2(S; \mathbb{S})$ such that

$$\frac{1}{h}q_{V^h} \rightharpoonup B_w$$
 weakly in $L^2(S; \mathbb{S})$.

(ii) (lower bound). Defining $I_0^{2,c}$ by (104) we have

$$\liminf_{h \to 0} \frac{1}{h^4} E^h(u^h) \ge I_0^{2,c}(V, \tilde{B}_w),$$

(iii) (recovery sequence) For any infinitesimal bending $V \in H^2(S^1, \mathbb{R}^3)$ and $B_w \in L^2(S; \mathbb{S})$ there exists $u^h \in H^1(S^h; \mathbb{R}^3)$ satisfying (109) and such that conclusions of part (i) are true with $Q^h = I$ and $c^h = 0$. Moreover, equality holds in (ii).

Proof. We will only give the sketch of the proof since it is analogous to the previous cases. Since V is an infinitesimal bending, we have

(110)
$$\partial_{\tau} V(x) = A(x)\tau$$
, for all $\tau \in T_x S$ for some $A \in H^1(S; so(3))$,

Let us assume as in Proposition 4.5 that

$$\nabla_{\tan} \nabla_{\tan} (V_s^h \cdot n) \xrightarrow{2,\gamma} \nabla_{\tan} \nabla_{\tan} (Vn) + \sum_{i,j=1}^2 (\partial_{y_i y_j}^2 \varphi) \tau^i \otimes \tau^j,$$

for some $\varphi \in L^2(S; \dot{H}^2(\mathcal{Y}))$. Using (iii) of Proposition 4.2 as well as Lemma 6.2 we conclude that $\varphi = 0$. Thus from Proposition 4.5 we conclude that for E^h defined in (56) we have $E^h \xrightarrow{2,\gamma} E(x,y)$ where

(111)
$$E = B_w + \dot{B} + \frac{1}{2}(dV)^2 - tb_V + U,$$

for some $U \in L^2(S; L_0(I \times \mathcal{Y}))$ and $\dot{B} \in L^2(S; \dot{L}^2(\mathcal{Y}; \mathbb{S}))$. The lower bound easily follows from Lemma 4.6 and the definition of the functional $I_0^{2,c}$.

To prove the upper bound we follow the proof of Proposition 5.4, the case $\gamma = 0$. Namely, let us again take $A^{\delta} \in C^2(S; \mathrm{so}(3))$ such that $\lim_{\delta \to 0} ||A^{\delta} - A||_{H^1} = 0$ and \dot{B}^{δ} such that for every $i, j = 1, 2, (\dot{B}^{\delta})_{ij} \in FL(S; \dot{C}^{\infty}(\mathcal{Y})), g^{\delta} \in C^1(S; C^1(I \times \mathcal{Y}; \mathbb{R}^3))$ and

(112)
$$\lim_{\delta \to 0} \left\| \mathcal{U}_0^{2,c}(\dot{B}^{\delta}, g^{\delta}) - \Pi_0^{2,c} \left(B_w + \frac{1}{2} (dV)^2, -b_V \right) \right\|_{L^2(S; \mathbb{R}^{3 \times 3})} = 0.$$

By Lemma 6.2 there exist $z^{\delta} \in (FL(S; \dot{C}^{\infty}(\mathcal{Y})))^2$ and $\varphi^{\delta} \in FL(S; \dot{C}^{\infty}(\mathcal{Y})$ solving the system

(113)
$$\sum_{i,j=1,2} \left(\operatorname{sym} \nabla_y z^{\delta} \right)_{ij} \tau^i \otimes \tau^j + \varphi^{\delta} \mathbf{S}_{ij} = \dot{B}^{\delta}.$$

We define

$$\begin{split} \tilde{v}^{\delta,h} &= v^{\delta,h}, \\ \tilde{w}^{\delta,h} &= w^{\delta,h} + \varphi^{\delta}(\cdot, r/\varepsilon)n + \varepsilon_{(h)} \big(z_1^{\delta}(\cdot, r/\varepsilon)\tau^1 + z_2^{\delta}(\cdot, r/\varepsilon)\tau^2 \big), \\ p^{\delta}(x,y) &= t(x) \left(\frac{|A_{\delta}(x)n(x)|^2}{2} I + A_{\delta}^2(x) \right) n(x) + 2 \int_0^{t(x)} g_{\alpha}^{\delta}(\pi(x), s, y) \ ds \ \tau^{\alpha}(x) \\ &+ \int_0^{t(x)} g_3^{\delta}(\pi(x), s, y) \ ds \ \tau^3(x). \end{split}$$

where $v^{\delta,h}$ is defined in (82) and $w^{\delta,h}$ is defined by the property (77). Notice that, similarly as before:

(114)
$$\varepsilon \left(\nabla \widetilde{w}^{\delta,h} - \nabla w^{\delta,h}\right) \text{ bounded in } L^{\infty}(S)$$

(115)
$$\varepsilon \left(\Omega_{w^{\delta,h}} - \Omega_{\tilde{w}^{\delta,h}}\right) \text{ bounded in } L^{\infty}(S)$$

(116)
$$\varepsilon \left(b_{\tilde{w}^{\delta,h}} - \left(b_{w}^{\delta,h} + \frac{1}{\varepsilon^2} \sum_{i,j=1}^{2} \partial_{y_i y_j} \varphi^{\delta} \tau^i \otimes \tau^j \right) \right)$$
 bounded in $L^{\infty}(S)$

(117)
$$\|q_{\tilde{w}^{\delta,h}}\|_{L^2}$$
 bounded independately of δ, h

(118)
$$\frac{1}{\varepsilon} \left(q_{\tilde{w}^{\delta,h}} - \left(q_{w^{\delta,h}} + \dot{B}^{\delta} \right) \right) \text{ bounded in } L^{\infty}(S).$$

Now we continue as in the proof of Proposition 5.4, after concluding that $\tilde{w}^{\delta,h}$ satisfies the condition (80). Boundedness of the right hand side of (86) follows easily. It can be easily seen that (80), (81) are valid. Instead of (89) we have (118).

A Auxiliary results

In the sequel we consider the sequence $\varepsilon(h) \to 0$ as $h \to 0$ and $\Omega \subset \mathbb{R}^3$ a Lipschitz domain and $\tau_i = \mathbf{e}_i$, for i = 1, 2, 3, where \mathbf{e}_i are standard coordinate vectors. The set \mathcal{Y} can be considered as the set $[0, 1)^3$ i.e. $[0, 1)^2$ with the topology of torrus. The claims can be trivially extended to \mathbb{R}^n . For the proofs see e.g. [All92]. For the last claim see [Velb, Lemma 3].

Lemma A.1. (i) Any sequence that is bounded in $L^2(\Omega)$ admits a two-scale convergence subsequence.

- (ii) Let $f \in L^2(\Omega \times \mathcal{Y})$ and $\{f^h\}_{h>0} \subset L^2(\Omega)$ be such that $f^h \xrightarrow{2,\gamma} f(x,y)$ weakly. Then $f^h \rightharpoonup \int_Y f(\cdot,y) \, dy$ weakly in $L^2(\Omega)$.
- (iii) Let $f^0 \in L^2(\Omega)$ and $\{f^h\}_{h>0} \subset L^2(\Omega)$ be such that $f^h \to f^0$ weakly in L^2 . Then (after passing to subsequences) we have $f^h \xrightarrow{2,\gamma} f^0(x) + \tilde{f}(x,y)$ for some $\tilde{f} \in L^2(\Omega \times Y)$ with $\int_Y \tilde{f}(\cdot, y) \, dy = 0$ almost everywhere in Ω .
- (iv) Let $f^0 \in L^2(\Omega)$ and $\{f^h\}_{h>0} \subset L^2(\Omega)$ be such that $f^h \to f^0$ strongly in L^2 . Then $f^h \xrightarrow{2,\gamma} f^0(x)$.

(v) Let $f^0 \in H^1(\Omega)$ and $\{f^h\}_{h>0} \subset H^1(\Omega)$ be such that $f^h \rightharpoonup f^0$ weakly in H^1 . Then (after passing to subsequences)

$$\nabla f^h \xrightarrow{2,\gamma} \nabla f^0 + \nabla_y \phi(x,y)$$

for some $\phi \in L^2(\Omega, \dot{H}^1(\mathcal{Y})).$

(vi) Let $f^0 \in H^2(\Omega)$ and $\{f^h\}_{h>0} \subset H^2(\Omega)$ be such that $f^h \rightharpoonup f^0$ weakly in H^2 . Then (after passing to subsequences)

$$\nabla^2 f^h \xrightarrow{2,\gamma} \nabla^2 f^0 + \nabla^2_y \phi(x,y)$$

for some $\phi \in L^2(\Omega, \dot{H}^2(\mathcal{Y}))$.

For brevity we shall write ε instead of $\varepsilon(h)$. At several places in our argument we are only interested in the oscillatory part of the two-scale limit. In the following, we introduce as in [HNV] the special notation $\xrightarrow{osc,\gamma}$ for that purpose. As a motivation consider a sequence $\{f^h\}_{h>0} \subset L^2(\Omega)$ with weak two-scale limit $f \in L^2(\Omega \times Y)$. Consider

$$f^0(x) := \int_Y f(x,y) \, dy$$
 and $\widetilde{f}(x,y) = f(x,y) - f^0(x)$

According to Lemma A.1 the function f^0 is the weak limit of f^h . We call $\tilde{f}(x, y)$ the oscillatory part of f. Evidently we have

(119)
$$\lim_{h \to 0} \int_{\Omega} f^{h}(x)\varphi(x)g(\frac{x'}{\varepsilon}) \, dx = \iint_{\Omega \times Y} \widetilde{f}(x,y)\varphi(x)g(y) \, dy \, dx$$

for all $\varphi \in C_{0}^{\infty}(\Omega)$ and $g \in C^{\infty}(\mathcal{Y})$ with $\int_{Y} g \, dy = 0$.

Motivated by that we introduce the following vocabulary (see [HNV]):

Definition A.2. For a sequence $(f^h)_{h>0} \subset L^2(\Omega)$ and $\tilde{f} \in L^2(\Omega \times Y)$ with $\int_Y \tilde{f}(\cdot, y) dy = 0$ almost everywhere in Ω we write

 $f^h \xrightarrow{osc,\gamma} \widetilde{f}(x,y),$

if (119) holds for all $\varphi \in C_0^{\infty}(\Omega)$ and $g \in C^{\infty}(\mathcal{Y})$ with $\int_Y g \, dy = 0$.

Lemma A.3. Let f^0 and $f^h \in L^2(\Omega)$ be such that $f^h \rightharpoonup f^0$ weakly in $L^2(\Omega)$ and $f^h \xrightarrow{osc,\gamma} \widetilde{f}(x,y)$. Then $f^h \xrightarrow{2,\gamma} f^0(x) + \widetilde{f}(x,y)$ weakly two-scale.

Proof. Straightforward.

The following Lemma was needed in the proof of Proposition 4.2.

Lemma A.4. (i) Let f^0 and $f^h \in H^1(\Omega)$ be such that $f^h \rightharpoonup f^0$ weakly in $H^1(\Omega)$ and assume that

$$\nabla f^h \xrightarrow{2,\gamma} \nabla f^0 + \nabla_y \phi(x,y)$$

for some $\phi \in L^2(\Omega; \dot{H}^1(\mathcal{Y}))$. Then

$$\frac{f^h}{\varepsilon} \xrightarrow{osc,\gamma} \phi.$$

(ii) Let f^0 and $f^h \in H^1(\Omega)$ be such that $f^h \rightharpoonup f^0$ weakly in $H^2(\Omega)$ and assume that

$$\nabla^2 f^h \xrightarrow{2,\gamma} \nabla^2 f^0 + \nabla^2_y \phi(x,y)$$

for some $\phi \in L^2(\Omega; \dot{H}^2(\mathcal{Y}))$. Then

$$\frac{f^h}{\varepsilon^2} \xrightarrow{osc,\gamma} \phi.$$

Proof. The proof of (i) is given in [HNV, Lemma 3.7]. Here we prove (ii) which goes in an analogous way. Let G denote the unique solution in $C^{\infty}(\mathcal{Y})$ to

$$-\Delta_y G = g, \qquad \int_Y G \, dy = 0.$$

We put $G^h(x) = g(\frac{x'}{\varepsilon})$. Then $\Delta_y G^h(x) = \frac{1}{\varepsilon^2} g(\frac{x}{\varepsilon})$ and $\nabla G^h(x) = \frac{1}{\varepsilon} \nabla_y G(\frac{x}{\varepsilon})$.

$$(120) \frac{1}{\varepsilon^2} \int_{\Omega} f^h(x) g(\frac{x}{\varepsilon^2}) \psi(x) \, dx = \int_{S} f^h \Delta G^h \psi \, dx$$

$$= -\int_{\Omega} \nabla f^h \cdot \nabla (G^h \psi) \, dx - 2 \int_{\Omega} f^h (\nabla G^h \cdot \nabla \psi) \, dx$$

$$-\int_{\Omega} f^h G^h \Delta \psi \, dx$$

$$= \int_{\Omega} \Delta f^h (G^h \psi) \, dx - 2 \int_{\Omega} f^h (\nabla G^h \cdot \nabla \psi) \, dx$$

$$-\int_{\Omega} f^h G^h \Delta \psi \, dx$$

$$\rightarrow \iint_{\Omega \times Y} \Delta_y \phi(x, y) G(y) \psi \, dx \, dy$$

$$= \iint_{\Omega \times Y} \phi(x, y) g(y) \psi \, dx \, dy,$$

where we have used (i) for the claim $\frac{f^h}{\varepsilon} \xrightarrow{osc,\gamma} 0$ that is used to conclude $\int_{\Omega} f^h (\nabla G^h \cdot \nabla \psi) \, dx \to 0$.

The following proposition can be found in [FJM06, Proposition 2]

Proposition A.5. Let Ω be a bounded Lipschitz domain in \mathbb{R}^n and let $1 , <math>k \in \mathbb{N}$ and $\lambda > 0$. Suppose that $u \in W^{k,p}(\Omega)$ and let

$$|u|_k(x) := \sum_{|\alpha| \le k} |\nabla^{\alpha} u(x)|.$$

Then there exists $u^{\lambda} \in W^{k,\infty}$ such that

$$\begin{aligned} \|u^{\lambda}\|_{W^{k,\infty}} &\leq C(p,k,\Omega)\lambda, \\ \left| \{x \in \Omega : u^{\lambda}(x) \neq u(x)\} \right| &\leq \frac{C(p,k)}{\lambda^{p}} \int_{|u|_{k} \geq \lambda/2} |u|_{k}^{p} \,\mathrm{d}x, \\ \|u^{\lambda}\|_{W^{k,p}} \leq C(p,k,\Omega) \|u\|_{W^{k,p}}. \end{aligned}$$

In particular

$$\lim_{\lambda \to \infty} \lambda^p \left| \{ x \in \Omega : u^{\lambda}(x) \neq u(x) \} \right| = 0,$$

and

$$\lim_{\lambda \to \infty} \|u^{\lambda} - u\|_{W^{k,p}} = 0.$$

The following diagonalization lemma is due to [Att84, Corollary 1.16]:

Lemma A.6. Let $g: [0,\infty) \times [0,\infty) \rightarrow [0,\infty)$ and suppose that

$$\limsup_{\delta \to 0} \limsup_{h \to 0} g(\delta, h) = 0$$

Then there is a monotone function $(0,\infty) \ni h \mapsto \delta(h) \in (0,\infty)$ with $\lim_{h\to 0} \delta(h) = 0$ and $\limsup_{h\to 0} g(\delta(h), h) = 0$.

Acknowledgement. Both authors were supported by Deutsche Forschungsgemeinschaft grant no. HO-4697/1-1. Part of this work was completed while the second author was affiliated with BCAM.

References

- [ABP91] Emilio Acerbi, Giuseppe Buttazzo, and Danilo Percivale. A variational definition of the strain energy for an elastic string. J. Elasticity, 25(2):137–148, 1991.
- [All92] Grégoire Allaire. Homogenization and two-scale convergence. SIAM J. Math. Anal., 23(6):1482–1518, 1992.
- [Att84] Hedy Attouch. Variational convergence for functions and operators. Applicable Mathematics Series. Pitman (Advanced Publishing Program), Boston, MA, 1984.
- [BB06] Jean-François Babadjian and Margarida Baía. 3D–2D analysis of a thin film with periodic microstructure. *Proc. Roy. Soc. Edinburgh Sect. A*, 136(2):223– 243, 2006.
- [BFF00] Andrea Braides, Irene Fonseca, and Gilles Francfort. 3D-2D asymptotic analysis for inhomogeneous thin films. *Indiana Univ. Math. J.*, 49(4):1367–1404, 2000.
- [Bra85] Andrea Braides. Homogenization of some almost periodic coercive functional. Rend. Accad. Naz. Sci. XL Mem. Mat. (5), 9(1):313–321, 1985.
- [Cho97] Daniel Choi. On geometrical rigidity of surfaces. Application to the theory of thin linear elastic shells. *Math. Models Methods Appl. Sci.*, 7(4):507–555, 1997.
- [Cia00] Philippe G. Ciarlet. Mathematical elasticity. Vol. III, volume 29 of Studies in Mathematics and its Applications. North-Holland Publishing Co., Amsterdam, 2000. Theory of shells.

- [CL96] Philippe G. Ciarlet and Véronique Lods. Asymptotic analysis of linearly elastic shells. I. Justification of membrane shell equations. Arch. Rational Mech. Anal., 136(2):119–161, 1996.
- [CLM96] Philippe G. Ciarlet, Véronique Lods, and Bernadette Miara. Asymptotic analysis of linearly elastic shells. II. Justification of flexural shell equations. Arch. Rational Mech. Anal., 136(2):163–190, 1996.
- [FJM02] Gero Friesecke, Richard D. James, and Stefan Müller. A theorem on geometric rigidity and the derivation of nonlinear plate theory from three-dimensional elasticity. *Comm. Pure Appl. Math.*, 55(11):1461–1506, 2002.
- [FJM06] Gero Friesecke, Richard D. James, and Stefan Müller. A hierarchy of plate models derived from nonlinear elasticity by gamma-convergence. Arch. Ration. Mech. Anal., 180(2):183–236, 2006.
- [FJMM03] Gero Friesecke, Richard D. James, Maria Giovanna Mora, and Stefan Müller. Derivation of nonlinear bending theory for shells from three-dimensional nonlinear elasticity by Gamma-convergence. C. R. Math. Acad. Sci. Paris, 336(8):697–702, 2003.
- [GSP95] Giuseppe Geymonat and Enrique Sánchez-Palencia. On the rigidity of certain surfaces with folds and applications to shell theory. *Arch. Rational Mech. Anal.*, 129(1):11–45, 1995.
- [HNV] Peter Hornung, Stefan Neukamm, and Igor Velčić. Derivation of the homogenized bending plate model from 3D nonlinear elasticity. *Preprint: http://www.mis.mpg.de/preprints/2013/preprint2013_23.pdf*.
- [Hor] Peter Hornung. Continuation of infinitesimal bendings on developable surfaces and equilibrium equations for nonlinear bending theory of plates. to appear in Comm. Part. Differ. Equat.
- [Hor12] Peter Hornung. The Willmore functional on isometric immersions. *MIS MPG Preprint*, 2012.
- [LDR95] Hervé Le Dret and Annie Raoult. The nonlinear membrane model as variational limit of nonlinear three-dimensional elasticity. J. Math. Pures Appl. (9), 74(6):549–578, 1995.
- [LDR96] H. Le Dret and A. Raoult. The membrane shell model in nonlinear elasticity: a variational asymptotic derivation. J. Nonlinear Sci., 6(1):59–84, 1996.
- [LMP10] Marta Lewicka, Maria Giovanna Mora, and Mohammad Reza Pakzad. Shell theories arising as low energy Γ-limit of 3d nonlinear elasticity. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), 9(2):253–295, 2010.
- [Lut85] Adam Lutoborski. Homogenization of thin elastic shell. *Journal of Elasticity*, 15(1):69–87, 1985.
- [MM03] Maria Giovanna Mora and Stefan Müller. Derivation of the nonlinear bendingtorsion theory for inextensible rods by Γ-convergence. Calc. Var. Partial Differential Equations, 18(3):287–305, 2003.

- [MM04] Maria Giovanna Mora and Stefan Müller. A nonlinear model for inextensible rods as a low energy Γ-limit of three-dimensional nonlinear elasticity. Ann. Inst. H. Poincaré Anal. Non Linéaire, 21(3):271–293, 2004.
- [Mül87] Stefan Müller. Homogenization of nonconvex integral functionals and cellular elastic materials. *Arch. Rational Mech. Anal.*, 99(3):189–212, 1987.
- [Neu10] Stefan Neukamm. Homogenization, linearization and dimension reduction in elasticity with variational methods. Phd thesis, Tecnische Universität München, 2010.
- [Neu12] Stefan Neukamm. Rigorous derivation of a homogenized bending-torsion theory for inextensible rods from three-dimensional elasticity. Arch. Ration. Mech. Anal., 206(2):645–706, 2012.
- [Ngu89] Gabriel Nguetseng. A general convergence result for a functional related to the theory of homogenization. SIAM J. Math. Anal., 20(3):608–623, 1989.
- [NV] Stefan Neukamm and Igor Velčić. Derivation of a homogenized von Kármán plate theory from 3D elasticity. to appear in M3AS; Preprint: http://www.mis.mpg.de/preprints/2012/preprint2012_61.pdf.
- [Pan01] Olivier Pantz. Une justification partielle du modèle de plaque en flexion par Γ-convergence. C. R. Acad. Sci. Paris Sér. I Math., 332(6):587–592, 2001.
- [Sch07] Bernd Schmidt. Plate theory for stressed heterogeneous multilayers of finite bending energy. J. Math. Pures Appl. (9), 88(1):107–122, 2007.
- [SP89a] Évariste Sanchez-Palencia. Statique et dynamique des coques minces. I. Cas de flexion pure non inhibée. C. R. Acad. Sci. Paris Sér. I Math., 309(6):411–417, 1989.
- [SP89b] Évariste Sanchez-Palencia. Statique et dynamique des coques minces. II. Cas de flexion pure inhibée. Approximation membranaire. C. R. Acad. Sci. Paris Sér. I Math., 309(7):531–537, 1989.
- [Vela] Igor Velčić. A note on the derivation of homogenized bending plate model. *Preprint: http://arxiv.org/abs/1212.2594.*
- [Velb] Igor Velčić. Periodically wrinkled plate of Föppl von Kármán type. to appear in Ann. Sc. Norm. Super. Pisa Cl. Sci., preprint: http://arxiv.org/abs/1104.0661.
- [Vis06] Augusto Visintin. Towards a two-scale calculus. ESAIM Control Optim. Calc. Var., 12(3):371–397 (electronic), 2006.
- [Vis07] Augusto Visintin. Two-scale convergence of some integral functionals. Calc. Var. Partial Differential Equations, 29(2):239–265, 2007.