

NEW BOUNDS FOR BILINEAR CALDERÓN-ZYGMUND OPERATORS AND APPLICATIONS

WENDOLÍN DAMIÁN, MAHDI HORMOZI AND KANGWEI LI

ABSTRACT. In this work we extend Lacey’s domination theorem to prove the pointwise control of bilinear Calderón–Zygmund operators with Dini–continuous kernel by sparse operators. The precise bounds are carefully tracked following the spirit in a recent work of Hytönen, Roncal and Tapiola. We also derive new mixed weighted estimates for a general class of bilinear dyadic positive operators using multiple A_∞ constants inspired in the Fujii–Wilson and Hrusčev classical constants. These estimates have many new applications including mixed bounds for multilinear Calderón–Zygmund operators and their commutators with BMO functions, square functions and multilinear Fourier multipliers.

1. INTRODUCTION

In the last decades, several advances have been carried out in the fruitful area of weighted inequalities concerning the precise determination of the optimal bounds of the weighted operator norm of Calderón–Zygmund operators in terms of the A_p constant of the weights. It has been a long journey from the proof of the linear dependence on the A_2 constant of w of the $L^2(w)$ norm of the Ahlfors–Beurling transform [28] leading to the full proof of the A_2 theorem due to T. Hytönen [12], plenty of previous partial attempts by others. We refer the interested reader to [12, 20] and the references therein for a survey on the advances on the topic.

After Hytönen’s proof, A. Lerner [20] gave an alternative proof of the A_2 theorem which showed that Calderón–Zygmund operators can be controlled in norm from above by a very special dyadic type operators defined by means of the concept of *sparseness*. More precisely, if \mathcal{S} is a collection of dyadic

Date: November 24, 2016.

2010 Mathematics Subject Classification. Primary: 42B20, Secondary: 42B25.

Key words and phrases. domination theorem, Dini condition, multilinear Calderón–Zygmund operators, commutators, square functions, Fourier multipliers.

W.D. and K.L. were supported by the European Union through T. Hytönen’s ERC Starting Grant “Analytic-probabilistic methods for borderline singular integrals”. They are members of the Finnish Centre of Excellence in Analysis and Dynamics Research. W.D. is also supported by the Spanish Ministry of Economy and Competitiveness through the project MTM2015-63699-P “Análisis Complejo, Espacios de Banach y Operadores”. K.L. is also supported by the Basque Government through the BERC 2014-2017 program and by Spanish Ministry of Economy and Competitiveness MINECO: BCAM Severo Ochoa excellence accreditation SEV-2013-0323.

cubes within a dyadic grid \mathcal{D} (see Sect. 2 for the definition), we say that the operator $\mathcal{A}_{\mathcal{S}}$ is *sparse* if

$$(1.1) \quad \mathcal{A}_{\mathcal{S},\mathcal{D}}f(x) = \sum_{Q \in \mathcal{S}} \langle f \rangle_Q \mathbf{1}_Q(x),$$

where $\mathbf{1}_Q$ is the characteristic function of the cube Q and the collection \mathcal{S} satisfies that there exists some $\gamma \in (0, 1)$ such that for each $Q \in \mathcal{S}$,

$$\sum_{S' \in \text{ch}_{\mathcal{S}}(S)} |S'| \leq \gamma |S|,$$

for every $S \in \mathcal{S}$. Here $\text{ch}_{\mathcal{S}}(S)$ denotes the set of the \mathcal{S} -children of a dyadic cube S . Namely, the set of the maximal cubes $S' \in \mathcal{D}$ such that $S' \subsetneq S$. One remarkable aspect from Lerner's proof is its flexibility to be adapted to the multilinear setting. In fact, in [7] the first author, A. Lerner and C. Pérez proved that multilinear Calderón–Zygmund operators can be controlled from above in norm by a supremum of sparse operators. More precisely, if X is a Banach function space over \mathbb{R}^n equipped with the Lebesgue measure, it holds that for any appropriate \vec{f} ,

$$(1.2) \quad \|T(\vec{f})\|_X \leq C_{T,n,m} \sup_{\mathcal{D},\mathcal{S}} \|\mathcal{A}_{\mathcal{D},\mathcal{S}}(\vec{f})\|_X,$$

where

$$(1.3) \quad \mathcal{A}_{\mathcal{D},\mathcal{S}}(\vec{f}) := \sum_{Q \in \mathcal{S}} \prod_{i=1}^m \langle f_i \rangle_Q \mathbf{1}_Q,$$

and the supremum is taken over arbitrary dyadic grids \mathcal{D} and sparse families $\mathcal{S} \in \mathcal{D}$. As an application of this result, it was derived a multilinear analogue of the A_2 theorem, proving that in this more general scenario, a linear bound on the corresponding multiple weight constant also holds. Lately, this result was extended by the third author, K. Moen and W. Sun [24] who proved the sharp bounds for the class of multilinear sparse operators from which follows the sharp bounds for Calderón–Zygmund operators. More precisely, if $\vec{w} = (w_1, \dots, w_m)$ are weights, $1 < p_1, \dots, p_m < \infty$ and p are numbers verifying that $\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m}$ and we denote $\vec{P} = (p_1, \dots, p_m)$,

$$(1.4) \quad \|\mathcal{A}_{\mathcal{S},\mathcal{D}}(\vec{f})\|_{L^p(\nu_{\vec{w}})} \lesssim [\vec{w}]_{A_{\vec{P}}}^{\max(1, \frac{p'_1}{p}, \dots, \frac{p'_m}{p})} \prod_{i=1}^m \|f_i\|_{L^{p_i}(w_i)}.$$

Here $\nu_{\vec{w}} = \prod_{i=1}^m w_i^{p/p_i}$ and the multiple $A_{\vec{P}}$ constant is defined as follows,

$$(1.5) \quad [\vec{w}]_{A_{\vec{P}}} = \sup_Q \left(\frac{1}{|Q|} \int_Q \nu_{\vec{w}} \right) \prod_{i=1}^m \left(\frac{1}{|Q|} \int_Q w_i^{1-p'_i} \right)^{p/p'_i} < \infty.$$

However, the problem of finding the sharp bounds in the multilinear setting for the full range of exponents was still open, since (1.2) does not apply if $1/m < p < 1$, in which case $L^p(\nu_{\vec{w}})$ is not a Banach function space.

Later on, this problem was solved independently by A. Lerner and F. Nazarov [21] and J.M. Conde–Alonso and G. Rey [6]. The main idea in both works was a pointwise control of multilinear Calderón–Zygmund operators by sparse operators avoiding the use of the adjoint operators and duality, which was the key point in Lerner’s original proof.

Another remarkable improvement in [6, 21] was considering weaker regularity conditions on the kernels of Calderón–Zygmund operators. In fact, in both works it was considered the case of log-Dini continuous kernels. Notwithstanding, this pointwise control also holds in the linear setting under the weaker Dini condition, as recently shown by M. Lacey [19] in a qualitative way or, shortly after, by T. Hytönen, L. Roncal and O. Tapiola [16] tracking the precise dependence on the constants.

The aim of this note is two-fold. On one hand, we prove the pointwise control by sparse operators of bilinear Calderón–Zygmund operators with Dini-continuous kernels taking care of the precise constants.

On the other hand, we prove three different mixed bounds for a general class of bilinear dyadic positive operators using the parallel stopping cubes technique. The first bound (see Theorem 4.1) follows the spirit in the work of the third author and W. Sun [25], combining a product of the $A_{\vec{p}}$ and A_{∞} linear constants of the weights involved. The other two mixed weighted bounds combine the multiple $A_{\vec{p}}$ constant with natural extensions of the linear Hrusčev and Fujii–Wilson A_{∞} constants (Theorems 4.2 and 4.4).

As a consequence, we are able to extend these weighted bounds to multilinear Calderón–Zygmund operators with Dini-continuous kernels and obtain new precise weighted bounds for their commutators with BMO functions, square functions and Fourier multipliers in the multiple scenario.

For the sake of simplicity, throughout this paper we are mainly going to consider the bilinear case. Notwithstanding, a similar argument can be used to obtain the general multilinear case. Observe that in the section concerning commutators we give the general proof since it is more convenient.

The organization of this paper is as follows. In Section 2 we give some background and definitions which will be useful to prove our main results. In Section 3 we prove the pointwise control of multilinear Calderón–Zygmund operators by sparse operators whereas in Section 4 we obtain three quantitative bounds for a general class of positive dyadic operators. In Section 5, main results in the previous section are applied to derive mixed weighted bounds for commutators of multilinear Calderón–Zygmund operators as well as for multilinear square functions and Fourier multipliers. Finally, in Section 6 we prove quantitative versions of some classical boundedness results in the multilinear setting.

Throughout this paper, we will denote the average of a function f over a cube Q as

$$(1.6) \quad \langle f \rangle_Q = \int_Q f = \int_Q f(x)dx = \frac{1}{|Q|} \int_Q f(x)dx,$$

where $|Q|$ denotes the Lebesgue measure of Q . If w is a weight, i.e. a measurable locally integrable function defined in \mathbb{R}^n taking values in $(0, \infty)$ for almost every point, we will denote $w(Q) := \int_Q w(x)dx$ and $w\mathbf{1}_Q(x) := w(x)\mathbf{1}_Q(x)$. We will use the notation $A \lesssim B$ to indicate that there is a constant c , independent of the weight constant, such that $A \leq cB$.

2. PRELIMINARIES

2.1. ω -bilinear Calderón–Zygmund operators. We say that T is a ω -bilinear Calderón–Zygmund operator if it is a bilinear operator originally defined on the product of Schwartz spaces and taking values into the space of tempered distributions,

$$(2.1) \quad T : \mathcal{S}(\mathbb{R}^n) \times \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n),$$

and for some $1 \leq q_1, q_2 < \infty$ it extends to a bounded bilinear operator from $L^{q_1} \times L^{q_2}$ to L^q , where $1/q_1 + 1/q_2 = 1/q$, and if there exists a function K , defined off the diagonal $x = y = z$ in $(\mathbb{R}^n)^3$, satisfying

$$(2.2) \quad T(f_1, f_2)(x) = \iint_{(\mathbb{R}^n)^2} K(x, y, z) f_1(y) f_2(z) dy dz,$$

for all $x \notin \text{supp } f_1 \cap \text{supp } f_2$. The kernel K must also satisfy, for some constants $C_K > 0$ and $\tau \in (0, 1)$, the following size condition

$$(2.3) \quad |K(x, y, z)| \leq \frac{C_K}{(|x - y| + |x - z|)^{2n}},$$

and, the smoothness estimate

$$\begin{aligned} & |K(x + h, y, z) - K(x, y, z)| + |K(x, y + h, z) - K(x, y, z)| \\ & + |K(x, y, z + h) - K(x, y, z)| \\ & \leq \frac{1}{(|x - y| + |x - z|)^{2n}} \omega \left(\frac{|h|}{|x - y| + |x - z|} \right), \end{aligned}$$

whenever $|h| \leq \tau \max(|x - y|, |x - z|)$.

If $\omega : [0, \infty) \rightarrow [0, \infty)$ is a modulus of continuity (i.e. it is increasing, subadditive ($\omega(t + s) \leq \omega(t) + \omega(s)$) and $\omega(0) = 0$), the kernel K is said to be a log-Dini-continuous kernel if ω satisfies the following condition

$$(2.4) \quad \|\omega\|_{\log\text{-Dini}} := \int_0^1 \omega(t) \left(1 + \log \left(\frac{1}{t} \right) \right) \frac{dt}{t} < \infty.$$

We are mostly interested in the weaker case when K is a *Dini*(a)-continuous kernel. Namely, when ω satisfies the following condition:

$$(2.5) \quad \|\omega\|_{\text{Dini}(a)} := \int_0^1 \omega^a(t) \frac{dt}{t} < \infty.$$

In the case $a = 1$, we will denote $\|\omega\|_{\text{Dini}(a)}$ simply as $\|\omega\|_{\text{Dini}}$.

Given a bilinear Calderón-Zygmund operator T , the maximal truncation of T is defined as the operator T_{\sharp} given by

$$(2.6) \quad T_{\sharp}(f_1, f_2)(x) = \sup_{\varepsilon > 0} |T_{\varepsilon}(f_1, f_2)(x)|,$$

where T_{ε} is the ε -truncation of T

$$(2.7) \quad T_{\varepsilon}(f_1, f_2)(x) = \int_{|x-y|^2 + |x-z|^2 > \varepsilon^2} K(x, y, z) f_1(y) f_2(z) dy dz.$$

2.2. Dyadic cubes, adjacent systems and sparse operators. The standard system of dyadic cubes in \mathbb{R}^n is the collection \mathcal{D} ,

$$(2.8) \quad \mathcal{D} := \{2^{-k}([0, 1]^n + m) : k \in \mathbb{Z}, m \in \mathbb{Z}^n\},$$

consisting of simple half-open cubes of different length scales with sides parallel to the coordinate axes. These cubes satisfy the following properties:

- (1) for any $Q \in \mathcal{D}$, the sidelength $\ell(Q)$ is of the form 2^k , $k \in \mathbb{Z}$.
- (2) $Q \cap R \in \{Q, R, \emptyset\}$, for any $Q, R \in \mathcal{D}$.
- (3) the cubes of fixed sidelength 2^k form a partition of \mathbb{R}^n .

Since given a ball $B(x, r)$, there does not always exist a cube $Q \in \mathcal{D}$ such that $B(x, r) \subset Q$ and $\ell(Q) \approx r$, a finite number of adjacent dyadic systems \mathcal{D}^u can be used to overcome this problem. More precisely, these dyadic systems are the following

$$(2.9) \quad \mathcal{D}^u := \{2^{-k}([0, 1]^u + m + (-1)^k \frac{1}{3}u) : k \in \mathbb{Z}, m \in \mathbb{Z}^n\}, \quad u \in \{0, 1, 2\}^n.$$

The next two lemmas will be quite useful in the following. The first result can be found in [13, Lemma 2.5] in an stronger version whereas the second result is in [16].

Lemma 2.10. *For any ball $B := B(x, r) \subset \mathbb{R}^n$, there exists a cube $Q_B \in \mathcal{D}^u$ for some $u \in \{0, 1, 2\}^n$ such that $B \subset Q_B$ and $6r < \ell(Q_B) < 12r$.*

Observe that, as a consequence of [13, Lemma 2.5], the collection $\mathcal{D}_0 := \cup_{u \in \{0, 1, 2\}^n} \mathcal{D}^u$ can be seen as a countable approximation of the collection of all balls in \mathbb{R}^n . This family satisfies (1) and (3) listed above, but it satisfies (2) only in various weaker forms. We slightly abuse of the common terminology and say that Q is a dyadic cube if $Q \in \mathcal{D}_0$.

Lemma 2.11. *If $Q_0 \in \cup_{u \in \{0, 1, 2\}^n} \mathcal{D}^u$, then for any ball $B := B(x, r) \subset Q_0$ there exists a cube $Q_B \in \cup_{u \in \{0, 1, 2\}^n} \mathcal{D}^u$ such that $B \subset Q_B \subseteq Q_0$ and $\ell(Q_B) \leq 12r$.*

2.3. Multiple weights. Along this section we recall some basic concepts related to some constants involved in the multiple theory of weights.

First, let us define the central object in the multiple weight theory introduced in [22]. Given $\vec{f} = (f_1, f_2)$, we define the multilinear maximal operator \mathcal{M} by

$$\mathcal{M}(\vec{f})(x) = \sup_{Q \ni x} \prod_{i=1}^m \frac{1}{|Q|} \int_Q |f_i(y_i)| dy_i,$$

where the supremum is taken over all cubes containing x .

Next, let us recall some useful definitions of the basic multiple weight constants that we are using throughout this paper. Consider numbers $1 < p_1, \dots, p_m < \infty$ and p such that $\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m}$ and denote $\vec{P} = (p_1, \dots, p_m)$. Now define

$$(2.12) \quad [w, \vec{\sigma}]_{A_{\vec{P}}} := \sup_Q \langle w \rangle_Q \prod_{i=1}^m \langle \sigma_i \rangle_Q^{\frac{p}{p_i}}.$$

Notice that this definition is more general than that presented in [22], since when $\sigma_i = w_i^{1-p'_i}$, $i = 1, \dots, m$, and $w = \nu_{\vec{w}}$ we recover the $A_{\vec{P}}$ condition in (1.5) if $[w, \vec{\sigma}]_{A_{\vec{P}}} < \infty$.

In [4], Chen and the first author introduced the following multilinear analogue of the A_∞ constant, which was defined by Fujii in [9] and later rediscovered by J.M Wilson [30]. We say that \vec{w} satisfies the $W_{\vec{P}}^\infty$ condition if

$$(2.13) \quad [\vec{w}]_{W_{\vec{P}}^\infty} = \sup_Q \left(\int_Q \prod_{i=1}^m M(w_i \mathbf{1}_Q)^{\frac{p}{p_i}} dx \right) \left(\int_Q \prod_{i=1}^m w_i^{\frac{p}{p_i}} dx \right)^{-1} < \infty.$$

We can also define a more natural multilinear A_∞ constant extending the classical Hruscev A_∞ constant in [11] as follows. We say that \vec{w} satisfies the $H_{\vec{P}}^\infty$ condition if

$$(2.14) \quad [\vec{w}]_{H_{\vec{P}}^\infty} := \sup_Q \prod_{i=1}^m \langle w_i \rangle_Q^{\frac{p}{p_i}} \exp \left(\int_Q \log w_i^{-1} \right)^{\frac{p}{p_i}}.$$

3. DOMINATION THEOREM FOR BILINEAR CZOS

In this section we will prove an extension of the domination theorem due to M. Lacey [19] for bilinear Calderón–Zygmund operators following the scheme of proof in [16] to track the precise constants.

3.1. Some auxiliary operators and a related lemma. Let T be a bilinear Calderón–Zygmund operator with Dini-continuous kernel. For every cube $P \subset \mathbb{R}^n$, we defined the P -localized maximal truncation of T as the operator

$$(3.1) \quad T_{\#,P}(f_1, f_2)(x) := \sup_{0 < \varepsilon < \delta < \frac{1}{2} \text{dist}(x, \partial P)} |T_{\varepsilon, \delta}(f_1, f_2)(x)| \mathbf{1}_P(x),$$

where $T_{\varepsilon, \delta}$ is defined as follows

$$(3.2) \quad T_{\varepsilon, \delta}(f_1, f_2)(x) := \iint_{\varepsilon^2 < |x-y|^2 + |x-z|^2 < \delta^2} K(x, y, z) f_1(y) f_2(z) dz dy.$$

We also need to define a truncated centered bilinear maximal function $\mathcal{M}_{\varepsilon,\delta}^c$ in the following way,

$$(3.3) \quad \mathcal{M}_{\varepsilon,\delta}^c(f_1, f_2)(x) := \sup_{\varepsilon < r < \delta} \prod_{i=1}^2 \langle |f_i| \rangle_{B(x,r)}.$$

We have the following relationship between the truncations $T_{\varepsilon,\delta}$ and $\mathcal{M}_{\varepsilon,\delta}^c$.

Lemma 3.4. *Suppose that $|x - x'| \leq \frac{1}{4}\varepsilon$. Then*

$$(3.5) \quad |T_{\varepsilon,\delta}(f_1, f_2)(x) - T_{\varepsilon,\delta}(f_1, f_2)(x')| \leq c_n(C_K + \|\omega\|_{\text{Dini}})\mathcal{M}_{\varepsilon,2\delta}^c(f_1, f_2)(x).$$

Proof. First observe that

$$\begin{aligned} & |T_{\varepsilon,\delta}(f_1, f_2)(x) - T_{\varepsilon,\delta}(f_1, f_2)(x')| \\ &= \left| \iint_{\varepsilon^2 < |x-y|^2 + |x-z|^2 < \delta^2} K(x, y, z) f_1(y) f_2(z) dz dy \right. \\ &\quad \left. - \iint_{\varepsilon^2 < |x'-y|^2 + |x'-z|^2 < \delta^2} K(x', y, z) f_1(y) f_2(z) dz dy \right| \\ &= \left| \iint_{\varepsilon^2 < |x-y|^2 + |x-z|^2 < \delta^2} (K(x, y, z) - K(x', y, z)) f_1(y) f_2(z) dz dy \right. \\ &\quad \left. + \left(\iint_{\varepsilon^2 < |x-y|^2 + |x-z|^2 < \delta^2} K(x', y, z) f_1(y) f_2(z) dz dy \right. \right. \\ &\quad \left. \left. - \iint_{\varepsilon^2 < |x'-y|^2 + |x'-z|^2 < \delta^2} K(x', y, z) f_1(y) f_2(z) dz dy \right) \right| \\ &:= |I + II|. \end{aligned}$$

For the first term, using the smoothness of the kernel and the properties of the modulus of continuity ω , we get

$$\begin{aligned} |I| &\lesssim \iint_{\varepsilon^2 < |x-y|^2 + |x-z|^2 < \delta^2} \omega\left(\frac{|x-x'|}{|x-y| + |x-z|}\right) \frac{|f_1(y)||f_2(z)|}{(|x-y| + |x-z|)^{2n}} dy dz \\ &\lesssim \sum_{k:\varepsilon^2 \leq (2^k\varepsilon)^2 < \delta^2} \iint_{(2^k\varepsilon)^2 < |x-y|^2 + |x-z|^2 \leq (2^{k+1}\varepsilon)^2} \omega\left(\frac{|x-x'|}{2^k\varepsilon}\right) \frac{|f_1(y)||f_2(z)|}{(2^k\varepsilon)^{2n}} dy dz \\ &\leq \sum_{k=0}^{\infty} \omega\left(\frac{|x-x'|}{2^k\varepsilon}\right) \iint_{B(x, 2^{k+1}\varepsilon)} \frac{|f_1(y)||f_2(z)|}{(2^k\varepsilon)^{2n}} dy dz \\ &\leq c'_n \mathcal{M}_{\varepsilon,2\delta}^c(f_1, f_2)(x) \sum_{k=0}^{\infty} \int_{|x-x'|/2^k\varepsilon}^{|x-x'|/2^{k-1}\varepsilon} \omega(t) \frac{dt}{t} \end{aligned}$$

$$\leq c'_n \mathcal{M}_{\varepsilon, 2\delta}^c(f_1, f_2)(x) \int_0^1 \omega(t) \frac{dt}{t}.$$

For the second term, we make a similar decomposition as in [16], namely

$$II = II_\varepsilon - II_\delta,$$

where

$$\begin{aligned} II_r &:= \left(\iint_{|x-y|^2+|x-z|^2>r^2} - \iint_{|x'-y|^2+|x'-z|^2>r^2} K(x', y, z) f_1(y) f_2(z) dz dy \right) \\ &= \iint_{|x-y|^2+|x-z|^2>r^2 \geq |x'-y|^2+|x'-z|^2} K(x', y, z) f_1(y) f_2(z) dz dy \\ &\quad - \iint_{|x'-y|^2+|x'-z|^2>r^2 \geq |x-y|^2+|x-z|^2} K(x', y, z) f_1(y) f_2(z) dz dy. \end{aligned}$$

Since $|x - x'| \leq \frac{\varepsilon}{4} \leq \frac{r}{4}$, for the first integral, $|x - y| + |x - z| \leq |x' - y| + |x' - z| + 2|x - x'| \leq \sqrt{2}(|x' - y|^2 + |x' - z|^2)^{1/2} + \frac{\varepsilon}{2} \leq \sqrt{2}r + \frac{r}{2} \leq 2r$, and $|x' - y| + |x' - z| \geq |x - y| + |x - z| - 2|x - x'| \geq (|x - y|^2 + |x - z|^2)^{\frac{1}{2}} - 2|x - x'| \geq \frac{r}{2}$, we have

$$\begin{aligned} &\left| \iint_{|x-y|^2+|x-z|^2>r^2 \geq |x'-y|^2+|x'-z|^2} K(x', y, z) f_1(y) f_2(z) dz dy \right| \\ &\leq \iint_{|x-y|+|x-z| \leq 2r} \frac{C_K}{(r/2)^{2n}} |f_1(y)| |f_2(z)| dy dz \\ &\leq c_n C_K \langle |f_1| \rangle_{B(x, 2r)} \langle |f_2| \rangle_{B(x, 2r)} \leq c_n C_K \mathcal{M}_{\varepsilon, 2\delta}^c(f_1, f_2)(x). \end{aligned}$$

For the second integral, we have $|x' - y| + |x' - z| \geq (|x' - y|^2 + |x' - z|^2)^{\frac{1}{2}} \geq r$. Therefore,

$$\begin{aligned} &\left| \iint_{|x'-y|^2+|x'-z|^2>r^2 \geq |x-y|^2+|x-z|^2} K(x', y, z) f_1(y) f_2(z) dz dy \right| \\ &\leq \iint_{|x-y|^2+|x-z|^2 \leq r^2} \frac{C_K}{r^{2n}} |f_1(y)| |f_2(z)| dy dz \\ &\leq c_n C_K \langle |f_1| \rangle_{B(x, r)} \langle |f_2| \rangle_{B(x, r)} \leq c_n C_K \mathcal{M}_{\varepsilon, 2\delta}^c(f_1, f_2)(x). \end{aligned}$$

Consequently,

$$II \leq 4c_n C_K \mathcal{M}_{\varepsilon, 2\delta}^c(f_1, f_2)(x).$$

which shows the desired result. \square

The following result is an extension of the pointwise domination of the maximal truncation of T by a sum of sparse operators in the bilinear setting.

Theorem 3.6. *Let T be a bilinear Calderón–Zygmund operator with Dini continuous kernel. Then for any pair of compactly supported functions*

$f_1, f_2 \in L^1(\mathbb{R}^n)$, there exist sparse collections $\mathcal{S}^u \subset \mathcal{D}^u$, $u = 1, 2, \dots, 3^n$, such that

$$(3.7) \quad T_{\sharp}(f_1, f_2)(x) \leq c_n(\|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q} + C_K + \|\omega\|_{\text{Dini}}) \sum_{u=1}^{3^n} \mathcal{A}_{\mathcal{S}^u}(f_1, f_2)(x),$$

for almost every $x \in \mathbb{R}^n$, where the constant c_n depends only on the dimension and $\|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q}$ denotes the norm of the operator.

The proof of the previous theorem follows exactly the same scheme of proof of [16, Thm. 2.4] with slight modifications. Notwithstanding, since the key ingredient for the proof of this theorem is essentially the next lemma, we are only going to give the details of its proof here for the sake of completeness.

Lemma 3.8. *Let f_1, f_2 be integrable functions. Then, for every $Q_0 \in \mathcal{D}_0$, there exists a collection $\mathcal{D}(Q_0)$ of dyadic cubes $Q \subset Q_0$ such that the following three conditions hold:*

- (1) $\sum_{Q \in \mathcal{D}(Q_0)} |Q| \leq \varepsilon_n |Q_0|$.
- (2) if $Q' \subset Q$, and $Q', Q \in \mathcal{D}(Q_0)$, then $Q' = Q$.
- (3) we have

$$(3.9) \quad T_{\sharp, Q_0}(f_1, f_2) \leq C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0} \mathbf{1}_{Q_0} + \max_{Q \in \mathcal{D}(Q_0)} T_{\sharp, Q}(f_1, f_2),$$

where $C_T^0 := c_n(\|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q} + C_K + \|\omega\|_{\text{Dini}})$ and $\varepsilon_n \searrow 0$ as $c_n \nearrow \infty$.

Proof. We want to prove that for any constant $C_T^0 > 0$ we can cover the set E_0 ,

$$E_0 := \left\{ x \in Q_0 : T_{\sharp, Q_0}(f_1, f_2)(x) > C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0} \right\},$$

with countably many cubes $Q_i \in \mathcal{D}_0$ that satisfying conditions (2) and (3) and if the constant C_T^0 is of the form $c_n(\|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q} + C_K + \|\omega\|_{\text{Dini}})$, then the cubes also satisfy condition (1).

Let $x \in E_0$. Since the function $(\varepsilon, \delta) \mapsto T_{\varepsilon, \delta}(f_1, f_2)(x)$ is continuous, we can choose such radii $0 < \sigma_x < \tau_x \leq \frac{1}{2} \cdot \text{dist}(x, \partial Q_0)$ that

$$|T_{\sigma_x, \tau_x}(f_1, f_2)(x)| \geq C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0}$$

and

$$|T_{\sigma, \tau}(f_1, f_2)(x)| \leq C_T^0 \prod_{i=1}^2 \langle |f_i| \rangle_{Q_0} \quad \text{if } \sigma_x \leq \sigma \leq \tau \leq \frac{1}{2} \cdot \text{dist}(x, Q_0).$$

For simplicity, we drop the conditions $\varepsilon > 0$ and $\delta \leq \frac{1}{2} \cdot \text{dist}(x, Q_0)$ from the notation. Now the maximality of σ_x implies the following:

$$\begin{aligned} T_{\sharp, Q_0}(f_1, f_2)(x) &= \sup_{\varepsilon \leq \delta} |T_{\varepsilon, \delta}(f_1, f_2)(x)| \\ &= \sup_{\varepsilon \leq \delta \leq \sigma_x} |T_{\varepsilon, \delta}(f_1, f_2)(x)| \\ &\vee \sup_{\sigma_x \leq \varepsilon \leq \delta} |T_{\varepsilon, \delta}(f_1, f_2)(x)| \\ &\vee \sup_{\varepsilon \leq \sigma_x \leq \delta} |T_{\varepsilon, \delta}(f_1, f_2)(x)| \\ &=: I \vee II \vee III, \end{aligned}$$

where

$$III = \sup_{\varepsilon \leq \sigma_x \leq \delta} |T_{\varepsilon, \sigma_x}(f_1, f_2)(x) + T_{\sigma_x, \delta}(f_1, f_2)(x)| \leq I + II,$$

and $II \leq C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0}$ by definition. So altogether we find that

$$(3.10) \quad T_{\sharp, Q_0}(f_1, f_2)(x) \leq \sup_{\varepsilon \leq \delta \leq \sigma_x} |T_{\varepsilon, \delta}(f_1, f_2)(x)| + C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0} \quad \forall x \in E_0,$$

which is a preliminary version of the pointwise domination result we are proving. Now we can use Lemma 2.11 to get from the preliminary version to the desired estimate. Since $B(x, 2\sigma_x) \subset Q_0$ for every $x \in E_0$, there exists a cube $Q_x \in \mathcal{D}_0$ such that $B(x, 2\sigma_x) \subset Q_x \subset Q_0$ and $\ell(Q_x) \leq 12 \cdot 2\sigma_x$ for every $x \in E_0$. Let $(Q_i)_i$ be the sequence of such cubes Q_x that are maximal with respect to inclusion, that is, for each Q_i there does not exist $R \in \mathcal{D}_0$ such that $Q_i \subsetneq R \subseteq Q_0$. Then for every $x \in E_0$ we have

$$\begin{aligned} T_{\sharp, Q_0}(f_1, f_2)(x) &\stackrel{(3.10)}{\leq} \sup_{0 < \varepsilon \leq \delta \leq \sigma_x} |T_{\varepsilon, \delta}(f_1, f_2)(x)| + C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0} \\ &\leq \sup_{0 < \varepsilon \leq \delta \leq \frac{1}{2} \cdot \text{dist}(x, \partial Q_x)} |T_{\varepsilon, \delta}(f_1, f_2)(x)| + C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0} \\ &= T_{\sharp, Q_x}(f_1, f_2)(x) + C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0} \\ &\leq \max_i T_{\sharp, Q_i}(f_1, f_2)(x) + C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0} \end{aligned}$$

and for every $x \in Q_0 \setminus E_0$ we have $T_{\sharp, Q_0}(f_1, f_2)(x) \leq C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0}$ by definition. Thus, the cubes Q_i satisfy Lacey's conditions (2) and (3) and to complete the proof, we only need to show that with a suitable choice of C_T^0

the cubes also satisfy property (1). Let us split the set E_0 into two parts:

$$E_1 := \{x \in E_0 : \mathcal{M}_{\sigma_x, 2\tau_x}(f_1, f_2)(x) \leq C_T^1 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0}\}, \quad E_2 := E_0 \setminus E_1,$$

where C_T^1 is a constant whose value we will fix in the next step. Then, for $x \in E_1$ and $x' \in B(x, \frac{1}{4}\sigma_x)$, we have

$$\begin{aligned} |T_{\sigma_x \tau_x}(f_1, f_2)(x') - T_{\sigma_x \tau_x}(f_1, f_2)(x)| &\stackrel{3.4}{\leq} c_n(C_K + \|\omega\|_{\text{Dini}}) \mathcal{M}_{\sigma_x, 2\tau_x}^c(f_1, f_2)(x) \\ &\leq c_n(C_K + \|\omega\|_{\text{Dini}}) C_T^1 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0} \\ &= \frac{1}{2} C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0}, \end{aligned}$$

provided that we choose

$$C_T^1 := \frac{C_T^0}{2c_n(C_K + \|\omega\|_{\text{Dini}})}.$$

Then, since $x \in E_1 \subseteq E_0$, it follows that

$$\begin{aligned} T_{\sharp}(\mathbf{1}_{Q_0} f_1, \mathbf{1}_{Q_0} f_2)(x') &\geq |T_{\sigma_x, \tau_x}(f_1, f_2)(x')| \\ &\geq |T_{\sigma_x, \tau_x}(f_1, f_2)(x)| - \frac{1}{2} C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0} \\ &> \frac{1}{2} C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0} \end{aligned}$$

for all $x' \in B(x, \frac{1}{4}\sigma_x)$. In particular,

$$\begin{aligned} \left| \bigcup_{x \in E_1} B(x, \frac{1}{4}\sigma_x) \right|^2 &\leq \left| \{T_{\sharp}(\mathbf{1}_{Q_0} f_1, \mathbf{1}_{Q_0} f_2) > \frac{1}{2} C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0}\} \right|^2 \\ &\leq \frac{\|T_{\sharp}\|_{L^1 \times L^1 \rightarrow L^{1/2, \infty}}^m}{\frac{1}{2} C_T^0 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0}} \prod_{i=1}^m \|\mathbf{1}_{Q_0} f_i\|_{L^1} \\ &= \frac{2\|T_{\sharp}\|_{L^1 \times L^1 \rightarrow L^{1/2, \infty}}}{C_T^0} |Q_0|^2 \end{aligned}$$

by the weak inequality of T_{\sharp} .

Let us then show that with this choice of C_T^1 and a suitable choice of C_T^0 the size of E_2 is controlled. Let $x \in E_2$. By definition, we can choose some $\rho_x \in [\sigma_x, 2\tau_x]$ such that

$$\prod_{j=1}^2 \int_{B(x, \rho_x)} |f_j(y_j)| dy_j > C_T^1 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0}.$$

Since $\tau_x \leq \frac{1}{2} \cdot \text{dist}(x, \partial Q_0)$, we know that $B(x, 2\rho_x) \subset Q_0$. In particular,

$$\mathcal{M}(\mathbf{1}_{Q_0} f_1, \mathbf{1}_{Q_0} f_2)(x') > C_T^1 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0}$$

for all $x' \in B(x, \rho_x)$, where \mathcal{M} is the noncentered bilinear maximal operator

$$\mathcal{M}(f_1, f_2)(x) := \sup_{B \ni x} \prod_{j=1}^2 \int_B |f_j| dx.$$

Thus

$$\begin{aligned} & \left| \bigcup_{x \in E_2} B(x, \frac{1}{4}\sigma_x) \right|^2 \\ & \leq \left| \bigcup_{x \in E_2} B(x, \rho_x) \right|^2 \leq |\{ \mathcal{M}(\mathbf{1}_{Q_0} f_1, \mathbf{1}_{Q_0} f_2) > C_T^1 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0} \}|^2 \\ & \leq \frac{c_n}{C_T^1 \prod_{j=1}^2 \langle |f_j| \rangle_{Q_0}} \prod_{j=1}^2 \|\mathbf{1}_{Q_0} f_j\|_{L^1} = \frac{c'_n (C_K + \|\omega\|_{\text{Dini}})}{C_T^0} |Q_0|^2. \end{aligned}$$

by the weak inequality of the bilinear maximal operator.

Finally, let us combine all the previous calculations. For every maximal cube Q_i , let $x_i \in E_0$ be a point such that $Q_i = Q_{x_i}$. Then, since $\ell(Q_x) \leq 12 \cdot 2\sigma_x$ for each $x \in E_0$, we have $|Q_{x_i}| \leq c_n |B(x_i, \frac{1}{4}\sigma_{x_i})|$ for every i . In particular, since the cubes in the collection $\{Q_{x_i} : Q_{x_i} \in \mathcal{D}^u\}$ are pairwise disjoint for a fixed $u \in \{0, 1, 2\}^n$ and $B(x_i, 2\sigma_{x_i}) \subset Q_{x_i}$, $B(x_i, \frac{1}{4}\sigma_{x_i})$ are pairwise disjoint and therefore,

$$\begin{aligned} \sum_i |Q_{x_i}| &= \sum_{u \in \{0, 1, 2\}^n} \sum_{i: Q_{x_i} \in \mathcal{D}^u} |Q_{x_i}| \\ &\leq c_n \sum_{u \in \{0, 1, 2\}^n} \sum_{i: Q_{x_i} \in \mathcal{D}^u} |B(x_i, \frac{1}{4}\sigma_{x_i})| \\ &= c_n \sum_{u \in \{0, 1, 2\}^n} \left| \bigcup_{i: Q_{x_i} \in \mathcal{D}^u} B(x_i, \frac{1}{4}\sigma_{x_i}) \right| \\ &\leq 3^n c_n \left(\left| \bigcup_{x \in E_1} B(x, \frac{1}{4}\sigma_x) \right| + \left| \bigcup_{x \in E_2} B(x, \rho_x) \right| \right) \\ &\leq c'_n \left(\frac{\|T_\# \|_{L^{q_1} \times L^{q_2} \rightarrow L^q} + C_K + \|\omega\|_{\text{Dini}}}{C_T^0} \right)^{\frac{1}{2}} |Q_0|. \end{aligned}$$

Hence, if

$$C_T^0 = c_n (C_K + \|\omega\|_{\text{Dini}} + \|T_\# \|_{L^{q_1} \times L^{q_2} \rightarrow L^q}),$$

then the cubes Q_i satisfy property (1). □

4. QUANTITATIVE BOUNDS FOR BILINEAR SPARSE OPERATORS

In this section we establish three different bounds for the family of bi-sublinear sparse operators $\mathcal{A}_{p_0, \gamma, \mathcal{S}}$. As a consequence of the domination theorem proved in the previous section, we will obtain the same bounds for bilinear Calderón-Zygmund operators or any other class of operators which can be controlled by this class of positive dyadic operators. For $\gamma > 0$, $p_0 \geq 1$, we define $\mathcal{A}_{p_0, \gamma, \mathcal{S}}$ as follows,

$$\mathcal{A}_{p_0, \gamma, \mathcal{S}}(\vec{f})(x) := \left(\sum_{Q \in \mathcal{S}} \left[\prod_{i=1}^2 \langle f_i \rangle_{Q, p_0} \right]^\gamma \mathbf{1}_Q(x) \right)^{1/\gamma},$$

where for any cube Q ,

$$\langle f \rangle_{Q, p_0} := \left(\frac{1}{|Q|} \int_Q |f(x)|^{p_0} dx \right)^{\frac{1}{p_0}}.$$

Throughout this section we will use the following notation, $\vec{P} = (\frac{p_1}{p_0}, \frac{p_2}{p_0})$.

Let us state our main results in this section. Our first bound is a mixed $A_{\vec{P}}-A_\infty$ estimate.

Theorem 4.1. *Let $\gamma > 0$. Suppose that $p_0 < p_1, p_2 < \infty$ with $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$. Let w and $\vec{\sigma}$ be weights satisfying that $[w, \vec{\sigma}]_{A_{\vec{P}/p_0}} < \infty$ and $w, \sigma_i \in A_\infty$ for $i = 1, 2$. If $\gamma \geq p_0$, then*

$$\begin{aligned} & \|\mathcal{A}_{p_0, \gamma, \mathcal{S}}(\cdot \sigma_1, \cdot \sigma_2)\|_{L^{p_1}(\sigma_1) \times L^{p_2}(\sigma_2) \rightarrow L^p(w)} \\ & \lesssim [w, \vec{\sigma}]_{A_{\vec{P}/p_0}}^{\frac{1}{p}} \left(\prod_{i=1}^2 [\sigma_i]_{A_\infty}^{\frac{1}{p_i}} + [w]_{A_\infty}^{(\frac{1}{\gamma} - \frac{1}{p})_+} \sum_{j=1}^2 \prod_{i \neq j} [\sigma_i]_{A_\infty}^{\frac{1}{p_i}} \right), \end{aligned}$$

where

$$\left(\frac{1}{\gamma} - \frac{1}{p} \right)_+ := \max \left\{ \frac{1}{\gamma} - \frac{1}{p}, 0 \right\}.$$

If $\gamma < p_0$, then the above result still holds for all $p > \gamma$.

Our second result is a mixed bound combining the $A_{\vec{P}}$ constant and a generalization of the Fujii-Wilson A_∞ constant to the bilinear setting which was introduced in [4].

Theorem 4.2. *Let $\gamma > 0$. Suppose that $p_0 < p_1, p_2 < \infty$ with $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$ and set $q = p/\gamma$. Let w and $\vec{\sigma}$ be weights satisfying that $[w, \vec{\sigma}]_{A_{\vec{P}/p_0}} < \infty$. If $\gamma \geq p_0$, then*

$$(4.3) \quad \begin{aligned} & \|\mathcal{A}_{p_0, \gamma, \mathcal{S}}(\cdot \sigma_1, \cdot \sigma_2)\|_{L^{p_1}(\sigma_1) \times L^{p_2}(\sigma_2) \rightarrow L^p(w)} \\ & \leq [w, \vec{\sigma}]_{A_{\vec{P}/p_0}}^{1/p} \left([\vec{\sigma}]_{W_{\vec{P}}^\infty}^{1/p} + \sum_{i=1}^2 [\vec{\sigma}^i]_{W_{\vec{P}^i}^\infty}^{1/\gamma(\frac{p_i}{\gamma})'} \right), \end{aligned}$$

where $[\bar{\sigma}^i]_{W_{\bar{P}^i}^\infty} = 1$ if $p \leq \gamma$ and otherwise,

$$[\bar{\sigma}^i]_{W_{\bar{P}^i}^\infty} = \sup_Q \left(\int_Q M(\mathbf{1}_Q w)^{\frac{(p_i/\gamma)'}{q'}} \prod_{j \neq i} M(\mathbf{1}_Q \sigma_j)^{\frac{(p_i/\gamma)'}{p_j/\gamma}} dx \right) \times \left(\int_Q w^{\frac{(p_i/\gamma)'}{q'}} \prod_{j \neq i} \sigma_j^{\frac{(p_i/\gamma)'}{p_j/\gamma}} dx \right)^{-1}.$$

If $\gamma < p_0$, then the above result still holds for all $p > \gamma$.

Finally, we give a mixed bound combining the $A_{\bar{P}}$ constant and a generalization of the Hrusčev A_∞ constant to the bilinear setting.

Theorem 4.4. *Let $\gamma > 0$. Suppose that $p_0 < p_1, p_2 < \infty$ with $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$ and set $q = p/\gamma$. Let w and $\bar{\sigma}$ be weights satisfying that $[w, \bar{\sigma}]_{A_{\bar{P}/p_0}} < \infty$. If $\gamma \geq p_0$, then*

(4.5)

$$\|\mathcal{A}_{p_0, \gamma, \mathcal{S}}(\cdot \sigma_1, \cdot \sigma_2)\|_{L^{p_1}(\sigma_1) \times L^{p_2}(\sigma_2) \rightarrow L^p(w)} \leq [w, \bar{\sigma}]_{A_{\bar{P}/p_0}}^{\frac{1}{p}} \left([\bar{\sigma}]_{H_{\bar{P}}^\infty}^{1/p} + \sum_{i=1}^2 [\bar{\sigma}^i]_{H_{\bar{P}^i}^\infty}^{1/p'_i} \right),$$

where $[\bar{\sigma}^i]_{H_{\bar{P}^i}^\infty} = 1$ if $p \leq \gamma$ and otherwise,

$$[\bar{\sigma}^i]_{H_{\bar{P}^i}^\infty} = \sup_Q \langle w \rangle_Q^{p'_i(\frac{1}{\gamma} - \frac{1}{p})_+} \exp \left(\int_Q \log w^{-1} \right)^{p'_i(\frac{1}{\gamma} - \frac{1}{p})_+} \times \prod_{j \neq i} \langle \sigma_i \rangle_Q^{p'_i/p_j} \exp \left(\int_Q \log \sigma_i^{-1} \right)^{p'_i/p_j}.$$

If $\gamma < p_0$, then the above result still holds for all $p > \gamma$.

Before proving Theorems 4.1, 4.2 and 4.4, we need the following two results. The first proposition can be found in [3, Proposition 2.2].

Proposition 4.7. *Let $1 < s < \infty$, σ be a positive Borel measure and*

$$\phi = \sum_{Q \in \mathcal{D}} \alpha_Q \mathbf{1}_Q, \quad \phi_Q = \sum_{Q' \subset Q} \alpha_{Q'} \mathbf{1}_{Q'}.$$

Then

$$\|\phi\|_{L^s(\sigma)} \sim \left(\sum_{Q \in \mathcal{D}} \alpha_Q (\langle \phi_Q \rangle_Q^\sigma)^{s-1} \sigma(Q) \right)^{1/s}.$$

The following proposition follows the same spirit as that in [14] and it allows us to avoid the ‘‘slicing’’ argument. Namely, the separate consideration of families of cubes with the $A_{\bar{P}}$ characteristic ‘‘frozen’’ to a certain value $\langle w \rangle_Q \prod_{i=1}^2 \langle \sigma_i \rangle_Q^{p/p'_i} \sim 2^k$.

By using Proposition 4.7, it is also possible to give an alternative proof of our main results by using the outer measure theory studied in [8, 29]. Notice that here the stopping cubes method provides a more direct proof.

Proposition 4.8. *Let \mathcal{S} be a sparse family and $0 \leq \gamma, \eta < 1$ satisfying $\gamma + \eta < 1$. Then*

$$(4.9) \quad \sum_{\substack{Q \in \mathcal{S} \\ Q \subset R}} \langle u \rangle_Q^\gamma \langle v \rangle_Q^\eta |Q| \lesssim \langle u \rangle_R^\gamma \langle v \rangle_R^\eta |R|.$$

Proof. Indeed, set $1/r := \gamma + \eta$, $1/s := \gamma + (1 - 1/r)/2$ and $1/s' := 1 - 1/s$. By sparseness and Kolmogorov's inequality, we have

$$\begin{aligned} \sum_{\substack{Q \in \mathcal{S} \\ Q \subset R}} \langle u \rangle_Q^\gamma \langle v \rangle_Q^\eta |Q| &\leq 2 \sum_{\substack{Q \in \mathcal{S} \\ Q \subset R}} \langle u \rangle_Q^\gamma \langle v \rangle_Q^\eta |E_Q| \\ &\leq 2 \int_R M(u \mathbf{1}_R)^\gamma M(v \mathbf{1}_R)^\eta dx \\ &\leq 2 \left(\int_R M(u \mathbf{1}_R)^{s\gamma} \right)^{1/s} \left(\int_R M(v \mathbf{1}_R)^{s'\eta} \right)^{1/s'} \\ &\lesssim \langle u \rangle_R^\gamma |R|^{1/s} \langle v \rangle_R^\eta |R|^{1/s'} = \langle u \rangle_R^\gamma \langle v \rangle_R^\eta |R|. \end{aligned}$$

□

Our first observation is that we can reduce the problem to study the case of $p_0 = 1$. Indeed, consider the two weight norm inequality

$$(4.10) \quad \|\mathcal{A}_{p_0, \gamma, \mathcal{S}}(f_1, f_2)\|_{L^p(w)} \leq \mathcal{N}(\vec{P}, p_0, \gamma, w, \vec{\sigma}) \|f_1\|_{L^{p_1}(w_1)} \|f_2\|_{L^{p_2}(w_2)},$$

where we use $\mathcal{N}(\vec{P}, p_0, \gamma, w, \vec{\sigma})$ to denote the best constant such that (4.10) holds. Rewrite (4.10) as

$$\|\mathcal{A}_{p_0, \gamma, \mathcal{S}}(f_1^{p_0}, f_2^{p_0})\|_{L^p(w)}^{p_0} \leq \mathcal{N}(\vec{P}, p_0, \gamma, w, \vec{\sigma})^{p_0} \|f_1^{p_0}\|_{L^{p_1}(w_1)}^{p_0} \|f_2^{p_0}\|_{L^{p_2}(w_2)}^{p_0},$$

which is equivalent to the following

$$\|\mathcal{A}_{1, \frac{\gamma}{p_0}, \mathcal{S}}(f_1, f_2)\|_{L^{p/p_0}(w)} \leq \mathcal{N}(\vec{P}, p_0, \gamma, w, \vec{\sigma})^{p_0} \|f_1\|_{L^{p_1/p_0}(w_1)} \|f_2\|_{L^{p_2/p_0}(w_2)}.$$

Therefore, if we denote by $\mathcal{N}(\vec{P}, \gamma, w, \sigma)$ the best constant for the case $p_0 = 1$, then the best constant for general p_0 would be $\mathcal{N}(\vec{P}/p_0, \gamma/p_0, w, \sigma)^{1/p_0}$. Therefore, it suffices to study the case of $p_0 = 1$.

Our second observation can be stated as follows, as it was done in [14, 23].

Lemma 4.11. *Suppose that $p > \gamma$. Let \mathcal{N} denote the best constant such that the following inequality holds*

$$(4.12) \quad \|\mathcal{A}_{1, \gamma, \mathcal{S}}(f_1 \sigma_1, f_2 \sigma_2)\|_{L^p(w)} \leq \mathcal{N} \|f_1\|_{L^{p_1}(\sigma_1)} \|f_2\|_{L^{p_2}(\sigma_2)}.$$

Then (4.12) is equivalent to the following inequality with $\mathcal{N}' \simeq \mathcal{N}^\gamma$

$$(4.13) \quad \left\| \left(\sum_{Q \in \mathcal{S}} \langle f_1 \rangle_Q^{\sigma_1} \langle f_2 \rangle_Q^{\sigma_2} \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma \mathbf{1}_Q \right)^{\frac{1}{\gamma}} \right\|_{L^p(w)}^\gamma \leq \mathcal{N}' \|f_1\|_{L^{\frac{p_1}{\gamma}}(\sigma_1)} \|f_2\|_{L^{\frac{p_2}{\gamma}}(\sigma_2)}.$$

Proof. On one hand, if (4.13) holds, we have

$$\begin{aligned}
& \|\mathcal{A}_{1,\gamma,\mathcal{S}}(f_1\sigma_1, f_2\sigma_2)\|_{L^p(w)} \\
& \leq \left\| \left(\sum_{Q \in \mathcal{S}} \langle M_{\mathcal{D}}^{\sigma_1}(f_1)^\gamma \rangle_Q^{\sigma_1} \langle M_{\mathcal{D}}^{\sigma_2}(f_2)^\gamma \rangle_Q^{\sigma_2} \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma \mathbf{1}_Q \right)^{\frac{1}{\gamma}} \right\|_{L^p(w)} \\
& \lesssim \mathcal{N} \|M_{\mathcal{D}}^{\sigma_1}(f_1)^\gamma\|_{L^{p_1/\gamma}(\sigma_1)}^{1/\gamma} \|M_{\mathcal{D}}^{\sigma_2}(f_2)^\gamma\|_{L^{p_2/\gamma}(\sigma_2)}^{1/\gamma} \\
& \leq \mathcal{N} \|f_1\|_{L^{p_1}(\sigma_1)} \|f_2\|_{L^{p_2}(\sigma_2)},
\end{aligned}$$

where $M_{\mathcal{D}}^\sigma$ denotes the dyadic weighted maximal function, namely

$$(4.14) \quad M_{\mathcal{D}}^\sigma(f) = \sup_{Q \in \mathcal{D}} \frac{1}{\sigma(Q)} \int_Q |f(x)| \sigma dx,$$

which is bounded from $L^p(\sigma)$ into itself for every $p > 1$. On the other hand, if (4.12) holds, we have

$$\begin{aligned}
& \left\| \left(\sum_{Q \in \mathcal{S}} \langle f_1 \rangle_Q^{\sigma_1} \langle f_2 \rangle_Q^{\sigma_2} \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma \mathbf{1}_Q \right)^{\frac{1}{\gamma}} \right\|_{L^p(w)} \\
& \leq \left\| \left(\sum_{Q \in \mathcal{S}} (\langle M_{\gamma,\mathcal{D}}^{\sigma_1}(f_1^{1/\gamma}) \rangle_Q^{\sigma_1})^\gamma (\langle M_{\gamma,\mathcal{D}}^{\sigma_2}(f_2^{1/\gamma}) \rangle_Q^{\sigma_2})^\gamma \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma \mathbf{1}_Q \right)^{\frac{1}{\gamma}} \right\|_{L^p(w)} \\
& \leq \mathcal{N} \|M_{\gamma,\mathcal{D}}^{\sigma_1}(f_1^{1/\gamma})\|_{L^{p_1}(\sigma_1)} \|M_{\gamma,\mathcal{D}}^{\sigma_2}(f_2^{1/\gamma})\|_{L^{p_2}(\sigma_2)} \\
& \lesssim \mathcal{N} \|f_1^{1/\gamma}\|_{L^{p_1}(\sigma_1)} \|f_2^{1/\gamma}\|_{L^{p_2}(\sigma_2)},
\end{aligned}$$

where $M_{\gamma,\mathcal{D}}^\sigma(f) = (M_{\mathcal{D}}^\sigma(f^\gamma))^{1/\gamma}$ and we have used in the last step that $p > \gamma$, which implies $p_1, p_2 > \gamma$ and consequently, the boundedness of the maximal functions. \square

Therefore, we further reduce the problem to study (4.13). Finally, we give the following lemma, which is the key to prove Theorems 4.1, 4.2 and 4.4.

Lemma 4.15. *Let $\gamma > 0$. Suppose that $1 < p_1, p_2 < \infty$ with $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$. Let w and $\vec{\sigma}$ be weights. Then for any sparse collection \mathfrak{S} ,*

$$(4.16) \quad \left\| \left(\sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma \mathbf{1}_Q \right)^{\frac{1}{\gamma}} \right\|_{L^p(w)} \leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{1}{p}} \left(\sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^{\frac{p}{p_1}} \langle \sigma_2 \rangle_Q^{\frac{p}{p_2}} |Q| \right)^{1/p},$$

and if $p > \gamma$, then there holds

$$(4.17) \quad \begin{aligned} & \left\| \sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^{\gamma-1} \langle w \rangle_Q \mathbf{1}_Q \right\|_{L^{(\frac{p_2}{\gamma})'}(\sigma_2)} \\ & \leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} \left(\sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^{\frac{\gamma(\frac{p_2}{\gamma})'}{p_1}} \langle w \rangle_Q^{(\frac{p_2}{\gamma})'(1-\frac{\gamma}{p})} |Q| \right)^{\frac{1}{(\frac{p_2}{\gamma})'}} \end{aligned}$$

and

$$(4.18) \quad \left\| \sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^{\gamma-1} \langle \sigma_2 \rangle_Q^\gamma \langle w \rangle_Q \mathbf{1}_Q \right\|_{L^{(\frac{p_1}{\gamma})'(\sigma_1)}} \\ \leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} \left(\sum_{Q \in \mathfrak{S}} \langle \sigma_2 \rangle_Q^{\frac{\gamma(\frac{p_1}{\gamma})'}{p_2}} \langle w \rangle_Q^{(\frac{p_1}{\gamma})'(1-\frac{\gamma}{p})} |Q| \right)^{\frac{1}{(\frac{p_1}{\gamma})'}}.$$

Proof. We start proving (4.16). Following the spirit in [14], observe that the right hand side of (4.16) is independent of γ . Therefore, it suffices to study the problem for small γ . More precisely, for fixed \vec{P} , we reduce the problem to study the case $\gamma < \min\{p, 1\}$ with $(p/\gamma)' < \max\{p_1, p_2\}$. Without loss of generality, we may assume that $(p/\gamma)' < p_1 = \max\{p_1, p_2\}$. Then it is easy to check that

$$(4.19) \quad 0 \leq \gamma - \frac{\gamma p_1'}{p_2'} < 1, \quad 0 \leq 1 - \frac{\gamma p_1'}{p} < 1,$$

and

$$(4.20) \quad \gamma - \frac{\gamma p_1'}{p_2'} + 1 - \frac{\gamma p_1'}{p} < 1.$$

By Proposition 4.7, we have

$$\begin{aligned} & \left\| \left(\sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma \mathbf{1}_Q \right)^{1/\gamma} \right\|_{L^p(w)} \\ & \approx \left(\sum_{Q \in \mathfrak{S}} \lambda_Q \left(\frac{1}{w(Q)} \sum_{Q' \subset Q} \langle \sigma_1 \rangle_{Q'}^\gamma \langle \sigma_2 \rangle_{Q'}^\gamma w(Q') \right)^{\frac{p-1}{\gamma}} \right)^{\frac{1}{p}} \\ & \lesssim [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{(p-\gamma)p_1'}{p^2}} \left(\sum_{Q \in \mathfrak{S}} \lambda_Q \left(\frac{1}{w(Q)} \sum_{Q' \subset Q} \langle \sigma_2 \rangle_{Q'}^{\gamma(1-\frac{p_1'}{p_2'})} \langle w \rangle_{Q'}^{1-\frac{p_1'\gamma}{p}} |Q'| \right)^{\frac{p-1}{\gamma}} \right)^{\frac{1}{p}} \\ & \stackrel{(4.9)}{\lesssim} [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{(p-\gamma)p_1'}{p^2}} \left(\sum_{Q \in \mathfrak{S}} \lambda_Q \left(\frac{1}{w(Q)} \langle \sigma_2 \rangle_Q^{\gamma(1-\frac{p_1'}{p_2'})} \langle w \rangle_Q^{1-\frac{p_1'\gamma}{p}} |Q| \right)^{\frac{p-1}{\gamma}} \right)^{\frac{1}{p}} \\ & = [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{(p-\gamma)p_1'}{p^2}} \left(\sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^{\gamma+(1-\frac{p_1'}{p_2'})(p-\gamma)} \langle w \rangle_Q^{1-\frac{p_1'(p-\gamma)}{p}} |Q| \right)^{1/p} \\ & \lesssim [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{(p-\gamma)p_1'}{p^2} + \frac{1}{p} - \frac{(p-\gamma)p_1'}{p^2}} \left(\sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^{\frac{p}{p_1}} \langle \sigma_2 \rangle_Q^{\frac{p}{p_2}} |Q| \right)^{1/p} \\ & = [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{1}{p}} \left(\sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^{\frac{p}{p_1}} \langle \sigma_2 \rangle_Q^{\frac{p}{p_2}} |Q| \right)^{1/p}, \end{aligned}$$

where $\lambda_Q = \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma w(Q)$. By symmetry, we only need to prove (4.17). Let us consider the case $(p/\gamma)' \geq \max\{p_1, p_2\}$ and $(p/\gamma)' < \max\{p_1, p_2\}$ separately. For the case $(p/\gamma)' < \max\{p_1, p_2\}$, without loss of generality, we

may assume that $p_1 > p_2$. Again, having into account (4.19) and (4.20) and using Proposition 4.7, we obtain

$$\begin{aligned}
& \left\| \sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^{\gamma-1} \langle w \rangle_Q \mathbf{1}_Q \right\|_{L^{(\frac{p_2}{\gamma})'}(\sigma_2)} \\
& \simeq \left(\sum_{Q \in \mathfrak{S}} \lambda_Q \left(\frac{1}{\sigma_2(Q)} \sum_{Q' \subset Q} \langle \sigma_1 \rangle_{Q'}^\gamma \langle \sigma_2 \rangle_{Q'}^\gamma w(Q') \right)^{(\frac{p_2}{\gamma})'-1} \right)^{\frac{1}{(\frac{p_2}{\gamma})'}} \\
& \leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{p_1' \gamma^2}{pp_2}} \left(\sum_{Q \in \mathfrak{S}} \lambda_Q \left(\frac{1}{\sigma_2(Q)} \sum_{Q' \subset Q} \langle \sigma_2 \rangle_{Q'}^{\gamma(1-\frac{p_1'}{p_2})} \langle w \rangle_{Q'}^{1-\frac{\gamma p_1'}{p}} |Q'| \right)^{(\frac{p_2}{\gamma})'-1} \right)^{\frac{1}{(\frac{p_2}{\gamma})'}} \\
& \stackrel{(4.9)}{\lesssim} [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{p_1' \gamma^2}{pp_2}} \left(\sum_{Q \in \mathfrak{S}} \lambda_Q \left(\frac{1}{\sigma_2(Q)} \langle \sigma_2 \rangle_Q^{\gamma(1-\frac{p_1'}{p_2})} \langle w \rangle_Q^{1-\frac{\gamma p_1'}{p}} |Q| \right)^{(\frac{p_2}{\gamma})'-1} \right)^{\frac{1}{(\frac{p_2}{\gamma})'}} \\
& = [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{p_1' \gamma^2}{pp_2}} \left(\sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^{\gamma(\frac{p_2}{\gamma})' - (\frac{\gamma p_1'}{p_2} + 1)((\frac{p_2}{\gamma})'-1)} \langle w \rangle_Q^{(\frac{p_2}{\gamma})' - \frac{\gamma p_1'}{p}((\frac{p_2}{\gamma})'-1)} |Q| \right)^{\frac{1}{(\frac{p_2}{\gamma})'}} \\
& \leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} \left(\sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^{\frac{\gamma(\frac{p_2}{\gamma})'}{p_1}} \langle w \rangle_Q^{(\frac{p_2}{\gamma})'(1-\frac{\gamma}{p})} |Q| \right)^{\frac{1}{(\frac{p_2}{\gamma})'}},
\end{aligned}$$

where $\lambda_Q = \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma w(Q)$. It remains to consider when $(p/\gamma)' \geq \max\{p_1, p_2\}$. In this case,

$$\gamma - \frac{p}{p_1'} \geq 0, \quad \gamma - \frac{p}{p_2'} \geq 0.$$

Moreover, since we are considering the case $p > \gamma$,

$$\gamma - \frac{p}{p_1'} + \gamma - \frac{p}{p_2'} = 2\gamma - 2p + 1 < 1.$$

Applying Proposition 4.7 again, we have

$$\begin{aligned}
& \left\| \sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^{\gamma-1} \langle w \rangle_Q \mathbf{1}_Q \right\|_{L^{(\frac{p_2}{\gamma})'}(\sigma_2)} \\
& \simeq \left(\sum_{Q \in \mathfrak{S}} \lambda_Q \left(\frac{1}{\sigma_2(Q)} \sum_{Q' \subset Q} \langle \sigma_1 \rangle_{Q'}^\gamma \langle \sigma_2 \rangle_{Q'}^\gamma w(Q') \right)^{(\frac{p_2}{\gamma})'-1} \right)^{\frac{1}{(\frac{p_2}{\gamma})'}} \\
& \leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p_2}} \left(\sum_{Q \in \mathfrak{S}} \lambda_Q \left(\frac{1}{\sigma_2(Q)} \sum_{Q' \subset Q} \langle \sigma_1 \rangle_{Q'}^{\gamma-\frac{p}{p_1'}} \langle \sigma_2 \rangle_{Q'}^{\gamma-\frac{p}{p_2'}} |Q'| \right)^{(\frac{p_2}{\gamma})'-1} \right)^{\frac{1}{(\frac{p_2}{\gamma})'}} \\
& \stackrel{(4.9)}{\lesssim} [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p_2}} \left(\sum_{Q \in \mathfrak{S}} \lambda_Q \left(\frac{1}{\sigma_2(Q)} \langle \sigma_1 \rangle_Q^{\gamma-\frac{p}{p_1'}} \langle \sigma_2 \rangle_Q^{\gamma-\frac{p}{p_2'}} |Q| \right)^{(\frac{p_2}{\gamma})'-1} \right)^{\frac{1}{(\frac{p_2}{\gamma})'}} \\
& = [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p_2}} \left(\sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^{\frac{\gamma}{p_2-\gamma}(p_2-\frac{p}{p_1'})} \langle \sigma_2 \rangle_Q^{\frac{1}{p_2-\gamma}(p_2(1-\gamma)-\frac{p}{p_2})} \langle w \rangle_Q |Q| \right)^{\frac{1}{(\frac{p_2}{\gamma})'}}
\end{aligned}$$

$$\leq [w, \vec{\sigma}]_{A_{\vec{F}}}^{\frac{\gamma}{p}} \left(\sum_{Q \in \mathfrak{S}} \langle \sigma_1 \rangle_Q^{\frac{\gamma(\frac{p_2}{\gamma})'}{p_1}} \langle w \rangle_Q^{(\frac{p_2}{\gamma})'(1-\frac{\gamma}{p})} |Q| \right)^{\frac{1}{(\frac{p_2}{\gamma})'}}.$$

where again $\lambda_Q = \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma w(Q)$. \square

Now we are ready to prove our main results.

Proof of Theorem 4.1. First we consider the case $p > \gamma$, and we denote $q = p/\gamma$. By Lemma 4.11, we have

$$\begin{aligned} & \left\| \left(\sum_{Q \in \mathfrak{S}} \langle f_1 \rangle_Q^{\sigma_1} \langle f_2 \rangle_Q^{\sigma_2} \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma \mathbf{1}_Q \right)^{\frac{1}{\gamma}} \right\|_{L^p(w)}^\gamma \\ &= \sup_{\|h\|_{L^{q'}(w)}=1} \sum_{Q \in \mathfrak{S}} \langle f_1 \rangle_Q^{\sigma_1} \langle f_2 \rangle_Q^{\sigma_2} \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma \int_Q h dw \\ &= \sup_{\|h\|_{L^{q'}(w)}=1} \sum_{Q \in \mathfrak{S}} \langle f_1 \rangle_Q^{\sigma_1} \langle f_2 \rangle_Q^{\sigma_2} \langle h \rangle_Q^w \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma w(Q) \end{aligned}$$

For each $i = 1, 2$, let \mathcal{F}_i be the stopping family starting at Q_0 and defined by the stopping condition

$$\text{ch}_{\mathcal{F}_i}(F_i) := \{F'_i \in \mathfrak{S} : F'_i \subset F_i \text{ maximal such that } \langle f_i \rangle_{F'_i}^{\sigma_i} > 2\langle f_i \rangle_{F_i}^{\sigma_i}\}.$$

Each collection \mathcal{F}_i is σ_i -sparse, since

$$\sum_{F'_i \in \text{ch}_{\mathcal{F}_i}(F_i)} \sigma_i(F'_i) \leq \frac{1}{2} \frac{\sum_{F'_i \in \text{ch}_{\mathcal{F}_i}(F_i)} \int_{F'_i} f d\sigma}{\int_{F_i} f d\sigma} \sigma_i(F_i) \leq \frac{1}{2} \sigma_i(F_i).$$

The \mathcal{F}_i -stopping parent $\pi_{\mathcal{F}_i}(Q)$ of a cube Q is defined by $\pi_{\mathcal{F}_i}(Q) := \{F_i \in \mathcal{F}_i : F_i \text{ minimal such that } F_i \supseteq Q\}$. By the stopping condition, for every cube Q we have $\langle f_i \rangle_Q^{\sigma_i} \leq 2\langle f_i \rangle_{\pi_{\mathcal{F}_i}(Q)}^{\sigma_i}$. Let \mathcal{H} be the analogue stopping family associated with h and the weight w , verifying the corresponding properties.

By rearranging the summation according to the stopping parents and removing the supremum, we obtain

$$\begin{aligned} & \sum_{Q \in \mathfrak{S}} \langle f_1 \rangle_Q^{\sigma_1} \langle f_2 \rangle_Q^{\sigma_2} \langle h \rangle_Q^w \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma w(Q) \\ &= \left(\sum_{\substack{F_1 \in \mathcal{F}_1 \\ F_2 \subset F_1}} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ H \subset F_2}} \sum_{\substack{H \in \mathcal{H} \\ H \subset F_2}} \sum_{\substack{Q \in \mathfrak{S} \\ \pi(Q)=(F_1, F_2, H)}} + \sum_{F_2 \in \mathcal{F}_2} \sum_{\substack{F_1 \in \mathcal{F}_1 \\ F_1 \subset F_2}} \sum_{\substack{H \in \mathcal{H} \\ H \subset F_2}} \sum_{\substack{Q \in \mathfrak{S} \\ \pi(Q)=(F_1, F_2, H)}} \right) \\ &+ \sum_{F_1 \in \mathcal{F}_1} \sum_{\substack{H \in \mathcal{H} \\ H \subset F_1}} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ F_2 \subset H}} \sum_{\substack{Q \in \mathfrak{S} \\ \pi(Q)=(F_1, F_2, H)}} + \sum_{F_2 \in \mathcal{F}_2} \sum_{\substack{H \in \mathcal{H} \\ H \subset F_2}} \sum_{\substack{F_1 \in \mathcal{F}_1 \\ F_1 \subset H}} \sum_{\substack{Q \in \mathfrak{S} \\ \pi(Q)=(F_1, F_2, H)}} \end{aligned}$$

$$\begin{aligned}
& + \sum_{H \in \mathcal{H}} \sum_{\substack{F_1 \in \mathcal{F}_1 \\ F_1 \subset H}} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ F_2 \subset F_1}} \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = (F_1, F_2, H)}} + \sum_{H \in \mathcal{H}} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ F_2 \subset H}} \sum_{\substack{F_1 \in \mathcal{F}_1 \\ F_1 \subset F_2}} \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = (F_1, F_2, H)}} \\
& \times \langle f_1 \rangle_Q^{\sigma_1} \langle f_2 \rangle_Q^{\sigma_2} \langle h \rangle_Q^w \lambda_Q \\
& := I + I' + II + II' + III + III',
\end{aligned}$$

where $\pi(Q)$ means that $\pi_{\mathcal{F}_i}(Q) = F_i$, for all $i = 1, 2$ and $\pi_{\mathcal{H}}(Q) = H$ and

$$\lambda_Q := \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma w(Q).$$

By symmetry, it suffices to give an estimate for I . We have

$$\begin{aligned}
I & \leq \sum_{F_1 \in \mathcal{F}_1} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ F_2 \subset F_1}} \sum_{\substack{H \in \mathcal{H} \\ H \subset F_2}} \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = (F_1, F_2, H)}} \langle f_1 \rangle_Q^{\sigma_1} \langle f_2 \rangle_Q^{\sigma_2} \langle h \rangle_Q^w \lambda_Q \\
& \leq 8 \sum_{F_1 \in \mathcal{F}_1} \langle f_1 \rangle_{F_1}^{\sigma_1} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ F_2 \subset F_1}} \langle f_2 \rangle_{F_2}^{\sigma_2} \sum_{\substack{H \in \mathcal{H} \\ H \subset F_2}} \langle h \rangle_H^w \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = (F_1, F_2, H)}} \lambda_Q \\
& \lesssim \sum_{F_1 \in \mathcal{F}_1} \langle f_1 \rangle_{F_1}^{\sigma_1} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ F_2 \subset F_1}} \langle f_2 \rangle_{F_2}^{\sigma_2} \int \left(\sup_{\substack{H' \in \mathcal{H} \\ \pi_{\mathcal{F}_2}(H') = F_2}} \langle h \rangle_{H'}^w \mathbf{1}_{H'} \right) \\
& \times \sum_{\substack{H \in \mathcal{H} \\ H \subset F_2}} \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = (F_1, F_2, H)}} \frac{\lambda_Q}{w(Q)} \mathbf{1}_Q dw \\
& \leq \sum_{F_1 \in \mathcal{F}_1} \langle f_1 \rangle_{F_1}^{\sigma_1} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ \pi_{\mathcal{F}_1}(F_2) = F_1}} \langle f_2 \rangle_{F_2}^{\sigma_2} \left\| \sum_{\substack{H \in \mathcal{H} \\ \pi_{\mathcal{F}_2}(H) = F_2}} \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = (F_1, F_2, H)}} \frac{\lambda_Q}{w(Q)} \mathbf{1}_Q \right\|_{L^q(w)} \\
& \quad \times \left\| \sup_{\substack{H' \in \mathcal{H} \\ \pi_{\mathcal{F}_2}(H') = F_2}} \langle h \rangle_{H'}^w \mathbf{1}_{H'} \right\|_{L^{q'}(w)} \\
& \leq \left(\sum_{F_1 \in \mathcal{F}_1} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ \pi_{\mathcal{F}_2}(F_2) = F_1}} (\langle f_1 \rangle_{F_1}^{\sigma_1} \langle f_2 \rangle_{F_2}^{\sigma_2})^q \left\| \sum_{\substack{H \in \mathcal{H} \\ \pi_{\mathcal{F}_2}(H) = F_2}} \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = (F_1, F_2, H)}} \frac{\lambda_Q}{w(Q)} \mathbf{1}_Q \right\|_{L^q(w)}^q \right)^{1/q} \\
& \quad \times \left(\sum_{F_1 \in \mathcal{F}_1} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ \pi_{\mathcal{F}_1}(F_2) = F_1}} \sum_{\substack{H' \in \mathcal{H} \\ \pi_{\mathcal{F}_2}(H') = F_2}} (\langle h \rangle_{H'}^w)^{q'} w(H') \right)^{1/q'} \\
& \lesssim \left(\sum_{F_1 \in \mathcal{F}_1} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ \pi_{\mathcal{F}_1}(F_2) = F_1}} (\langle f_1 \rangle_{F_1}^{\sigma_1} \langle f_2 \rangle_{F_2}^{\sigma_2})^q \left\| \sum_{\substack{H \in \mathcal{H} \\ \pi_{\mathcal{F}_2}(H) = F_2}} \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = (F_1, F_2, H)}} \frac{\lambda_Q}{w(Q)} \mathbf{1}_Q \right\|_{L^q(w)}^q \right)^{1/q}.
\end{aligned}$$

By (4.16), we have

$$\left\| \sum_{\substack{Q \in \mathcal{S} \\ \pi_{\mathcal{F}_2}(Q) = F_2}} \frac{\lambda_Q}{w(Q)} \mathbf{1}_Q \right\|_{L^q(w)} \leq [w, \vec{\sigma}]_{A\vec{P}}^\gamma \left(\sum_{\substack{Q \in \mathcal{S} \\ \pi_{\mathcal{F}_2}(Q) = F_2}} \langle \sigma_1 \rangle_Q^{\frac{p}{p_1}} \langle \sigma_2 \rangle_Q^{\frac{p}{p_2}} |Q| \right)^{\gamma/p}.$$

Therefore,

$$\begin{aligned}
I &\leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} \left(\sum_{F_1 \in \mathcal{F}_1} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ \pi_{\mathcal{F}_1}(F_2) = F_1}} (\langle f_1 \rangle_{F_1}^{\sigma_1} \langle f_2 \rangle_{F_2}^{\sigma_2})^q \sum_{\substack{Q \in \mathcal{S} \\ \pi_{\mathcal{F}_2}(Q) = F_2}} \langle \sigma_1 \rangle_Q^{\frac{p}{p_1}} \langle \sigma_2 \rangle_Q^{\frac{p}{p_2}} |Q| \right)^{1/q} \\
&\leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} \left(\sum_{F_1 \in \mathcal{F}_1} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ \pi_{\mathcal{F}_1}(F_2) = F_1}} (\langle f_1 \rangle_{F_1}^{\sigma_1} \langle f_2 \rangle_{F_2}^{\sigma_2})^q \right. \\
&\quad \times \left(\sum_{\substack{Q \in \mathcal{S} \\ \pi_{\mathcal{F}_2}(Q) = F_2}} \langle \sigma_1 \rangle_Q |Q| \right)^{\frac{p}{p_1}} \left(\sum_{\substack{Q \in \mathcal{S} \\ \pi_{\mathcal{F}_2}(Q) = F_2}} \langle \sigma_2 \rangle_Q |Q| \right)^{\frac{p}{p_2}} \Big)^{1/q} \\
&\leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} \left(\sum_{F_1 \in \mathcal{F}_1} (\langle f_1 \rangle_{F_1}^{\sigma_1})^{p_1/\gamma} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ \pi_{\mathcal{F}_1}(F_2) = F_1}} \sum_{\substack{Q \in \mathcal{S} \\ \pi_{\mathcal{F}_2}(Q) = F_2}} \langle \sigma_1 \rangle_Q |Q| \right)^{\frac{\gamma}{p_1}} \\
&\quad \times \left(\sum_{F_2 \in \mathcal{F}_2} (\langle f_2 \rangle_{F_2}^{\sigma_2})^{p_2/\gamma} \sum_{\substack{Q \in \mathcal{S} \\ \pi_{\mathcal{F}_2}(Q) = F_2}} \langle \sigma_2 \rangle_Q |Q| \right)^{\frac{\gamma}{p_2}} \\
&\leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} \left(\prod_{i=1}^2 [\sigma_i]_{A_{\infty}}^{\frac{\gamma}{p_i}} \right) \|f_1\|_{L^{p_1/\gamma}(\sigma_1)} \|f_2\|_{L^{p_2/\gamma}(\sigma_2)}.
\end{aligned}$$

It remains to consider the case $p \leq \gamma$. By Lemma 4.11, we have

$$\begin{aligned}
&\left\| \left(\sum_{Q \in \mathcal{S}} \langle f_1 \rangle_Q^{\sigma_1} \langle f_2 \rangle_Q^{\sigma_2} \langle \sigma_1 \rangle_Q^{\gamma} \langle \sigma_2 \rangle_Q^{\gamma} \mathbf{1}_Q \right)^{\frac{1}{\gamma}} \right\|_{L^p(w)}^{\gamma} \\
&\lesssim \left\| \left(\sum_{F_1 \in \mathcal{F}_1} \langle f_1 \rangle_{F_1}^{\sigma_1} \sum_{F_2 \in \mathcal{F}_2} \langle f_2 \rangle_{F_2}^{\sigma_2} \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = (F_1, F_2)}} \langle \sigma_1 \rangle_Q^{\gamma} \langle \sigma_2 \rangle_Q^{\gamma} \mathbf{1}_Q \right)^{\frac{1}{\gamma}} \right\|_{L^p(w)}^{\gamma} \\
&\leq \left(\sum_{F_1 \in \mathcal{F}_1} (\langle f_1 \rangle_{F_1}^{\sigma_1})^q \sum_{F_2 \in \mathcal{F}_2} (\langle f_2 \rangle_{F_2}^{\sigma_2})^q \right\| \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = (F_1, F_2)}} \langle \sigma_1 \rangle_Q^{\gamma} \langle \sigma_2 \rangle_Q^{\gamma} \mathbf{1}_Q \Big\|_{L^q(w)}^q \Big)^{\frac{1}{q}} \\
&\lesssim \left(\sum_{F_1 \in \mathcal{F}_1} (\langle f_1 \rangle_{F_1}^{\sigma_1})^q \sum_{\substack{F_2 \in \mathcal{F}_2 \\ F_2 \subset F_1}} (\langle f_2 \rangle_{F_2}^{\sigma_2})^q \right\| \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = (F_1, F_2)}} \langle \sigma_1 \rangle_Q^{\gamma} \langle \sigma_2 \rangle_Q^{\gamma} \mathbf{1}_Q \Big\|_{L^q(w)}^q \Big)^{\frac{1}{q}} \\
&+ \left(\sum_{F_2 \in \mathcal{F}_2} (\langle f_2 \rangle_{F_2}^{\sigma_2})^q \sum_{\substack{F_1 \in \mathcal{F}_1 \\ F_1 \subset F_2}} (\langle f_1 \rangle_{F_1}^{\sigma_1})^q \right\| \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = (F_1, F_2)}} \langle \sigma_1 \rangle_Q^{\gamma} \langle \sigma_2 \rangle_Q^{\gamma} \mathbf{1}_Q \Big\|_{L^q(w)}^q \Big)^{\frac{1}{q}}.
\end{aligned}$$

Then by the previous arguments, the desired estimate follows. This completes the proof. \square

The proof of Theorem 4.2 follows the same idea as the proof of the previous theorem.

Proof of Theorem 4.2. We only check the estimate for I , since the other terms are similar. By (4.16), we have

$$\begin{aligned} \left\| \sum_{\substack{Q \in \mathcal{S} \\ Q \subset F_2}} \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma \mathbf{1}_Q \right\|_{L^q(w)} &\leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} \left(\sum_{\substack{Q \in \mathcal{S} \\ Q \subset F_2}} \langle \sigma_1 \rangle_Q^{\frac{p}{p_1}} \langle \sigma_2 \rangle_Q^{\frac{p}{p_2}} |Q| \right)^{\gamma/p} \\ &\lesssim [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} \left(\int_{F_2} \prod_{i=1}^2 M(\mathbf{1}_{F_2} \sigma_i)^{p/p_i} dx \right)^{\gamma/p} \\ &\leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} [\vec{\sigma}]_{W_{\vec{P}}}^{\gamma/p} \left(\int_{F_2} \prod_{i=1}^2 \sigma_i^{p/p_i} dx \right)^{\gamma/p}. \end{aligned}$$

Therefore,

$$\begin{aligned} I &\leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} [\vec{\sigma}]_{W_{\vec{P}}}^{\gamma/p} \left(\sum_{F_1 \in \mathcal{F}_1} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ \pi_{\mathcal{F}_1}(F_2) = F_1}} (\langle f_1 \rangle_{F_1}^{\sigma_1} \langle f_2 \rangle_{F_2}^{\sigma_2})^q \int_{F_2} \prod_{i=1}^2 \sigma_i^{p/p_i} dx \right)^{1/q} \\ &\leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} [\vec{\sigma}]_{W_{\vec{P}}}^{\gamma/p} \left(\int_{F_1 \in \mathcal{F}_1} (\langle f_1 \rangle_{F_1}^{\sigma_1})^q \mathbf{1}_{F_1} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ \pi_{\mathcal{F}_1}(F_2) = F_1}} (\langle f_2 \rangle_{F_2}^{\sigma_2})^q \mathbf{1}_{F_2} \prod_{i=1}^2 \sigma_i^{p/p_i} dx \right)^{1/q} \\ &\leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} [\vec{\sigma}]_{W_{\vec{P}}}^{\gamma/p} \left(\int M_{\mathcal{D}}^{\sigma_1}(f_1)^q M_{\mathcal{D}}^{\sigma_2}(f_2)^q \prod_{i=1}^2 \sigma_i^{p/p_i} dx \right)^{1/q} \\ &\leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} [\vec{\sigma}]_{W_{\vec{P}}}^{\gamma/p} \|M_{\mathcal{D}}^{\sigma_1}(f_1)\|_{L^{p_1/\gamma}(\sigma_1)} \cdot \|M_{\mathcal{D}}^{\sigma_2}(f_2)\|_{L^{p_2/\gamma}(\sigma_2)} \\ &\lesssim [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} [\vec{\sigma}]_{W_{\vec{P}}}^{\gamma/p} \|f_1\|_{L^{p_1/\gamma}(\sigma_1)} \cdot \|f_2\|_{L^{p_2/\gamma}(\sigma_2)}. \end{aligned}$$

□

Again, the proof of Theorem 4.4 also follows the same idea as the proof of the previous theorem.

Proof of Theorem 4.4. Likewise, we only study the estimate of I . By (4.16) again, we have

$$\begin{aligned} &\left\| \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = F_2}} \langle \sigma_1 \rangle_Q^\gamma \langle \sigma_2 \rangle_Q^\gamma \mathbf{1}_Q \right\|_{L^q(w)} \\ &\leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} \left(\sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = F_2}} \langle \sigma_1 \rangle_Q^{\frac{p}{p_1}} \langle \sigma_2 \rangle_Q^{\frac{p}{p_2}} |Q| \right)^{\gamma/p} \\ &\leq [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{\gamma}{p}} [\vec{\sigma}]_{H_{\vec{P}}}^{\frac{\gamma}{p}} \left(\sum_{\substack{Q \in \mathcal{S} \\ \pi(Q) = F_2}} \prod_{i=1}^2 \exp \left(\int_Q \log \sigma_i \right)^{\frac{p}{p_i}} |Q| \right)^{\gamma/p} \end{aligned}$$

$$\begin{aligned}
&\leq [w, \vec{\sigma}]_{A\bar{P}}^{\frac{\gamma}{p}} [\vec{\sigma}]_{H\bar{P}}^{\frac{\gamma}{p}} \prod_{i=1}^2 \left(\sum_{\substack{Q \in \mathcal{S} \\ \pi(Q)=F_2}} \exp \left(\int_Q \log \sigma_i \right) |Q| \right)^{\gamma/p_i} \\
&\lesssim [w, \vec{\sigma}]_{A\bar{P}}^{\frac{\gamma}{p}} [\vec{\sigma}]_{H\bar{P}}^{\frac{\gamma}{p}} \left(\sum_{\substack{Q \in \mathcal{S} \\ \pi(Q)=F_2}} \exp \left(\int_Q \log \sigma_1 \right) |Q| \right)^{\gamma/p_1} \|M_0(\mathbf{1}_{F_2} \sigma_2)\|_{L^1}^{\frac{\gamma}{p_2}} \\
&\leq [w, \vec{\sigma}]_{A\bar{P}}^{\frac{\gamma}{p}} [\vec{\sigma}]_{H\bar{P}}^{\frac{\gamma}{p}} \sigma_2(F)^{\frac{\gamma}{p_2}} \left(\sum_{\substack{Q \in \mathcal{S} \\ \pi(Q)=F_2}} \exp \left(\int_Q \log \sigma_1 \right) |Q| \right)^{\gamma/p_1},
\end{aligned}$$

where

$$(4.21) \quad M_0(f) := \sup_Q \exp \left(\int_Q \log |f| \right) \mathbf{1}_Q,$$

is the logarithmic maximal function. Here we have used the fact that this maximal function is bounded from L^p into itself for $p \in (0, \infty)$ with bound independent of the dimension in the dyadic case as proved in [15, Lemma 2.1]. Hence,

$$\begin{aligned}
I &\leq [w, \vec{\sigma}]_{A\bar{P}}^{\frac{\gamma}{p}} [\vec{\sigma}]_{H\bar{P}}^{\frac{\gamma}{p}} \left(\sum_{F_1 \in \mathcal{F}_1} (\langle f_1 \rangle_{F_1}^{\sigma_1})^q \sum_{\substack{F_2 \in \mathcal{F}_2 \\ \pi_{\mathcal{F}_1}(F_2)=F_1}} (\langle f_2 \rangle_{F_2}^{\sigma_2})^q \sigma_2(F)^{\frac{p}{p_2}} \right. \\
&\quad \times \left. \left(\sum_{\substack{Q \in \mathcal{S} \\ \pi(Q)=F_2}} \exp \left(\int_Q \log \sigma_1 \right) |Q| \right)^{p/p_1} \right)^{\frac{\gamma}{p}} \\
&\leq [w, \vec{\sigma}]_{A\bar{P}}^{\frac{\gamma}{p}} [\vec{\sigma}]_{H\bar{P}}^{\frac{\gamma}{p}} \left(\sum_{F_1 \in \mathcal{F}_1} (\langle f_1 \rangle_{F_1}^{\sigma_1})^q \left(\sum_{\substack{F_2 \in \mathcal{F}_2 \\ \pi_{\mathcal{F}_1}(F_2)=F_1}} (\langle f_2 \rangle_{F_2}^{\sigma_2})^{p_2/\gamma} \sigma_2(F) \right)^{\frac{p}{p_2}} \right. \\
&\quad \times \left. \left(\sum_{\substack{F_2 \in \mathcal{F}_2 \\ \pi_{\mathcal{F}_1}(F_2)=F_1}} \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q)=F_2}} \exp \left(\int_Q \log \sigma_1 \right) |Q| \right)^{p/p_1} \right)^{\frac{\gamma}{p}} \\
&\leq [w, \vec{\sigma}]_{A\bar{P}}^{\frac{\gamma}{p}} [\vec{\sigma}]_{H\bar{P}}^{\frac{\gamma}{p}} \left(\sum_{F_1 \in \mathcal{F}_1} (\langle f_1 \rangle_{F_1}^{\sigma_1})^{p_1/\gamma} \left(\sum_{\substack{F_2 \in \mathcal{F}_2 \\ \pi_{\mathcal{F}_1}(F_2)=F_1}} \sum_{\substack{Q \in \mathcal{S} \\ \pi(Q)=F_2}} \exp \left(\int_Q \log \sigma_1 \right) |Q| \right) \right)^{\frac{\gamma}{p_1}} \\
&\quad \times \left(\sum_{F_1 \in \mathcal{F}_1} \sum_{\substack{F_2 \in \mathcal{F}_2 \\ \pi_{\mathcal{F}_1}(F_2)=F_1}} (\langle f_2 \rangle_{F_2}^{\sigma_2})^{p_2/\gamma} \sigma_2(F) \right)^{\frac{\gamma}{p_2}} \\
&\leq [w, \vec{\sigma}]_{A\bar{P}}^{\frac{\gamma}{p}} [\vec{\sigma}]_{H\bar{P}}^{\frac{\gamma}{p}} \|f_1\|_{L^{p_1/\gamma}(\sigma_1)} \|f_2\|_{L^{p_2/\gamma}(\sigma_2)}.
\end{aligned}$$

□

5. APPLICATIONS

5.1. Mixed A_p - A_∞ estimate for commutators of multilinear Calderón–Zygmund operators. Throughout this section, we will work with commutators of multilinear Calderón–Zygmund operators with symbols in BMO . Recall that BMO consists of all locally integrable functions b with $\|b\|_{BMO} < \infty$, where

$$\|b\|_{BMO} := \sup_Q \frac{1}{|Q|} \int_Q |b(y) - \langle b \rangle_Q| dy,$$

and the supremum in the above definition is taken over all cubes $Q \in \mathbb{R}^n$ with sides parallel to the axes.

Given a multilinear Calderón–Zygmund operator T and $\vec{b} \in BMO^m$, we consider the following commutators with \vec{b} ,

$$[\vec{b}, T] = \sum_{i=1}^m [\vec{b}, T]_i,$$

where

$$[\vec{b}, T]_i(\vec{f}) := b_i T(\vec{f}) - T(f_1, \dots, f_{i-1}, b_i f_i, f_{i+1}, \dots, f_m).$$

Our aim in this section is to prove the following mixed estimate for commutators of multilinear Calderón–Zygmund operators following the same spirit as in [5].

Theorem 5.1. *Let T be a multilinear Calderón–Zygmund operator and $\vec{b} \in BMO^m$. If we assume that $[w, \vec{\sigma}]_{A_{\vec{b}}} < \infty$, then*

$$\begin{aligned} & \|[\vec{b}, T]\|_{L^{p_1}(w_1) \times \dots \times L^{p_m}(w_m) \rightarrow L^p(w)} \\ & \leq [w, \vec{\sigma}]_{A_{\vec{b}}}^{\frac{1}{p}} \left(\prod_{i=1}^m [\sigma_i]_{A_\infty}^{\frac{1}{p_i}} + [w]_{A_\infty}^{\frac{1}{p'}} \sum_{j=1}^m \prod_{i \neq j} [\sigma_i]_{A_\infty}^{\frac{1}{p_i}} \right) \\ & \quad \times \left([w]_{A_\infty} + \sum_{i=1}^m [\sigma_i]_{A_\infty} \right) \left(\sum_{i=1}^m \|b_i\|_{BMO} \right), \end{aligned}$$

where $\sigma_i = w_i^{1-p'_i}$, $i = 1, \dots, m$.

Before proving our main result in this section we need to recall some basic properties about BMO functions and A_∞ weights that we are going to use in the sequel. Recall that a key property of BMO functions is the celebrated John–Nirenberg inequality [17].

Proposition 5.2. [18, pp. 31–32] *There are dimensional constants $0 < \alpha_n < 1 < \beta_n < \infty$ such that*

$$(5.3) \quad \sup_Q \frac{1}{|Q|} \int_Q \exp\left(\frac{\alpha_n}{\|b\|_{BMO}} |b(y) - \langle b \rangle_Q|\right) dy \leq \beta_n.$$

In fact, we can take $\alpha_n = \frac{1}{2^{n+2}}$.

It is well-known that if $w \in A_\infty$, then $\log w \in BMO$. Using the John-Nirenberg inequality, Chung, Pereyra, and Pérez [5] proved the following bound.

Proposition 5.4. *Let $b \in BMO$ and let $0 < \alpha_n < 1 < \beta_n < \infty$ be the dimensional constants from (5.3). Then*

$$s \in \mathbb{R}, |s| \leq \frac{\alpha_n}{\|b\|_{BMO}} \min\left\{1, \frac{1}{p-1}\right\} \Rightarrow e^{sb} \in A_p \text{ and } [e^{sb}]_{A_p} \leq \beta_n^p.$$

In [15], Hytönen and Pérez also showed the following bound for the Fujii-Wilson A_∞ constant of a particular family of weights.

Proposition 5.5. *There are dimensional constants ε_n and c_n such that*

$$[e^{\text{Re}z b} w]_{A_\infty} \leq c_n [w]_{A_\infty} \quad \text{if } |z| \leq \frac{\varepsilon_n}{\|b\|_{BMO} [w]_{A_\infty}}.$$

For our purpose, we need to show the following variation of the previous lemmas.

Lemma 5.6. *Suppose that $[w, \vec{\sigma}]_{A_{\vec{p}}} < \infty$ and $w, \sigma_i \in A_\infty$, $i = 1, 2, \dots, m$. Then for any $1 \leq j \leq m$,*

$$[w e^{pb\text{Re}z}, \sigma_1, \dots, \sigma_j e^{-p'_j b\text{Re}z}, \dots, \sigma_m]_{A_{\vec{p}}} \leq c_{n, \vec{p}} [w, \vec{\sigma}]_{A_{\vec{p}}},$$

provided that

$$|z| \leq \frac{\alpha_n \min\left\{1, \frac{p'_1}{p}, \dots, \frac{p'_m}{p}\right\}}{p(1 + \max\{[w]_{A_\infty}, [\sigma_1]_{A_\infty}, \dots, [\sigma_m]_{A_\infty}\}) \|b\|_{BMO}}.$$

To prove the previous lemma, we need to recall this sharp version of the reverse Hölder's inequality proved in [15].

Proposition 5.7. *Let $w \in A_\infty$. Then for any $0 \leq r \leq 1 + \frac{1}{c_n [w]_{A_\infty}}$, we have*

$$\left(\frac{1}{|Q|} \int_Q w(x)^r dx\right)^{\frac{1}{r}} \leq 2 \frac{1}{|Q|} \int_Q w(x) dx.$$

Proof of Lemma 5.6. Set

$$r = 1 + \frac{1}{c_n \max\{[w]_{A_\infty}, [\sigma_j]_{A_\infty}\}}.$$

By definition of the $A_{\vec{p}}$ constant, Hölder's inequality and Proposition 5.7, we have

$$\begin{aligned} & [w e^{pb\text{Re}z}, \sigma_1, \dots, \sigma_j e^{-p'_j b\text{Re}z}, \dots, \sigma_m]_{A_{\vec{p}}} \\ &= \sup_Q \langle w e^{pb\text{Re}z} \rangle_Q \langle \sigma_j e^{-p'_j b\text{Re}z} \rangle_Q^{\frac{p}{p'_j}} \prod_{i \neq j} \langle \sigma_i \rangle_Q^{\frac{p}{p'_i}} \\ &\leq \sup_Q \langle w^r \rangle_Q^{\frac{1}{r}} \langle e^{pb\text{Re}z} \rangle_Q^{\frac{1}{r'}} \langle \sigma_j^r \rangle_Q^{\frac{p}{r p'_j}} \langle e^{-p'_j b\text{Re}z} \rangle_Q^{\frac{p}{r' p'_j}} \prod_{i \neq j} \langle \sigma_i \rangle_Q^{\frac{p}{p'_i}} \end{aligned}$$

$$\begin{aligned}
&\leq 4 \sup_Q \langle w \rangle_Q \langle e^{pbr' \operatorname{Re} z} \rangle_Q^{\frac{1}{r'}} \langle \sigma_j \rangle_Q^{\frac{p}{p_j'}} \langle e^{-p_j' br' \operatorname{Re} z} \rangle_Q^{\frac{p}{r' p_j'}} \prod_{i \neq j} \langle \sigma_i \rangle_Q^{\frac{p}{p_i'}} \\
&\leq 4 [w, \vec{\sigma}]_{A_{\vec{P}}} [e^{pbr' \operatorname{Re} z}]_{A_{1+\frac{p}{p_j'}}} \\
&\leq c_{n, \vec{P}} [w, \vec{\sigma}]_{A_{\vec{P}}},
\end{aligned}$$

where Proposition 5.4 is used in the last step. \square

Now we are ready to prove the main result in this section.

Proof of Theorem 5.1. It suffices to study the boundedness of $[b, T]_i$. Without loss of generality, we just consider the case $i = 1$. Using the same trick as that in [5, Thm. 3.1], for any complex number z , we define

$$T_z^1(\vec{f}) = e^{zb} T(e^{-zb} f_1, f_2, \dots, f_m).$$

Then by using the Cauchy integral theorem, we get for “nice” functions,

$$[b, T]_1(\vec{f}) = \frac{d}{dz} T_z^1(\vec{f}) \Big|_{z=0} = \frac{1}{2\pi i} \int_{|z|=\varepsilon} \frac{T_z^1(\vec{f})}{z^2} dz, \quad \varepsilon > 0.$$

Next, using Minkowski’s inequality, for $p \geq 1$,

$$(5.8) \quad \|[b, T]_1(\vec{f})\|_{L^p(w)} \leq \frac{1}{2\pi\varepsilon^2} \int_{|z|=\varepsilon} \|T_z^1(\vec{f})\|_{L^p(w)} |dz|.$$

Notice that

$$(5.9) \quad \|T_z^1(\vec{f})\|_{L^p(w)} = \|T(e^{-zb} f_1, f_2, \dots, f_m)\|_{L^p(we^{pb \operatorname{Re} z})}.$$

Therefore, applying the boundedness properties for Calderón–Zygmund operators in Theorem 4.1 for weights $(we^{pb \operatorname{Re} z}, w_1 e^{p_1 b \operatorname{Re} z}, w_2, \dots, w_m)$ with $p_0 = \gamma = 1$, we get

$$\begin{aligned}
(5.10) \quad &\|T(e^{-zb} f_1, f_2, \dots, f_m)\|_{L^p(we^{pb \operatorname{Re} z})} \lesssim [e^{pb \operatorname{Re} z} w, e^{-p_1' b \operatorname{Re} z} \sigma_1, \sigma_2, \dots, \sigma_m]_{A_{\vec{P}}}^{1/p} \\
&\times \left([e^{-p_1' b \operatorname{Re} z} \sigma_1]_{A_\infty}^{1/p_1} \prod_{i=2}^m [\sigma_i]_{A_\infty}^{1/p_i} + [e^{pb \operatorname{Re} z} w]_{A_\infty}^{1/p'} \left(\prod_{i=2}^m [\sigma_i]_{A_\infty}^{1/p_i} + \right. \right. \\
&\left. \left. + \sum_{i'=2}^m [\sigma_{i'} e^{-p_1' b \operatorname{Re} z}]_{A_\infty}^{1/p_1} \prod_{\substack{i \neq i' \\ i > 1}} [\sigma_i]_{A_\infty}^{1/p_i} \right) \right) \|f_1 e^{-zb}\|_{L^{p_1}(e^{b p_1 \operatorname{Re} z} w_1)} \prod_{i=2}^m \|f_i\|_{L^{p_i}(w_i)}.
\end{aligned}$$

Combining (5.8), (5.9) and (5.10) and using Proposition 5.5 and Lemma 5.6, we arrive at

$$(5.11) \quad \begin{aligned} & \| [b, T]_1(\vec{f}) \|_{L^p(w)} \\ & \leq \frac{1}{2\pi\varepsilon} [w, \vec{\sigma}]_{A_{\vec{P}}}^{1/p} \left(\prod_{i=1}^m [\sigma_i]_{A_\infty}^{1/p_i} + [w]_{A_\infty}^{1/p'} \sum_{i'=1}^m \prod_{i' \neq i} [\sigma_{i'}]_{A_\infty}^{1/p_{i'}} \right) \prod_{i=1}^m \|f_i\|_{L^{p_i}(w_i)}. \end{aligned}$$

Now taking

$$\varepsilon = \frac{c_{n, \vec{P}}}{([w]_{A_\infty} + \sum_{i=1}^m [\sigma_i]_{A_\infty}) \|b_1\|_{BMO}},$$

where $c_{n, \vec{P}}$ is sufficiently small such that it satisfies the hypotheses in Proposition 5.5 and Lemma 5.6. Then, we obtain

$$\begin{aligned} \| [b, T]_1(\vec{f}) \|_{L^p(w)} & \lesssim [w, \vec{\sigma}]_{A_{\vec{P}}}^{\frac{1}{p}} \left(\prod_{i=1}^m [\sigma_i]_{A_\infty}^{\frac{1}{p_i}} + [w]_{A_\infty}^{\frac{1}{p'}} \sum_{j=1}^m \prod_{i \neq j} [\sigma_i]_{A_\infty}^{\frac{1}{p_i}} \right) \\ & \quad \times ([w]_{A_\infty} + \sum_{i=1}^m [\sigma_i]_{A_\infty}) \|b_1\|_{BMO} \prod_{i=1}^m \|f_i\|_{L^{p_i}(w_i)}. \end{aligned}$$

The general result follows immediately combining the estimates for all the commutators in the different variables. \square

5.2. Mixed A_p - A_∞ estimates for multilinear square functions and multilinear Fourier multipliers. The results obtained in Section 4 can be applied to different instances of operators which can be reduced to the simpler dyadic operators $\mathcal{A}_{p_0, \gamma, \mathcal{S}}$.

Firstly, observe that the mixed weighted bounds obtained in the main theorems in Section 4 can be extended to the case of multilinear square functions taking into account [2, Prop. 4.2] and choosing $p_0 = 1$ and $\gamma = 2$.

These mixed bounds can also be extended to multilinear Fourier multipliers, which are a particular example of a general class of operators whose kernels satisfy weaker regularity conditions than the usual Hölder continuity. To obtain the corresponding mixed bounds, it is sufficient to consider the results in [1] together with the main theorems in Section 4 for $\gamma = 1$. It is worth mentioning that these mixed bounds for Fourier multipliers seem to be new in the multilinear scenario.

6. APPENDIX

In this appendix we state and prove some well-known boundedness results for bilinear Calderón–Zygmund operators and their maximal truncations, which also hold in the multilinear setting. It is worth mentioning that the novelty of these results is not only that they are stated in a quantitative way that will be useful for our purposes, but also that some of these results are proved under weaker regularity conditions on the kernels than those results in the literature.

Lemma 6.1. *Let T be a bilinear Dini-continuous Calderón-Zygmund operator. Then T is bounded from $L^1 \times L^1$ to $L^{1/2, \infty}$ and*

$$(6.2) \quad \|T\|_{L^1 \times L^1 \rightarrow L^{1/2, \infty}} \lesssim \|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q} + \|\omega\|_{\text{Dini}},$$

where $\|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q}$ denotes the norm of the operator as in its definition.

This result was proved under the $Dini(\frac{1}{2})$ condition in [26]. Observe that $Dini(1/2)$ condition is an stronger condition than Dini condition, which is also referred to as $Dini(1)$. In [27], Pérez and Torres studied the problem under the $BGHC$ condition. Namely, we say that a bilinear operator with kernel K satisfies the bilinear geometric Hörmander condition ($BGHC$) if there exists a fixed constant C such that and for any family of disjoint dyadic cubes D_1 and D_2 ,

$$(6.3) \quad \int_{\mathbb{R}^n} \sup_{y \in Q} \int_{\mathbb{R} \setminus Q^*} |K(x, y, z) - K(x, y_Q, z)| dx dz \leq C,$$

$$(6.4) \quad \int_{\mathbb{R}^n} \sup_{z \in P} \int_{\mathbb{R} \setminus P^*} |K(x, y, z) - K(x, y, z_P)| dx dy \leq C,$$

and

$$(6.5) \quad \sum_{(P, Q) \in D_1 \times D_2} |P||Q| \sup_{(y, z) \in P \times Q} \int_{\mathbb{R}^n \setminus (\cup_{R \in D_1} R) \cup (\cup_{S \in D_2} S)} |K(x, y, z) - K(x, y_P, z_Q)| dx \leq C(|\cup_{P \in D_1} P| + |\cup_{Q \in D_2} Q|).$$

Here Q^* is the cube with the same center as Q and sidelength $10\sqrt{n}\ell(Q)$. This condition, which is actually stated here in an equivalent way, was shown to be weaker than the Dini condition in [27, Prop. 2.3]). Thus, Lemma 6.1 follows immediately from the mentioned result. Here we give the proof with the precise constants.

Proof of Lemma 6.1. Suppose that T is bounded from $L^{q_1} \times L^{q_2}$ to L^q , where $\frac{1}{q_1} + \frac{1}{q_2} = \frac{1}{q}$. We shall dominate the bound $\|T\|_{L^1 \times L^1 \rightarrow L^{1/2, \infty}}$ by $\|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q} + \|\omega\|_{\text{Dini}}$. Indeed, fix $\lambda > 0$ and consider without loss of generality functions $f_i \geq 0$, $i = 1, 2$. Let $\alpha_i > 0$ be numbers to be determined later. Apply the Calderón-Zygmund decomposition to f_i at height $\alpha_i \lambda$, to obtain its good and bad parts g_i and b_i , respectively, and families of cubes $\{Q_k^i\}_k$ with disjoint interiors such that $f_i = g_i + b_i$ and $b_i = \sum_k b_k^i$ verifying the properties in [10, Thm. 4.3.1].

Next, set $\Omega_i = \cup_k 4nQ_k^i$. We have

$$\begin{aligned} |\{x : |T(f_1, f_2)(x)| > \lambda\}| &\leq |\Omega_1| + |\Omega_2| \\ &\quad + |\{x \in (\Omega_1 \cup \Omega_2)^c : |T(g_1, g_2)(x)| > \frac{\lambda}{4}\}| \\ &\quad + |\{x \in (\Omega_1 \cup \Omega_2)^c : |T(g_1, b_2)(x)| > \frac{\lambda}{4}\}| \end{aligned}$$

$$\begin{aligned}
& + |\{x \in (\Omega_1 \cup \Omega_2)^c : |T(b_1, g_2)(x)| > \frac{\lambda}{4}\}| \\
& + |\{x \in (\Omega_1 \cup \Omega_2)^c : |T(b_1, b_2)(x)| > \frac{\lambda}{4}\}|.
\end{aligned}$$

It is easy to see that

$$|\Omega_1| + |\Omega_2| \leq C_n \left(\frac{1}{\alpha_1 \lambda} \|f_1\|_{L^1} + \frac{1}{\alpha_2 \lambda} \|f_2\|_{L^1} \right).$$

For the third term, using Chebychev's inequality and the boundedness properties of T and g_i , we have

$$\begin{aligned}
& |\{x \in (\Omega_1 \cup \Omega_2)^c : |T(g_1, g_2)(x)| > \frac{\lambda}{4}\}| \\
& \leq \frac{4^q}{\lambda^q} \|T(g_1, g_2)\|_{L^q}^q \\
& \leq \frac{4^q}{\lambda^q} \|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q}^q \|g_1\|_{L^{q_1}}^q \|g_2\|_{L^{q_2}}^q \\
& \leq \frac{4^q}{\lambda^q} C_{n,q,q_1,q_2} \|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q}^q (\alpha_1 \lambda)^{q/q_1'} (\alpha_2 \lambda)^{q/q_2'} \|f_1\|_{L^1}^{q/q_1} \|f_2\|_{L^1}^{q/q_2}.
\end{aligned}$$

For the fourth term, if c_k denotes the center of the cube Q_k^2 , we have

$$\begin{aligned}
& |\{x \in (\Omega_1 \cup \Omega_2)^c : |T(g_1, b_2^k)(x)| > \frac{\lambda}{4}\}| \\
& \leq \frac{4}{\lambda} \int \left| \sum_k \int \int_{Q_k^2} (K(x, y, z) - K(x, y, c_k)) g_1(y) b_2^k(z) dz dy \right| dx \\
& \leq \frac{4}{\lambda} \sum_k \int \int \int_{Q_k^2} |K(x, y, z) - K(x, y, c_k)| \cdot |g_1(y)| \cdot |b_2^k(z)| dz dy dx \\
& \leq \frac{4}{\lambda} \sum_k \int_{Q_k^2} \int \int \omega \left(\frac{\sqrt{n} \ell(Q_k^2)}{2(|x-y| + |x-z|)} \right) \frac{|g_1(y)| \cdot |b_2^k(z)|}{(|x-y| + |x-z|)^{2n}} dy dx dz \\
& \leq C_n \alpha_1 \sum_k \int_{Q_k^2} \int \int \omega \left(\frac{\sqrt{n} \ell(Q_k^2)}{2(|y| + |x-z|)} \right) \frac{|b_2^k(z)|}{(|y| + |x-z|)^{2n}} dy dx dz \\
& \leq C_n \alpha_1 \sum_k \int_{Q_k^2} \int \int \omega \left(\frac{\sqrt{n} \ell(Q_k^2)}{2|x-z|} \right) \frac{|b_2^k(z)|}{(|y| + |x-z|)^{2n}} dy dx dz \\
& \leq C_n \alpha_1 \sum_k \int_{Q_k^2} \int_{|x-z| \geq n \ell(Q_k^2)} \omega \left(\frac{\sqrt{n} \ell(Q_k^2)}{2|x-z|} \right) \frac{|b_2^k(z)|}{|x-z|^n} dx dz \\
& \leq C_n' \alpha_1 \|\omega\|_{\text{Dini}} \|f_2\|_{L^1},
\end{aligned}$$

where we have used the cancellation properties of b_2^k , the regularity condition on the third variable of K (since $|z - c_k| < \tau \max(|x - y|, |x - z|)$ for $x \notin \Omega_1 \cup \Omega_2$), the fact that ω is increasing, the Dini condition, $\|g_1\|_{L^\infty} \leq c_n \alpha_1 \lambda$ and $\sum_k \|b_2^k\|_{L^1} \leq c_n \|f_2\|_{L^1}$.

Since the estimate of the fifth term is symmetric to the previous estimate, it remains to estimate the last term. If we denote as c_l and c_k the center of

the cubes Q_l^1 and Q_k^2 , respectively, proceeding similarly as in the previous estimate, we obtain

$$\begin{aligned}
& |\{x \in (\Omega_1 \cup \Omega_2)^c : |T(b_1, b_2)(x)| > \frac{\lambda}{4}\}| \\
& \leq \frac{4}{\lambda} \int \left| \sum_{k,l} \int_{Q_l^1} \int_{Q_k^2} (K(x, y, z) - K(x, y, c_k)) b_1^l(y) b_2^k(z) dz dy \right| dx \\
& \leq \frac{4}{\lambda} \sum_{k,l} \int_{(\Omega_1 \cup \Omega_2)^c} \int_{Q_l^1} \int_{Q_k^2} |K(x, y, z) - K(x, y, c_k)| |b_1^l(y)| |b_2^k(z)| dx dy dz \\
& \leq \frac{4}{\lambda} \sum_{k,l} \int_{Q_k^2} \int_{Q_l^1} \int_{(\Omega_1 \cup \Omega_2)^c} \omega\left(\frac{\sqrt{n}\ell(Q_k^2)}{2(|x-y|+|x-z|)}\right) \frac{|b_1^l(y)| |b_2^k(z)| dx dy dz}{(|x-y|+|x-z|)^{2n}} \\
& \leq \frac{C_n}{\lambda} \sum_{k,l} \int_{Q_k^2} \int_{Q_l^1} \int_{(\Omega_1 \cup \Omega_2)^c} \omega\left(\frac{\sqrt{n}\ell(Q_k^2)}{2(|x-c_l|+|x-c_k|)}\right) \frac{|b_1^l(y)| |b_2^k(z)| dx dy dz}{(|x-c_l|+|x-c_k|)^{2n}} \\
& \leq C_n \sum_{k,l} |Q_l^1| |Q_k^2| \alpha_1 \alpha_2 \lambda \int_{(\Omega_1 \cup \Omega_2)^c} \omega\left(\frac{\sqrt{n}(\ell(Q_k^2) + \ell(Q_l^1))}{2(|x-c_l|+|x-c_k|)}\right) \frac{dx}{(|x-c_l|+|x-c_k|)^{2n}} \\
& \leq C'_n \sum_{k,l} \alpha_1 \alpha_2 \lambda \int_{Q_k^2} \int_{Q_l^1} \int_{(\Omega_1 \cup \Omega_2)^c} \omega\left(\frac{\sqrt{n}(\ell(Q_k^2) + \ell(Q_l^1))}{2(|x-y|+|x-z|)}\right) \frac{dx dy dz}{(|x-y|+|x-z|)^{2n}} \\
& = C'_n \sum_{k,l} \alpha_1 \alpha_2 \lambda \left(\int_{\ell(Q_k^2) \geq \ell(Q_l^1)} + \int_{\ell(Q_l^1) \geq \ell(Q_k^2)} \right) \\
& \leq I + II.
\end{aligned}$$

By symmetry, it suffices to estimate I . We have

$$\begin{aligned}
I & \leq C'_n \sum_k \alpha_1 \alpha_2 \lambda \int_{Q_k^2} \int_{(\Omega_1 \cup \Omega_2)^c} \int_{\mathbb{R}^n} \omega\left(\frac{\sqrt{n}\ell(Q_k^2)}{|x-z|}\right) \frac{dy dx dz}{(|x-y|+|x-z|)^{2n}} \\
& = C'_n \sum_k \alpha_1 \alpha_2 \lambda \int_{Q_k^2} \int_{(\Omega_1 \cup \Omega_2)^c} \omega\left(\frac{\sqrt{n}\ell(Q_k^2)}{|x-z|}\right) \frac{1}{|x-z|^n} dx dz \\
& \leq C_n \alpha_1 \|\omega\|_{\text{Dini}} \|f_2\|_{L^1}.
\end{aligned}$$

Combining the arguments above, we have

$$\begin{aligned}
& |\{x : |T(f_1, f_2)(x)| > \lambda\}| \\
& \lesssim \frac{1}{\alpha_1 \lambda} \|f_1\|_{L^1} + \frac{1}{\alpha_2 \lambda} \|f_2\|_{L^1} \\
& \quad + \|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q}^q (\alpha_1)^{q/q'_1} (\alpha_2)^{q/q'_2} \lambda^{q-1} \|f_1\|_{L^1}^{q/q_1} \|f_2\|_{L^1}^{q/q_2} \\
& \quad + \alpha_1 \|\omega\|_{\text{Dini}} \|f_2\|_{L^1} + \alpha_2 \|\omega\|_{\text{Dini}} \|f_1\|_{L^1}
\end{aligned}$$

Take

$$\alpha_1 = \lambda^{-\frac{1}{2}} \frac{\|f_1\|_{L^1}^{\frac{1}{2}}}{\|f_2\|_{L^1}^{\frac{1}{2}} \left(\|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q} + \|\omega\|_{\text{Dini}} \right)^{\frac{1}{2}}}$$

$$\alpha_2 = \lambda^{-\frac{1}{2}} \frac{\|f_2\|_{L^1}^{\frac{1}{2}}}{\|f_1\|_{L^1}^{\frac{1}{2}}} \frac{1}{(\|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q} + \|\omega\|_{\text{Dini}})^{\frac{1}{2}}},$$

we get

$$\lambda \left| \{x : |T(f_1, f_2)(x)| > \lambda\} \right|^2 \leq (\|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q} + \|\omega\|_{\text{Dini}}) \|f_1\|_{L^1} \|f_2\|_{L^1}.$$

□

We also need to show that the maximal truncated operator T_{\sharp} is bounded from $L^1 \times L^1$ to $L^{\frac{1}{2}, \infty}$. Therefore, we need to check first that Cotlar's inequality holds for this class of operators.

Theorem 6.6. *Let T be a bilinear Dini-continuous Calderón-Zygmund operator with kernel K . Then, for all $\eta \in (0, \frac{1}{2})$, there exists a constant C such that*

$$(6.7) \quad T_{\sharp}(\vec{f}) \leq c_{\eta, n} (C_K + \|\omega\|_{\text{Dini}} + \|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q}) \mathcal{M}(\vec{f}) + M_{\eta}(|T(\vec{f})|).$$

In this proof we combine the strategies used in [26, Thm 6.4] and [16, Lemma 5.3] to determine the precise constants involved in the inequality.

Proof of Theorem 6.6. Let us begin defining the following maximal truncation

$$\tilde{T}_{\sharp}(f_1, f_2)(x) = \sup_{\varepsilon > 0} \left| \tilde{T}_{\varepsilon}(f_1, f_2)(x) \right|,$$

where

$$\tilde{T}_{\varepsilon}(f_1, f_2)(x) = \int_{\max\{|x-y|, |x-z|\} > \varepsilon} K(x, y, z) f_1(y) f_2(z) dy dz.$$

Since

$$(6.8) \quad \sup_{\varepsilon > 0} \left| \int_{\substack{\max\{|x-y|, |x-z|\} \leq \varepsilon \\ |x-y|^2 + |x-z|^2 > \varepsilon^2}} K(x, y, z) f_1(y) f_2(z) dy dz \right| \lesssim C_K \mathcal{M}(f_1, f_2)(x),$$

it suffices to show (6.7) with T_{\sharp} replaced by \tilde{T}_{\sharp} . Notice that we can write for $x' \in B(x, \varepsilon/2)$,

$$(6.9) \quad \begin{aligned} \tilde{T}_{\varepsilon}(f_1, f_2)(x) &= \int_{\max\{|x-y|, |x-z|\} > \varepsilon} (K(x, y, z) - K(x', y, z)) f_1(y) f_2(z) dy dz \\ &\quad + T(f_1, f_2)(x') + T(f_1^0, f_2^0)(x'), \end{aligned}$$

where $f_i^0 = f_i \mathbf{1}_{B(x, \varepsilon)}$. For the first term in (6.9), using the regularity assumptions on the kernel, we get

$$\left| \int_{\max\{|x-y|, |x-z|\} > \varepsilon} (K(x, y, z) - K(x', y, z)) f_1(y) f_2(z) dy dz \right|$$

$$\begin{aligned}
&\leq \int_{\max\{|x-y|, |x-z|\} > \varepsilon} \omega\left(\frac{|x-x'|}{|x-y|+|x-z|}\right) \frac{|f_1(y)||f_2(z)| dy dz}{(|x-y|+|x-z|)^{2n}} \\
&= \sum_{k=0}^{\infty} \int_{2^k \varepsilon < \max\{|x-y|, |x-z|\} \leq 2^{k+1} \varepsilon} \omega\left(\frac{|x-x'|}{2^k \varepsilon}\right) \frac{1}{(2^k \varepsilon)^{2n}} |f_1(y)||f_2(z)| dy dz \\
&\lesssim \mathcal{M}(f_1, f_2)(x) \sum_{k=0}^{\infty} \omega\left(\frac{|x-x'|}{2^k \varepsilon}\right) \\
&\lesssim \mathcal{M}(f_1, f_2)(x) \sum_{k=0}^{\infty} \int_{2^{k-1}}^{2^k} \omega\left(\frac{|x-x'|}{\varepsilon t}\right) \frac{dt}{t} \\
&= \mathcal{M}(f_1, f_2)(x) \sum_{k=0}^{\infty} \int_{\frac{|x-x'|}{2^k \varepsilon}}^{\frac{|x-x'|}{2^{k-1} \varepsilon}} \omega(u) \frac{du}{u} \\
&= \mathcal{M}(f_1, f_2)(x) \int_0^{\frac{2|x-x'|}{\varepsilon}} \omega(u) \frac{du}{u} \\
&\leq \|\omega\|_{\text{Dini}} \mathcal{M}(f_1, f_2)(x),
\end{aligned}$$

where the last step holds since $|x-x'| \leq \varepsilon/2$. Next, taking the L^η average over $x' \in B(x, \varepsilon/2)$, we arrive at

$$\begin{aligned}
|\tilde{T}_\varepsilon(f_1, f_2)(x)| &\lesssim \|\omega\|_{\text{Dini}} \mathcal{M}(f_1, f_2)(x) + M_\eta(|T(f_1, f_2)|)(x) \\
&\quad + \left(\frac{1}{|B(x, \varepsilon/2)|} \int_{B(x, \varepsilon/2)} |T(f_1^0, f_2^0)(x')|^\eta dx' \right)^{1/\eta}.
\end{aligned}$$

For the last term, using Kolmogorov's inequality to relate the L^η and $L^{1/2, \infty}$ norms and the boundedness of T from $L^1 \times L^1$ to $L^{1/2, \infty}$, we obtain for any $\eta \in (0, \frac{1}{2})$,

$$\begin{aligned}
&\left(\frac{1}{|B(x, \varepsilon/2)|} \int_{B(x, \varepsilon/2)} |T(f_1^0, f_2^0)(x')|^\eta dx' \right)^{1/\eta} \\
&= \|T(f_1^0, f_2^0)\|_{L^\eta(B(x, \frac{\varepsilon}{2}), \frac{dx}{|B(x, \frac{\varepsilon}{2})|})} \\
&\leq C_\eta \|T(f_1^0, f_2^0)\|_{L^{1/2, \infty}(B(x, \frac{\varepsilon}{2}), \frac{dx}{|B(x, \frac{\varepsilon}{2})|})} \\
&\leq C_\eta \|T\|_{L^1 \times L^1 \rightarrow L^{1/2, \infty}} \mathcal{M}(f_1, f_2)(x).
\end{aligned}$$

Combining all the terms, we finally arrive at

$$\begin{aligned}
|\tilde{T}_\varepsilon(f_1, f_2)(x)| &\leq c_n (\|\omega\|_{\text{Dini}} + C_\eta \|T\|_{L^1 \times L^1 \rightarrow L^{1/2, \infty}}) \mathcal{M}(f_1, f_2)(x) \\
&\quad + M_\eta(|T(f_1, f_2)|)(x),
\end{aligned}$$

which taking into account (6.8) and (6.2) leads to the desired result. \square

As a corollary of the previous result follows the weak boundedness of the maximal truncation of T .

Corollary 6.10. *Let T be a bilinear Calderón–Zygmund operator with Dini-continuous kernel K . Then*

$$(6.11) \quad \|T_{\sharp}\|_{L^1 \times L^1 \rightarrow L^{1/2, \infty}} \lesssim (C_K + \|\omega\|_{\text{Dini}} + \|T\|_{L^{q_1} \times L^{q_2} \rightarrow L^q}).$$

Proof. Fix $\eta \in (0, 1/2)$ and use the previous result together with the weak boundedness of the multilinear maximal function and bilinear Calderón–Zygmund operators and the fact that $M_{\eta} \circ T : L^1 \times L^1 \rightarrow L^{1/2, \infty}$. To prove the latter, notice that for the Hardy–Littlewood maximal function using Lemma 2.11, we can write

$$M(f) \approx \sum_{u=1}^{3^n} M_u(f),$$

where

$$M_u(f) := \sup_{\substack{Q \ni x \\ Q \in \mathcal{D}^u}} \frac{1}{|Q|} \int_Q |f(y)| dy.$$

Therefore,

$$\left| \{x : M(|T(f_1, f_2)|^{\eta})(x)^{\frac{1}{\eta}} > \lambda\} \right| \leq \sum_{u=1}^{3^n} \left| \{x : M_u(|T(f_1, f_2)|^{\eta})(x)^{\frac{1}{\eta}} > \lambda/3^n\} \right|.$$

Denote

$$E_u := \{x \in \mathbb{R}^n : M_u(|T(f_1, f_2)|^{\eta})(x)^{\frac{1}{\eta}} > \lambda/3^n\}.$$

We can find a collection of maximal dyadic cubes $\{Q_j\}_j$ such that $E_u = \cup_j Q_j$ and

$$\frac{1}{|Q_j|} \int_{Q_j} |T(f_1, f_2)|^{\eta} > \lambda^{\eta} (3^n)^{-\eta},$$

which means that

$$|E_u| \leq (3^n)^{\eta} \lambda^{-\eta} \int_{E_u} |T(f_1, f_2)|^{\eta}, \quad u = 1, \dots, 3^n.$$

Now using Kolmogorov’s inequality and the fact that $T : L^1 \times L^1 \rightarrow L^{1/2, \infty}$, and assuming that $\eta < 1/2$, we get

$$\int_{E_u} |T(f_1, f_2)|^{\eta} \lesssim \|T(f_1, f_2)\|_{L^{\frac{1}{2}, \infty}(E_u, \frac{dx}{|E_u|})}^{\eta} |E_u| \leq \|f_1\|_1^{\eta} \|f_2\|_1^{\eta} |E_u|^{1-2\eta}$$

Combining both estimates, it follows that

$$|E_u| \leq \lambda^{-\eta} (3^n)^{\eta} \|f_1\|_1^{\eta} \|f_2\|_1^{\eta} |E_u|^{1-2\eta},$$

which is exactly,

$$\lambda |E_u|^2 \leq c_{n, \eta} \|f_1\|_1 \|f_2\|_1.$$

□

7. ACKNOWLEDGEMENTS

The authors would like to thank Prof. Tuomas Hytönen for suggesting this problem and for carefully reading this manuscript as well as many helpful comments which have improved the quality of this paper.

REFERENCES

- [1] T. A. Bui, J. M. Conde-Alonso, X. T. Duong, and M. Hormozi, Weighted bounds for multilinear operators with non-smooth kernels, *Studia Math.*, to appear. Available at [1506.07752](#), (2015).
- [2] T. A. Bui and M. Hormozi, Weighted bounds for multilinear square functions, *Potential Anal.*, to appear. Available at [1502.05490](#), (2015).
- [3] C. Cascante, J.M. Ortega, and I.E. Verbitsky, Nonlinear potentials and two weight trace inequalities for general dyadic and radial kernels, *Indiana Univ. Math. J.*, 53(3) (2004), 845–882.
- [4] W. Chen and W. Damián, Weighted estimates for the multisublinear maximal function, *Rendiconti del Circolo Matematico di Palermo*, 62(2013), 379–391.
- [5] D. Chung, M. Cristina Pereyra, and C. Pérez, Sharp bounds for general commutators on weighted Lebesgue spaces, *Trans. Amer. Math. Soc.*, Vol. 364 (3), (2012), 1163–1177.
- [6] J.M. Conde-Alonso, and G. Rey, A pointwise estimate for positive dyadic shifts and some applications, *Math. Ann.* 365 (2016), no. 3-4, 1111–1135.
- [7] W. Damián, A.K. Lerner, and C. Pérez, Sharp weighted bounds for multilinear maximal functions and Calderón–Zygmund operators, *J. Fourier Anal. Appl.* 21 (1) (2015), 161–181.
- [8] Do, Yen, and Thiele, Christoph. L^p theory for outer measures and two themes of Lennart Carleson united. *Bull. Amer. Math. Soc. (N.S.)* 52 (2015), no. 2, 249–296.
- [9] N. Fujii, Weighted bounded mean oscillation and singular integrals, *Math. Japon.* 22(5) (1977), 529–534.
- [10] L. Grafakos, *Classical Fourier Analysis*, Graduate Texts in Mathematics, Vol. 249, Springer, New York, (2014).
- [11] S. Hrusčev, A description of weights satisfying the A_∞ condition of Muckenhoupt, *Proc. Amer. Math. Soc.* 90(2) (1984), 253–257.
- [12] T. Hytönen, *The sharp weighted bound for general Calderón–Zygmund operators*, *Ann. of Math. (2)* 175 (2012), no. 3, 1473–1506.
- [13] T. Hytönen, M.T. Lacey, and C. Pérez, Sharp weighted bounds for the q -variation of singular integrals, *Bull. Lond. Math. Soc.*, 45 (3) (2013), 529–540.
- [14] T. Hytönen, and K. Li, Weak and strong $A_p - A_\infty$ estimates for square functions and related operators, preprint. Available at [1509.00273](#), (2015).
- [15] T. Hytönen and C. Pérez, Sharp weighted bounds involving A_∞ , *Anal. & P.D.E.*, 6(2013), 777–818.
- [16] T. Hytönen, L. Roncal, and O. Tapiola, Quantitative weighted estimates for homogeneous singular integrals, *Israel J. Math.*, to appear. Available at [1510.05789](#), (2015).
- [17] F. John and L. Nirenberg, On functions of bounded mean oscillation, *Comm. Pure Appl. Math.* 14(1961), 415–426.
- [18] J. L. Journé, Calderón–Zygmund operators, pseudo-differential operators and the Cauchy integral of Calderón, *Lect. Notes Math.* 994, Springer Verlag, (1983).
- [19] M.T. Lacey, An elementary proof of the A_2 bound, *Israel J. Math.*, to appear. Available at [1501.05818](#), (2015).
- [20] A.K. Lerner, A simple proof of the A_2 conjecture, *Int. Math. Res. Not. IMRN*, 14 (2013), 3159–3170.

- [21] A.K. Lerner, and F. Nazarov, Intuitive dyadic calculus: the basics, preprint, [1508.05639](#).
- [22] A.K. Lerner, S. Ombrosi, C. Pérez, R.H. Torres, and R. Trujillo-González, New maximal functions and multiple weights for the multilinear Calderón–Zygmund theory, *Adv. Math.*, 220 (2009), 1222–1264.
- [23] K. Li, Two weight inequalities for bilinear forms, *Collect. Math.*, to appear. Available at [1511.07250](#), (2015).
- [24] K. Li, K. Moen and W. Sun, The sharp weighted bound for multilinear maximal functions and Calderón–Zygmund operators, *J. Fourier Anal. Appl.* 20 (4) (2014), 751–765.
- [25] K. Li and W. Sun, Weak and strong type weighted estimates for multilinear Calderón–Zygmund operators, *Adv. Math.*, 254(2014), 736–771.
- [26] D. Maldonado and V. Naibo, Weighted norm inequalities for paraproducts and bilinear pseudodifferential operators with mild regularity, *J. Fourier Anal. Appl.* 15(2009), no. 2, 218–261.
- [27] C. Pérez and R. Torres, Minimal regularity conditions for the end-point estimate of bilinear Calderón–Zygmund operators, *Proc. Amer. Math. Soc., Series B*, Volume 1 (2014), 1–13.
- [28] S. Petermichl, and A. Volberg, *Heating of the Ahlfors–Beurling operator: weakly quasiregular maps on the plane are quasiregular*, *Duke Math. J.* 112 (2002), 281–305.
- [29] Thiele, Christoph, and Treil, Sergei, and Volberg, Alexander . Weighted martingale multipliers in the non-homogeneous setting and outer measure spaces. *Adv. Math.* 285 (2015), 1155–1188.
- [30] J.M. Wilson, Weighted inequalities for the dyadic square function without dyadic A_∞ , *Duke Math. J.*, 55(1) (1987), 19–50.

(W. Damián) DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O.B. 68 (GUSTAF HÄLLSTRÖMIN KATU 2B), FI-00014 UNIVERSITY OF HELSINKI, FINLAND
E-mail address: wendolin.damian.gonzalez@gmail.com

(K. Li) DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O.B. 68 (GUSTAF HÄLLSTRÖMIN KATU 2B), FI-00014 UNIVERSITY OF HELSINKI, FINLAND.
Current address: BCAM–BASQUE CENTER FOR APPLIED MATHEMATICS, MAZARREDO, 14. 48009 BILBAO BASQUE COUNTRY, SPAIN
E-mail address: kli@bcamath.org

(M. Hormozi) UNIVERSITY OF GOTHENBURG & CHALMERS UNIVERSITY OF TECHNOLOGY SE-412 96 GÖTEBORG, SWEDEN
E-mail address: me.hormozi@gmail.com