

A Lower Bound on Local Energy of Partial Sum of Eigenfunctions for Laplace-Beltrami Operators*

Qi Lü [†]

Abstract

In this paper, a lower bound is established for the local energy of partial sum of eigenfunctions for Laplace-Beltrami operators (in Riemannian manifolds with low regularity data) with general boundary condition. This result is a consequence of a new pointwise and weighted estimate for Laplace-Beltrami operators, a construction of some nonnegative function with arbitrary given critical point location in the manifold, and also two interpolation results for solutions of elliptic equations with lateral Robin boundary conditions.

2010 Mathematics Subject Classification. Primary 93B07.

Key Words. Lower bound, local energy, partial sum of eigenfunctions, Laplace-Beltrami operator, Robin boundary condition.

1 Introduction and Main Result

Let M be a d ($d \in \mathbb{N}$) dimensional connected compact C^1 -smooth Riemannian manifold with an C^2 -smooth boundary Γ , and ω a nonempty open subset of M . Denote by g the C^1 -smooth Riemannian metric tensor on M ; by \mathcal{D}_g the Levi-Civita connection on M induced by g ; by ∇_M , div_M and Δ_M the gradient operator, the divergence operator and the Laplace-Beltrami operator (on M) given by \mathcal{D}_g , respectively; by $(\cdot, \cdot)_g$ and $|\cdot|_g$ the inner product and the norm for the tangent vector of M with respect to g , respectively; by $d_g x$ the volume element of M with respect to g ; and by $d_g \Gamma$ the volume element of Γ induced by g . We refer to [3] for more details on the notation/tool used in this paper, say Sobolev spaces on Riemannian manifold. Fix any $T > 0$, and put $Q = (0, T) \times M$ and $\Sigma = (0, T) \times \Gamma$. Throughout this paper, we use $C = C(M, \omega, d, g, T)$ to denote a generic positive constant, which may change from one place to another.

We define an unbounded operator A on $L^2(M)$ by

$$\begin{cases} D(A) = \left\{ u \in H^2(M) : \tilde{l} \frac{\partial_M u}{\partial \nu} + lu = 0 \text{ on } \Gamma \right\}, \\ Au = -\Delta_M u, \quad \forall u \in D(A), \end{cases} \quad (1.1)$$

*This work is partially supported by the NSF of China under grants 10831007, 11101070 and 60974035. This paper is an improved version of one chapter of the author's Ph D thesis ([13]) accomplished at Sichuan University under the guidance of Professor Xu Zhang. The author would like to take this opportunity to thank him deeply for his help.

[†]School of Mathematical Sciences, University of Electronic Science and Technology of China, Chengdu 610054, China; and Basque Center for Applied Mathematics (BCAM), Mazarredo, 14. 48009 Bilbao Basque Country - Spain. *E-mail:* luqi59@163.com.

where $\nu = \nu(x)$ is the unit outward normal vector of M at $x \in \Gamma$ with respect to the metric g , $\frac{\partial_M u}{\partial \nu} \Big|_{\Gamma} = (\nabla_M u, \nu)_g|_{\Gamma}$, both \tilde{l} and l belong to $L^\infty(\Gamma)$ and satisfy $\tilde{l} = 1$, $l \geq 0$ or $\tilde{l} = 0$, $l > 0$. Let $\{\lambda_i\}_{i=1}^\infty$ be the eigenvalues of A , and $\{e_i\}_{i=1}^\infty$ the corresponding eigenfunctions satisfying $\|e_i\|_{L^2(M)} = 1$. It is easy to show that $0 \leq \lambda_1 \leq \lambda_2 \leq \dots$, and $\{e_i\}_{i=1}^\infty$ constitutes an orthonormal basis of $L^2(M)$.

One can find the following result from [4, 7, 9].

Theorem 1.1 *If both Γ and g are C^∞ , $\tilde{l} = 0$ and $l = 1$, then it holds that*

$$\sum_{\lambda_i \leq r} |a_i|^2 \leq C e^{C\sqrt{r}} \int_{\omega} \left| \sum_{\lambda_i \leq r} a_i e_i(x) \right|^2 d_g x, \quad (1.2)$$

for every $r > 0$ and every choice of the coefficients $\{a_i\}_{\lambda_i \leq r}$ with $a_i \in \mathbb{C}$.

This result provides a delicate lower bound estimate for the local energy of partial sum of eigenfunctions for Laplace-Beltrami operators (in C^∞ -smooth Riemannian manifolds) with Dirichlet boundary condition. As remarked in [22], the power $\frac{1}{2}$ in the above $e^{C\sqrt{r}}$ is sharp. In terms of the control theory language, inequality (1.2) can be viewed as an observability estimate for partial sum of eigenfunctions for operator A . Besides its obviously independent interest, this inequality has many applications in control theory. In [7], by means of a time iteration approach, Lebeau and Robbiano used (1.2) to obtain null controllability of the heat equation with homogeneous Dirichlet boundary condition. In [9], inequality (1.2) was addressed by Lebeau and Zuazua, and via which null controllability of a linear system of thermoelasticity was analyzed. Further applications of this inequality to controllability problems can be found in [11, 15, 16, 21]. On the other hand, in [19], Wang used (1.2) to establish an L^∞ -null controllability for the heat equation, and especially, via which he solved a long-standing open problem in control theory for infinite dimensional systems, i.e., the Bang-Bang principle for time optimal control problem for the heat equation with a locally distributed controller. His results was recently extended to fractional order parabolic equations, see [12].

We remark that, in Theorem 1.1, both Γ and g are assumed to be C^∞ -smooth. L. Escauriaza pointed out that the C^∞ -regularity for Γ can be weakened to be C^2 but his proof was not published (see Remark 1.1 in [11]). In this paper, we shall address the sharp result in this respect and, in particular, consider a similar problem but with more general boundary conditions.

The main result of this paper can be stated as follows:

Theorem 1.2 *The conclusion in Theorem 1.1 still holds when the additional assumptions on Γ , g , \tilde{l} and l therein are dropped.*

Noting that the time iteration method developed in [7] does not depend on the boundary condition. Therefore, using Theorem 1.2 and this method, it is easy to obtain the corresponding controllability/optimal control results for equations with Robin boundary condition. On the other hand, Theorem 1.2 can also be employed to prove the null/approximate controllability of forward stochastic heat equations ([14]), which is, to the best of the author's knowledge, the first controllability result for forward stochastic partial differential equations with control acts only on the drift term.

Theorem 1.2 needs much lower regularities for both Γ and g than Theorem 1.1. Furthermore, Theorem 1.2 is for general Robin type boundary condition while Theorem 1.1 addresses only the homogeneous Dirichlet boundary condition.

In [4, 7, 9], the authors employed a local Carleman estimate to establish Theorem 1.1. The homogeneous Dirichlet boundary condition plays an important role in their proof. However, it seems to be quite difficult to prove Theorem 1.2 by using the same method. Instead, in this paper, we shall use a global (in space) Carleman estimate to overcome the difficulties introduced by the general boundary condition. On the other hand, it deserves to point out that, although a related global Carleman estimate was established in [2] addressing observability estimates for quite general parabolic equations, the approach therein does not seem to be able to provide the desired sharp estimate “ $e^{C\sqrt{r}}$ ” in Theorem 1.2. Indeed, in order to prove Theorem 1.2, we need to derive first a new pointwise and weighted estimate for Laplace-Beltrami operators (see Section 2), and then to prove the existence of a nonnegative function with arbitrary given critical point location in manifold M (see Section 3), and also to show some interpolation results for solutions of elliptic equations with lateral Robin boundary conditions in a cylinder (see Sections 4 and 5).

It is considerably easier to prove Theorem 1.2 with $\tilde{l} = 0$ and $l > 0$ than the case with $\tilde{l} = 1$ and $l \geq 0$. Noting that in both cases we can use the same method to obtain the desired inequalities. Therefore in the sequel we only prove Theorem 1.2 for the case that $\tilde{l} = 1$ and $l \geq 0$. The proof of this theorem will be given in Section 6. Note also that, even for the case of Dirichlet boundary condition, our method seems to be more elementary and also self-contained than that in [4, 7, 9].

2 A pointwise and weighted estimate for Laplace-Beltrami operators

In this section, we establish a pointwise weighted estimate for Laplace-Beltrami operators on a given Riemannian manifold, which will play a key role in the sequel.

Let N be a n -dimensional ($n \in \mathbb{N}$) Riemannian manifold with a C^1 -metric tensor b . The meaning of $(\cdot, \cdot)_b$, $|\cdot|_b$, ∇_N , div_N , Δ_N and so on can be understood similarly as mentioned at the very beginning of Section 1.

Let H , H_1 and H_2 be any given C^1 -vector fields on N . We recall the following well-known formulas which will be useful later (e.g. [3, Chapter 1], [5, Chapter 3]).

$$\operatorname{div}_N(hH) = (\nabla_N h, H)_b + h \operatorname{div}_N H, \quad \forall h \in C^1(N). \quad (2.1)$$

$$\nabla_N(H_1, H_2)_b = (\nabla_N H_1, H_2)_b + (\nabla_N H_2, H_1)_b + (\nabla_N b)(H_1, H_2), \quad (2.2)$$

where $(\nabla_N H_i, H_j)_b$ stands for the contraction of the tensor $b \otimes \nabla_N H_i \otimes H_j$ ($1 \leq i, j \leq 2, i + j = 3$), $(\nabla_N b)(H_1, H_2)$ stands for the contraction of the tensor $\nabla_N b \otimes H_1 \otimes H_2$. Also, for any $f \in C^1(N)$, we denote by $\nabla_N(\nabla_N f)$ the Hessian of f .

In the sequel, for arbitrary real function $\varphi \in C^2(N)$ and arbitrary positive real numbers s and λ , we choose functions α and θ as follows:

$$\alpha = e^{\lambda\varphi}, \quad \theta = e^{s\alpha}. \quad (2.3)$$

We have the following result:

Theorem 2.1 *Assume $v \in C^2(N)$ and put $w = \theta v$. Then it holds that*

$$\begin{aligned} & 2\theta^2 \left| \Delta_N v \right|_b^2 + D \\ & \geq B_1 \left| \nabla_N w \right|_b^2 + B_2 w^2 + 4s\lambda^2 \left(\nabla_N(\alpha \left| \nabla_N \varphi \right|_b^2), \nabla_N w \right)_b w \\ & \quad + 4s\lambda^2 \alpha \left(\nabla_N w, \nabla_N \varphi \right)_b^2 + 4s\lambda \alpha \left(\nabla_N w, \left(\left[\nabla_N \left(\nabla_N \varphi \right) \right], \nabla_N w \right)_b \right)_b \end{aligned}$$

$$+4s\lambda\alpha\left(\nabla_N w, (\nabla_N b)(\nabla_N w, \nabla_N \varphi)\right)_b - 2s\lambda\alpha\left((\nabla_N b)(\nabla_N w, \nabla_N w), \nabla_N \varphi\right)_b, \quad (2.4)$$

where

$$\left\{ \begin{array}{l} D = 2s\lambda\operatorname{div}_N \left[2\lambda\alpha|\nabla_N \varphi|_b^2 w \nabla_N w + s^2\lambda^2\alpha^3|\nabla_N \varphi|_b^2 w^2 \nabla_N \varphi \right. \\ \quad \left. + 2\alpha(\nabla_N \varphi, \nabla_N w)_b \nabla_N w - \alpha|\nabla_N w|_b^2 \nabla_N \varphi \right] \\ B_1 = 2s\lambda^2\alpha|\nabla_N \varphi|_b^2 - 2s\lambda\alpha\Delta_N \varphi|\nabla_N \varphi|_b^2 - s\lambda\alpha\left((\nabla_N b)(\nabla_N w, \nabla_N w), \nabla_N \varphi\right)_b \\ \quad = 2s\lambda^2\alpha|\nabla_N \varphi|_b^2 - s\alpha O(\lambda), \\ B_2 = 2s^3\lambda^4\alpha^3|\nabla_N \varphi|_b^4 + 2s^3\lambda^3\alpha^3\operatorname{div}_N(|\nabla_N \varphi|_b^2 \nabla_N \varphi) - 4s^2\lambda^2\alpha^2|\Delta_N \varphi|^2 - 4s^2\lambda^4\alpha^2|\nabla_N \varphi|_b^2 \\ \quad = 2s^3\lambda^4\alpha^3|\nabla_N \varphi|_b^4 - s^3\alpha^3 O(\lambda^3) - s^2\alpha^2 O(\lambda^4). \end{array} \right. \quad (2.5)$$

Remark 2.1 *There exist several pointwise and weighted estimates for second order partial differential operators in the literature (e.g., [1, 6, 10, 18, 20]). These estimates are quite useful in control theory and inverse problems for partial differential equations. In [18, Theorem 2.2], one can find an estimate similar to (2.4). The main advantage of our estimate (2.4) consists in that it is more convenient to deal with the Robin boundary condition, as shown in the proof of Theorem 4.1.*

Proof of Theorem 2.1: By the definition of v and w , we have that

$$\nabla_N v = \nabla_N(\theta^{-1}w) = w\nabla_N(\theta^{-1}) + \theta^{-1}\nabla_N w = -s\lambda\theta^{-1}\alpha w \nabla_N \varphi + \theta^{-1}\nabla_N w. \quad (2.6)$$

Hence, by (2.1), it follows that

$$\begin{aligned} -\theta\operatorname{div}_N(\nabla_N v) &= -\theta\operatorname{div}_N(-s\lambda\theta^{-1}\alpha w \nabla_N \varphi + \theta^{-1}\nabla_N w) \\ &= -\Delta_N w + 2s\lambda\alpha(\nabla_N \varphi, \nabla_N w)_b + s\lambda^2\alpha|\nabla_N \varphi|_b^2 w \\ &\quad - s^2\lambda^2\alpha^2|\nabla_N \varphi|_b^2 w + s\lambda\alpha w \Delta_N \varphi. \end{aligned} \quad (2.7)$$

Put

$$\left\{ \begin{array}{l} I_1 = -\Delta_N w - s^2\lambda^2\alpha^2|\nabla_N \varphi|_b^2 w, \\ I_2 = 2s\lambda\alpha(\nabla_N \varphi, \nabla_N w)_b + 2s\lambda^2\alpha|\nabla_N \varphi|_b^2 w, \\ I_3 = -\theta\Delta_N v - s\lambda\alpha w \Delta_N \varphi + s\lambda^2\alpha|\nabla_N \varphi|_b^2 w. \end{array} \right. \quad (2.8)$$

By (2.7)–(2.8), we see that $I_1 + I_2 = I_3$. Hence

$$2I_1 I_2 \leq |I_3|^2. \quad (2.9)$$

We estimate $|I_3|^2$ first.

$$\begin{aligned} |I_3|^2 &= \left| -\theta\Delta_N v - s\lambda\alpha w \Delta_N \varphi + s\lambda^2\alpha|\nabla_N \varphi|_b^2 w \right|^2 \\ &\leq 2\theta^2|\Delta_N v|^2 + 4s^2\lambda^2\alpha^2|\Delta_N \varphi|^2|w|^2 + 4s^2\lambda^4\alpha^2\left|\nabla_N \varphi\right|_b^4|w|^2. \end{aligned} \quad (2.10)$$

Next, let us estimate $I_1 I_2$. By (2.8), it follows that

$$\begin{aligned} I_1 I_2 &= 2s\lambda\alpha\left(-\Delta_N w - s^2\lambda^2\alpha^2|\nabla_N \varphi|_b^2 w\right)\left((\nabla_N \varphi, \nabla_N w)_b + \lambda|\nabla_N \varphi|_b^2 w\right) \\ &= 2s\lambda^2\alpha\left(-\Delta_N w - s^2\lambda^2\alpha^2|\nabla_N \varphi|_b^2 w\right)|\nabla_N \varphi|_b^2 w \\ &\quad - 2s^3\lambda^3\alpha^3|\nabla_N \varphi|_b^2(\nabla_N \varphi, \nabla_N w)_b w - 2s\lambda\alpha\Delta_N w(\nabla_N \varphi, \nabla_N w)_b. \end{aligned} \quad (2.11)$$

We need to compute the terms in the right-hand side of (2.11) one by one. By formula (2.1), we find that

$$\begin{aligned}
& 2s\lambda^2\alpha\left(-\Delta_N w - s^2\lambda^2\alpha^2|\nabla_N\varphi|_b^2 w\right)|\nabla_N\varphi|_b^2 w \\
&= -2s^3\lambda^4\alpha^3|\nabla_N\varphi|_b^4 w^2 - \operatorname{div}_N\left(2s\lambda^2\alpha|\nabla_N\varphi|_b^2 w\nabla_N w\right) \\
& \quad + 2s\lambda^2\left(\nabla_N(\alpha|\nabla_N\varphi|^2), \nabla_N w\right)_b w + 2s\lambda^2\alpha|\nabla_N\varphi|_b^2|\nabla_N w|_b^2. \tag{2.12}
\end{aligned}$$

Further,

$$\begin{aligned}
& -2s^3\lambda^3\alpha^3|\nabla_N\varphi|_b^2(\nabla_N\varphi, \nabla_N w)_b w \\
&= -\operatorname{div}_N\left(s^3\lambda^3\alpha^3|\nabla_N\varphi|_b^2 w^2\nabla_N\varphi\right) + 3s^3\lambda^4\alpha^3|\nabla_N\varphi|_b^4 w^2 \\
& \quad + s^3\lambda^3\alpha^3\operatorname{div}_N\left(|\nabla_N\varphi|_b^2\nabla_N\varphi\right)w^2. \tag{2.13}
\end{aligned}$$

Further,

$$\begin{aligned}
& -2s\lambda\alpha\Delta_N w(\nabla_N\varphi, \nabla_N w)_b \\
&= -\operatorname{div}_N\left(2s\lambda\alpha(\nabla_N\varphi, \nabla_N w)_b\nabla_N w\right) + 2s\lambda^2\alpha(\nabla_N\varphi, \nabla_N w)_b^2 \\
& \quad + 2s\lambda\alpha\left(\nabla_N w, \nabla_N(\nabla_N w, \nabla_N\varphi)_b\right)_b.
\end{aligned}$$

By formula (2.2),

$$\begin{aligned}
\left(\nabla_N w, \nabla_N(\nabla_N w, \nabla_N\varphi)_b\right)_b &= \left(\nabla_N w, \left([\nabla_N(\nabla_N w)], \nabla_N\varphi\right)_b\right)_b + \left([\nabla_N(\nabla_N\varphi)], \nabla_N w\right)_b^2 \\
& \quad + \left(\nabla_N w, (\nabla_N b)(\nabla_N w, \nabla_N\varphi)_b\right)_b.
\end{aligned}$$

Noting that

$$2\left(\nabla_N w, \left([\nabla_N(\nabla_N w)], \nabla_N\varphi\right)_b\right)_b = \left(\nabla_N|\nabla_N w|_b^2, \nabla_N\varphi\right)_b - \left((\nabla_N b)(\nabla_N w, \nabla_N w), \nabla_N\varphi\right)_b,$$

we arrive at

$$\begin{aligned}
& 2s\lambda\alpha\left(\nabla_N w, \nabla_N(\nabla_N w, \nabla_N\varphi)_b\right)_b \\
&= s\lambda\alpha\left(\nabla_N|\nabla_N w|_b^2, \nabla_N\varphi\right)_b - s\lambda\alpha\left((\nabla_N b)(\nabla_N w, \nabla_N w), \nabla_N\varphi\right)_b \\
& \quad + 2s\lambda\alpha\left(\nabla_N w, \left([\nabla_N(\nabla_N\varphi)], \nabla_N w\right)_b\right)_b + 2s\lambda\alpha\left(\nabla_N w, (\nabla_N b)(\nabla_N w, \nabla_N\varphi)_b\right)_b \\
&= s\lambda\operatorname{div}_N\left(\alpha|\nabla_N w|_b^2\nabla_N\varphi\right) - s\lambda^2\alpha|\nabla_N\varphi|_b^2|\nabla_N w|_b^2 - s\lambda\alpha\Delta_N\varphi|\nabla_N w|_b^2 \\
& \quad - s\lambda\alpha\left((\nabla_N b)(\nabla_N w, \nabla_N w), \nabla_N\varphi\right)_b + 2s\lambda\alpha\left(\nabla_N w, \left([\nabla_N(\nabla_N\varphi)], \nabla_N w\right)_b\right)_b \\
& \quad + 2s\lambda\alpha\left(\nabla_N w, (\nabla_N b)(\nabla_N w, \nabla_N\varphi)_b\right)_b.
\end{aligned}$$

Therefore it holds

$$\begin{aligned}
& -2s\lambda\alpha\Delta_N w(\nabla_N\varphi, \nabla_N w)_b \\
&= -\operatorname{div}_N\left(2s\lambda\alpha(\nabla_N\varphi, \nabla_N w)_b\nabla_N w\right) + 2s\lambda^2\alpha(\nabla_N\varphi, \nabla_N w)_b^2 + \operatorname{div}_N\left(s\lambda\alpha|\nabla_N w|_b^2\nabla_N\varphi\right) \\
& \quad - s\lambda\alpha\left((\nabla_N b)(\nabla_N w, \nabla_N w), \nabla_N\varphi\right)_b - s\lambda^2\alpha|\nabla_N\varphi|_b^2|\nabla_N w|_b^2 - s\lambda\alpha\Delta_N\varphi|\nabla_N w|_b^2 \\
& \quad + 2s\lambda\alpha\left(\nabla_N w, \left([\nabla_N(\nabla_N\varphi)], \nabla_N w\right)_b\right)_b + 2s\lambda\alpha\left(\nabla_N w, (\nabla_N b)(\nabla_N w, \nabla_N\varphi)_b\right)_b. \tag{2.14}
\end{aligned}$$

Finally, by (2.9)–(2.14), we obtain (2.4). \square

3 A nonnegative function with an arbitrary given critical point location in the manifold

In this section, we prove the existence of a nonnegative function with an arbitrary given critical point location in manifold M . This result is a modification of the corresponding result in [2] for flat spaces. In the sequel, this construction will play a key role in the choice of the weight function in our global Carleman estimate.

Our result is stated as follows:

Theorem 3.1 *There exists a function $\psi \in C^2(M)$ such that $\psi > 0$ in M , $\psi = 0$ on Γ and*

$$|\nabla_M \psi|_g^2 > 0, \quad \forall x \in \overline{M} \setminus \omega_0, \quad (3.1)$$

where ω_0 is an arbitrary fixed nonempty open subset of M such that $\overline{\omega_0} \subset \omega$.

Proof of Theorem 3.1: We borrow some idea from [2]. Choose a function $p \in C^2(M)$ such that

$$p > 0 \text{ in } M, \quad p = 0 \text{ and } |\nabla_M p|_g > 0 \text{ on } \Gamma. \quad (3.2)$$

By the density of Morse functions in $C^2(M)$ (see [17, Chapter 1]), there exists a sequence of Morse functions $\{p^k(x)\}_{k=1}^\infty$ such that

$$p^k \rightarrow p \text{ in } C^2(M), \text{ as } k \rightarrow \infty. \quad (3.3)$$

Denote by $M_1 = \{x \in M \mid \nabla_M p(x) = 0\}$ the set of critical points of function p . Since $|\nabla_M p|_g > 0$ on ∂M , there exist a positive number $\xi_1 > 0$ and an open set $M_2 \subset M$ such that

$$|\nabla_M p|_g > \xi_1 > 0 \text{ in } M_2, \quad M_1 \cap \overline{M_2} = \emptyset, \quad \Gamma \subset \overline{M_2}. \quad (3.4)$$

Let $f \in C^\infty(M)$ such that

$$f = 1 \text{ on } \Gamma, \quad f = 0 \text{ in } M \setminus \overline{M_2}. \quad (3.5)$$

Put $q^k(x) = p^k(x) + f(x)[p(x) - p^k(x)]$. By the definition of q^k , we know

$$q^k = 0 \text{ on } \Gamma, \quad \nabla_M q^k = \nabla_M p^k \text{ in } M \setminus \overline{M_2} \quad (3.6)$$

and

$$\nabla_M q^k(x) = \nabla_M p^k(x) + f(x)[\nabla_M p(x) - \nabla_M p^k(x)] + \nabla_M f(x)[p(x) - p^k(x)]. \quad (3.7)$$

By (3.3), we know that there exists a $\bar{k} \in \mathbb{N}$ such that for any integer $k > \bar{k}$, we have

$$f(x)[\nabla_M p(x) - \nabla_M p^k(x)] + \nabla_M f(x)[p(x) - p^k(x)] < \frac{\xi_1}{2}. \quad (3.8)$$

From (3.4), (3.7) and (3.8), for any integer $k_1 > \bar{k}$, it follows that

$$|\nabla_M q^{k_1}|_g > 0 \text{ in } \overline{M_2}. \quad (3.9)$$

Letting $q(x) = q^{k_1}(x)$, we know that q is a Morse function satisfying $|\nabla_M q|_g > 0$ in $\overline{M_2}$.

Denote by CP_1 the set of critical points of function q , i.e., $CP_1 = \{x \in M \mid \nabla_M q(x) = 0\}$. Hence CP_1 is a finite set. Assume $CP_1 = \{x_1, x_2, \dots, x_m\}$. Consider a sequence of functions $\{\rho^i\}_{i=1}^m \subset C^\infty([0, 1]; M)$ such that

$$\begin{cases} \rho^i(t) \in M, \forall t \in [0, 1], \rho^i(t_1) \neq \rho^i(t_2), \forall t_1, t_2 \in [0, 1], t_1 \neq t_2, i = 1, \dots, m, \\ \rho^i(1) = x_i, \rho^i(0) \in \omega_1, i = 1, \dots, m, \\ \rho^i(t_1) \neq \rho^j(t_2), \forall i \neq j, \forall t_1, t_2 \in [0, 1], \end{cases} \quad (3.10)$$

where ω_1 is a nonempty open set such that $\overline{\omega_1} \subset \omega_0$. By (3.10), there exists a sequence of C^2 -vector fields $\{\eta^i\}_{i=1}^m$ on M and a sequence of C^∞ -functions $\{\gamma^i\}_{i=1}^m$ on M such that

$$\frac{d\rho^i(t)}{dt} = \eta^i(\rho^i(t)), \quad \text{in } [0, 1], i = 1, \dots, m, \quad (3.11)$$

$$\text{supp } \gamma^i \subset M, \quad i = 1, \dots, m, \quad (3.12)$$

$$\text{supp } \gamma^i \cap \text{supp } \gamma^j = \emptyset, \quad \forall i \neq j, \quad (3.13)$$

$$\gamma^i(\rho^i(t)) = 1, \quad \forall t \in [0, 1], i = 1, \dots, m. \quad (3.14)$$

Let $V^i(x) = \gamma^i(x)\eta^i(x)$. Consider the system of the ordinary differential equations on manifold M as follows:

$$\begin{cases} \frac{dx}{dt} = V^i(x), \\ x(0) = x_0. \end{cases} \quad (3.15)$$

Denote by $S_t^i : M \rightarrow M$ ($i = 1, \dots, m$) the operator such that $S_t^i(x_0) = x(t)$, where $x(t)$ is the solution of equation (3.15). Hence S_t^i ($i = 1, \dots, m$) are diffeomorphisms on M .

By (3.10), (3.11) and (3.14), we have

$$S_1^i(\rho^i(0)) = x_i, \quad i = 1, \dots, m. \quad (3.16)$$

Put $S(x) = S_1^1 \circ S_1^2 \circ \dots \circ S_1^m$ and $\psi(x) = q(S(x))$. By (3.12), there exists a domain $M_3 \subset M$ such that $\Gamma \subset \overline{M_3}$ and

$$S_1^i(x) = x, \quad \forall x \in M_3, i = 1, \dots, m. \quad (3.17)$$

Therefore $\psi(x) = q(x)$, $\forall x \in M_3$. Hence $\psi(x) = 0$, $\forall x \in \partial M$. Denote by CP_2 the critical points of ψ . Since the mapping S is a diffeomorphism, we have

$$CP_2 = \left\{ x \in M \mid S(x) \in CP_1 \right\}. \quad (3.18)$$

By (3.13), we have

$$S(\rho^i(0)) = x_i, \quad i = 1, \dots, m. \quad (3.19)$$

It follows from (3.18) and (3.19) that $CP_2 \subset \omega_0$, which completes the proof. \square

4 Interpolation inequality I

This section is devoted to showing an interpolation result for solutions to the following elliptic equation:

$$\begin{cases} u_{tt} + \Delta_M u = 0 & \text{in } Q, \\ \frac{\partial_M u}{\partial \nu} + l(x)u = 0 & \text{on } \Sigma. \end{cases} \quad (4.1)$$

Our result reads:

Theorem 4.1 *Let $0 < \gamma < \frac{T}{2}$ and $2\gamma < T' < T'' < T - \gamma$. Then there exists a constant $\mu \in (0, 1)$ such that any solution $u \in H^2(Q)$ of (4.1) satisfies*

$$|u|_{L^2(M \times (T', T''))} \leq C |u|_{L^2(\omega \times (\gamma, T - \gamma))}^\mu |u|_{H^1(Q)}^{1-\mu}. \quad (4.2)$$

This sort of interpolation estimate has already appeared in the framework of boundary control and stabilization for hyperbolic equations (e.g.[8]) and also for inverse problems (e.g.[18]).

Proof of Theorem 4.1: We borrow some ideas from [18]. The key is to use Theorem 2.1. The proof is divided into five steps.

Step 1. Firstly, we will explain the construction of the weight function θ appeared in Theorem 2.1. By (3.1), we have

$$h \triangleq \frac{1}{|\psi|_{L^\infty(M)}} \min_{x \in M \setminus \omega_0} |\nabla \psi(x)|_g > 0. \quad (4.3)$$

Without loss of generality, let us assume that $T' \leq T - T''$. Let

$$a = \frac{T}{2} - 2\gamma, \quad a_0 = \frac{T - T' - 2\gamma}{2}, \quad a_1 = \frac{T}{2} - \gamma \quad (4.4)$$

It is easy to check that

$$\frac{T}{2} - T' < a_0 < a < a_1 < \frac{T}{2}.$$

We choose

$$\varphi(x, t) = (c_1 - c_2) \frac{\psi(x)}{|\psi|_{L^\infty(M)}} + a^2 - \left(t - \frac{T}{2}\right)^2 + \kappa \quad (4.5)$$

and

$$\tilde{\varphi}(x, t) = -(c_1 - c_2) \frac{\psi(x)}{|\psi|_{L^\infty(M)}} + a^2 - \left(t - \frac{T}{2}\right)^2 + \kappa, \quad (4.6)$$

where $c_1 = a^2 - \left(\frac{T}{2} - T'\right)^2$, $c_2 = a^2 - \frac{1}{2} \left(\frac{T}{2} - T'\right)^2 - a_0^2$ and κ is chosen to be large enough to make $\tilde{\varphi} > 0$. It is easy to check that $c_1 > c_2$.

These give the functions $\alpha(x, t) = e^{\lambda\varphi(x, t)}$, $\tilde{\alpha}(x, t) = e^{\lambda\tilde{\varphi}(x, t)}$, $\theta = e^{s\alpha}$ and $\tilde{\theta} = e^{s\tilde{\alpha}}$. It is obvious that $0 < \tilde{\varphi} \leq \varphi$, $1 < \tilde{\alpha} \leq \alpha$ and $1 < \tilde{\theta} \leq \theta$.

By the definition of α , it is easy to check that

$$\begin{cases} \alpha(\cdot, t) \geq e^{c_1\lambda + \lambda\kappa}, & \left|t - \frac{T}{2}\right| \leq \frac{T}{2} - T', \\ \alpha(\cdot, t) \leq e^{(c_1 - c_2)\lambda + \lambda\kappa}, & \left|t - \frac{T}{2}\right| \geq a. \end{cases} \quad (4.7)$$

Noting that equation (4.1) has only partial boundary condition. We need to reduce it into an equation with full boundary condition. For this, let us choose a cut-off function $\phi(t) \in C_0^\infty\left(\frac{T}{2} - a_1, \frac{T}{2} + a_1\right) = C_0^\infty(\gamma, T - \gamma)$ such that

$$\begin{cases} 0 \leq \phi(t) \leq 1, & t \in (\gamma, T - \gamma), \\ \phi(t) = 1, & \left|t - \frac{T}{2}\right| \leq \frac{T}{2} - a. \end{cases} \quad (4.8)$$

Let $u_1 = \phi u$, noticing that ϕ is independent of x , it follows by equation (4.1) that

$$\begin{cases} (u_1)_{tt} + \Delta_M u_1 = \phi_{tt} u + 2\phi_t u_t & \text{in } Q, \\ \frac{\partial_M u_1}{\partial \nu} + l(x)u_1 = 0 & \text{on } \Sigma, \\ u_1 = 0 & \text{on } (M \times \{0\}) \cup (M \times \{T\}). \end{cases} \quad (4.9)$$

By (4.8), we know that there is a $Q_0 \subset Q$ such that

$$\begin{cases} \text{supp}(u_1) \subset Q_0, \\ \partial Q_0 \text{ is } C^2. \end{cases} \quad (4.10)$$

Put $\Sigma_0 = \partial Q_0 \cap \Sigma$.

Step 2. We now apply Theorem 2.1 to equation (4.9) with $n = d + 1$, $N = Q_0$, $b = 1 \otimes g$, v being replaced by u_1 , φ is as (4.5) and $w = \theta u_1$.

Integrating equality (2.4) on Q_0 , we obtain that

$$\begin{aligned} & \int_{Q_0} 2\theta^2 \left| (u_1)_{tt} + \Delta_M u_1 \right|^2 d_g x dt + \int_{Q_0} D d_g x dt \\ & \geq \int_{Q_0} B_1 |\nabla_N w|_b^2 d_g x dt + \int_{Q_0} B_2 w^2 d_g x dt + 4s\lambda^2 \int_{Q_0} \left(\nabla_N (\alpha |\nabla_N \varphi|_b^2), \nabla_N w \right)_b w d_g x dt \\ & \quad + 4s\lambda^2 \int_{Q_0} \alpha (\nabla_N w, \nabla_N \varphi)_b^2 d_g x dt + 4s\lambda \int_{Q_0} \alpha \left(\nabla_N w, \left[\nabla_N (\nabla_N \varphi) \right] \nabla_N w \right)_b d_g x dt \\ & \quad + 4s\lambda \int_{Q_0} \alpha \left(\nabla_N w, (\nabla_N b) (\nabla_N w, \nabla_N \varphi) \right)_b d_g x dt - 2s\lambda \int_{Q_0} \alpha \left((\nabla_N b) (\nabla_N w, \nabla_N w), \nabla_N \varphi \right)_b d_g x dt, \end{aligned} \quad (4.11)$$

Let us estimate the right-hand side of (4.11). By Cauchy-Schwarz inequality and noting that $\varphi \in C^2(Q_0)$, we have the following estimates:

$$4s\lambda^2 \left| \left(\nabla_N (\alpha |\nabla_N \varphi|_b^2), \nabla_N w \right)_b w \right| \leq C \left(s^2 \lambda^4 \alpha w^2 + \lambda^2 |\nabla_M w|_g^2 + \lambda^2 |w_t|^2 \right), \quad (4.12)$$

$$4s\lambda \left| \alpha \left(\nabla_N w, \left[\nabla_N (\nabla_N \varphi) \right] \nabla_N w \right)_b \right| \leq Cs\lambda \alpha \left(|\nabla_M w|_g^2 + |w_t|^2 \right), \quad (4.13)$$

$$4s\lambda \left| \alpha \left(\nabla_N w, (\nabla_N b) (\nabla_N w, \nabla_N \varphi) \right)_b \right| \leq Cs\lambda \alpha \left(|\nabla_M w|_g^2 + |w_t|^2 \right). \quad (4.14)$$

$$s\lambda \left| \alpha \left((\nabla_N b)(\nabla_N w, \nabla_N w), \nabla_N \varphi \right)_b \right| \leq Cs\lambda \alpha \left(|\nabla_M w|_g^2 + |w_t|^2 \right). \quad (4.15)$$

By the definition of B_1 , we have that

$$B_1 |\nabla_N w|_b^2 = \left[2s\lambda^2 \alpha |\nabla_N \varphi|_b^2 - s\alpha O(\lambda) \right] (|\nabla_M w|_g^2 + |w_t|^2). \quad (4.16)$$

By the definition of B_2 , we have that

$$B_2 w^2 = \left[2s^3 \lambda^4 \alpha^3 |\nabla_N \varphi|_b^4 - s^3 \alpha^3 O(\lambda^3) - s^2 \alpha^2 O(\lambda)^4 \right] w^2. \quad (4.17)$$

Recalling (4.5) for the definition of φ and (4.3) for the positive constant h , we conclude that there is a constant $\lambda_0 > 1$ such that for any $\lambda \geq \lambda_0$, one can find a constant $s_0 > 1$ so that for any $s \geq s_0$, the following estimates hold uniformly for $(x, t) \in \overline{M} \times (2-a, 2+a) \setminus \omega_0 \times (2-a_0, 2+a_0)$:

$$\begin{cases} B_1 |\nabla_N w|_b^2 - C(s\lambda \alpha + \lambda^2 \alpha) (|\nabla_M w|_g^2 + |w_t|^2) \geq (c_1 - c_2)^2 h^2 s \lambda^2 \alpha (|\nabla_M w|_g^2 + |w_t|^2), \\ B_2 w^2 - Cs^2 \lambda^4 \alpha w^2 \geq (c_1 - c_2)^4 h^4 s^3 \lambda^4 \alpha^3 |w|^2. \end{cases} \quad (4.18)$$

From (4.11) and (4.18), we conclude that

$$\begin{aligned} & s\lambda^2 \int_{Q_0} \alpha \left(|\nabla_M w|_g^2 + |w_t|^2 \right) d_g x dt + s^3 \lambda^4 \int_{Q_0} \alpha^3 |w|^2 d_g x dt \\ & \leq C \left\{ \int_{Q_0} \theta^2 |(u_1)_{tt} + \Delta_M u_1|^2 d_g x dt + \int_{Q_0} D d_g x dt \right. \\ & \quad \left. + s\lambda^2 \int_0^T \int_{\omega_0} \alpha \left(|\nabla_M w|_g^2 + |w_t|^2 \right) d_g x dt + s^3 \lambda^4 \int_0^T \int_{\omega_0} \alpha^3 |w|^2 d_g x dt \right\}. \end{aligned} \quad (4.19)$$

Step 3. We now get rid of the boundary term $\int_{Q_0} D d_g x dt$ in (4.19).

Using the divergence theorem and the boundary condition of equation (4.9), the first term in $\int_{Q_0} D d_g x dt$ reads

$$4s\lambda^2 \int_{\Sigma_0} \alpha |\nabla_N \varphi|_b^2 w \frac{\partial_M w}{\partial \nu} d\Gamma_g dt = 4s\lambda^2 \int_{\Sigma_0} \alpha (|\nabla_M \varphi|_g^2 + |\varphi_t|^2) \left(s\lambda \alpha \frac{\partial_M \varphi}{\partial \nu} w^2 - lw^2 \right) d_g \Gamma dt. \quad (4.20)$$

The second one is

$$2s^3 \lambda^3 \int_{\Sigma_0} \alpha^3 (|\nabla_M \varphi|_g^2 + |\varphi_t|^2) \frac{\partial_M \varphi}{\partial \nu} w^2 d_g \Gamma dt. \quad (4.21)$$

The third one is

$$\begin{aligned} & 4s\lambda \int_{\Sigma_0} \alpha (\nabla_N \varphi, \nabla_N w)_b \frac{\partial_M w}{\partial \nu} d_g \Gamma dt \\ & = 4s\lambda \int_{\Sigma_0} \alpha [(\nabla_M \varphi, \nabla_M w)_g + \varphi_t w_t] \left(s\lambda \alpha w \frac{\partial_M \varphi}{\partial \nu} + \theta \frac{\partial_M u_1}{\partial \nu} \right) d_g \Gamma dt. \end{aligned} \quad (4.22)$$

By the boundary condition of u_1 , we have that $\frac{\partial_M u_1}{\partial \nu} = -lu_1$. Especially, noting that $\psi|_{\Gamma} = 0$, we have that $\nabla_M \varphi|_{\Gamma} = \frac{\partial_M \varphi}{\partial \nu} \nu|_{\Gamma}$. Hence from (4.22), we get that

$$\begin{aligned} & 4s\lambda \int_{\Sigma_0} \alpha(\nabla_N \varphi, \nabla_N w)_b \frac{\partial_M w}{\partial \nu} d_g \Gamma dt \\ &= \int_{\Sigma_0} \left(4s^3 \lambda^3 \alpha^3 |\nabla_M \varphi|_g^2 \frac{\partial_M \varphi}{\partial \nu} w^2 - 8s^2 \lambda^2 \alpha^2 |\nabla_M \varphi|_g^2 l w^2 + 4s\lambda \alpha \frac{\partial_M \varphi}{\partial \nu} l^2 w^2 \right) d_g \Gamma dt \\ & \quad + 4s^2 \lambda^2 \int_{\Sigma_0} \alpha^2 \varphi_t w_t w \frac{\partial_M \varphi}{\partial \nu} d_g \Gamma dt - 4s\lambda \int_{\Sigma_0} \alpha \varphi_t l w_t w d_g \Gamma dt. \end{aligned} \quad (4.23)$$

By integration by parts, we get that

$$\begin{aligned} & -4s\lambda \int_{\Sigma_0} \alpha \varphi_t w_t \theta l u_1 d_g \Gamma dt \\ &= -4s\lambda \int_{\Sigma_0} \theta \alpha \varphi_t (s\lambda \alpha \varphi_t \theta u_1 + \theta(u_1)_t) l u_1 d_g \Gamma dt \\ &= 2s\lambda \int_{\Sigma_0} \left\{ \lambda \alpha \varphi_t^2 l w^2 + \alpha \varphi_{tt} l w^2 \right\} d_g \Gamma dt. \end{aligned} \quad (4.24)$$

Therefore, we obtain that

$$\begin{aligned} & 4s\lambda \int_{\Sigma_0} \alpha(\nabla_N \varphi, \nabla_N w)_b \frac{\partial_M w}{\partial \nu} d_g \Gamma dt \\ &= \int_{\Sigma_0} \left(4s^3 \lambda^3 \alpha^3 |\nabla_M \varphi|_g^2 \frac{\partial_M \varphi}{\partial \nu} w^2 - 8s^2 \lambda^2 \alpha^2 |\nabla_M \varphi|_g^2 l w^2 + 4s\lambda \alpha \frac{\partial_M \varphi}{\partial \nu} l^2 w^2 \right) d_g \Gamma dt \\ & \quad + 4s^2 \lambda^2 \int_{\Sigma_0} \alpha^2 \varphi_t w_t w \frac{\partial_M \varphi}{\partial \nu} d_g \Gamma dt + \int_{\Sigma_0} \left\{ 2s^2 \lambda^2 \alpha^2 \varphi_t^2 l w^2 - 2s\lambda^2 \alpha \varphi_t^2 l w^2 - 2s\lambda \alpha \varphi_{tt} l w^2 \right\} d_g \Gamma dt. \end{aligned} \quad (4.25)$$

The fourth one is

$$\begin{aligned} & -2 \int_{\Sigma_0} s\lambda \alpha |\nabla_N w|_b^2 \frac{\partial_M \varphi}{\partial \nu} d_g \Gamma dt \\ &= -2 \int_{\Sigma_0} s\lambda \alpha (|\nabla_M w|_g^2 + w_t^2) \frac{\partial_M \varphi}{\partial \nu} d_g \Gamma dt \\ &= -2 \int_{\Sigma_0} \left[s^3 \lambda^3 \alpha^3 |\nabla_M \varphi|_g^2 \frac{\partial_M \varphi}{\partial \nu} w^2 - 2s^2 \lambda^2 \alpha^2 |\nabla_M \varphi|_g^2 l w^2 + s\lambda \alpha \theta^2 |\nabla_M u_1|_g^2 \frac{\partial_M \varphi}{\partial \nu} \right] d_g \Gamma dt \\ & \quad - 2 \int_{\Sigma_0} s\lambda \alpha w_t^2 \frac{\partial_M \varphi}{\partial \nu} d_g \Gamma dt. \end{aligned} \quad (4.26)$$

Therefore we have

$$\begin{aligned} \int_{Q_0} D d_g x dt &= \int_{\Sigma_0} \left[4s^2 \lambda^3 \alpha^2 (|\nabla_M \varphi|_g^2 + |\varphi_t|^2) w^2 + 4s^3 \lambda^3 \alpha^3 |\nabla_M \varphi|_g^2 w^2 + 2s^3 \lambda^3 \alpha^3 |\varphi_t|^2 w^2 \right. \\ & \quad \left. + 4s\lambda \alpha l^2 w^2 + 4s^2 \lambda^2 \alpha^2 \varphi_t w w_t - 2s\lambda \alpha \theta^2 |\nabla_M u_1|_g^2 - 2s\lambda \alpha l w_t^2 \right] \frac{\partial_M \varphi}{\partial \nu} d_g \Gamma dt \\ & \quad - \int_{\Sigma_0} \left[4s\lambda^2 \alpha |\nabla_M \varphi|_g^2 l (1 + s\alpha) - 2s\lambda^2 \alpha l |\varphi_t|^2 + 2s\lambda \alpha l \varphi_{tt} \right] w^2 d_g \Gamma dt. \end{aligned} \quad (4.27)$$

Since $|\nabla_M \varphi|_g|_\Gamma > 0$, we know that there exists an $s_1 > 0$ such that for all $s > s_1$, we have that

$$\int_{\Sigma_0} \left[4s\lambda^2 \alpha |\nabla_M \varphi|_g^2 l (1 + s\alpha) - 2s\lambda^2 \alpha l |\varphi_t|^2 + 2s\lambda \alpha l \varphi_{tt} \right] w^2 d_g \Gamma dt \geq 0. \quad (4.28)$$

Hence the right-hand side of (4.27) could be divided into two parts. The second integral is negative and has the property we expect. We need only to deal with the first integral in the right hand side of (4.27). We now choose another weight function $\tilde{\theta}$. By (4.5), (4.6) and noting that ψ vanishes on Γ , we have the following equalities:

$$\varphi|_\Sigma = \tilde{\varphi}|_\Sigma, \quad \frac{\partial_M \varphi}{\partial \nu}|_\Sigma = -\frac{\partial_M \tilde{\varphi}}{\partial \nu}|_\Sigma, \quad \alpha|_\Sigma = \tilde{\alpha}|_\Sigma, \quad w|_\Sigma = \tilde{w}|_\Sigma, \quad w_t|_\Sigma = \tilde{w}_t|_\Sigma. \quad (4.29)$$

Similar to (4.19) and (4.27), we deduce the following inequality:

$$\begin{aligned} & s\lambda^2 \int_{Q_0} \tilde{\alpha} \left(|\nabla_M \tilde{w}|_g^2 + |\tilde{w}_t|^2 \right) d_g x dt + s^3 \lambda^4 \int_{Q_0} \tilde{\alpha}^3 |\tilde{w}|^2 d_g x dt \\ & \leq C \left\{ \int_{Q_0} \tilde{\theta}^2 \left| (u_1)_{tt} + \Delta_M u_1 \right|^2 d_g x dt + \int_{Q_0} \tilde{D} d_g x dt \right. \\ & \quad \left. + s\lambda^2 \int_0^T \int_{\omega_0} \tilde{\alpha} \left(|\nabla_M \tilde{w}|_g^2 + |\tilde{w}_t|^2 \right) d_g x dt + s^3 \lambda^4 \int_0^T \int_{\omega_0} \tilde{\alpha}^3 |\tilde{w}|^2 d_g x dt \right\}, \quad (4.30) \end{aligned}$$

where $\tilde{w} = \tilde{\theta} u_1$ and

$$\begin{aligned} \int_{Q_0} \tilde{D} d_g x dt & = - \int_{\Sigma_0} \left[4s^2 \lambda^3 \alpha^2 \left(|\nabla_M \varphi|_g^2 + |\varphi_t|^2 \right) w^2 + 6s^3 \lambda^3 \alpha^3 |\nabla_M \varphi|_g^2 w^2 + 4s^3 \lambda^3 \alpha^3 |\varphi_t|^2 w^2 \right. \\ & \quad \left. + 4s\lambda \alpha l_1^2 w^2 + 4s^2 \lambda^2 \alpha^2 \varphi_t w w_t - 2s\lambda \alpha \theta^2 |\nabla_M u_1|_g^2 - 2s\lambda \alpha l w_t^2 \right] \frac{\partial_M \varphi}{\partial \nu} d_g \Gamma dt \\ & \quad - \int_{\Sigma_0} \left[4s\lambda^2 \alpha |\nabla_M \varphi|_g^2 l (1 + s\alpha) - 2s\lambda^2 \alpha l |\varphi_t|^2 - 2s\lambda \alpha l \varphi_{tt} \right] w^2 d_g \Gamma dt. \quad (4.31) \end{aligned}$$

From (4.27) and (4.31), we know that for $\forall s > s_1$, we have that

$$\int_{Q_0} \left(D + \tilde{D} \right) d_g x dt = - \int_{\Sigma_0} \left[8s\lambda^2 \alpha |\nabla_M \varphi|_g^2 l (1 + s\alpha) - 4s\lambda^2 \alpha l |\varphi_t|^2 - 4s\lambda \alpha l \varphi_{tt} \right] w^2 d_g \Gamma dt \leq 0. \quad (4.32)$$

By (4.19), (4.30) and noting (4.32), we arrive at

$$\begin{aligned} & s\lambda^2 \int_{Q_0} \left[\alpha \left(|\nabla_M w|_g^2 + |w_t|^2 \right) + \tilde{\alpha} \left(|\nabla_M \tilde{w}|_g^2 + |\tilde{w}_t|^2 \right) \right] d_g x dt \\ & + s^3 \lambda^4 \int_{Q_0} \left(\alpha^3 |w|^2 + \tilde{\alpha}^3 |\tilde{w}|^2 \right) d_g x dt \\ & \leq C \left\{ \int_{Q_0} \left(\theta^2 \left| (u_1)_{tt} + \Delta_M u_1 \right|^2 + \tilde{\theta}^2 \left| (u_1)_{tt} + \Delta_M u_1 \right|^2 \right) d_g x dt + s^3 \lambda^4 \int_0^T \int_{\omega_0} \left(\alpha^3 |w|^2 + v \tilde{\alpha}^3 |\tilde{w}|^2 \right) d_g x dt \right. \\ & \quad \left. + s\lambda^2 \int_0^T \int_{\omega_0} \left[\alpha \left(|\nabla_M w|_g^2 + |w_t|^2 \right) + \tilde{\alpha} \left(|\nabla_M \tilde{w}|_g^2 + |\tilde{w}_t|^2 \right) \right] d_g x dt \right\}. \quad (4.33) \end{aligned}$$

Step 4. We now return both w and \tilde{w} in (4.33) to u_1 . Recalling that $w = \theta u_1$ and $\tilde{w} = \tilde{\theta} u_1$, we obtain that

$$\begin{aligned} \frac{1}{C} \theta^2 \left(|\nabla_M u_1|_g^2 + |(u_1)_t|^2 + s^2 \lambda^2 \alpha^2 |u_1|^2 \right) &\leq |\nabla_M w|_g^2 + |w_t|^2 + s^2 \lambda^2 \alpha^2 w^2 \\ &\leq C \theta^2 \left(|\nabla_M u_1|_g^2 + |(u_1)_t|^2 + s^2 \lambda^2 \alpha^2 |u_1|^2 \right) \end{aligned} \quad (4.34)$$

and

$$\begin{aligned} \frac{1}{C} \tilde{\theta}^2 \left(|\nabla_M u_1|_g^2 + |(u_1)_t|^2 + s^2 \lambda^2 \tilde{\alpha}^2 |u_1|^2 \right) &\leq |\nabla_M \tilde{w}|_g^2 + |\tilde{w}_t|^2 + s^2 \lambda^2 \tilde{\alpha}^2 \tilde{w}^2 \\ &\leq C \tilde{\theta}^2 \left(|\nabla_M u_1|_g^2 + |(u_1)_t|^2 + s^2 \lambda^2 \alpha^2 |u_1|^2 \right). \end{aligned} \quad (4.35)$$

By the definition of α , $\tilde{\alpha}$, θ and $\tilde{\theta}$, we know $\alpha \geq \tilde{\alpha} > 1$ and $\theta \geq \tilde{\theta} > 1$. Hence, by (4.33)-(4.35), we end up with the following inequality:

$$\begin{aligned} &s^3 \lambda^4 \int_{Q_0} \alpha^3 \theta^2 |u_1|^2 d_g x dt \\ &\leq C \left\{ \int_{Q_0} \theta^2 \left| \Delta_M u_1 + (u_1)_{tt} \right|^2 d_g x dt + s^3 \lambda^4 \int_0^T \int_{\omega_0} \alpha^3 \theta^2 |u_1|^2 d_g x dt \right. \\ &\quad \left. + s \lambda^2 \int_0^T \int_{\omega_0} \alpha \theta^2 \left(|\nabla_M u_1|_g^2 + |(u_1)_t|^2 \right) d_g x dt \right\}. \end{aligned} \quad (4.36)$$

Recalling that u_1 is the solution of equation (4.9), we know

$$\left| \Delta_M u_1 + (u_1)_{tt} \right|^2 \leq \left| \phi_{tt} u + 2\phi_t u_t \right|^2. \quad (4.37)$$

Choose a cut-off function $g \in C_0^\infty(\omega)$ with $g = 1$ in ω_0 and $0 \leq g \leq 1$ in ω . Multiplying equation (4.9) by $g\theta^2 \alpha u_1$ and integrating it in Q_0 , using integration by parts, we get

$$\begin{aligned} &\int_0^T \int_{\omega_0} \alpha \theta^2 \left(|\nabla_M u_1|_g^2 + |(u_1)_t|^2 \right) d_g x dt \\ &\leq C \left[s^2 \lambda^2 \int_0^T \int_{\omega} \theta^2 \alpha^2 |u_1|^2 d_g x dt + \int_{Q_0} \theta^2 |\phi_{tt} u + 2\phi_t u_t|^2 d_g x dt \right]. \end{aligned} \quad (4.38)$$

From (4.36)-(4.38), we obtain

$$\begin{aligned} &s^3 \lambda^4 \int_{Q_0} \alpha^3 \theta^2 |u_1|^2 d_g x dt \\ &\leq C \left\{ s \lambda^2 \int_{Q_0} \theta^2 |\phi_{tt} u + 2\phi_t u_t|^2 d_g x dt + s^3 \lambda^4 \int_0^T \int_{\omega} \alpha^3 \theta^2 |u_1|^2 d_g x dt \right\}. \end{aligned} \quad (4.39)$$

Step 5. Finally, we shall drop the weight functions in the integrands of (4.39) to get the desired result. Noting that α satisfies (4.7) and φ satisfies (4.8), we have the following inequalities:

$$\int_{Q_0} s^3 \lambda^4 \alpha^3 \theta^2 u_1^2 d_g x dt \geq s^3 \lambda^4 e^{3(c_1 \lambda + \lambda \kappa)} e^{2s(e^{c_1 \lambda + \lambda \kappa})} \int_{T'}^{T''} \int_M |u|^2 d_g x dt, \quad (4.40)$$

$$\begin{aligned}
& \int_{Q_0} \theta^2 |\phi_{tt}u + 2\phi_t u_t|^2 d_g x dt \\
&= \int_{\frac{T}{2}-a_1}^{\frac{T}{2}+a_1} \int_M \theta^2 |\phi_{tt}u + 2\phi_t u_t|^2 d_g x dt \\
&\leq C e^{2se^{(c_1-c_2)\lambda+\lambda\kappa}} \left\{ \int_{\frac{T}{2}-a_1}^{\frac{T}{2}-a} \int_M (|u|^2 + |u_t|^2) d_g x dt + \int_{\frac{T}{2}+a}^{\frac{T}{2}+a_1} \int_M (|u|^2 + |u_t|^2) d_g x dt \right\} \\
&\leq C e^{2se^{(c_1-c_2)\lambda+\lambda\kappa}} |u|_{H^1(Q)}^2,
\end{aligned} \tag{4.41}$$

and

$$\begin{aligned}
& \int_0^T \int_\omega s^3 \lambda^4 \alpha^3 \theta^2 |u_1|^2 d_g x dt \\
&= \int_{\frac{T}{2}-a_1}^{\frac{T}{2}+a_1} \int_\omega s^3 \lambda^4 \alpha^3 \theta^2 |u_1|^2 d_g x dt \\
&\leq s^3 \lambda^4 e^{3\lambda(\frac{T^2}{4}+c_1-c_2+\kappa)} e^{2se^{\lambda(\frac{T^2}{4}+c_1-c_2+\kappa)}} \int_\gamma^{T-\gamma} \int_\omega |u|^2 d_g x dt.
\end{aligned} \tag{4.42}$$

From (4.39)-(4.42), we obtain

$$\begin{aligned}
& s^3 \lambda^4 e^{3(c_1\lambda+\lambda\kappa)} e^{2s(e^{c_1\lambda+\lambda\kappa})} \int_{T'}^{T''} \int_M |u|^2 d_g x dt \\
&\leq s^3 \lambda^4 e^{3\lambda(\frac{T^2}{4}+c_1-c_2+\kappa)} e^{2se^{\lambda(\frac{T^2}{4}+c_1-c_2+\kappa)}} \int_\gamma^{T-\gamma} \int_\omega |u|^2 d_g x dt + C e^{2se^{(c_1-c_2)\lambda+\lambda\kappa}} |u|_{H^1(Q)}^2.
\end{aligned} \tag{4.43}$$

Recalling that $c_1 > c_2 > 0$, hence we know that $e^{2s(e^{c_1\lambda+\lambda\kappa})} > e^{2s(e^{(c_1-c_2)\lambda+\lambda\kappa})}$. Let $\lambda = \lambda_0$,

$$\varepsilon = \frac{e^{2s(e^{(c_1-c_2)\lambda_0+\kappa\lambda_0})}}{\lambda_0^4 s^3 e^{3\lambda_0(c_1+\kappa)} e^{2se^{c_1\lambda_0+\kappa\lambda_0}}}, \quad k = \frac{e^{\lambda_0(\frac{T^2}{4}+c_1-c_2+\kappa)} - e^{c_1\lambda_0+\kappa\lambda_0}}{e^{c_1\lambda_0+\kappa\lambda_0} - e^{(c_1-c_2)\lambda_0+\kappa\lambda_0}},$$

and

$$\varepsilon_0 = \frac{e^{2s_0(e^{(c_1-c_2)\lambda_0+\kappa\lambda_0})}}{\lambda_0^4 s_0^3 e^{3\lambda_0(c_1+\kappa)} e^{2s_0 e^{c_1\lambda_0+\kappa\lambda_0}}}.$$

From (4.43), we know that for any $\varepsilon \in (0, \varepsilon_0]$, it holds

$$|u|_{L^2(M \times (T', T''))}^2 \leq \varepsilon^{-k} |u|_{L^2(\omega \times (\gamma, T-\gamma))}^2 + C\varepsilon |u|_{H^1(Q)}^2, \tag{4.44}$$

which in turn implies that the above inequality holds for any $\varepsilon > 0$.

Let $\mu = \frac{1}{1+k}$, $\varepsilon = \left(\frac{|u|_{L^2(\omega \times (\gamma, T-\gamma))}}{|u|_{H^1(Q)}} \right)^{2\mu}$, by inequality (4.44), we get

$$|u|_{L^2(M \times (T', T''))} \leq C |u|_{L^2(\omega \times (\gamma, T-\gamma))}^\mu |u|_{H^1(M \times (0, T))}^{1-\mu}. \tag{4.45}$$

□

5 Interpolation inequality II

This section is devoted to showing another interpolation result for solutions to equation (4.1). Our result is stated as follows:

Theorem 5.1 *Let $0 < \gamma < \frac{T}{2}$. Then there exists a constant $\delta \in (0, 1)$ such that any solution $u \in H^2(Q)$ of (4.1) satisfies*

$$|u|_{H^1(\omega \times (\gamma, T-\gamma))} \leq C(|u(0)|_{L^2(\omega)} + |u_t(0)|_{L^2(\omega)} + |\nabla_M u(0)|_{L^2(\omega)})^\delta |u|_{H^1(Q)}^{1-\delta}. \quad (5.1)$$

Proof of Theorem 5.1 : We divide the proof into three steps.

Step 1. Let $\omega_2 \subset\subset \omega$. Denote by $\text{dist}((x, t), \omega_2 \times \{0\})$ the distance from (x, t) to $\omega_2 \times \{0\}$. Put $N(\tau) = \{(x, t) \in Q \mid \text{dist}((x, t), \omega_2 \times \{0\}) < \tau\}$. Let $0 < \tau_1 < \tau_2 < \tau_3$ such that $N(\tau_3) \subset Q$ and $N(\tau_3) \cap (M \times \{0\}) \subset \omega \times \{0\}$.

Let h be an C^2 -function such that

$$\begin{cases} 3 < h < 4 & \text{if } (x, t) \in N(\tau_1), \\ 0 < h < 1 & \text{if } (x, t) \in N(\tau_3) \setminus N(\tau_2), \\ |\nabla_M h| > 0 & \text{for all } (x, t) \in N(\tau_3). \end{cases}$$

The construction of h is very easy. For example, we can choose a smooth function $h_1 : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\begin{cases} h_1' < 0 \text{ and } 3 < h_1(s) < 4 & \text{if } 0 < s < \tau_1^2, \\ 0 < h_1(s) < 1 & \text{if } \tau_2^2 < s < \tau_3^2. \end{cases}$$

Then $h(x, t) = h_1(\text{dist}^2((x, t), \omega_2 \times \{0\}))$ is the desired function.

In what follows, we shall use Theorem 2.1 (with φ replaced by h) to prove Theorem 5.1. For simplicity of the notations, we still use θ to denote the weight function if there is no confusion.

Denote by $d_b \partial N(\tau_3)$ the volume element of $\partial N(\tau_3)$ in its Riemannian metric and by $\bar{\nu} = \bar{\nu}(x)$ the unit outward normal vector of $N(\tau_3)$ at $x \in \partial N(\tau_3)$ with its Riemannian metric.

For ε small enough, define

$$N_\varepsilon(\partial(N(\tau_3)) \setminus (\omega_2 \times \{0\})) \triangleq \{x : x \in N(\tau_3), \text{dist}((x, t), \partial(N(\tau_3)) \setminus (\omega_2 \times \{0\})) < \varepsilon\}.$$

Choose a function $\chi \in C^\infty(N(\tau_3))$ such that $0 \leq \chi \leq 1$ and that

$$\chi = \begin{cases} 1 & \text{if } (x, t) \in N(\tau_2), \\ 0 & \text{if } (x, t) \in N(\tau_3) \cap N_\varepsilon(\partial(N(\tau_3)) \setminus (\omega_2 \times \{0\})). \end{cases}$$

Put $\bar{u} = \chi u$ where u is the solution of equation (4.1). Then, \bar{u} satisfies the following equation:

$$\begin{cases} \bar{u}_{tt} + \Delta_M \bar{u} = \chi_{tt} u + 2\chi_t u_t + u \Delta_M \chi + 2(\nabla_M u, \nabla_M \chi)_g & \text{in } N(\tau_3), \\ |\nabla_M \bar{u}|_g = \bar{u} = 0 & \text{on } \partial N(\tau_3) \setminus (\omega \times \{0\}). \end{cases} \quad (5.2)$$

Apply Theorem 2.1 to equation (5.2) with $b = 1 \otimes g$, v replaced by \bar{u} and $w = \theta \bar{u}$.

Proceeding as in (4.12)-(4.18), similar to (4.19), and noting that h has no critical point in $N(\tau_3)$, we obtain that

$$\begin{aligned} & s\lambda^2 \int_{N(\tau_3)} \alpha \left(|\nabla_M w|_g^2 + |w_t|^2 \right) d_g x dt + s^3 \lambda^4 \int_{N(\tau_3)} \alpha^3 |w|^2 d_g x dt \\ & \leq C \left\{ \int_{N(\tau_3)} \theta^2 \left| \bar{u}_{tt} + \Delta_M \bar{u} \right|^2 d_g x dt + \int_{N(\tau_3)} D_1 d_g x dt \right\}, \end{aligned} \quad (5.3)$$

where

$$\begin{aligned} D_1 = 2s\lambda \operatorname{div}_N & \left[2\lambda\alpha |\nabla_N h|_b^2 w \nabla_N w + s^2 \lambda^2 \alpha^3 |\nabla_N h|_b^2 w^2 \nabla_N h \right. \\ & \left. + 2\alpha (\nabla_N h, \nabla_N w)_b \nabla_N w - \alpha |\nabla_N w|_b^2 \nabla_N h \right]. \end{aligned} \quad (5.4)$$

By the divergence theorem, $\int_{N(\tau_3)} D_1 d_g x dt$ is the boundary term.

For the first term therein, we have

$$\begin{aligned} & \int_{\partial N(\tau_3)} 2s\lambda^2 \alpha |\nabla_N h|_b^2 w \frac{\partial w}{\partial \bar{\nu}} d_b \partial N(\tau_3) \\ & \leq C \int_{\partial N(\tau_3)} \left\{ s\lambda\alpha (|\nabla_M w|_g^2 + w_t^2) + s\lambda^3 \alpha w^2 \right\} d_b \partial N(\tau_3). \end{aligned}$$

Due to the definition of w , we know

$$\begin{cases} w|_{\partial N(\tau_3) \setminus (\omega \times \{0\})} = \bar{u}|_{\partial N(\tau_3) \setminus (\omega \times \{0\})} = 0, \\ \nabla_N w|_{\partial N(\tau_3) \setminus (\omega \times \{0\})} = \nabla_N \bar{u}|_{\partial N(\tau_3) \setminus (\omega \times \{0\})} = 0. \end{cases}$$

Hence we know that

$$\begin{aligned} & \int_{\partial N(\tau_3)} 2s\lambda^2 \alpha |\nabla_N h|_b^2 w \frac{\partial w}{\partial \bar{\nu}} d_b \partial N(\tau_3) \\ & \leq C \int_{\omega \times \{0\}} \left\{ s\lambda\alpha (|\nabla_M w|_g^2 + w_t^2) + s\lambda^3 \alpha w^2 \right\} d_g x. \end{aligned}$$

By the same argument, we obtain the estimates for the remainder terms. Therefore, it follows that

$$\int_Q D_1 d_g x dt \leq C \int_{\omega \times \{0\}} \left[s\lambda\alpha (|\nabla_M w|_g^2 + w_t^2) + s^3 \lambda^3 \alpha^3 w^2 \right] d_g x. \quad (5.5)$$

Combining (5.3) and (5.5), we obtain that

$$\begin{aligned} & s\lambda^2 \int_{N(\tau_3)} \alpha \left(|\nabla_M w|_g^2 + |w_t|^2 \right) d_g x dt + s^3 \lambda^4 \int_{N(\tau_3)} \alpha^3 |w|^2 d_g x dt \\ & \leq C \left\{ \int_{N(\tau_3)} \theta^2 \left| \bar{u}_{tt} + \Delta_M \bar{u} \right|^2 d_g x dt + \int_{\omega \times \{0\}} \left[s\lambda\alpha (|\nabla_M w|_g^2 + |w_t|^2) + s^3 \lambda^3 \alpha^3 |w|^2 \right] d_g x \right\}. \end{aligned} \quad (5.6)$$

Step 2. We now return the w in (5.6) to \bar{u} . Recalling $w = \theta \bar{u}$, it is clear that

$$\begin{aligned} \frac{1}{C} \theta^2 \left(|\nabla_M \bar{u}|_g^2 + |\bar{u}_t|^2 + s^2 \lambda^2 \alpha^2 |\bar{u}|^2 \right) & \leq |\nabla_M w|_g^2 + |w_t|^2 + s^2 \lambda^2 \alpha^2 w^2 \\ & \leq C \theta^2 \left(|\nabla_M \bar{u}|_g^2 + |\bar{u}_t|^2 + s^2 \lambda^2 \alpha^2 |\bar{u}|^2 \right). \end{aligned} \quad (5.7)$$

From (5.6)-(5.7), and noting the first equation in (5.2), we obtain that

$$\begin{aligned}
& s\lambda^2 \int_{N(\tau_3)} \alpha\theta^2 \left(|\nabla_M \bar{u}|_g^2 + |\bar{u}_t|^2 \right) d_g x dt + s^3 \lambda^4 \int_{N(\tau_3)} \alpha^3 \theta^2 |\bar{u}|^2 d_g x dt \\
& \leq C \left\{ \int_{N(\tau_3)} \theta^2 \left| \chi_{tt} u + 2\chi_t u_t + u \Delta_M \chi + 2(\nabla_M u, \nabla_M \chi)_g \right|^2 d_g x dt \right. \\
& \quad \left. + \int_{\omega \times \{0\}} \left[s\lambda \alpha (|\nabla_M \bar{u}|_g^2 + \bar{u}_t^2) + s^3 \lambda^3 \alpha^3 \bar{u}^2 \right] d_g x \right\}. \tag{5.8}
\end{aligned}$$

By the definition of \bar{u} , we know that $\bar{u} = u$ in $N(\tau_1)$. By the definition of h , we know that

$$\begin{cases} \alpha \geq e^{3\lambda} \text{ and } \theta \geq e^{se^{3\lambda}} & \text{if } (x, t) \in N(\tau_1), \\ \alpha \leq e^\lambda \text{ and } \theta \leq e^{se^\lambda} & \text{if } (x, t) \in N(\tau_3) \setminus N(\tau_2). \end{cases}$$

By the definition of χ we know that

$$\chi_t = 0 \text{ and } \nabla_M \chi = 0 \text{ if } (x, t) \in N(\tau_2).$$

Therefore we have the following inequalities:

$$\int_{N(\tau_3)} s\lambda^2 \alpha \theta^2 \left(|\nabla_M \bar{u}|_g^2 + |\bar{u}_t|^2 \right) d_g x dt \geq s\lambda^2 e^{3\lambda} e^{2se^{3\lambda}} \int_{N(\tau_1)} \left(|\nabla_M \bar{u}|_g^2 + |\bar{u}_t|^2 \right) d_g x dt, \tag{5.9}$$

$$\int_{N(\tau_3)} s^3 \lambda^4 \alpha^3 \theta^2 \bar{u}^2 d_g x dt \geq s^3 \lambda^4 e^{9\lambda} e^{2se^{3\lambda}} \int_{N(\tau_1)} |u|^2 d_g x dt, \tag{5.10}$$

$$\begin{aligned}
& \int_{N(\tau_3)} \theta^2 \left| \chi_{tt} u + 2\chi_t u_t + u \Delta_M \chi + 2(\nabla_M u, \nabla_M \chi)_g \right|^2 d_g x dt \\
& \leq C e^{2se^\lambda} \int_{N(\tau_3)} \left(|u|^2 + |\nabla_M u|_g^2 + |u_t|^2 \right) d_g x dt, \tag{5.11}
\end{aligned}$$

$$\int_{\omega \times \{0\}} s^3 \lambda^3 \alpha^3 \theta^2 |\bar{u}|^2 d_g x \leq s^3 \lambda^3 e^{12\lambda} e^{2se^{4\lambda}} \int_{\omega \times \{0\}} |u|^2 d_g x, \tag{5.12}$$

and

$$\int_{\omega \times \{0\}} s\lambda \alpha \theta^2 \left(|\nabla_M \bar{u}|_g^2 + \bar{u}_t^2 \right) d_g x \leq s\lambda e^{4\lambda} e^{2se^{4\lambda}} \int_{\omega \times \{0\}} \left(|\nabla_M u|_g^2 + u_t^2 \right) d_g x. \tag{5.13}$$

From (5.8) to (5.13), we know that

$$\begin{aligned}
& s\lambda^2 e^{3\lambda} e^{2se^{3\lambda}} \int_{N(\tau_1)} \left(|\nabla_M \bar{u}|_g^2 + |\bar{u}_t|^2 \right) d_g x dt + s^3 \lambda^4 e^{9\lambda} e^{2se^{3\lambda}} \int_{N(\tau_1)} s^3 \lambda^4 |u|^2 d_g x dt \\
& \leq C \left\{ e^{2se^\lambda} \int_{N(\tau_3)} \left(|u|^2 + |\nabla_M u|_g^2 + |u_t|^2 \right) d_g x dt + s^3 \lambda^3 e^{12\lambda} e^{2se^{4\lambda}} \int_{\omega \times \{0\}} |u|^2 d_g x \right. \\
& \quad \left. + s\lambda e^{4\lambda} e^{2se^{4\lambda}} \int_{\omega \times \{0\}} \left(|\nabla_M u|_g^2 + u_t^2 \right) d_g x \right\}. \tag{5.14}
\end{aligned}$$

Similar to (4.44), by (5.14), we obtain that there exist a $\beta > 0$ and an $\epsilon_0 > 0$ such that for any $\epsilon \in (0, \epsilon_0]$, we have

$$|u|_{H^1(N(\tau_1))}^2 \leq \epsilon^{-\beta} \left(|u_t(0)|_{L^2(\omega)}^2 + |\nabla_M u(0)|_{L^2(\omega)}^2 + |u(0)|_{L^2(\omega)}^2 \right) + C\epsilon |u|_{H^1(Q)}^2, \quad (5.15)$$

which in turn implies that the above inequality holds for any $\epsilon > 0$.

Noting that $\tau_1 > 0$, hence there is an open ball $B \subset N(\tau_1)$. Then we know that

$$|u|_{H^1(B)}^2 \leq \epsilon^{-\beta} \left(|u_t(0)|_{L^2(\omega)}^2 + |\nabla_M u(0)|_{L^2(\omega)}^2 + |u(0)|_{L^2(\omega)}^2 \right) + C\epsilon |u|_{H^1(Q)}^2. \quad (5.16)$$

Put $\delta' = \frac{1}{1+\beta}$ and let $\epsilon = \left(\frac{|u_t(0)|_{L^2(\omega)}^2 + |\nabla_M u(0)|_{L^2(\omega)}^2 + |u(0)|_{L^2(\omega)}^2}{|u|_{H^1(Q)}^2} \right)^{\delta'}$ in (5.16), we get

$$\begin{aligned} |u|_{H^1(B)} &\leq C \left(|u_t(0)|_{L^2(\omega)}^2 + |\nabla_M u(0)|_{L^2(\omega)}^2 + |u(0)|_{L^2(\omega)}^2 \right)^{\frac{\delta'}{2}} |u|_{H^1(Q)}^{1-\delta'} \\ &\leq C \left(|u_t(0)|_{L^2(\omega)} + |\nabla_M u(0)|_{L^2(\omega)} + |u(0)|_{L^2(\omega)} \right)^{\delta'} |u|_{H^1(Q)}^{1-\delta'}. \end{aligned} \quad (5.17)$$

Step 3. To complete the proof, it suffices to show that the following proposition: For any given open set $L \subset\subset Q$, there exists a constant $0 < \delta'' < 1$ such that

$$|u|_{H^1(L)} \leq C |u|_{H^1(B)}^{\delta''} |u|_{H^1(Q)}^{1-\delta''}. \quad (5.18)$$

Firstly, we admit this claim and continue our proof. After that, we prove this proposition.

By inequality (5.17) and (5.18), we deduce that for any given subset $L \subset\subset Q$, we have

$$|u|_{H^1(L)} \leq C \left(|u_t(0)|_{L^2(\omega)} + |\nabla_M u(0)|_{L^2(\omega)} + |u(0)|_{L^2(\omega)} \right)^{\delta} |u|_{H^1(Q)}^{1-\delta} \quad (5.19)$$

where $\delta = \delta' \delta''$. Now we choose $L = \omega \times (\gamma, T - \gamma)$ to get Theorem 5.1.

Now we prove the above proposition. Let B_1, B_2 and B_3 be three open balls in Q such that $B_1 \subset\subset B_2 \subset\subset B_3 \subset\subset Q$. Choose a cut-off function $\eta \in C_0^\infty(Q)$ such that $\eta = 1$ in B_3 and $0 < \eta < 1$. Let $y = \eta u$. Then, y solves

$$\begin{cases} y_{tt} + \Delta_M y = \eta_{tt} u + 2\eta_t u_t + u \Delta_M \eta + 2(\nabla_M u, \nabla_M \eta)_g & \text{in } Q, \\ |\nabla_M y| = \bar{y} = 0 & \text{on } \partial Q. \end{cases} \quad (5.20)$$

Denote by P the center of B_1 . Let $r(x, t) = \text{dist}^2((x, t), P)$. Replace the above φ (in θ) by r . By the same argument as the proof of Theorem 4.1, we conclude that there exists a constant $0 < \tilde{\delta} < 1$ such that

$$|u|_{H^1(B_2)} \leq C |u|_{H^1(B_1)}^{\tilde{\delta}} |u|_{H^1(Q)}^{1-\tilde{\delta}}. \quad (5.21)$$

For any ball $B' \subset\subset Q$, we can find a finite number $m \in \mathbb{N}$ and two sequences of balls $\{B^i\}_{i=1}^m$ and $\{\tilde{B}^i\}_{i=1}^m$ such that

$$\begin{cases} B' \subset\subset B^1, \\ \tilde{B}^i \subset\subset B^i \cap B^{i+1} \text{ for } i = 1, \dots, m-1, \\ \tilde{B}^m \subset\subset B^m, \\ \tilde{B}^m = B'. \end{cases} \quad (5.22)$$

By inequality (5.21), we know that there exists a sequences $\{\tilde{\delta}_i\}_{i=1}^m$ satisfying $0 < \tilde{\delta}_i < 1$ for $i = 1, \dots, m$, such that

$$\begin{aligned} |u|_{H^1(B')} &\leq |u|_{H^1(B^1)} \leq C|u|_{H^1(\tilde{B}^1)}^{\tilde{\delta}_1} |u|_{H^1(Q)}^{1-\tilde{\delta}_1} \leq C|u|_{H^1(B^2)}^{\tilde{\delta}_1} |u|_{H^1(Q)}^{1-\tilde{\delta}_1} \\ &\leq C|u|_{H^1(\tilde{B}^2)}^{\tilde{\delta}_1 \tilde{\delta}_2} |u|_{H^1(Q)}^{1-\tilde{\delta}_1 \tilde{\delta}_2} \leq \dots \leq C|u|_{H^1(\tilde{B}^n)}^{\tilde{\delta}_1 \tilde{\delta}_2 \dots \tilde{\delta}_m} |u|_{H^1(Q)}^{1-\tilde{\delta}_1 \tilde{\delta}_2 \dots \tilde{\delta}_m}. \end{aligned} \quad (5.23)$$

Put $\tilde{\delta} = \tilde{\delta}_1 \tilde{\delta}_2 \dots \tilde{\delta}_m$, then we know that

$$|u|_{H^1(B')} \leq C|u|_{H^1(B)}^{\tilde{\delta}} |u|_{H^1(Q)}^{1-\tilde{\delta}}. \quad (5.24)$$

For any given $L \subset\subset Q$, we can find finite balls contained in the internal of Q to cover it. Hence from inequality (5.24), we know that there exist a constant $0 < \delta'' < 1$ such that (5.18) holds.

$$|u|_{H^1(L)} \leq C|u|_{H^1(B)}^{\delta''} |u|_{H^1(Q)}^{1-\delta''}. \quad (5.25)$$

This completes the proof of Theorem 5.1. \square

6 Proof of Theorem 1.2

We are now in a position to prove Theorem 1.2.

Proof of Theorem 1.2: For simplicity, choose $T = 4$, $T' = 1$ and $T'' = 3$ in inequalities (4.2) and (5.1). From Theorem 4.1, we get

$$|u|_{L^2(M \times (1,3))} \leq C|u|_{L^2(\omega \times (\gamma, 4-\gamma))}^\mu |u|_{H^1(Q)}^{1-\mu}. \quad (6.1)$$

By Theorem 5.1, we obtain

$$|u|_{L^2(M \times (1,3))} \leq C \left(|u_t(0)|_{L^2(\omega)}^2 + |\nabla_M u(0)|_{L^2(\omega)}^2 + |u(0)|_{L^2(\omega)}^2 \right)^{\mu\delta} |u|_{H^1(Q)}^{1-\mu\delta}, \quad (6.2)$$

where $u \in H^2(Q)$ is any solution of equation (4.1).

For any $\{a_i\}_{\lambda_i \leq r}$ with $a_i \in \mathbb{C}$, set

$$y(x, t) = \sum_{\lambda_j \leq r} \frac{\text{sh}(t\sqrt{\lambda_j})}{\sqrt{\lambda_j}} a_j e_j \quad (6.3)$$

with $\frac{\text{sh}(tb)}{b} = t$ if $b = 0$. Then, both the real part and the imaginary part of y are solutions of (4.1) and $\text{Re } y = \text{Im } y = 0$ on $M \times \{0\}$. Therefore $\text{Re } y$ and $\text{Im } y$ satisfy inequality (6.2). For the left term of (6.2), we have

$$\begin{aligned} |\text{Re } y|_{L^2(M \times (1,3))}^2 &= \int_1^3 \int_M \left| \sum_{\lambda_j \leq r} \frac{\text{sh}(t\sqrt{\lambda_j})}{\sqrt{\lambda_j}} (\text{Re } a_j) e_j \right|^2 dx dt = \sum_{\lambda_j \leq r} |\text{Re } a_j|^2 \int_1^3 \left| \frac{\text{sh}(t\sqrt{\lambda_j})}{\sqrt{\lambda_j}} \right|^2 dt \\ &\geq \sum_{\lambda_j \leq r} |\text{Re } a_j|^2 \int_1^3 t^2 dt = \frac{8}{3} \sum_{\lambda_j \leq r} |\text{Re } a_j|^2. \end{aligned} \quad (6.4)$$

For the right term of (6.2), we have $\partial_t \operatorname{Re} y(x, 0) = \sum_{\lambda_j \leq r} \operatorname{Re} a_j e_j$ and

$$|\operatorname{Re} y|_{H^1(Q)}^2 \leq C e^{8\sqrt{r}} (1+r) \sum_{\lambda_j \leq r} |\operatorname{Re} a_j|^2 \leq C e^{9\sqrt{r}} \sum_{\lambda_j \leq r} |\operatorname{Re} a_j|^2. \quad (6.5)$$

Therefore we get

$$\sum_{\lambda_j \leq r} |\operatorname{Re} a_j|^2 \leq C \left(\int_{\omega} \left| \sum_{\lambda_j \leq r} \operatorname{Re} a_j e_j \right|^2 dx \right)^{\mu\delta} \left(e^{9\sqrt{r}} \sum_{\lambda_j \leq r} |\operatorname{Re} a_j|^2 \right)^{1-\mu\delta}. \quad (6.6)$$

Hence we have

$$\sum_{\lambda_j \leq r} |\operatorname{Re} a_j|^2 \leq C e^{C\sqrt{r}} \int_{\omega} \left| \sum_{\lambda_i \leq r} \operatorname{Re} a_i e_i \right|^2 dx. \quad (6.7)$$

By the same argument, we can get

$$\sum_{\lambda_j \leq r} |\operatorname{Im} a_j|^2 \leq C e^{C\sqrt{r}} \int_{\omega} \left| \sum_{\lambda_i \leq r} \operatorname{Im} a_i e_i \right|^2 dx. \quad (6.8)$$

From (6.7) and (6.8), we obtain

$$\sum_{\lambda_j \leq r} |a_j|^2 \leq C e^{C\sqrt{r}} \int_{\omega} \left| \sum_{\lambda_i \leq r} a_i e_i \right|^2 dx. \quad (6.9)$$

□

Acknowledgments. The author acknowledges the anonymous referees for valuable and helpful comments.

References

- [1] X. Fu, *A weighted identity for partial differential operators of second order and its applications*, *C. R. Acad. Sci. Paris Série I*, **342**(2006), 579–584.
- [2] A. V. Fursikov and O. Yu. Imanuvilov, *Controllability of Evolution Equations*, Lectures Notes Series, vol. 34, Seoul National University, Seoul, 1996.
- [3] E. Hebey, *Nonlinear Analysis on Manifolds: Sobolev Spaces and Inequalities*, Courant Lecture Notes in Mathematics, vol. 5, New York University Courant Institute of Mathematical Sciences, New York, 1999.
- [4] D. Jerison and G. Lebeau, *Nodal sets of sums of eigenfunctions*, in *Harmonic Analysis and Partial Differential Equations (Chicago, IL, 1996)*, 223–239, Chicago Lectures in Math., Univ. Chicago Press, Chicago, IL, 1999.
- [5] J. Jost, *Riemann Geometry and Geometric Analysis*, Springer-Verlag, Berlin, Heidelberg, 2005.
- [6] M. M. Lavrent'ev, V. G. Romanov and S. P. Shishat'skii, *Ill-posed Problems of Mathematical Physics and Analysis*, American Mathematical Society, Providence. 64, 1986.
- [7] G. Lebeau and L. Robbiano, *Contrôle exact de l'équation de la chaleur*, *Commun. Part. Diff. Eq.*, **20**(1995), 335–356.
- [8] G. Lebeau and L. Robbiano, *Stabilization de l'équation des ondes par le bord*, *Duke Math. J.*, **86**(1997), 465–491.
- [9] G. Lebeau and E. Zuazua, *Null controllability of a system of linear thermoelasticity*, *Arch. Rational Mech. Anal.*, **141**(1998), 297–329.
- [10] X. Liu and X. Zhang, *On the local controllability of a class of multidimensional quasilinear parabolic equations*, *C. R. Math. Acad. Sci. Paris*, **347**(2009), 1379–1384.

- [11] A. López, X. Zhang and E. Zuazua, *Null controllability of the heat equation as singular limit of the exact controllability of dissipative wave equations*, *J. Math. Pure. Appl.*, **79**(2000), 741–808.
- [12] Q. Lü, *Bang-Bang principle of time optimal controls and null controllability of fractional order parabolic equations*, *Acta Math. Sin.*, **26**(2010), 2377–2386.
- [13] Q. Lü, *Control and Observation of Stochastic Partial Differential Equations*, Ph D Thesis, Sichuan University, 2010.
- [14] Q. Lü, *Some results on the controllability of forward stochastic heat equations with control on the drift*, *J. Funct. Anal.*, **260**(2011), 832–851.
- [15] Q. Lü and G. Wang, *On the existence of time optimal controls with constraints of the rectangular type for heat equations*, *SIAM J. Control Optim.*, **49**(2011), 1124–1149.
- [16] L. Miller, *How violent are fast controls for Schrödinger and plate vibrations?*, *Arch. Rational Mech. Anal.*, **172**(2004), 429–456.
- [17] J. Milnor, *Morse Theory*, Ann. of Math. Studies, Princeton Univ. Press, Princeton, N. J., 1963.
- [18] K.-D. Phung and X. Zhang, *Time reversal focusing of the initial state for Kirchhoff plate*, *SIAM J. Appl. Math.*, **68**(2008), 1535–1556.
- [19] G. Wang, *L^∞ -null controllability for the heat equation and its consequences for the time optimal control problem*, *SIAM J. Control Optim.*, **47**(2008), 1701–1720.
- [20] X. Zhang, *Explicit observability estimate for the wave equation with lower order terms by means of Carleman inequalities*, *SIAM J. Control Optim.*, **39**(2001), 812–834.
- [21] C. Zheng, *Controllability of the time discrete heat equation*, *Asymptot. Anal.*, **59**(2008), 139–177.
- [22] E. Zuazua, *Controllability and observability of partial differential equations: Some results and open problems*, In *Handbook of Differential Equations: Evolutionary Differential Equations*, vol. 3, Elsevier Science, 2006, 527–621.