



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Paediatrics and Child Health

Division of Woman and Child Health

3-2020

A descriptive study of double burden of malnutrition in mothers of children with severe acute malnutrition admitted in children hospital and institute of child health, Multan

Saadia Khan

Children Hospital and Institute of Child Health, Multan, Pakistan

Umme Ammara

Children Hospital and Institute of Child Health, Multan, Pakistan

Reema Arshad

Aga Khan University, reema.arshad@aku.edu

Farah Naz

Children Hospital and Institute of Child Health, Multan, Pakistan

Kamran Ishfaq

Bahauddin Zakarya University, Multan, Pakistan

Follow this and additional works at: <https://ecommons.aku.edu/>

[pakistan_fhs_mc_women_childhealth_paediatr](https://ecommons.aku.edu/pakistan_fhs_mc_women_childhealth_paediatr)



Part of the [Maternal and Child Health Commons](#), and the [Pediatrics Commons](#)

Recommended Citation

Khan, S., Ammara, U., Arshad, R., Naz, F., Ishfaq, K. (2020). A descriptive study of double burden of malnutrition in mothers of children with severe acute malnutrition admitted in children hospital and institute of child health, Multan. *JPMA. The Journal of the Pakistan Medical Association*, 70(3), 417-420. Available at: https://ecommons.aku.edu/pakistan_fhs_mc_women_childhealth_paediatr/822

A descriptive study of double burden of malnutrition in mothers of children with severe acute malnutrition admitted in Children Hospital and Institute of Child Health, Multan

Saadia Khan¹, Umme Ammara², Reema Arshad³, Farah Naz⁴, Kamran Ishfaq⁵

Abstract

Objective: To assess the nutritional status of mothers of severely malnourished children, and to evaluate the factors associated with inadequate caloric intake of children with severe acute malnutrition versus sufficient caloric intake by mothers.

Methods: The cross-sectional descriptive study was conducted from January to June 2016 at Children Hospital and Institute of Child Health, Multan, Pakistan, and comprised mothers of severely malnourished children admitted for treatment. Data was collected using a questionnaire administered through interviews. Elements noted included body mass index, 24-hour recall, dietary restriction during pregnancy, lactation and complimentary feeding patterns of the children. The mothers were also examined for clinical signs of iron deficiency anaemia.

Results: Of the 100 women, 20(20%) were malnourished, 42(42%) were of normal weight, and 38(38%) were either overweight, pre-obese or obese. The caloric intake of 26(26%) mothers was less than 1500 kcal per day, while 42(42%) were taking between 2000-2500 kcal per day and 32(32%) were taking >2500 kcal daily. Also, 67(67%) women were suffering from anaemia and 80(80%) women had some myths related with dietary restrictions during pregnancy, lactation and complimentary feeding of children. Overall, 23(23%) women breastfed their babies.

Keywords: SAM, Severe acute malnutrition, Double burden, Obesity, BMI. (JPMA 70: 417; 2020).

<https://doi.org/10.5455/JPMA.290318>

Introduction

There is a recent emergence of 'double burden of malnutrition', which consists of under-nutrition among children and over-nutrition in adults, especially women. This alarming shift from under-nutrition in childhood to over-nutrition in adulthood is the result of speedy economic development, globalisation, urbanisation, lifestyle changes, high-calory diets and reduced physical activity. Thus, over-nutrition and under-nutrition simultaneously occur among different age groups within the same population.¹ This double burden of malnutrition has been noticed at country, household, and even individual levels. The more pronounced and visible pattern is an overweight or obese mother with a malnourished and underweight child. Although poverty is linked with under-nutrition among children, it can also cause obesity among adults. Moreover, under-nutrition

in childhood also escalates the possibility of obesity in adulthood if the environmental factors are obesogenic. Both under-nutrition and over-nutrition has the potential to lead to chronic diseases.² Low socioeconomic status (SES) and poverty contribute to the double burden of malnutrition.³

Obesity is a dangerous factor for many chronic diseases, such as respiratory illnesses, osteoporosis, diabetes mellitus (DM), hypertension (HTN), and cardiovascular diseases.^{4,5} Obesity is also a risk factor for the top four preventable diseases related to child mortality which are acute respiratory diseases, diarrhoea, neonatal sepsis and malaria.⁶ Almost 35% of child mortality is linked to macro- and -nutrient deficiencies.⁷ In addition to its effect on mortality, under-nutrition also affects human development in many aspects. The World Bank (WB) estimates that the annual cost burden of malnutrition to the global economy is approximately \$80 billion.⁸

Similarly, obesity also drastically affects health and it is an independent factor for diseases like cardiovascular

^{1,2,4}Children Hospital and Institute of Child Health, Multan, Pakistan; ³Aga Khan University, Karachi, Pakistan; ⁵Bahauddin Zakarya University, Multan, Pakistan.

Correspondence: Reema Arshad. e-mail: reemaarshad0@gmail.com

diseases, HTN, type 2 diabetes mellitus (T2DM), hyperlipidaemia, osteoarthritis, and a few types of cancer.^{9,10}

There are a rapid rise in obesity in adult female general population of Pakistan and yet wasting in children is still rampant. According to the National Nutritional Survey 2011, Pakistan is also facing double burden of malnutrition. Female obesity is on the rise and on the other hand 24% children are severely stunted.¹¹ On the one hand, the average female is consuming more calories than required while on the other, they are still facing significant micro-nutrient deficiencies such as iron, calcium and vitamin s A and D. Almost half of the adult women population is anaemic.¹² The obese female population indicates the availability of sufficient food and caloric resources, but malnourished children indicate lack of food given to children or wrong feeding practices. It is an indication that although there are sufficient food sources present at home to meet energy requirements of the mothers, but the children are being underfed. These underfed children may be due to dietary restrictions and myths related to complementary feeding despite having enough food at home.

The current study was planned to assess the nutritional status of the mothers of severely malnourished children, and to evaluate the factors associated with inadequate caloric intake of children with severe acute malnutrition (SAM) versus sufficient caloric intake by the mothers.

Subjects and Methods

The cross-sectional descriptive study was conducted from January to June 2016 at Children Hospital and Institute of Child Health, Multan, Pakistan, and comprised mothers of severely malnourished children admitted for treatment. Non purposive simple selection technique was used to raise the sample. All the mothers of patients admitted during the study period who furnished consent were included. The mothers of children with other causes of malnutrition, like coeliac disease and chronic illnesses, and those who refused to give consent were excluded. Body mass index (BMI) was used to assess the nutritional status of the mothers and they were classified using the World Health Organisation (WHO) classification of BMI for South Asian populations.¹³

A questionnaire was designed and data was collected through detailed interviews with the mothers (Annexure). A pre-designed questionnaire was used to evaluate

Annexure: Questionnaire used for data collection.

Name: _____		Age: _____		Contact no. _____	
Education: _____		Income level: _____		Address: _____	
Height:		Weight:		BMI:	
Clinical signs for iron deficiency					
Eye examination					
Palm pallor					
Tongue examination					
24 Hour Recall					
Meal	Food item	Quantity	Kcal		
Breakfast					
Snack 1					
Lunch					
Snack 2					
Dinner					
Illness of child during past 6 months					
Disease	No	Once	Twice	>3	
Pneumonia					
Diarrhoea					
Baby feed					
Breastfeeding	Yes	No			
Complimentary feeding	Food item	Quantity	Frequency		
Foods avoided	Food item	Reasons to avoid			
During pregnancy					
During lactation					

dietary restriction during pregnancy, lactation, breastfeeding and complimentary feeding patterns of children. A 24-hour dietary recall was also taken to evaluate the daily caloric intake of the subjects. The mothers were asked to recall every single item they had eaten in the preceding 24 hours. The quantity of food was calculated by showing the mothers standard cups and spoons. The 24-hour recall was evaluated using Pakistan Food Composition Tables.¹⁴ As the mothers were not literate. It was not possible to obtain a 3-day written recall for dietary evaluation.

Mothers were also examined for clinical signs of iron deficiency anaemia i.e., palm paleness, eye and tongue examination. Risk factors associated with SAM children were further evaluated in percentages according to underweight and overweight women.

Results

Of the 100 women, 20(20%) were malnourished, 42(42%) were of normal weight, 17(17%) were overweight, 14(14%) were pre-obese and 7(7%) were obese. The overall mean age was 25.7±3.4 years. The caloric intake

Table: Risk factors leading to SAM in children with underweight and overweight/obese mothers.

Risk factors leading to SAM in children	n	SAM children of underweight mothers	SAM children of overweight and obese mothers
Lack of breastfeeding	77	32 (32%)	45 (45%)
Lack of adequate complimentary feeding	68	28 (28%)	40 (40%)
Food restrictions due to myths/taboo	33	33 (33%)	47 (47%)
Frequent infections i.e. Diarrhoea	16	16 (16%)	24 (24%)
Pneumonia	30	12.2 (12.2%)	17.7 (17.7%)

"Severe acute malnutrition (SAM) is defined by a very low weight for height (below -3z scores of the median World Health Organisation [WHO] growth standards), by visible severe wasting, or by the presence of nutritional oedema."²⁴

of 26(26%) mothers was less than 1500 kcal per day, while 42(42%) were taking between 2000-2500 kcal per day and 32(32%) were taking >2500 kcal daily. Also, 67(67%) women were suffering from anaemia and 80(80%) women had some myths related with dietary restrictions during pregnancy, lactation and complimentary feeding of children (Table). Overall, 23(23%) women breastfed their babies.

Despite the availability of adequate diet at home, all children were suffering from SAM although 29(29%) of the women were overweight and obese.

Discussion

Pakistan is a developing country facing a double burden' of malnutrition. The current findings that 29% mothers of SAM children were themselves overweight or obese is indicative of the double burden of malnutrition which leads to disorders linked with under-development, poor quality of life because of infectious diseases and nutrition deficiencies, and preventable diseases secondary to urbanisation or rapid industrialisation.²

In Pakistan, adult obesity and overweight has become a public health hazard in the last two decades. A study in Thailand has also shown similar results regarding alarming magnitudes of overweight and obesity in urban populations where females tend to have higher proportion of overweight and obesity than males of similar population.¹⁴ Researches from Bangladesh and Indonesia have observed low physical activity and high-calorie diets mainly composed of junk foods and less consumptions of fruits and vegetables further contribute to obesity.^{15,16} Studies in Pakistan also found results similar to this study.^{5,17}

In a study conducted in several low-income countries, a growing problem of under-nutrition and over-nutrition was also found, like in Pakistan. The children are dying because malnutrition and being underfed, and the adults, especially women, are becoming more overweight and facing complications related to obesity.¹⁸ A research in Brazil slums also showed this contradictory co-occurrence of under-nutrition and over-nutrition in the same family.¹⁹

According to a study, pregnancy and early childhood is a crucial time to provide a good base for adequate growth and development. The poor nutritional status in children under the age of five specifies the nutritional status and dietary intake of mothers during pregnancy and complimentary feeding after 6 months. The trend of obese mothers and SAM children indicates that although there is sufficient availability of food resources at home, the children are still not fed enough calories to provide adequate growth and development. The insufficient calorie intake of children is mainly due to lack of breastfeeding and intake of diluted formula via bottle.²⁰ Similar trends were reported by this study.

Related findings were observed in a report by the United Nations International Children's Emergency Fund (UNICEF) that children's diet is also lacking in vital nutrients due to certain myths and taboos regarding food intake during pregnancy, lactation and complementary food. Nutritious foods like meat, eggs, nuts and seeds are prohibited to pregnant and lactating women because of myths of miscarriage and complications associated with these foods. The children are often given commercially prepared expensive formulas and cereals in diluted forms because their concentrated forms are believed to cause diarrhoea. Due to these food myths and taboos, the child's diet is deficient in calories and vital nutrients.^{21,22} According to studies, there is also an increase in the threat of overweight and obesity among children despite the persistently high burden of under-nutrition. Over nutrition phenomenon is age-related and related to the simultaneous decrease in physical activity. The strategies must have interventions for the prevention of obesity among children by promoting physical activity and healthy dietary intake, especially in schools.^{23,24}

Another study also stated that it is fundamental to recognise that both infant / child malnutrition and maternal obesity may have a common aetiology, both of which are significantly associated with poverty and

adverse conditions of their environment.^{25,26}

There is a threat of emerging burden of malnutrition in Pakistan. Further studies are recommended to find out the reasons for inadequate calorie intake in children whose mothers are either of normal weight, overweight or obese.

Conclusion

Only about one-fourth of the women were undernourished and most of them were consuming adequate calories or more than what was required for them. On the other hand, their children were severely malnourished. The breastfeeding rates were also low and diluted bottle feeding and insufficient complementary feeding was common. There were many myths and taboos associated with dietary intake during pregnancy, lactation and complimentary feeding of children.

Disclaimer: None.

Conflict of Interest: None.

Source of Funding: None.

References

- Kolcic I. Double burden of malnutrition: A silent driver of double burden of disease in low-and middle-income countries. *J Glob Health* 2012; 2: 020303.
- Ravishankar A. Is India shouldering a double burden of malnutrition? *J Health Manag* 2012; 14: 313-28.
- Delisle H, Batal M. The double burden of malnutrition associated with poverty. *Lancet* 2016; 387: 2504-5.
- Must A, Spadano J, Coakley EH, Field AE, Colditz G, Dietz WH. The disease burden associated with overweight and obesity. *JAMA* 1999; 282: 1523-9.
- Burton BT, Foster WR, Hirsch J, Van Itallie TB. Health implications of obesity: an NIH Consensus Development Conference. *Int J Obes* 1985; 9: 155-70.
- World Health Organization. World malaria report 2009. Geneva, Switzerland: WHO Press, 2009.
- Black RE, Allen LH, Bhutta ZA, Caulfield LE, De Onis M, Ezzati M, et al. Maternal and child undernutrition: global and regional exposures and health consequences. *Lancet* 2008; 371: 243-60.
- Subramanian SV, Kawachi I, Smith GD. Income inequality and the double burden of under-and overnutrition in India. *J Epidemiol Community Health* 2007; 61: 802-9.
- Swinburn BA, Sacks G, Hall KD, McPherson K, Finegood DT, Moodie ML, et al. The global obesity pandemic: shaped by global drivers and local environments. *Lancet* 2011; 378: 804-14.
- Campos P, Saguy A, Ernsberger P, Oliver E, Gaesser G. The epidemiology of overweight and obesity: public health crisis or moral panic? *Int J Epidemiol* 2006; 35: 55-60.
- Di Cesare M, Bhatti Z, Soofi SB, Fortunato L, Ezzati M, Bhutta ZA. Geographical and socioeconomic inequalities in women and children's nutritional status in Pakistan in 2011: an analysis of data from a nationally representative survey. *Lancet Glob Health* 2015; 3: e229-39.
- Butchart SH, Walpole M, Collen B, Van Strien A, Scharlemann JP, Almond RE, et al. Global biodiversity: indicators of recent declines. *Science* 2010; 328: 1164-8.
- WHO EC. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 2004; 363: 157-63.
- UNICEF. Food Composition Table for Pakistan (revised 2001). Islamabad, Pakistan. [Online] 2001 [Cited 2014 September 02]. Available from URL: http://www.fao.org/fileadmin/templates/food_composition/documents/regional/Book_Food_Composition_Table_for_Pakistan_.pdf
- Aekplakorn W, Kosulwat V, Suriyawongpaisal P. Obesity indices and cardiovascular risk factors in Thai adults. *Int J Obes (Lond)*. 2006; 30: 1782-90.
- Boutayeb A, Boutayeb S. The burden of non communicable diseases in developing countries. *Int J Equity Health* 2005; 4: 2.
- Oddo VM, Rah JH, Semba RD, Sun K, Akhter N, Sari M, et al. Predictors of maternal and child double burden of malnutrition in rural Indonesia and Bangladesh. *Am J Clin Nutr* 2012; 95: 951-8.
- Bhutta ZA, Hafeez A, Rizvi A, Ali N, Khan A, Ahmad F, et al. Reproductive, maternal, newborn, and child health in Pakistan: challenges and opportunities. *Lancet* 2013; 381: 2207-18.
- Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet* 2013; 382: 427-51.
- Silveira KB, Alves JF, Ferreira HS, Sawaya AL, Florêncio TM. Association between malnutrition in children living in favelas, maternal nutritional status, and environmental factors. *J Pediatr (Rio J)* 2010; 86: 215-20.
- Khan S, Iqbal I, Arshad R, Ishfaq K. Breast feeding; awareness and complementary feeding in mothers of children with severe acute malnutrition at nutrition stabilization centre children hospital and institute of child health Multan. *Professional Med J* 2015; 22: 1531-34.
- UNICEF. Tracking progress on child and maternal nutrition: a survival and development priority. New York, USA: UNICEF, 2009.
- Vitolo MR, Gama CM, Bortolini GA, Campagnolo PD, Drachler Mde L. Some risk factors associated with overweight, stunting and wasting among children under 5 years old. *J Pediatr (Rio J)* 2008; 84: 251-7.
- Jafar T, Qadri Z, Islam M, Hatcher J, Bhutta ZA, Chaturvedi N. Rise in childhood obesity with persistently high rates of undernutrition among urban school-aged Indo-Asian children. *Arch Dis Child* 2008; 93: 373-8.
- Collaboration APCS. The burden of overweight and obesity in the Asia-Pacific region. *Obes Rev* 2007; 8: 191-6.
- Abrahams Z, Mchiza Z, Steyn NP. Diet and mortality rates in Sub-Saharan Africa: stages in the nutrition transition. *BMC Public Health* 2011; 11: 801.