

Thomas Jefferson University
Jefferson Digital Commons

Phase 1

Class of 2022

1-2020

Effects of Rocking on Attention and Arousal

Jonathan Lubeck

Thomas Jefferson University, jonathan.lubeck@jefferson.edu


Roseann C. Schaaf, PhD, OTR/L, FAOTA

Thomas Jefferson University, Roseann.Schaaf@jefferson.edu

Rachel L. Dumont, MS, OTS

Thomas Jefferson University, rachel.dumont@jefferson.edu

Follow this and additional works at: https://jdc.jefferson.edu/si_ctr_2022_phase1

 Part of the [Occupational Therapy Commons](#), and the [Translational Medical Research Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Lubeck, Jonathan; Schaaf, PhD, OTR/L, FAOTA, Roseann C.; and Dumont, MS, OTS, Rachel L., "Effects of Rocking on Attention and Arousal" (2020). *Phase 1*. Paper 43.

https://jdc.jefferson.edu/si_ctr_2022_phase1/43

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Effects of Rocking on Attention and Arousal

Jonathan Lubeck, Roseann C. Schaaf, PhD., OTR/L, FAOTA*,

Rachel L. Dumont, MS, OTR/L

(*) indicates primary project advisor

Introduction: Difficulties regulating attention and arousal underlie common developmental conditions and impact performance in academic and work tasks. Sensory-based activities such as sitting in a rocker chair may help regulate attention and arousal by providing sensory stimulation. This pilot study assessed autonomic nervous system activity (ANS) while seated in a specially designed rocking chair providing vestibular sensation during a performance task.

Methods: Thirty-four adults (aged 20-35 years) were randomized to either a rocker chair (Virco Zuma, n = 17) or static chair (n =17) while performing cognitive tasks in a counterbalanced design. ANS measures of skin conductance level (SCL) and respiratory sinus arrhythmia (RSA) were taken at 3-minute periods at baseline, performance, and recovery to assess sympathetic and parasympathetic activation respectively.

Results: For the group using the stationary chair first, sympathetic activity increased from baseline to task ($p=.000$) and baseline to recovery ($p=.002$). Likewise with the Zuma chair first group, sympathetic activity increased from baseline to task ($p=.000$) and baseline to recovery ($p=.000$). For the Zuma chair group only, sympathetic activity decreased from task to recovery ($p=.000$). RSA did not have any statistically significant changes for either group.

Discussion: While these findings were not consistent with the hypothesis that chair type impacts a change in ANS during performance, only the Zuma chair first group had a significant decrease in sympathetic activity from task to recovery. This suggests that using a rocker chair may help decrease arousal level after the task. Further research with larger samples sizes and longer duration tasks is needed to further validate this finding.