

1-2020

Characteristics and Outcomes of Patients Discharged Directly Home from a Medical Intensive Care Unit

Michele Fiorella, MS

Thomas Jefferson University, michele.fiorella@jefferson.edu

Preeyal M. Patel

Thomas Jefferson University, preeyal.patel@jefferson.edu

Ann Zheng

Thomas Jefferson University, ann.zheng@jefferson.edu

Lauren McDonnell, MSN, ACNP

Thomas Jefferson University, lauren.mcdonnell@jefferson.edu

Mina Yasuoka, MSN

*Thomas Jefferson University, mina.yasuoka@jefferson.edu*Follow this and additional works at: https://jdc.jefferson.edu/si_ctr_2022_phase1 Part of the [Translational Medical Research Commons](#)*See next page for additional authors***[Let us know how access to this document benefits you](#)**

Recommended Citation

Fiorella, MS, Michele; Patel, Preeyal M.; Zheng, Ann; McDonnell, MSN, ACNP, Lauren; Yasuoka, MSN, Mina; and Yoo, MD, Erika J., "Characteristics and Outcomes of Patients Discharged Directly Home from a Medical Intensive Care Unit" (2020). *Phase 1*. Paper 71.

https://jdc.jefferson.edu/si_ctr_2022_phase1/71

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Michele Fiorella, MS; Preeyal M. Patel; Ann Zheng; Lauren McDonnell, MSN, ACNP; Mina Yasuoka, MSN; and Erika J. Yoo, MD

Authors: Preeyal M. Patel, BS**; Michele A. Fiorella, MS; Ann Zheng, BS**; Lauren McDonnell, MSN, ACNP; Mina Yasuoka, MSN Erika J. Yoo, MD*

Division of Pulmonary, Allergy, & Critical Care Medicine, Jane and Leonard Korman Respiratory Institute, Sidney Kimmel Medical College, Thomas Jefferson University

Title: Characteristics and Outcomes of Patients Discharged Directly Home from a Medical Intensive Care Unit

RATIONALE: Discharging patients directly from ICUs is an increasingly common practice, largely due to decreased availability of ward beds. The purpose of this study was to describe the population and evaluate the outcomes of patients discharged directly from the MICU.

METHODS: We conducted a retrospective chart review of direct discharges to home from June 2018 to June 2019 from two MICUs. Patients were separated into two groups based on wait time (<24 hours or \geq 24 hours) between ward transfer order and actual discharge. The primary outcome was 30-day hospital readmission. Risk was adjusted using Mortality Probability Model (MPM-III); ICU workload at admission and discharge was estimated using the nine equivalents of nursing manpower use score (NEMS). Patient characteristics were compared using t-test and Fisher exact or X^2 .

RESULTS: There was no difference in severity-of-illness or admission NEMS between the two groups. Patients who waited <24 hours for discharge were more likely to be admitted from home. Patients who waited \geq 24 hours prior to discharge had significantly longer mean hospital LOS compared to those who waited <24 hours (4.63 days vs. 2.65 days, $p < 0.001$). There was no significant difference in 30-day readmission between patients who were discharged after waiting <24 hours vs. waiting \geq 24 hours ($p = 0.70$).

CONCLUSION: Patients who returned directly home from the MICU without any discharge delay were not readmitted to the hospital more frequently within 30 days than

those discharged to home after a delay exceeding 24 hours. Further investigation into identifying those patients for whom early discharge planning directly to home from the ICU is viable and safe may aid in reducing unnecessary critical care resource utilization.