

Thomas Jefferson University
Jefferson Digital Commons

Phase 1

Class of 2022

1-2020

Evaluating the Utility of Diagnostic Workups for Biliary Atresia in Neonates with Cholestatic Jaundice Following Prolonged TPN


Melissa Austin

Thomas Jefferson University, melissa.austin@jefferson.edu

Zubair H. Aghai

Thomas Jefferson University, zubairul.aghai@jefferson.edu

Follow this and additional works at: https://jdc.jefferson.edu/si_ctr_2022_phase1

 Part of the [Obstetrics and Gynecology Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Austin, Melissa and Aghai, Zubair H., "Evaluating the Utility of Diagnostic Workups for Biliary Atresia in Neonates with Cholestatic Jaundice Following Prolonged TPN" (2020). *Phase 1*. Paper 99.

https://jdc.jefferson.edu/si_ctr_2022_phase1/99

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

SI/CTR Abstract

Word count: 250 words

Evaluating the Utility of Diagnostic Workups for Biliary Atresia in Neonates with Cholestatic Jaundice Following Prolonged TPN

Melissa, Zubair Aghai*

(*) indicates primary project advisor

Introduction: Parenteral nutrition associated cholestasis (PNAC) develops in 40-60% of premature infants following TPN for 2-4+ weeks. The incidence of biliary atresia is low and there is a 60-day, post-term window for corrective surgery. There is limited data on both the natural history of PNAC in premature infants following prolonged TPN, as well as the utility of diagnostic tools investigating biliary atresia in these patients.

Methods: A retrospective chart review using EMR data from the Intensive Care Nursery was performed with the following criteria: premature babies diagnosed with cholestasis, born at <1500 grams, and received TPN for ≥ 14 days. Ultimately 61 babies met criteria and data was collected and pooled to produce descriptive statistics and graphs describing laboratory trends.

Results: Median gestational age was 26 [IQR 25, 28] weeks, birth weight was 732 [650, 930] grams and 60% (36/60) were male. After being on TPN for a median of 51 [38, 73] days and developing cholestasis, 12/61 (19.7%) babies underwent hepatobiliary scintigraphy, three of which also underwent repeat scans, 29/61 (47.5%) received GI consults and 32/61 (52.5%) underwent abdominal ultrasounds. No babies were

diagnosed with biliary atresia. Graphical depiction of laboratory trends demonstrates an initial spike in direct bilirubin after TPN cessation, followed by a gradual decline 3-4 weeks later.

Discussion: Despite many diagnostic procedures and consults, PNAC was the only observed diagnosis at discharge for this cohort of babies. Based on the laboratory trends, delaying the investigation of elevated bilirubin until 3-4 weeks after ceasing TPN might prevent unnecessary diagnostics and improve resource allocation.