

Thomas Jefferson University Jefferson Digital Commons

Phase 1

Class of 2022

1-2020

Effect of Contrast-Enhanced Echocardiograms on the Prognosis of Infective Endocarditis

D. Arias Thomas Jefferson University, diego.arias@jefferson.edu

E. Halpern Thomas Jefferson University, ethan.halpern@jefferson.edu

F. Forsberg Thomas Jefferson University, flemming.forsberg@jefferson.edu

J. R. Eisenbrey Thomas Jefferson University, john.eisenbrey@jefferson.edu

Follow this and additional works at: https://jdc.jefferson.edu/si_ctr_2022_phase1

Part of the Cardiology Commons, and the Radiology Commons
<u>Let us know how access to this document benefits you</u>

Recommended Citation

Arias, D.; Halpern, E.; Forsberg, F.; and Eisenbrey, J. R., "Effect of Contrast-Enhanced Echocardiograms on the Prognosis of Infective Endocarditis" (2020). *Phase 1.* Paper 100. https://jdc.jefferson.edu/si_ctr_2022_phase1/100

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's Center for Teaching and Learning (CTL). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson Scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Effect of Contrast-Enhanced Echocardiograms on the Prognosis of Infective Endocarditis

D. Arias, E. Halpern, F. Forsberg, J.R. Eisenbrey

Abstract (\leq 300 words)

Objective

Infective endocarditis (IE) is an infectious disease of the cardiac valves where bacteria colonize the valves; typically, via the formation of vegetations. Recent research has shown that the microbubbles in a contrast-enhanced ultrasound (CEUS) examination can move and dislodge bacterial vegetations *in vitro*. This study investigated whether CEUS resulted in faster resolution of IE in vivo by dislodging the vegetations.

Methods

This IRB approved retrospective study reviewed 36 patients who were diagnosed with IE via echocardiography. Data was sourced from patients within the Jefferson University Hospital's Cardiology EMR system by searching for contrast and vegetation from January 1^{st} , 2013 – January 1^{st} , 2018. Fifteen patients were not given contrast, whereas 21 patients were given contrast via agitated saline (n=16) or an ultrasound contrast agent (n=5). All patients received an echocardiogram after blood cultures confirmed an infection, but before resolution of infection (defined by negative blood cultures). A student's t-test was used for analyses.

Results

The study population was heterogeneous in terms of sex (67.5% male) and race (70% Caucasian, 25% African American, and 5% Asian), with an average age of 51 ± 20 years, and an average BMI of 29.65 ± 7.43 in the contrast group and 27.67 ± 3.16 in the non-contrast group (p=0.37). Following ultrasound, no patients had documented stroke, pulmonary embolism, or systemic blood clot, which physicians could have attributed to a thrombus resulting from dislodging of bacterial vegetation. Overall, blood cultures did not clear faster in patients receiving CEUS compared to those undergoing standard echocardiography, (2.63 ± 2.69 days vs. 1.34 ± 1.11 days, p=0.09). CEUS also did not shorten the admission length in patients with IE, (16.9 ± 7.7 days vs. 19.9 ± 12.1 days; p=0.36).

Conclusion

Based on this limited sample size, patients who underwent CEUS did not have a different prognosis when compared to patients who received a non-contrast echocardiogram.