



Thomas Jefferson University
Jefferson Digital Commons

Phase 1

Class of 2022

1-2020

The Effects of Centering Pregnancy on Attitudes towards Breastfeeding among Women with Opioid Use Disorder

Francesca Mancuso

Thomas Jefferson University, francesca.mancuso@jefferson.edu

Vanessa Short, PhD, MPH

Thomas Jefferson University, vanessa.short@jefferson.edu

Follow this and additional works at: https://jdc.jefferson.edu/si_phr_2022_phase1

 Part of the [Maternal and Child Health Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Mancuso, Francesca and Short, PhD, MPH, Vanessa, "The Effects of Centering Pregnancy on Attitudes towards Breastfeeding among Women with Opioid Use Disorder" (2020). *Phase 1*. Paper 23.

https://jdc.jefferson.edu/si_phr_2022_phase1/23

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

**The Effects of Centering Pregnancy on Attitudes towards Breastfeeding among Women
with Opioid Use Disorder**

Francesca Mancuso, Vanessa Short*

Introduction: Breastfeeding has been found to decrease severity of neonatal abstinence syndrome (NAS), however, breastfeeding rates among women with opioid use disorder (OUD) are significantly lower than national averages. CenteringPregnancy, a group prenatal care model, increases exposure to education regarding breastfeeding, making it more likely that patients breastfeed in traditional populations.

Objective: This project's purpose was to identify and understand current infant feeding attitudes among mothers with OUD and explore whether participation in the CenteringPregnancy "Breastfeeding Session" positively impacted knowledge and attitudes towards breastfeeding.

Methods: Data were collected from 12 women in treatment for OUD who participated in CenteringPregnancy. A survey was given prior to and after the CenteringPregnancy "Breastfeeding Session," and after completion of CenteringPregnancy. Descriptive statistics, including means for continuous data and count and percent frequencies for categorical data used to describe data.

Results: Concern for transmission of methadone and hepatitis C through breast milk, travel schedules to the OUD treatment center, and desire to smoke cigarettes were identified as barriers to breastfeeding. After the breastfeeding session, 58.3% of the women intended to breastfeed.

SI/PHR Abstract

Among those women, none indicated that they did not feel confident in their ability to breastfeed. Follow-up surveys revealed participants found the discussions surrounding breastfeeding helpful and played a role in their decision to breastfeed.

Conclusion: Breastfeeding education programs, such as PregnancyCentering, are needed to address deficits in knowledge of and attitudinal barriers towards breastfeeding among women with OUD. Due to small study size, further studies are needed.