



Thomas Jefferson University
Jefferson Digital Commons

Phase 1

Class of 2022

1-2020

Obesity and Cardiovascular Health Differences Between Urban and Suburban Philadelphia High School Athletes

Peri Levey

Thomas Jefferson University, peri.levey@jefferson.edu

Kyle Prochno

Thomas Jefferson University, kyle.prochno@jefferson.edu

Prashant Rao, MD

David Shipon, MD, FACC, FACP

Thomas Jefferson University, david.shipon@jefferson.edu

Jeremy Close, MD

Thomas Jefferson University, jeremy.close@jefferson.edu

Follow this and additional works at: https://jdc.jefferson.edu/si_phr_2022_phase1

 Part of the [Pediatrics Commons](#), and the [Public Health Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Levey, Peri; Prochno, Kyle; Rao, MD, Prashant; Shipon, MD, FACC, FACP, David; and Close, MD, Jeremy, "Obesity and Cardiovascular Health Differences Between Urban and Suburban Philadelphia High School Athletes" (2020). *Phase 1*. Paper 25.

https://jdc.jefferson.edu/si_phr_2022_phase1/25

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

SKMC Class of 2022

Word count: 245

Obesity and Cardiovascular Health Differences Between Urban and Suburban

Philadelphia High School Athletes

Peri Levey, Kyle Prochno, Prashant Rao, MD, David Shipon, MD, FACC, FACP,**

Jeremy Close,* MD

Introduction: The strongest risk factor for primary hypertension in children is obesity, and concern for pediatric hypertension rises with the rising prevalence of obesity in American children. Specifically, in urban Philadelphia schools from 2009 to 2012, preparticipation physical evaluations (PPEs) demonstrated a positive correlation between blood pressure measurements demonstrating stage 1 or 2 hypertension and elevated body mass index ($P < 0.00001$).

Objective: The purpose was to identify potential differences in cardiovascular health and prevalence of obesity between adolescents in the urban and suburban Philadelphia areas.

Methods: Data from PPEs administered to urban Philadelphia high school athletes in 2018 by the Athlete Health Organization (AHO) was analyzed and compared to data from the Simon's Heart Foundation's HeartBytes registry, which provided data regarding suburban Philadelphia students. Demographic information and basic physical examination data were collected. Each participant's body mass index (BMI) was categorized by percentile for age, and blood pressures were classified according to American Heart Association pediatric guidelines.

Results: Analysis of the AHO data shows that 44.9% of urban students who completed PPEs in 2018 were overweight or obese, and that 43.1% of these students had blood pressures that would

qualify as either stage 1 or 2 hypertension. Further comparison to the HeartBytes data is forthcoming.

Conclusion: Given results to date, it is expected that upon conclusion of the study, the data will support the hypothesis that rates of hypertension and obesity will be higher in the urban underserved population than in the suburban population.