



Thomas Jefferson University  
Jefferson Digital Commons

---

Phase 1

Class of 2022

---

1-2020

## Angiogenin mRNA Expression Levels in Prostate Cancer Tissue

Bowen Yao

Megumi Shigematsu, PhD

Kamlesh Ganesh Pawar, PhD

Venetia Pliatsika, MS

Yohei Kirino, PhD

Follow this and additional works at: [https://jdc.jefferson.edu/si\\_dh\\_2022\\_phase1](https://jdc.jefferson.edu/si_dh_2022_phase1)

 Part of the [Oncology Commons](#), and the [Urology Commons](#)

[Let us know how access to this document benefits you](#)

---

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

**SKMC Class of 2022: SI/DH Abstract**

**Word count: 221**

**Angiogenin mRNA Expression Levels in Prostate Cancer Tissue**

**Bowen Yao, Megumi Shigematsu PhD, Kamlesh Ganesh Pawar PhD, Venetia Pliatsika**

**MS, Yohei Kirino PhD\***

**Computational Medicine Center, Sidney Kimmel Medical College, Thomas Jefferson  
University**

**Introduction:** Prostate cancer is the most commonly diagnosed cancer in men and second leading cause of cancer deaths. Studies have shown that tRNA fragments are upregulated in prostate cancers and play important roles in carcinogenesis. This project looks at how tRNA cleaving enzyme angiogenin expression is regulated in prostate cancer tissues.

**Methods:** Clinical data and mRNA expression levels of selected tRNA cleaving enzymes were extracted from the TCGA website. mRNAs were sequenced using IlluminaGA\_RNASeqV2 at University of North Carolina.

**Results:** 546 samples from 494 patients, with normal tissue from 53 patients were collected. ANG mRNA levels were lower in patients with higher Gleason scores(Intercept=1321.787362, regression coefficient= -87.05499452,  $R^2=0.038$ ). ANG mRNA levels were inconclusive in different clinical T grade( $p=0.15$ ), but were lower in higher pathologic T grade(intercept=1100.484695, x variable=-166.9047227,  $R^2=0.038$ );

ANG expression was lower in patients with nodal involvement versus without(539.56 vs 673.58,  $p=0.005$ ).

**Discussion:** Overall trend we found from the results were ANG mRNA expression levels are down regulated in patients that have more advanced disease versus early disease. This supports the hypothesis that ANG expression plays an interesting role in prostate cancer biology. This trend might be due to the negative feedback due to high levels of tRNA fragments however there is no single theory available to answer this question.

**No references/tables/figures/appendices allowed.**