

RESEARCH ARTICLE

Knee sensorimotor control following anterior cruciate ligament reconstruction: A comparison between reconstruction techniques

Cristóbal San Martín-Mohr^{1,2}*, Iver Cristi-Sánchez^{2,3}, Patricio A. Pincheira^{3,4,5}, Alvaro Reyes⁶, Francisco José Berral⁷, Claudio Oyarzo^{2,8}

1 Carrera de Kinesiología, Departamento Ciencias de la salud, Facultad de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile, **2** Clínica MEDS, Santiago, Chile, **3** Escuela de Kinesiología, Facultad de Ciencias, Universidad Mayor, Santiago, Chile, **4** Laboratorio Integrativo Biomecánica y Fisiología del Ejercicio, Escuela de Kinesiología, Universidad de los Andes, Santiago, Chile, **5** School of Human Movement and Nutrition Sciences, The University of Queensland, Brisbane, Australia, **6** Facultad de Ciencias de la Rehabilitación, Universidad Andrés Bello, Santiago, Chile, **7** Universidad Pablo de Olavide ES-41013 Sevilla, España, **8** Universidad de Los Andes, Santiago, Chile

* These authors contributed equally to this work.

* csanmartino@uc.cl


 OPEN ACCESS

Citation: San Martín-Mohr C, Cristi-Sánchez I, Pincheira PA, Reyes A, Berral FJ, Oyarzo C (2018) Knee sensorimotor control following anterior cruciate ligament reconstruction: A comparison between reconstruction techniques. *PLoS ONE* 13 (11): e0205658. <https://doi.org/10.1371/journal.pone.0205658>

Editor: Riccardo Di Giminiani, University of L'Aquila, ITALY

Received: January 11, 2018

Accepted: September 29, 2018

Published: November 15, 2018

Copyright: © 2018 San Martín-Mohr et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The data was deposited in a stable public repository Figshare (<http://figshare.com/>) DOI [10.6084/m9.figshare.7170368](https://doi.org/10.6084/m9.figshare.7170368).

Funding: The authors received no specific funding for this work. Clínica MEDS provided support in the form of salaries for authors CSMM, ICS and CO, but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. The

Abstract

The sensorimotor system helps to maintain functional joint stability during movement. After anterior cruciate ligament (ACL) injury and reconstruction, several sensorimotor deficits may arise, including altered proprioception and changes in neuromuscular control. It is still unknown whether the type of autograft used in the reconstruction may influence knee sensorimotor impairments. The aim of this study was to comparatively assess the effects of the hamstring tendon (HT) and bone-patellar tendon-bone (BPTB) ACL reconstruction techniques on knee sensorimotor control 6–12 months post-operation. A total of 83 male subjects participated in this study: 27 healthy participants, 30 BPTB-operated patients and 26 HT-operated patients. Active joint position sense in 3 ranges of motion (90–60°, 60–30°, and 30–0° of knee flexion), isometric steadiness, and onset of muscle activation were used to compare sensorimotor system function between groups. Both operated groups had a small (< 5°) but significant joint position sense error in the 30–0° range when compared to the healthy group. No significant differences were found between the operated and the control groups for isometric steadiness or onset of muscle activation. The results of this study suggest that operated patients present knee proprioceptive deficits independently of surgical technique. Nevertheless, the clinical implications of this impairment are still unknown. It seems that selected surgical approach for ACL reconstruction do not affect functioning of the sensorimotor system to a large degree.

specific roles of these authors are articulated in the 'author contributions' section.

Competing interests: The authors have the following interests. Cristóbal San Martín-Mohr, Iver Cristi-Sánchez and Claudio Oyarzo are employed by Clínica MEDS. There are no patents, products in development or marketed products to declare. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials, as detailed online in the guide for authors.

Introduction

The sensorimotor system encompasses all the afferent, efferent, and central integration- processing components involved in maintaining functional joint stability during body movements [1, 2]. Inadequate functioning of this system may predispose joint damage [3, 4], a situation that commonly occurs after ligament injury. Following anterior cruciate ligament (ACL) injury, various sensorimotor impairments may occur, including proprioceptive deficit [5–10], decreased quadriceps and hamstrings strength [11–13], and alterations in muscle activation onset patterns [14]. Moreover, an ACL injury can produce alterations in ligament afferent signals that provoke altered brain activity patterns during knee movements [15]. While some deficits can be compensated by a greater flow of afferent information from other somatosensory pathways (e.g. visual feedback) [16], in some patients, these compensations are insufficient for recovering functional joint stability.

When the conservative treatment for ACL injury is unsuccessful in restoring knee joint functionality, ACL reconstruction (ACLR) may be indicated to restore joint stability. However, sensorimotor impairments and dynamic deficiencies may persist after surgery [17]. For instance, muscle recruitment patterns [18] and muscle strength [19] may remain altered in asymptomatic subjects even when some clinical indicators are within normal ranges. In an extensive literature review, Gokeler et al. [20] indicated that proprioceptive deficits detected by commonly used tests in ACLR patients are not conclusive in detecting sensorimotor impairments. As such, a comprehensive sensorimotor evaluation is needed to assess the function of the joint before returning to normal activities, a decision that in some cases, is based on subjective judgement or time-after-surgery criteria [21]. Assessments that evaluate the integrity and function of sensorimotor components by measuring variables along the afferent or efferent pathways may be used to obtain broader information about joint functionality after ACL injury and surgery [22].

In ACLR patients, reconstruction type may be an important factor related to impaired sensorimotor control [23]. For instance, during the post-operative stage, the graft used for the reconstruction contributes to tissue and mechanoreceptors regeneration [24, 25]. Nevertheless, the exact time needed for mechanoreceptors to appear is unclear, as is the time needed for other somatosensory pathways to compensate for the lower afferent signals coming from the ligament [25, 26]. This altered afferent information results in knee sensorimotor deficits, which may be dependent on the autograft used for ACLR [27]. While quadriceps muscle strength/activation deficits [28], knee proprioception impairment [8], and altered muscle activation patterns [29–31], among other alterations, are present in patients who receive either a bone patellar tendon bone (BPTB) graft or a hamstring tendon (HT) graft, some studies report differences in sensorimotor impairments between graft types [32, 33]. Therefore, existing knowledge on the effects of graft type on sensorimotor control of the knee is still conflicting.

The aim of this study was to comparatively assess the effects of the HT and BPTB ACL reconstruction techniques on knee sensorimotor control in patients 6–12 months post-operation. Essential methods to assess the components of the sensorimotor system [22] were used including: joint-position sense (JPS), muscle tension sense (steadiness), and onset time of electromyographic (EMG) activity of knee muscles during an unexpected perturbation. These methods evaluate the afferent and efferent components of functional joint stability [22] that may be affected differently in each type of graft. Therefore, this information may help to establish the status of individual sensorimotor components that contributes to joint stability during the post-operative period and recovery phase [22]. Due to the nature of the surgery, we hypothesized that the type of graft would differentially impact at least one dimension of sensorimotor control.

Materials and methods

Participants

Participants were recruited by public announcements within the clinic and by contacting ACLR patients listed in the clinic’s database. A total of 30 healthy participants and 89 ACLR patients were contacted within a timeframe of 18 months and were interviewed to assess clinical eligibility. Subject exclusion criteria included the following: any previous surgical intervention in the lower extremities (healthy subjects); acute or chronic pain in the lower extremities within the past six months (healthy subjects); more than one concomitant injury/repair during surgery (e.g. tibial collateral ligament injury plus meniscus repair; operated subjects); body mass index higher than 30 (both groups); chronic ankle instability (both groups). After checking for inclusion and exclusion criteria, 83 males were invited to participate in the study, and were divided into three groups: control (n = 27), BPTB (n = 30) and HT (n = 26). Anthropometric characteristics of each group are presented in Table 1. All operated patients were assessed 6 to 12 months post-surgery (8.6 ± 2.3 months). All study participants scored between four and seven on the Tegner scale for physical activity [34].

All reconstructed patients were rehabilitated at the same clinic with the same general aims for each rehabilitation phase [17]. However, specific precautions were taken for each type of graft, with excessive stress avoided over the anatomical zone where the graft was obtained. The aims of the first six weeks of rehabilitation were to reduce inflammation, restore the knee range of motion, favor muscle activation, and gait retraining. In the second stage, (six weeks post-surgery), the aims were to strengthen the knee musculature and increase neuromuscular control. In the third stage (12 weeks post-surgery), the aims were to normalize knee musculature strength and to start jogging. In the final stage (20 and 32 weeks post-surgery), the aim was to restore functional performance (e.g. in plyometric exercises, jumping, and changes in direction) [17]. All participants provided signed informed consent before participating in the study, which was approved by the Ethical Committee of the Faculty of Medicine, Pontificia Universidad Catolica de Chile and was conducted in accordance with the Declaration of Helsinki.

Measurements

The testing order was randomized by the investigators. Grouping assignments were sorted by the principal investigator, who was blinded to information regarding surgical method and the post-operation time of each patient. The sensorimotor control of the knee was measured through: (a) JPS in three distinct ranges of joint movement; (b) quadriceps muscle tension sensation using the isometric steadiness technique, and; (c) EMG onset of muscle activation of

Table 1. Anthropometric characteristics, time post-surgery and activity level of each group.

	Groups		
	Control (n = 27)	BPTB (n = 30)	HT (n = 26)
Age (years)	24.27 ± 3.28	25.77 ± 4.47	26.60 ± 5.74
Weight (kg)	75.90 ± 8.53	73.16 ± 7.46	78.08 ± 9.40
Height (m)	1.76 ± 0.06	1.72 ± 0.06	1.74 ± 0.06
BMI (kg/m ²)	24.28 ± 2.07	24.69 ± 1.80	25.48 ± 2.42
Time Post-Surgery (months)	—	7.77 ± 2.28	8.92 ± 2.30
Activity Level (Tegner Scale)	6.40 ± 1.71	5.88 ± 1.41	5.39 ± 1.99

Characteristics of each study group. Values are mean ± standard deviation. HT: hamstring tendon; BPTB: bone-tendon-bone.

<https://doi.org/10.1371/journal.pone.0205658.t001>

vastus medialis, vastus lateralis, semitendinosus, and biceps femoris muscles after an unexpected perturbation. The test side for ACLR subjects was the reconstructed leg, whereas the test side for control subjects was pseudorandomly selected to counterbalance the ACLR limbs. This resulted in three groups that were used for comparisons: (i) leg of the healthy subjects (control group); (ii) reconstructed leg of patients operated with the HT method; and (iii) reconstructed leg of patients operated with the BPTB method.

Joint-position sense. The aim of this test was to evaluate JPS, the ability of subjects to actively replicate a previously determined joint position [22, 35]. Angular measurements of the knee joint were taken with a uniaxial electrogoniometer (Kinetecnic, Santiago, Chile) using an incremental encoder (Hengstler, Aldingen, Germany) connected to a computer. The encoder was adapted for the patient with a rod, enabling knee angle assessment (Fig 1A). This device was shown to have a good resolution (5000 pulses, 0.072° of resolution) with a high reproducibility rate (intraclass correlation coefficient = 0.999). As previously reported [36], three ranges of movement (90–60°, 60–30°, and 30–0°) were used [5, 7], with 0° being knee full extension. These ranges were chosen because previous studies suggest that afferent discharge of capsule-ligamentous proprioceptors is influenced by joint angle, where different discharge rates occur within the knee range of motion [37]

For the test, each subject was seated on the edge of a stretcher, with the fulcrum of the electrogoniometer aligned with the axis of the knee (i.e. lateral femoral condyle). For each range, a target knee angle was first achieved using visual feedback on a computer screen to ensure that the subject selected the target within the ranges of movement pre-established for the evaluation. While the subject was extending their knee, a green light appeared on the screen if the knee angle was within the range of movement, and a red light appeared if the knee angle was out of range. Once a target angle was chosen, the subject pushed a button associated with the encoder system, which saved the selected target angle in the computer. In other words, each subject chose any angle within the range of movement as the target angle for the JPS assessment, angle that was used in the consecutive repetitions. This procedure was repeated for each




Fig 1. Knee sensorimotor assessment. A Joint-position sense test with the uniaxial electrogoniometer. B Steadiness test (green line) with the load cell and paradigm set to 15% maximum isometric voluntary contraction (red line). C Muscle activation onset test in the knee, showing the positions of the surface EMG sensors and platform.

<https://doi.org/10.1371/journal.pone.0205658.g001>

of the assessed ranges (i.e., 90–60°, 60–30°, and 30–0°), obtaining three different target angles for each of the ranges [36]. Later, without visual feedback, the subject tried to reproduce the target angle for each of the ranges. Independent of the evaluated range of movement, subjects started from 90° of knee flexion towards extension. This procedure was repeated 10 times for each extremity and for each of the defined ranges of joint movement. Therefore, a total of 30 repetitions were obtained. For each repetition, the absolute value of the difference between the achieved angle and the target angle was calculated (e.g. error angle). Then, the differences of the 10 repetitions were finally averaged for each range. The joint position signals from the electrogoniometer were processed on Igor Pro 6.0 software (WaveMetrics Inc, Lake Oswego, USA).

Muscle tension sensation: Steadiness. The purpose of this assessment was to evaluate the ability of subjects to maintain a constant force at 15% of maximum voluntary isometric contraction, which reflects fine muscle control [38, 39]. Isometric steadiness was represented as the coefficient of variation between the target and the force maintained by the subject. First, the maximum voluntary isometric contraction (MVIC) was assessed. Participants were seated in an instrumented chair with approximately 90° of knee flexion, with a load cell (Scottsdale, Arizona, USA) anchored to the distal end of the leg at the level of the ankle (Fig 1B). Participants were asked to exert an MVIC of the knee extensor muscles for 4 s. This procedure was repeated three times, with a rest period of 1 min between each repetition. The signal from the load cell was captured with a Trigno Wireless System amplifier (Delsys, Boston, USA) at a sample frequency of 2000 Hz, where the maximum force obtained between the three repetitions was selected for further analysis.

Knee isometric steadiness was evaluated with the same setup as the MVIC assessment. A computer screen projected a trapezoidal figure (Fig 1B), where the upper part represented the 15% of the previously measured MVIC. Each subject was asked to exert knee extensor force to reach this target, coinciding with the projected trapezoidal figure. Real-time feedback on the exerted force was provided so that the task could be sustained for 20 s and to match the trapezoidal target. Subjects performed four practice trials separated by 45 seconds of rest. After practice, subjects rested for two minutes and then performed three repetitions of the task, with a rest period of 1 min between each repetition. Isometric steadiness was quantified as the coefficient of variation between isometric strength fluctuations around the projected trapezoidal target [38, 39]. Stabilization of the exerted force occurs normally during the first 8 seconds of the contraction [40], therefore, a visual criteria was used to select the start of the stable signal. A 10-second window of analysis was used thereafter, beginning from the selected start of the stable signal. This method has been previously used to analyze the most stable part of the generated force and has been proven reliable [39, 41]. The final obtained result was the coefficient of variation (%) of the three repetitions, which is a measure of statistical dispersion that describes the degree of variability between measurements. Igor Pro 6.0 software was used to calculate isometric steadiness.

Onset time of muscle activation in the knee muscles. The onset of muscle activation in the knee muscles was estimated utilizing surface electromyography. EMG bipolar sensors (Delsys, Boston, USA) were positioned on the vastus medialis, vastus lateralis, semitendinosus, and biceps femoris muscles of each subject according to SENIAM recommendations [42]. EMG signals were pre-amplified in a simple differential manner, filtered in a bandwidth of 50–450 Hz, and recorded at a sampling frequency of 2000 Hz (Trigno Wireless System, Delsys, Boston, USA). Two destabilizing platforms elicited perturbations in both lower limbs as previously reported [22, 30, 43] (Fig 1C). A sudden fall of the platforms causes 20° of inversion at the ankle in a weight-bearing condition. This mechanical perturbation over the ankle causes a generalized lower limb destabilization, inducing neuromuscular responses to overcome a loss

in balance [35]. The aim of this method is to stimulate the lower limb in a position that ensures muscle activation, joint capsule compression, and stretching of the skin during the evaluation, all of which is crucial information for central proprioception and balance control [35, 44].

The fall of the platform was detected with a triaxial accelerometer (Delsys, Boston, USA) that was synchronized with the EMG signals. This procedure was performed six times for each extremity, with the side of perturbation being randomly selected. Blindfolds and earplugs were used with each participant to nullify the effects of vision and hearing in the evaluations. To calculate the onset of EMG activity, the signals were fully rectified using the Average Rectified Value (AVR) method. Then, an activation threshold was established as the average of the basal amplitude plus five standard deviations in a window of 500 ms measured 50 ms prior to the perturbation. Thus, the start of the muscle activation was defined as any EMG burst that exceeds the activation threshold [45, 46]. A computational script (Igor Pro 6.0, WaveMetrics Inc, Lake Oswego, USA) plus visual inspection allowed for the semiautomatic detection of muscle onset time, defined as the time between the start of perturbation (i.e., stop after the fall of the platform) and the start of EMG muscle activation. This procedure was performed for each of the muscles evaluated. Three randomly selected repetitions were averaged, obtaining the EMG onset latency, which was considered for further analysis.

Statistical analysis

Sample size was estimated based on a one-way analysis of variance (ANOVA) design. Using an alpha level of 0.05 adjusted by multiple comparison tests after ANOVA, a statistical power of 0.9 and an estimated effect size of at least 0.8 (Hedges' g) on the difference of average angle for joint reposition, a sample size of 10 participants per group was obtained [5]. The Shapiro-Wilk test was used to evaluate normality assumptions for all continuous variables. Thus, according to data distribution, a one-way analysis of variance (ANOVA) test or a Kruskal-Wallis test was used to compare JPS, steadiness, and EMG onset among study groups (i.e.: Control, HT, BPTB). A multiple comparison post hoc test (Scheffé or Dunn's) was used after ANOVA. All statistical analyses were performed using the STATA 9.1 software. The level of significance was set at $\alpha = 0.05$.

Results

Joint-position sense

The BPTB and HT groups showed significantly higher error values in comparison to the control group in the 30–0° range ($p = 0.017$ and $p = 0.039$ respectively) (Fig 2). No other significant differences were found when comparing JPS results between control, BPTB, and HT groups. Details of the JPS results are provided in Fig 2.

Muscle tension sense: Steadiness

The coefficient of variation was $2.91 \pm 0.45\%$ for the control, $3.16 \pm 1.32\%$ for the BPTB, and $2.41 \pm 1.21\%$ for the HT groups. A significant difference was found between the HT and BPTB groups ($p = 0.041$) but no difference was found between the HT ($p = 0.061$) and BPTB ($p = 0.99$) vs the control group (Fig 3).

Onset of muscle activation in the knee

No significant differences were found when comparing the onset of muscle activation for the vastus medialis, vastus lateralis, semitendinosus, and biceps femoris muscles between the Control, BPTB, and HT groups. Details of these results are presented in Table 2.




Fig 2. Joint position sense results. Comparative results for JPS between the control, HT, and BPTB groups for the three different angle ranges. Bars represent the mean ± standard deviation (error bars). * indicates significant differences ($p < 0.05$) between the assessed variables. Abbreviations: JPS, joint-position sense; HT, hamstrings tendon; BPTB, bone-patellar tendon-bone.

<https://doi.org/10.1371/journal.pone.0205658.g002>

Discussion

The aim of this study was to determine differences in knee sensorimotor control of ACLR patients operated with HT and BPTB techniques. Our results indicate that 6 to 12 months after surgery, steadiness and onset of muscle activation are similar between both operated groups and the control group. However, near full knee extension, small magnitude JPS errors can persist in operated patients independently of surgical approach. While this detriment in JPS is significant, the clinical implications of this finding remain unknown.

Joint-position sense

Both operated groups tended to present a higher JPS error than the control group near knee full extension. Previous evidence support these findings in athletic [9] and non-athletic populations [10]. Increased errors in JPS in the reconstructed groups may be explained by two theories. Firstly, an increased error in the capacity to reproduce specific joint positions may be interpreted as an alteration in the modulation of proprioceptive information by the muscle




Fig 3. Isometric steadiness results. Comparative results for isometric steadiness between the control, HT, and BPTB groups. Bars represent the mean ± standard deviation (error bars). Abbreviations: HT, hamstrings tendon; BPTB, bone-patellar tendon-bone.

<https://doi.org/10.1371/journal.pone.0205658.g003>

Table 2. Onset of muscle activation for the control and operated groups.

	Vastus medialis (ms)	Vastus lateralis (ms)	Semitendinosus (ms)	Biceps femoris (ms)
Control	97.54 ± 11.11	99.68 ± 13.70	96.85 ± 8.97	101.2 ± 9.35
BPTB	91.48 ± 11.01	90.95 ± 12.17	91.44 ± 12.12	98.75 ± 11.03
HT	93.41 ± 16.19	93.90 ± 17.29	91.38 ± 12.98	94.00 ± 13.66

Onset of muscle activation for the different muscles and groups. Values are mean ± standard deviation. HT: hamstring tendon; BPTB: bone-tendon-bone.

<https://doi.org/10.1371/journal.pone.0205658.t002>

spindle [35, 37]. It has been proposed that the muscle spindle is the final common input of sensory information that the central nervous system uses as feedback for joint position [47] and sensorimotor control. That is, most of the proprioceptive information provided by joint receptors go through the same afferent pathway of the muscle spindle. The spindle contributes to muscle force control and joint position sense through a spinal circuit known as the gamma (γ) loop. This circuit is formed by γ -motoneurons that transmit excitatory/inhibitory pulses to α -motoneurons via Ia afferents, reflex loop that can be modulated by afferent information [48]. Since mechanoreceptors in the ACL provide important afferent information on the relative position and movement of the knee joint [1, 9], ACL injury and reconstruction appears to impair proprioceptive ability through disruption of the transmission and modulation of this sensory information. Abnormal neurologic output from the articular capsule, the collateral and the posterior cruciate ligaments may also contribute to this abnormal spindle modulation. Furthermore, capsuloligamentous structures seems to be more sensitive to joint position changes near knee extension [37], which may be related with our findings of altered JPS just in the 30–0° range. Therefore, the alteration of JPS in the operated subjects may indicate a reduced joint position sense ability due to disruption in muscle spindles' modulation of proprioceptive information.

Another possible explanation for the altered JPS may be related with changes in central nervous system (CNS) after injury and reconstruction. It has been demonstrated that ACL reconstruction and subsequent rehabilitation process evoke neuroplastic changes at the level motor cortex that are not normalized after treatment or return to activity [49]. Indeed, these modifications may elicit supraspinal inhibition of voluntary muscle activation [50], and/or altered neuromuscular control during functional tasks [51]. Even simple cognitive and sensorimotor tasks present higher attentional demands [52] and neurocognitive overload [51] in this kind of patients. This supports the concept that somatosensory, neuroplastic, cognitive, and visual-motor changes can occur after ACL injury and reconstruction [15]. Therefore, the JPS errors in this study may result from CNS reorganization and subsequent altered processing of sensory information, which according to our results, seems to be similar for the HT and BPTB surgical techniques. Our finding of altered JPS in the operated groups can be interpreted from a different point of view. While significant, the recorded error values for JPS were small (~3°) and fall below values established for the clinical relevance (5°) of similar techniques [53]. Furthermore, it seems that there is no substantial evidence of a strong relationship between joint position sense ability and functional performance [20]. Previous studies using H-reflexes to evaluate neuromuscular control in ACLR patients, have found that muscle spindle function is restored within the first 6 months post-surgery [28]. Further, it has been suggested an increased neuromuscular excitability (increased H-reflex amplitude) in patients receiving HT grafts [54]. Therefore, our JPS findings could be considered circumstantial. However, JPS and H-reflex evaluations when used to infer muscle spindle function are methodologically different, thus the obtained results may be interpreted differently. For instance, it can be argued that

joint repositioning includes a higher proportion of descending activation/inhibition from supraspinal centers [1, 35], in comparison to H-reflexes which are considered in essence an estimation of spinal excitability [55]. Overall, the results of this study interpreted in light of the current literature suggest that the altered JPS found in the operated patients may not have clinical implications.

Muscle tension sense: Steadiness

Force steadiness was not altered in the HT or BPTP groups in comparison to the controls. To our knowledge, this is the first study comparing this variable between patients with different ACL surgical grafts. Force fluctuation is dependent on the interaction of multiple features of motor unit behavior, which change as a function of contraction intensity [56]. Alterations in motor unit recruitment and rate coding properties or adaptations in the activation pattern of the motor unit population (e.g., motor unit synchronization and coherence) would affect force variability [57]. Impairments in force steadiness have been associated with ACL injury and ACLR [12]. Considering this, the obtained steadiness results suggest that, by 6–12 months post-surgery, both assessed patient groups (HT and BPTP) had adequately adapted neuromotor control of the knee muscles and were, therefore, able to reduce force fluctuations during isometric contractions. While the exact mechanisms of this adaptation are unknown, several hypotheses can be postulated. First, the post-operative period (i.e., 6–12 months) may have been enough for muscle contraction to recover. Although ACL injury and the degree of graft regeneration are linked to muscle weakness [16, 28], quadriceps and hamstrings strength recovers early during the rehabilitation period after ACLR [58] regardless of the degree of graft regeneration [23, 59] probably due to neuromuscular adaptations such as enhanced ligament-muscular reflex arc excitability [60]. Second, neuromuscular adaptations such as increased antagonist coactivation may increase fine muscle control and force steadiness, thus increasing muscle stiffness and joint stability [11, 12]. Moreover, lower limb training such as within the rehabilitation process, may improve force steadiness directly [61] or indirectly through increased muscle coactivation [62]. Therefore, the results of the present study suggest that commonly used rehabilitation protocols (as used with the assessed patient groups [17]) may restore force steadiness 6 to 12 months after ACL surgery, independent of the type of graft used in the surgery. Future studies may use prospective designs to clarify the effect of the rehabilitation protocols in force steadiness.

Muscle onset timing after ACLR

The patients in both operated groups presented similar muscle onset values at the moment of evaluation. In humans, long latency muscle stretch reflex responses occur within 50–200 ms following an external stimulus [63]. This protective reflex is a rapid muscle response that keeps the joint stable against perturbations that put stability at risk [64]. Deficits in sensory information could alter the latency of a reflex response, thus increasing the risk for joint instability [1]. Prior research has reported that the onset of muscle activation can be delayed in patients with joint instability [52] and ACLR [63]. For instance, a recent study found that ACL reconstructed knees present altered muscle onset timing after ankle perturbations while standing [30]. The possible cause of these differences may be related with the magnitude of the destabilization (30° of inversion, 10° plantarflexion in their study compared to 20° of inversion in this study). However, our comprehensive rehabilitation process over a large sample size make us believe that the onset of muscle activation, and the other components of sensorimotor control evaluated here, may have improved 6 to 12 months post-surgery. Previous literature supports neuroplastic changes following simple exercise, balance and resistive training, and motor task

training [49]. These interventions may enhance neurogenesis, improve cognitive function, and modify nervous system excitability [49], thus enhancing neuromuscular control after ACL reconstruction. Indeed, supraspinal reorganization may contribute to the restoration of long latency muscle stretch reflex responses [60], compensating for possible deficits in afferent information from the joint. Specific muscle strengthening [18] and sensorimotor training can also increase knee joint stability via enhanced reflex excitability and/or an earlier recruitment of motoneurons [65]. Thus, the rehabilitation process that the ACLR group of this study underwent may have partly reestablishing normal functioning of the sensorimotor system after changes produced by injury and reconstruction. This may be related with the lack of differences found between ACL reconstructed groups and the controls. However, these assumptions remain as speculations due to the cross-sectional nature of our cohort. Prospective studies are needed to shed light on the effects of ACLR rehabilitation protocols on specific components of the sensorimotor system.

Study limitations and future research

The results of this study should be interpreted considering some limitations. As mentioned before, the JPS difference found between the control and HT groups was small and the clinical importance of this change remains undetermined. Since afferent information to the muscle spindle may be influenced in part by joint position, future studies should consider more sensitive methods to assess JPS and spindle function such as the H-reflex [66]. Regarding steadiness, the measurement of this variable at different joint positions might be desirable, since knee position may have an effect on muscle force production and muscle activation [67]. In relation with the measurement of muscle onset around the knee, ankle perturbations may be considered non-specific as compared to other methods proposed in the literature [63, 68]. Furthermore, the magnitude of the perturbation might not be enough to elicit a clear response in the knee muscles [30]. However, distal perturbations in a weight bearing situation includes the proprioceptive information of the ankle, which is crucial in the assessment of lower limb sensorimotor control [35] and might be important in the modulation of muscle spindle function. Therefore, we believe that this method contributes to assessing the efferent pathways of the sensorimotor system related to knee stability [22]. Whereas our preliminary results included women's in our sample, we were unable to have similar proportions of males and females between ACLR groups. As there is a possibility that the male response may drive the differences with no effect in females, or there can be interactions between gender and the dependent variables, we opted to exclude women from the final analysis. This means that our conclusions can only be drawn for male patients. Finally, since sensorimotor impairments after ACLR have been found at the motor cortex [16] and spine [54], future studies should assess these levels of the sensorimotor system. Measurement techniques such as functional MRI, transcranial magnetic stimulation and H-reflexes among others may be needed to explore central nervous system reorganization after ACLR.

Conclusions

ACLR individuals have an altered JPS near knee extension ($30-0^\circ$) when compared to control subjects, independently of the surgical approach utilized. This suggests altered modulation and/or processing of proprioceptive information. Because JPS impairment was small in magnitude ($\sim 3^\circ$ of error), and since no other altered sensorimotor components (e.g., force steadiness, muscle onset) were found, the impact of this finding on knee function is arguable. Therefore, the surgical approach used for ACLR appears to not affect the sensorimotor behavior of the knee 6 to 12 months after surgery.

Acknowledgments

The authors would like to thank Rony Silvestre for proof-reading the article.

Author Contributions

Conceptualization: Cristóbal San Martín-Mohr, Iver Cristi-Sánchez, Francisco José Berral, Claudio Oyarzo.

Formal analysis: Iver Cristi-Sánchez, Alvaro Reyes.

Methodology: Cristóbal San Martín-Mohr, Iver Cristi-Sánchez, Alvaro Reyes.

Project administration: Cristóbal San Martín-Mohr, Iver Cristi-Sánchez.

Supervision: Cristóbal San Martín-Mohr, Iver Cristi-Sánchez, Francisco José Berral, Claudio Oyarzo.

Validation: Francisco José Berral.

Visualization: Cristóbal San Martín-Mohr, Iver Cristi-Sánchez.

Writing – original draft: Cristóbal San Martín-Mohr, Iver Cristi-Sánchez, Patricio A. Pincheira.

Writing – review & editing: Cristóbal San Martín-Mohr, Iver Cristi-Sánchez, Patricio A. Pincheira.

References

1. Riemann BL, Lephart SM. The sensorimotor system, part I: the physiologic basis of functional joint stability. *J Athl Train.* 2002; 37(1):71. PMID: [16558670](#)
2. Riemann BL, Lephart SM. The sensorimotor system, part II: the role of proprioception in motor control and functional joint stability. *J Athl Train.* 2002; 37(1):80. PMID: [16558671](#)
3. Myers JB, Lephart SM. The role of the sensorimotor system in the athletic shoulder. *J Athl Train.* 2000; 35(3):351. PMID: [16558648](#)
4. Myers JB, Wassinger CA, Lephart SM. Sensorimotor contribution to shoulder stability: effect of injury and rehabilitation. *Man Ther.* 2006; 11(3):197–201. <https://doi.org/10.1016/j.math.2006.04.002> PMID: [16777465](#)
5. Angoules A, Mavrogenis A, Dimitriou R, Karzis K, Drakoulakis E, Michos J, et al. Knee proprioception following ACL reconstruction; a prospective trial comparing hamstrings with bone–patellar tendon–bone autograft. *Knee.* 2011; 18(2):76–82. <https://doi.org/10.1016/j.knee.2010.01.009> PMID: [20149662](#)
6. Lee DH, Lee JH, Ahn SE, Park MJ. Effect of Time after Anterior Cruciate Ligament Tears on Proprioception and Postural Stability. *PLoS One.* 2015; 10(9):e0139038. <https://doi.org/10.1371/journal.pone.0139038> PMID: [26422800](#). Pubmed Central PMCID: 4589391.
7. Arockiaraj J, Korula R, Oommen A, Devasahayam S, Wankhar S, Velkumar S, et al. Proprioceptive changes in the contralateral knee joint following anterior cruciate injury. *Bone Joint J.* 2013; 95(2):188–91. <https://doi.org/10.1302/0301-620X.95B2.30566> PMID: [23365027](#)
8. Cossich V, Mallrich F, Titonelli V, de Sousa EB, Velasques B, Salles JI. Proprioceptive deficit in individuals with unilateral tearing of the anterior cruciate ligament after active evaluation of the sense of joint position. *Rev Bras Ortop.* 2014; 49(6):607–12. <https://doi.org/10.1016/j.rboe.2013.07.003> PMID: [26229869](#)
9. Relph N, Herrington L. Knee joint position sense ability in elite athletes who have returned to international level play following ACL reconstruction: A cross-sectional study. *Knee.* 2016; 23(6):1029–34. <https://doi.org/10.1016/j.knee.2016.09.005> PMID: [27712856](#)
10. Relph N, Herrington L, Tyson S. The effects of ACL injury on knee proprioception: a meta-analysis. *Physiotherapy.* 2014; 100(3):187–95. <https://doi.org/10.1016/j.physio.2013.11.002> PMID: [24690442](#)
11. Telianidis S, Perraton L, Clark RA, Pua Y-H, Fortin K, Bryant AL. Diminished sub-maximal quadriceps force control in anterior cruciate ligament reconstructed patients is related to quadriceps and hamstring muscle dyskinesia. *J Electromyogr Kinesiol.* 2014; 24(4):513–9. <https://doi.org/10.1016/j.jelekin.2014.04.014> PMID: [24875460](#)

12. Bryant AL, Clark RA, Pua YH. Morphology of hamstring torque-time curves following acl injury and reconstruction: mechanisms and implications. *J Orthop Res*. 2011; 29(6):907–14. <https://doi.org/10.1002/jor.21306> PMID: 21259335
13. Bryant AL, Pua Y-H, Clark RA. Morphology of knee extension torque-time curves following anterior cruciate ligament injury and reconstruction. *J Bone Joint Surg Am*. 2009; 91(6):1424–31. <https://doi.org/10.2106/JBJS.H.01335> PMID: 19487521
14. Williams GN, Barrance PJ, Snyder-Mackler L, Buchanan TS. Altered quadriceps control in people with anterior cruciate ligament deficiency. *Med Sci Sports Exerc*. 2004; 36(7):1089–97. PMID: 15235310
15. Grooms DR, Page SJ, Nichols-Larsen DS, Chaudhari AM, White SE, Onate JA. Neuroplasticity associated with anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther*. 2017; 47(3):180–9. <https://doi.org/10.2519/jospt.2017.7003> PMID: 27817301
16. Nyland J, Gamble C, Franklin T, Caborn DNM. Permanent knee sensorimotor system changes following ACL injury and surgery. *Knee Surg Sports Traumatol Arthrosc*. 2017 May; 25(5):1461–74. <https://doi.org/10.1007/s00167-017-4432-y> PMID: 28154888.
17. Kvist J. Rehabilitation Following Anterior Cruciate Ligament Injury: Current Recommendations for Sports Participation. *Sports Med*. 2004 February 15, 2004; 34:269–80. <https://doi.org/10.2165/00007256-200434040-00006> PMID: 15049718
18. Crow J, Pizzari T, Buttifant D. Muscle onset can be improved by therapeutic exercise: a systematic review. *Phys Ther Sport*. 2011; 12(4):199–209. <https://doi.org/10.1016/j.ptsp.2010.12.002> PMID: 22085715
19. Hiemstra LA, Webber S, MacDonald PB, Kriellaars DJ. Knee strength deficits after hamstring tendon and patellar tendon anterior cruciate ligament reconstruction. *Med Sci Sports Exerc*. 2000; 32(8):1472–9. PMID: 10949014
20. Gokeler A, Benjaminse A, Hewett TE, Lephart SM, Engebretsen L, Ageberg E, et al. Proprioceptive deficits after ACL injury: are they clinically relevant? *Br J Sports Med*. 2012; 46(3):180–92. <https://doi.org/10.1136/bjism.2010.082578> PMID: 21511738
21. Barber-Westin SD, Noyes FR. Factors used to determine return to unrestricted sports activities after anterior cruciate ligament reconstruction. *Arthroscopy*. 2011; 27(12):1697–705. <https://doi.org/10.1016/j.arthro.2011.09.009> PMID: 22137326
22. Riemann BL, Myers JB, Lephart SM. Sensorimotor system measurement techniques. *J Athl Train*. 2002; 37(1):85. PMID: 16558672
23. Takeda Y, Kashiwaguchi S, Matsuura T, Higashida T, Minato A. Hamstring muscle function after tendon harvest for anterior cruciate ligament reconstruction. *Am J Sports Med*. 2006; 34(2):281–8. <https://doi.org/10.1177/0363546505279574> PMID: 16210579
24. MacLeod TD, Snyder-Mackler L, Axe MJ, Buchanan TS. Early regeneration determines long-term graft site morphology and function after reconstruction of the anterior cruciate ligament with semitendinosus-gracilis autograft: a case series. *Int J Sports Phys Ther*. 2013; 8(3):256–68. PMID: 23772342
25. Shimizu T, Takahashi T, Wada Y, Tanaka M, Morisawa Y, Yamamoto H. Regeneration process of mechanoreceptors in the reconstructed anterior cruciate ligament. *Arch Orthop Trauma Surg*. 1999; 119(7):405–9.
26. Barrack RL, Lund PJ, Munn BG, Wink C, Happel L. Evidence of reinnervation of free patellar tendon autograft used for anterior cruciate ligament reconstruction. *Am J Sports Med*. 1997; 25(2):196–202. <https://doi.org/10.1177/036354659702500210> PMID: 9079173
27. Ingersoll CD, Grindstaff TL, Pietrosimone BG, Hart JM. Neuromuscular consequences of anterior cruciate ligament injury. *Clin Sports Med*. 2008 Jul; 27(3):383–404, vii. <https://doi.org/10.1016/j.csm.2008.03.004> PMID: 18503874.
28. Lepley AS, Gribble PA, Thomas AC, Tevald MA, Sohn DH, Pietrosimone BG. Quadriceps neural alterations in anterior cruciate ligament reconstructed patients: A 6-month longitudinal investigation. *Scand J Med Sci Sports*. 2015 Dec; 25(6):828–39. <https://doi.org/10.1111/sms.12435> PMID: 25693627.
29. Dalton EC, Pfile KR, Weniger GR, Ingersoll CD, Herman D, Hart JM. Neuromuscular changes after aerobic exercise in people with anterior cruciate ligament–reconstructed knees. *J Athl Train*; 2011.
30. Pincheira PA, Silvestre R, Armijo-Olivo S, Guzman-Venegas R. Ankle perturbation generates bilateral alteration of knee muscle onset times after unilateral anterior cruciate ligament reconstruction. *Peer J*. 2018; 6:e5310. <https://doi.org/10.7717/peerj.5310> PMID: 30083454
31. Patras K, Ziogas G, Ristanis S, Tsepis E, Stergiou N, Georgoulis AD. ACL reconstructed patients with a BPTB graft present an impaired vastus lateralis neuromuscular response during high intensity running. *J Sci Med Sport*. 2010 Nov; 13(6):573–7. <https://doi.org/10.1016/j.jsams.2009.12.001> PMID: 20227341.
32. Aune AK, Holm I, Risberg MA, Jensen HK, Steen H. Four-strand hamstring tendon autograft compared with patellar tendon-bone autograft for anterior cruciate ligament reconstruction: a randomized study

- with two-year follow-up. *Am J Sports Med.* 2001; 29(6):722–8. <https://doi.org/10.1177/03635465010290060901> PMID: 11734484
33. Heijne A, Werner S. A 2-year follow-up of rehabilitation after ACL reconstruction using patellar tendon or hamstring tendon grafts: a prospective randomised outcome study. *Knee Surg Sports Traumatol Arthrosc.* 2010; 18(6):805–13. <https://doi.org/10.1007/s00167-009-0961-3> PMID: 19851754
 34. Tegner Y, Lysholm J. Rating systems in the evaluation of knee ligament injuries. *Clin Orthop Relat Res.* 1985; 198:42–9.
 35. Han J, Waddington G, Adams R, Anson J, Liu Y. Assessing proprioception: a critical review of methods. *J Sport Health Sci.* 2016; 5(1):80–90. <https://doi.org/10.1016/j.jshs.2014.10.004> PMID: 30356896
 36. Panics G, Tallay A, Pavlik A, Berkes I. Effect of proprioception training on knee joint position sense in female team handball players. *Br J Sports Med.* 2008; 42(6):472–6. <https://doi.org/10.1136/bjism.2008.046516> PMID: 18390919
 37. Proske U, Gandevia SC. The proprioceptive senses: their roles in signaling body shape, body position and movement, and muscle force. *Physiol Rev.* 2012; 92(4):1651–97. <https://doi.org/10.1152/physrev.00048.2011> PMID: 23073629
 38. Beck TW, DeFreitas JM, Stock MS, Dillon MA. Effects of resistance training on force steadiness and common drive. *Muscle Nerve.* 2011; 43(2):245–50. <https://doi.org/10.1002/mus.21836> PMID: 21254090
 39. Tracy BL, Enoka RM. Older adults are less steady during submaximal isometric contractions with the knee extensor muscles. *J Appl Physiol.* 2002; 92(3):1004–12. <https://doi.org/10.1152/jappphysiol.00954.2001> PMID: 11842033
 40. Seynnes O, Hue OA, Garrandes F, Colson SS, Bernard PL, Legros P, et al. Force steadiness in the lower extremities as an independent predictor of functional performance in older women. *J Aging Phys Act.* 2005; 13(4):395–408. PMID: 16301752
 41. Carville SF, Perry MC, Rutherford OM, Smith ICH, Newham DJ. Steadiness of quadriceps contractions in young and older adults with and without a history of falling. *Eur J Appl Physiol.* 2007; 100(5):527–33. <https://doi.org/10.1007/s00421-006-0245-2> PMID: 16983499
 42. Hermens HJ, Freriks B, Merletti R, Stegeman D, Blok J, Rau G, et al. European recommendations for surface electromyography. Roessingh Research and Development. 1999; 8(2):13–54.
 43. Hopkins JT, McLoda T, McCaw S. Muscle activation following sudden ankle inversion during standing and walking. *Eur J Appl Physiol.* 2007; 99(4):371–8. <https://doi.org/10.1007/s00421-006-0356-9> PMID: 17165055
 44. Han J, Anson J, Waddington G, Adams R, Liu Y. The Role of Ankle Proprioception for Balance Control in relation to Sports Performance and Injury. *Biomed Res Int.* 2015; 2015:842804. Pubmed Central PMCID: <https://doi.org/10.1155/2015/842804> PMID: 26583139.
 45. Hodges PW, Bui BH. A comparison of computer-based methods for the determination of onset of muscle contraction using electromyography. *Electroencephalogr Clin Neurophysiol.* 1996; 101(6):511–9. PMID: 9020824
 46. Staude G, Flachenecker C, Daumer M, Wolf W. Onset detection in surface electromyographic signals: a systematic comparison of methods. *EURASIP J Appl Signal Processing.* 2001; 2001(1):67–81.
 47. Needle AR, Charles BBS, Farquhar WB, Thomas SJ, Rose WC, Kaminski TW. Muscle spindle traffic in functionally unstable ankles during ligamentous stress. *J Athl Train.* 2013 Mar-Apr; 48(2):192–202. <https://doi.org/10.4085/1062-6050-48.1.09> PMID: 23672383. Pubmed Central PMCID: 3600921.
 48. Rice DA, McNair PJ, editors. Quadriceps arthrogenic muscle inhibition: neural mechanisms and treatment perspectives. *Semin Arthritis Rheum*; 2010: Elsevier.
 49. Needle AR, Lepley AS, Grooms DR. Central Nervous System Adaptation After Ligamentous Injury: a Summary of Theories, Evidence, and Clinical Interpretation. *Sports Med.* 2017 Jul; 47(7):1271–88. <https://doi.org/10.1007/s40279-016-0666-y> PMID: 28005191.
 50. Urbach D, Awiszus F. Impaired ability of voluntary quadriceps activation bilaterally interferes with function testing after knee injuries. A twitch interpolation study. *Int J Sports Med.* 2002; 23(04):231–6.
 51. Dingenen B, Janssens L, Claes S, Bellemans J, Staes FF. Lower extremity muscle activation onset times during the transition from double-leg stance to single-leg stance in anterior cruciate ligament reconstructed subjects. *Clin Biomech (Bristol, Avon).* 2016; 35:116–23.
 52. Negahban H, Ahmadi P, Salehi R, Mehravar M, Goharpey S. Attentional demands of postural control during single leg stance in patients with anterior cruciate ligament reconstruction. *Neurosci Lett.* 2013; 556:118–23. <https://doi.org/10.1016/j.neulet.2013.10.022> PMID: 24157849
 53. Burgess P, Wei JY, Clark F, Simon J. Signaling of kinesthetic information by peripheral sensory receptors. *Annu Rev Neurosci.* 1982; 5(1):171–88.

54. Rosenthal M, Moore J, Stoneman P, DeBerardino T. Neuromuscular excitability changes in the vastus medialis following anterior cruciate ligament reconstruction. *Electromyogr Clin Neurophysiol.* 2009; 49(1):43–51. PMID: [19280799](https://pubmed.ncbi.nlm.nih.gov/19280799/)
55. Burke D. Clinical uses of H reflexes of upper and lower limb muscles. *Clinical Neurophysiology Practice.* 2016; 1:9–17. <https://doi.org/10.1016/j.cnp.2016.02.003> PMID: [30214954](https://pubmed.ncbi.nlm.nih.gov/30214954/)
56. Taylor AM, Christou EA, Enoka RM. Multiple features of motor-unit activity influence force fluctuations during isometric contractions. *J Neurophysiol.* 2003; 90(2):1350–61. <https://doi.org/10.1152/jn.00056.2003> PMID: [12702706](https://pubmed.ncbi.nlm.nih.gov/12702706/)
57. Vila-Chã C, Hassanlouei H, Farina D, Falla D. Eccentric exercise and delayed onset muscle soreness of the quadriceps induce adjustments in agonist–antagonist activity, which are dependent on the motor task. *Exp Brain Res.* 2012; 216(3):385–95. <https://doi.org/10.1007/s00221-011-2942-2> PMID: [22094715](https://pubmed.ncbi.nlm.nih.gov/22094715/)
58. Harput G, Kilinc HE, Ozer H, Baltaci G, Mattacola CG. Quadriceps and Hamstring Strength Recovery During Early Neuromuscular Rehabilitation After ACL Hamstring-Tendon Autograft Reconstruction. *J Sport Rehabil.* 2015 Nov; 24(4):398–404. <https://doi.org/10.1123/jsr.2014-0224> PMID: [26633588](https://pubmed.ncbi.nlm.nih.gov/26633588/)
59. Janssen RP, van der Velden MJ, Pasmans HL, Sala HA. Regeneration of hamstring tendons after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2013; 21(4):898–905. <https://doi.org/10.1007/s00167-012-2125-0> PMID: [22763570](https://pubmed.ncbi.nlm.nih.gov/22763570/)
60. Mrachacz-Kersting N, Grey MJ, Sinkjær T. Evidence for a supraspinal contribution to the human quadriceps long-latency stretch reflex. *Exp Brain Res.* 2006; 168(4):529. <https://doi.org/10.1007/s00221-005-0120-0> PMID: [16240144](https://pubmed.ncbi.nlm.nih.gov/16240144/)
61. Hortobágyi T, Tunnel D, Moody J, Beam S, DeVita P. Low- or high-intensity strength training partially restores impaired quadriceps force accuracy and steadiness in aged adults. *J Gerontol A Biol Sci Med Sci.* 2001; 56(1):B38–B47. PMID: [11193224](https://pubmed.ncbi.nlm.nih.gov/11193224/)
62. Shultz R, Silder A, Malone M, Braun HJ, Dragoo JL. Unstable surface improves quadriceps: hamstring co-contraction for anterior cruciate ligament injury prevention strategies. *Sports Health.* 2015; 7(2):166–71. <https://doi.org/10.1177/1941738114565088> PMID: [25984263](https://pubmed.ncbi.nlm.nih.gov/25984263/)
63. Madhavan S, Shields RK. Neuromuscular responses in individuals with anterior cruciate ligament repair. *Clin Neurophysiol.* 2011; 122(5):997–1004. <https://doi.org/10.1016/j.clinph.2010.09.002> PMID: [20884289](https://pubmed.ncbi.nlm.nih.gov/20884289/)
64. Solomonow M. Sensory–motor control of ligaments and associated neuromuscular disorders. *J Electromyogr Kinesiol.* 2006; 16(6):549–67. <https://doi.org/10.1016/j.jelekin.2006.08.004> PMID: [17045488](https://pubmed.ncbi.nlm.nih.gov/17045488/)
65. Gruber M, Bruhn S, Gollhofer A. Specific adaptations of neuromuscular control and knee joint stiffness following sensorimotor training. *Int J Sports Med.* 2006; 27(08):636–41.
66. Hwang I-S, Lin Y-C, Ho K-Y. Modulation of Soleus H-Reflex Amplitude and Variance during Pretibial Contraction Effects of Joint Position and Effort Level. *Int J Neurosci.* 2002; 112(6):623–38. PMID: [12325309](https://pubmed.ncbi.nlm.nih.gov/12325309/)
67. Krishnan C, Allen EJ, Williams GN. Effect of knee position on quadriceps muscle force steadiness and activation strategies. *Muscle Nerve.* 2011; 43(4):563–73. <https://doi.org/10.1002/mus.21981> PMID: [21404288](https://pubmed.ncbi.nlm.nih.gov/21404288/)
68. Beard DJ, Kyberd PJ, O'Connor JJ, Fergusson CM, Dodd CA. Reflex hamstring contraction latency in anterior cruciate ligament deficiency. *J Orthop Res.* 1994; 12(2):219–28. <https://doi.org/10.1002/jor.1100120211> PMID: [8164095](https://pubmed.ncbi.nlm.nih.gov/8164095/)