

## Approaching ancient disease from a One Health perspective: interdisciplinary review for the investigation of zoonotic brucellosis

Robin Bendrey<sup>1</sup>, Joe Cassidy<sup>2</sup>, Guillaume Fournié<sup>3</sup>, Deborah C. Merrett<sup>4</sup>, Rebecca Oakes<sup>5</sup>, G. Michael Taylor<sup>6</sup>

<sup>1</sup> School of History, Classics and Archaeology, University of Edinburgh, William Robertson Wing, Old Medical School, Teviot Place, Edinburgh EH8 9AG, UK. [robin.bendrey@ed.ac.uk](mailto:robin.bendrey@ed.ac.uk)

<sup>2</sup> School of Veterinary Medicine, University College Dublin, Belfield, Dublin 4, Ireland. [joseph.cassidy@ucd.ie](mailto:joseph.cassidy@ucd.ie)

<sup>3</sup> Veterinary Epidemiology, Economics and Public Health group, Department of Production and Population Health, Royal Veterinary College, University of London, Hawkshead Lane, North Mymms, Hatfield AL9 7TA, UK. [gfournie@rvc.ac.uk](mailto:gfournie@rvc.ac.uk)

<sup>4</sup> Department of Archaeology, Simon Fraser University, Burnaby, British Columbia V5A 1S6, Canada. [dcmerrret@sfu.ca](mailto:dcmerrret@sfu.ca)

<sup>5</sup> Department of History, University of Winchester, Sparkford Road, Winchester, S022 4NR, UK. [rebecca.oakes@winchester.ac.uk](mailto:rebecca.oakes@winchester.ac.uk)

<sup>6</sup> Department of Microbial Sciences, Faculty of Health and Medical Sciences, University of Surrey, Guildford, UK. [gm.taylor@surrey.ac.uk](mailto:gm.taylor@surrey.ac.uk)

### *Corresponding author*

Robin Bendrey, School of History, Classics and Archaeology, University of Edinburgh, William Robertson Wing, Old Medical School, Teviot Place, Edinburgh EH8 9AG, UK. Telephone: +44 (0)131 6504562. Email: [robin.bendrey@ed.ac.uk](mailto:robin.bendrey@ed.ac.uk)

### *Abstract*

Today, brucellosis is the most common global bacterial zoonosis, bringing with it a range of significant health and economic consequences, yet it is rarely identified from the archaeological record. Detection and understanding of past zoonoses could be improved by triangulating evidence and proxies generated through different approaches. The complex socio-ecological systems that support zoonoses involve humans, animals, and pathogens interacting within specific environmental and cultural contexts, and as such there is a diversity of potential datasets that can be targeted. To capture this, in this paper we consider how to approach the study of zoonotic brucellosis in the past from a One Health perspective, one which explicitly acknowledges the health link between people, animals and environments (both physical and cultural). One Health research is explicitly interdisciplinary and conceptually moves away from an anthropocentric approach, allowing the component parts to be considered in holistic and integrated ways to deliver more comprehensive understanding. To this end, in this paper we review the methods, selected evidence and potential for past brucellosis identification and understanding, focussing on osteological markers in humans and animals, historical, biomolecular and epidemiological approaches. We also present an agenda and potential for future research.

### *Key words*

Zoonoses; brucellosis; *Brucella*; One Health; palaeopathology; biomolecular approaches

### *Suggested running title*

Approaching ancient brucellosis from a 'One Health' perspective

## 1 Introduction

Brucellosis is a disease of global significance today, with major human and animal health and economic impacts. It is the most common bacterial zoonosis, with >500,000 new cases reported each year (Pappas *et al.*, 2006), although the true incidence is estimated at 5-12.5 million cases annually (Hull and Schumaker, 2018). The causative agents are bacteria of the genus *Brucella*, which infect a range of mammalian hosts (Moreno, 2014). *Brucella* species are gram-negative, facultative aerobic, non-motile coccoid or rod-shaped aerobic bacteria which replicate within phagocytic cells of the host reticuloendothelial system (species and their preferred hosts are summarised in Appendix S1 and Table S1).

Of the *Brucella* species known, those infecting domestic animals are generally both zoonotic and most virulent (with the exception of *B. ovis*) compared to strains affecting wild animals (Moreno, 2014). Species that are generally pathogenic to humans are *B. melitensis*, *B. abortus* and biovars 1 and 3 of *B. suis*. They exploit the reproductive cycle of their infected hosts, multiplying in the placenta inducing abortion, in mammary glands following pregnancy and shedding through milk, then mainly infecting other animals through ingestion (Díaz Aparicio, 2013). Human infection normally occurs through ingestion of unpasteurized dairy products and direct contact with infected animals (Moreno, 2014).

Brucellosis is thus a very common zoonotic infection, yet it is rarely identified from the archaeological record. Detection and understanding of past zoonoses could be improved by triangulating evidence and proxies generated through different approaches. The complex socio-ecological systems that support zoonoses involve humans, animals, and pathogens interacting within specific environmental and cultural contexts, and as such there is a diversity of potential datasets that can be targeted. To capture this, in this paper we consider how to approach the study of zoonotic brucellosis in antiquity from a One Health perspective, one which explicitly acknowledges the health link between people, animals and environments (both physical and cultural). In epidemiology, the 'epidemiological triad' of hosts, pathogens and environment is commonly used to summarise the factors that influence infections (Figure 1A). Studies of archaeological human remains have long drawn on the biocultural paradigm to emphasize that these factors can only be fully understood when placed within the context of culture and the constructed human niche (Figure 1B). The One Health approach is explicitly interdisciplinary and emphasizes communication and collaboration across multiple sectors in the delivery of improved health outcomes (Figure 1C). It also ensures that the component factors influencing infections are considered in holistic and integrated ways to deliver more comprehensive understanding than siloed approaches by single disciplines (Lebov *et al.*, 2017). To this end, here we will review the methods, selected evidence and potential for past brucellosis identification, focussing on osteological markers in humans and animals, historical, biomolecular and epidemiological approaches (Figure 1D). We also present an agenda and potential for future research.

## 2 Brucellosis and human palaeopathology

*Brucella* causes systemic infections in humans; any organ of the body may be affected with bacteria localizing intracellularly in the immune system, transported through lymph and haematogenously spread to the spine and knee, sacroiliac and interphalangeal joints, amongst others (Buzgan *et al.*, 2010; Turan *et al.*, 2011). However, a mouse model suggests dispersal of bacteria throughout the body results directly in localization in osteoarticulations suggesting that intermediate tissues may not be necessary for development of skeletal brucellosis (Magnani *et al.*, 2013) as previously thought. Many patients experience joint pain that is often associated with joint swelling and development of septic arthritis, that is, infection of the joint and associated tissues. Although

brucellosis can affect any region of the spine, it most often affects the lumbar vertebrae with concomitant lower back pain (Buzgan et al., 2010; Madkour et al., 1988; Turan et al., 2011). Localization of the disease in the sacroiliac joint(s) is also often accompanied by sciatica and back pain (Corbel, 2006; Priest et al., 2008). Fatality rate in humans is very low, most often following development of brucellar endocarditis (Buzgan et al., 2010; Hull and Schumaker, 2018).

Skeletal manifestations of brucellosis are diverse and non-specific with potential for presentation of many atypical forms (Corbel 2006). The prevalence of complications varies among clinical studies ranging from ten to eighty per cent of brucellosis patients (Mehanic et al., 2012; Turan et al., 2011). Vertebral body lesions are resorptive in nature early in the disease process, with more bone deposition and sclerosis in healing stages (Lifeso et al. 1985) than is seen in tuberculosis, a disease with which brucellosis is often misdiagnosed (Buzgan et al., 2010; Glasgow, 1976). Vertebral destruction is usually less severe in brucellosis than in tuberculosis (Chelli Bouaziz et al., 2008). In addition, clinical manifestations vary with age of the individual: monoarthritis of knee and hip in children, sacroiliitis in children and young adults, and spondylitis in older adults (Chelli Bouaziz et al., 2008; Esmailnejad-Ganji and Esmailnejad-Ganji, 2019). Thus diseases included in the differential diagnosis may differ depending on age of the individual and osseous elements affected.

Monoarticular involvement of peripheral joints, seen mostly in children in the knee results in bone resorption starting at the joint capsule margins. However as complete joint destruction is unusual (al-Shahed et al., 1994), we might expect brucellar osteoarthritis to be less severe and thus less visible in the archaeological record and thus more difficult to diagnose definitively than diseases such as tuberculosis and osteomyelitis that can cause more extensive bone destruction. As with peripheral arthropathy, sacroiliitis is seen as irregular areas of resorption on articular facets. In clinical setting it is visualized radiographically and through CT and MR imaging as widening and blurring of the joint space (Ariza et al., 1993). However, similar lesions are also seen in gout and psoriatic arthropathy (Dayan et al., 2009). To our knowledge, sacroiliitis has not yet been linked specifically with brucellosis in archaeological contexts.

Brucellar spondyloarthropathy, seen in older adults, primarily affects the lumbar spine, although all regions are possible. Resorption is exhibited on the anterior superior margin of vertebral body below attachment site of the annulus fibrosus of the intervertebral disc. Resorptive lesions develop slowly, with sclerosis, increased bone density and thickening of trabeculae (Capasso, 1999; D'Anastasio et al., 2009, 2011). Brucellar lesions of vertebral body endplates (Madkour et al., 1988) have been noted clinically.

Although most of the changes are resorptive as seen above, bone deposition is seen in the development of anterior bone spurs, the 'parrot beak' osteophytes projecting from the inferior end of the vertebral body zone of resorption. For beak osteophytes and Schmorl's nodes, differential diagnosis should include herniation of intervertebral disc or trauma (Aufderheide and Rodríguez-Martín, 1998). Development of paravertebral abscesses (Ariza et al., 1985) including psoas abscess (Turan et al., 2011) similar to that seen in psoas fascia calcification of tuberculosis (Ortner 2003: 232; Roberts and Buikstra, 2019) have also been observed clinically but less frequently than in tuberculosis (Corbel, 2006). Deposition of new woven bone may be present on the visceral surfaces of ribs and the anterior and lateral surfaces of vertebral bodies. Presence of both resorption and deposition in the same bone counter-indicates tuberculosis, unless the new bone is solely for vertebral stabilisation following vertebral body collapse or joint destruction (Mehanic et al. 2012; Roberts and Buikstra, 2019; Waldron 2009). In a Turkish clinical study of 2018 cases, the most common laboratory result was anaemia (Buzgan et al., 2010). Thus, it may be useful to add indicators of anaemia (*cribra orbitalia* and porotic hyperostosis) to the list of potentially diagnostic criteria in palaeopathological studies.

In bone, the feature most commonly observed is Pedro-Pons' sign in vertebrae, which is resorption of the anterior superior margin of one or several vertebrae with underlying sclerosis (Aufderheide and Rodríguez-Martín, 1998; Glasgow 1976; Mehanic et al., 2012; Roberts and Buikstra, 2019; Waldron, 2009). Other skeletal changes include subperiosteal new bone on anterior surfaces of ribs and vertebral bodies (D'Anastasio et al., 2011) and resorption of vertebral endplates (Chelli Bouaziz et al., 2008; Madkour et al., 1988). Any resorptive lesions thought to be pathogen-related on sacroiliac joint surfaces (Dayan et al., 2009) must be distinguished from those that are age-related. Thus, for adults, methods of age estimation other than that of auricular surface morphology should be used in cases of suspected brucellosis. Skeletal region targeted varies with the age-at-death of the individual observed: peripheral arthritis and sacroiliitis in children and young adults, and spondyloarthropathy in older adults (Chelli Bouaziz et al., 2008; Esmailnejad-Ganji and Esmailnejad-Ganji, 2019) although length of chronic infection may also play a part in the predominance of spinal changes in older individuals. In his macroscopic analysis of vertebral lesions from medieval Wharram Percy, England, Mays (2007) takes a conservative approach, suggesting that without at least two categories of evidence (one being biomolecular) diagnosis must be tentative.

Good bone preservation and recovery of as much of each skeleton as possible are the ideal for securing a diagnosis. If death occurs early in disease progression, bony changes may not be sufficiently developed for accurate diagnosis. Because of the variability of disease expression we cannot assume that all cases of brucellosis will lead to advanced and pathognomonic changes in bone, nor that an infected individual, in life, exhibited skeletal brucellosis. Additionally, in a diseased individual, presence of one illness can decrease immune function increasing risk of co-infection. If the coinfecting organism also involves the skeletal system, bone morphology and pattern of lesion location may reflect neither of the infections adequately for definitive diagnosis (Christensen et al., 2013; de Boer et al., 2016). Thus, any estimate in the archaeological record must be considered an underestimate of the true prevalence of brucellosis. Even with the above caveats and difficulties, human skeletons from past populations have been identified as having evidence of *Brucella* infection (e.g. Table 1).

### **3 Brucellosis and animal palaeopathology**

In animals, brucellosis is a sub-acute or chronic disease (Corbel, 2006). The bacteria may enter the body via the gastrointestinal tract, inhalation or conjunctiva, and once they have accessed the circulatory system may spread and cause bacteraemia (Hull and Schumaker, 2018), then settle in the reproductive or musculoskeletal systems (Glynn and Lynn, 2008). In terms of the impact on reproductive tissues, in females infection frequently causes abortion, a key clinical sign of brucellosis, and in males it causes epididymitis and orchitis (Poester et al., 2013).

Of relevance to osteoarchaeological identification, *Brucella* organisms can also localise in bones, especially vertebrae, and synovial structures such as joints, bursae and tendon sheaths causing inflammation and spill-over of infection and inflammation from these sites. The latter can impact on adjacent bone resulting in characteristic focally extensive periosteal responses that can be identified in skeletal remains (Table 2). Localisation in and inflammation of bursae is well known in horses, in conditions such as 'fistulous withers' or 'poll evil' (Denny, 1973), in cattle and, less commonly, sheep and goats with carpal bursitis (carpal hygromas) (Ramadan et al., 1991). In goats and sheep, arthritis may also occur as a rare clinical sign of *B. melitensis* infection (Corbel, 2006). Arthritis affecting the larger limb joints as well as lumbar vertebral lesions and spondylitis are commonly described in pigs with *B. suis* infection (Schlafer and Foster, 2016).

Despite these well-documented impacts of brucellosis on the skeletal system of animals, the fact that these morphological responses simulate those of other bacterial infections within joints and bone (Lignereux and Peters 1999) presents a key challenge to zooarchaeological investigations. Limited modern comparative data on the skeletal manifestation of infectious diseases in domestic animals limits ability to provide definitive identifications, although some propositions have been forwarded in the palaeopathological literature on the separation of diseases, for example Baker and Brothwell (1980, 77) suggest that there is greater periosteal proliferation in brucellosis than tuberculosis. There are no definitive published cases of archaeological animal brucellosis (Table 3).

Palaeopathological approaches should focus on detailed description and development of differential diagnoses of possible infectious agents, something easier attempted in more complete articulating skeletons, with subsequent biomolecular analysis to refine the disease identification. Analysis of disarticulated material should focus on the identification and quantification of potential markers of infection across the skeleton, and assessment of their correlation with locations of known skeletal involvement (Table 3), again supported with biomolecular analysis. Brucellosis also causes late foetal abortion in some taxa, and foetal age estimation of very young remains may therefore give clues as to the potential presence of an infectious agent. Given the range of other pathogens that cause late term abortions in domestic livestock (Tables S2), however, other evidence (e.g. biomolecular) would be needed to confirm an infectious agent.

#### **4 Biomolecular evidence for *Brucella* species in the archaeological record**

Reports of confirmed *Brucella* species retrieved from archaeological remains are limited (Table 4). The earliest DNA evidence comes from the Early Bronze Age North Caucasus (c.3700-3300 BC). During a study of mitochondrial DNA haplogroups and human origins in this region conducted on burials of the Maikop culture, a human burial from Novosvobodnaya was found to have generated sequences from *Brucella abortus* (Sokolov et al., 2016). The authors used a high-throughput sequencing approach rather than individual PCR methods. The DNA fragments obtained were short, in the region of 51-75 bp long and exhibited many C to T transitions, observations consistent with degraded ancient DNA. The isolation of *B. abortus* from these individuals is consistent with the known farming practices of Maikop culture peoples, who kept predominantly pigs and cattle, with the latter being the preferential host of *B. abortus*.

Mutolo et al. (2011) successfully amplified *Brucella* genomic DNA from two human skeletons from medieval Butrint, Albania (Table 4). The PCR targets used were the multi-copy element *IS711* (formerly known as *IS6510*) (Ouahrani et al., 1993) and DNA coding for the 31kDa membrane protein *Bcsp31*. Short templates were targeted; 58bp in the case of *IS711* and 59 bp in the *Bcsp31* PCR. They found that burial 4015 was positive for both *loci* and that burial 2272 was positive only with the more sensitive *IS711* method, suggesting poorer DNA preservation in this individual. As infection with tuberculosis was part of the differential diagnosis, the remains were also tested for three MTB complex *loci* which have been used in ancient DNA (aDNA) studies for some years, namely *IS6110*, *mtp40* and the *oxyR* pseudogene (e.g. Mays et al., 2001). All were negative. The authors were careful to apply tuberculosis PCR methods which would detect similarly degraded templates (62-65 bp) so that this mycobacterial pathogen should also have been detected in bone extracts, if present.

Kay et al. (2014) also used a high-throughput approach (metagenomic shotgun sequencing) to study DNA extracted and amplified from a calcified abdominal nodule present in an adult male burial from Geridu, Sardinia (Table 4). The remains displayed lesions of DISH (diffuse idiopathic skeletal hyperostosis) but were without obvious morphological evidence of brucellosis other than the presence of multiple calcified nodules, which have sometimes been associated with chronic brucellosis, amongst other pathologies (Arcomano et al., 1977; Sevilla-López et al., 2011). Using this

unbiased approach (i.e. without target-specific amplification or capture) they managed to obtain 6.5 fold coverage of a strain of *B. melitensis* from this individual. Further analysis with SNP and deletion typing confirmed that the medieval Geridu-1 isolate belonged to the Ether clade of *B. melitensis*, a lineage considered basal to the phylogenetic tree of the species (Pisarenko et al., 2018).

In a recent publication using proteomic analysis, Greco et al. (2018) reported the presence of a specific peptide sequence associated with *B. melitensis* extracted from organic material preserved in a storage jar from an Egyptian site dating back to the 19<sup>th</sup> Dynasty (1295-1186 BC). Analysis of the cheese residues by UHPLC/high-resolution nanoESI-mass spectrometry showed this contained cow's milk and either sheep or goats milk proteins. The peptide sequence the authors described as indicative of *B. melitensis* (GSIKER) could conceivably have originated from another organism, *Coxiella burnetii*, a Gram-negative organism affecting ruminants. Unfortunately, aDNA analysis was not undertaken to validate the proteomic findings.

This author (GMT) has applied screening PCR methods for *Brucella* to all cases of human skeletal tuberculosis where lesions have suggested that it might be a differential diagnosis. Over the years, this has resulted in testing over 200 human cases and several dozen animal bones (e.g. Bendrey et al., 2008) but in only one instance has any evidence of *Brucella* DNA been detected. This observation was made in an adult female from Tyva, south Siberia (Table 4), where evidence of *Mycobacterium bovis* had already been found in specimens taken from lumbar vertebrae (L3/L4) displaying the classic spinal lesions of tuberculosis (Murphy et al., 2009). The identification of *Brucella* DNA was a late observation made after the completion of the main aDNA analyses which had focused on the typing of *M. bovis* isolates retrieved from four burials of nomadic pastoralists. These individuals spent their lives in close proximity to a number of herd species. The amplification of *Brucella* pathogen DNA was a reproducible observation, but we were not able to pursue this at the time as the sample had been returned for reburial. However, a gel run of the 144 bp amplicon from the *IS711* PCR product was subsequently published in a palaeopathology review (Donoghue, 2008).

## **5 Screening archaeological samples for *Brucella* species: points for consideration and future studies**

After the death of an individual or animal, *postmortem* action of endonucleases and microbial activity results in fragmentation of both host and pathogen DNA. Over time, the DNA may be further modified by chemical processes such as hydrolysis and oxidation (Lindahl and Nyberg, 1972). The soil environment often contains the presence of naturally occurring fixative acids and tannins. These can damage DNA over time or inhibit PCR reactions, if co-extracted (Sidstedt et al., 2015). Extracted aDNA may thus block PCR polymerases due to both intra and inter-strand nucleic acid cross-linking. Modification or loss of nucleotide bases, particularly depurination (Lindahl, 1993) may introduce errors which allow extension but then appear as nucleotide transitions when remnant DNA templates are amplified by PCR and used later for downstream validation measures like cloning and sequencing. The majority of miscoding changes involve C → T/U and G → A transitions (Taylor, 2014 and references therein).

The study of mycobacterial pathogens in the past has been a productive area of research. By protecting them from initial degradation, the waxy outer cell wall of mycobacterial species may be partly responsible for the number of reports of tuberculosis and leprosy in the literature. *Brucella* species have an atypical lipopolysaccharide (LPS) responsible for structural and functional integrity of the bacteria (Cardoso et al., 2006) but lack the mycolic acids and derivatives, which makes the mycobacterial cell membrane relatively impermeable and resilient (Brennan and Nikaido, 1995). The limited reports of *Brucella* in the bioarchaeological literature may thus be a consequence of a greater susceptibility to degradation. Further studies are needed to investigate this aspect in both

human and faunal remains. Information on the association of the pathogen to skeletal lesions and uninvolved or distant skeletal elements is also minimal. The recovery of *Brucella* DNA from the Russian burial lacking obvious osteoarticular lesions mentioned above (Sokolov et al., 2016), implies sampling skeletal remains without lesions might be productive. A factor possibly favouring detection is the faster doubling time (3-4 hours) of the *Brucella* species and hence potential higher bacterial load compared to the slower doubling times of pathogenic mycobacteria such as *M. bovis* (16-20 hours) *M. tuberculosis* (18-54 hrs) or *M. leprae* (14 days). Testing of skeletal lesions proposed as indicators of *Brucella* infection should also be undertaken. Some authors have suggested that lytic erosions on the anterior-superior aspect of the vertebral body are indicative of *Brucellosis* in archaeological cases (Exteberria, 1994; Curate, 2006). However, this is as yet unsupported by testing for the pathogen and an alternative cause, traumatic anterior disc herniation, has been suggested (Mays, 2007). Suggestions for aDNA studies are included in supporting information (Appendix S2).

## **6 Perspectives from historical records**

Sub-disciplines of history have explored questions relating to medical history, human health, demography, agriculture and socio-economic experiences. However, brucellosis remains under-examined and rarely mentioned in historiography. It is not until the development of microbiology that this disease was identified explicitly in the historical record, although given what we know of its transmission, clinical manifestations and impacts, its likely presence and significance to past societies cannot be denied. Re-examination of source materials and their interpretations derived from across the separate fields of historical research have the potential to contribute to both the identification of the disease in the past and the cultural contextualisation of human-animal-environment relationships (Figure 1); examples and future research potential are included in Appendix S3.

## **7 Exploring the dynamics of animal populations and *Brucella* transmission through epidemiological modelling**

Findings about the structure and management of domestic animal populations, the nature and intensity of animal-human interactions, and the trends in the consumption of animal products can also allow scientists to investigate the impact of those features on pathogens' transmission dynamics, and in particular whether epidemiological conditions were met to support disease emergence and endemicity. Such an approach was adopted by Fournié et al. (2017) to explore the potential impact of the origins of animal husbandry on the emergence of zoonotic brucellosis. The Early Neolithic of the Zagros mountains was chosen as a case study for investigating past brucellosis emergence associated with early goat husbandry as there is strong indirect contextual evidence and probable human osteological evidence for the disease (Merrett 2002; 2004). Moreover, previous archaeological investigations had dated a sequence of site assemblages, and, for each of those sites, characterised the age and sex structure of managed goat populations based on the fusion of post-cranial bone remains (Zeder, 2008). Mathematical models simulating the dynamics of domestic goat populations were developed and fitted to these reconstructed goat demographic profiles in each site. *Brucella* transmission was then modelled to assess the likely effect of changes in goat population structure (Fournié et al., 2017). The models indicate that the pathogen could have been sustained, even for low levels of transmission, in small domestic goat populations that lie within the likely ranges estimated for these early farming settlements. This resulted from the creation of dense domestic goat populations, but also the decisions made by early goat farmers on the demographic composition of their herds. As goat farming evolved, some communities began to preferentially retain domestic female goats into adulthood in herds, and selectively cull male goats at a younger age. In this way people inadvertently created population demographic structures which would have increased the transmission potential of the pathogen among goats, as the infectious material

excreted by females following abortion or full-term parturition is the main source of infection. Conditions were thus met for the maintenance of a permanent reservoir of zoonotic infection in close proximity to human settlements, exposing humans to greater risk of infection.

Such an approach has several limitations, due to the nature of the data on which the models rely, and the assumptions about the transmission of *Brucella*, which are based on current knowledge about the disease epidemiology. However, these models can be used to generate hypotheses about factors promoting the circulation and maintenance of pathogens within domestic animal populations and their zoonotic transfer; hypotheses which could then be tested by other disciplines.

## 8 Conclusions

Although brucellosis is today's most common bacterial zoonosis, it is only rarely identified in the past. It is notable, for example, that the *Brucella melitensis* sequenced draft genome from medieval Sardinia shows a close relationship with modern Italian strains indicating continuity of this disease on a regional basis (Kay *et al.*, 2014), however this continuum through time is currently not visible to us. This is due to the diverse challenges of positively identifying the disease using current approaches. Attempts to build up a picture of past human-animal-pathogen relationships must engage with a range of evidence. We propose taking a One Health approach and triangulating evidence and proxies generated through different methods to improve detection of past zoonoses (Figure 1). Such an approach – integrating studies that are typically performed independently – will help maximise understanding for different diseases for which there is differential ability to identify accessible records of their presence. This interdisciplinary review has identified potential for the advancement of methods and integration of datasets. For both human and animal skeletons, researchers should be aware of the potential distribution of brucellosis lesions to support investigations, from which to develop differential diagnoses, and where potential cases are identified, samples should be subject to biomolecular analyses. The anthropological and epidemiological contextualisation of palaeopathological and biomolecular investigations can help to move beyond the description of suspected and evidenced cases of infection in humans and their animals in the distant past, towards the analysis of the factors promoting zoonotic disease emergence. The description and conceptualisation, through epidemiological modelling, of the contexts within which domestic animal populations are structured, managed and humans exposed to these animal populations, can be used to generate hypotheses about zoonotic disease emergence drivers. These hypotheses could then inform the design of osteological and genetic research studies, i.e. the choice of sites, time period and samples, allowing the testing of these hypotheses.

### Supporting information

Appendix S1. Background to the genus *Brucella*

Appendix S2. Suggestions for aDNA studies

Appendix S3. Historical perspectives and potential

Table S1. *Brucella* species and their preferred hosts.

Table S2. Selected common infectious causes of abortion in domestic cattle, goats, sheep and pigs.

Table S3. Primers and FAM labelled probe used in our screening PCR method.

Table S4. Examples of historical approaches and evidence for potential case study examination of brucellosis in English history



## References

- al-Shahed SM, Sharif HS, Haddad MC, Aabed MY, Sammak BM, Mutairi MA. 1994. Imaging features of musculoskeletal brucellosis. *RadioGraphics* **14**: 333-348.
- Arcomano JP, Pizzolato NF, Singer R, Zucker SM. 1977. A unique type of calcification in chronic Brucellosis. *American Journal of Roentgenology* **128**: 135-137.
- Ariza J, Gudiol F, Valverde J, Pallarés R, Fernández-Viladrich P, Rufí G, Espadaler L, Fernández-Nogues. 1985. Brucellar Spondylitis: A detailed analysis based on current findings. *Reviews of Infectious Diseases* **7**(5): 656-664.
- Ariza J, Pujol M, Valverde J, Nolla JM, Rufí G, Viladrich PF, Corredoira JM, Gudiol F. 1993. Brucellar sacroiliitis: findings in 63 episodes and current relevance. *Clinical Infectious Diseases* **16**: 761-765.
- Aufderheide A, Rodríguez- Martín C. 1998. *Human Paleopathology*. Cambridge: Cambridge University Press.
- Baker J, Brothwell DR. 1980. *Animal diseases in archaeology*. London: Academic Press.
- Bendrey R, Cassidy JP, Bokovenko N, Lepetz S, Zaitseva GI. 2011. A possible case of 'poll-evil' in an early Scythian horse skull from Arzhan 1, Tuva Republic, Central Asia. *International Journal of Osteoarchaeology* **21**: 111-118.
- Bendrey R, Taylor GM, Bouwman AS, Cassidy JP. 2008. Suspected bacterial disease in two archaeological horse skeletons from southern England: palaeopathological and biomolecular studies. *Journal of Archaeological Science* **35**: 1581-1590.
- Brennan PJ, Nikaido H. 1995. The envelope of mycobacteria. *Annual Review Biochemistry* **64**: 29-63.
- Buzgan T, Karahocagil MK, Irmak H, Baran AI, Karsen H, Evirgen O, Akdeniz H. 2010. Clinical manifestations and complications in 1028 cases of brucellosis: A retrospective evaluation and review of the literature. *International Journal of Infectious Diseases* **14**: e469-e478.  
doi:10.1016/j.ijid.2009.06.031
- Capasso L. 1999. Brucellosis at Herculaneum (79 AD). *International Journal of Osteoarchaeology* **9**: 277-288.
- Cardoso PG, Macedo GC, Azevedo V Oliveira SC. 2006. Brucella spp. Noncanonical LPS: structure, biosynthesis and interaction with host immune system. *Microbial Cell Factories* **5**: 13.  
doi: 10.1186/1475-2859-5-13.
- Chelli Bouaziz M, Ladeb MF, Chakroun M, Chaabane S. 2008. Spinal brucellosis: A review. *Skeletal Radiology* **37**: 785-790. doi 10.1007/s00256-007-0371-x.
- Christensen T, Martínez-Lavín M, Peneda C. 2013. Periostitis and osteolysis in a Medieval skeleton from South-West Hungary: (Leprosy, treponematosi, tuberculosis or hypertrophic osteoarthropathy) A diagnostic challenge! *International Journal of Osteoarchaeology* **23**(1): 69-82.
- Corbel M. 2006. *Brucellosis in Humans and Animals*. WHO: Geneva.
- Craig LE, Dittmer KE, Thompson KG. 2016. Bones and joints. In, Maxie MG (ed.) *Jubb, Kennedy, and Palmer's Pathology of Domestic Animals*. Sixth edition. Volume 1. Elsevier: St. Louis; 16-163.

Curate F. 2006. Two possible cases of brucellosis from a Clarist monastery in Alácer do Sal, southern Portugal. *International Journal of Osteoarchaeology* **16**: 453–458.

D'Anastasio R, Zipfel B, Moggi-Cecchi J, Stanyon R, Capasso L. 2009. Possible brucellosis in an early hominin skeleton from Sterkfontein, South Africa. *PLoS ONE* **4**: e6439.

D'Anastasio R, Staniscia T, Milia ML, Manzoli L, Capasso L. 2011. Origin, evolution and paleoepidemiology of brucellosis. *Epidemiology and Infection* **139**: 149-156.

Dayan L, Deyev S, Palma L, Rozen N. 2009. Long-standing, neglected sacroiliitis with remarked sacroiliac degenerative changes as a result of *Brucella* spp. infection. *The Spine Journal* **9**: e1-e4. doi: 10.1016/j.spinee.2008.03.011.

de Boer H, Van der Merwe L. 2016. Diagnostic dry bone histology in human paleopathology. *Clinical Anatomy* **29**(7): 831-843.

Díaz Aparicio ED. 2013. Epidemiology of brucellosis in domestic animals caused by *Brucella melitensis*, *Brucella suis* and *Brucella abortus*. *Revue scientifique et technique - Office international des epizooties* **32**: 53-60.

Denny HR. 1973. A review of brucellosis in the horse. *Equine Veterinary Journal* **5**: 121-125.

Donoghue HD. 2008. Molecular palaeopathology of human infectious disease. In: Pinhasi, R and Mays, S, (eds.) *Advances in human palaeopathology*. John Wiley & Sons Ltd: Chichester; 147-176.

Esmailnejad-Ganji SM, Esmailnejad-Ganji SMR. 2019. Osteoarticular manifestations of human brucellosis: A review. *World Journal of Orthopedics* **18**(10): 54-62.

Etxeberria F. 1994. Vertebral epiphysitis: early signs of *Brucellar* disease. *Journal of Paleopathology* **6**: 41–49.

Fournié G, Pfeiffer DU, Bendrey R. 2017. Early animal farming and zoonotic disease dynamics: modelling brucellosis transmission in Neolithic goat populations. *Royal Society Open Science* **4**(2): 160943.

Glasgow MMS. 1976. Brucellosis of the spine. *British Journal of Surgery* **63**: 283-288.

Glynn MK, Lynn TV. 2008. Brucellosis. *Journal of the American Veterinary Medical Association* **233**: 900-908.

Greco E, El-Aguizy O, Ali MF, Foti S, Cunsolo V, Saletti R, Ciliberto E. 2018. Proteomic analyses on an ancient Egyptian cheese and biomolecular evidence of *Brucellosis*. *Analytical Chemistry* **90**: 9673-9676. doi: 10.1021/acs.analchem.8b02535.

Hull NC, Schumaker BA. 2018. Comparisons of brucellosis between human and veterinary medicine. *Infection ecology & epidemiology* **8**(1): 1500846.

Johnson-Walker YJ, Kaneene JB. 2018. Epidemiology: science as a tool to inform One Health policy. In: Herrmann, J.A., Johnson-Walker, Y.J. (Eds.), *Beyond One Health: From Recognition to Results*. Wiley Blackwell; Hoboken, NJ; 3–30.

Jones C. 2019. Brucellosis in an adult female from Fate Bell Rock Shelter, Lower Pecos, Texas (4000–1300 BP). *International Journal of Paleopathology* **24**: 252-264.

Kay GL, Sergeant MJ, Giuffra V, Bandiera P, Milanese M, Bramanti B, Bianucci R, Pallen MJ. 2014. Recovery of a medieval *Brucella melitensis* genome using shotgun metagenomics. *mBio* **5**: (4). e01337-14. doi:10.1128/mBio.01337-14.

Lebov J, Grieger K, Womack D, Zaccaro D, Whitehead N, Kowalczyk B, MacDonald PDM 2017. A framework for One Health research. *One Health* **3**: 44-50.

Lifeso RM, Harder E, McCorkell SJ. 1985. Spinal brucellosis. *The Journal of Bone and Joint Surgery* **16-B**(3)L 345-351.

Lignereux Y, Peters J. 1999. Elements for the retrospective diagnosis of tuberculosis on animal bones from archaeological sites. In, *Tuberculosis Past and Present*, G. Pálfi, O. Dutour, J. Deák, I. Hutás (eds.), Golden Book Publisher Ltd/Tuberculosis Foundation: Budapest; 339-348.

Lindahl T, Nyberg B. 1972. Rate of depurination of native deoxyribonucleic acid. *Biochemistry* **11**: 3610-3618. doi:10.1021/bi00769a018.

Lindahl T. 1993. Instability and decay of the primary structure of DNA. *Nature* **362**: 709-715.

Madkour MM, Sharif HS, Abed MY, Al-Fayez MA. 1988. Osteoarticular brucellosis: Results of bone scintigraphy in 140 patients. *American Journal of Radiology* **150**: 1101-1105.

Magnani DM, Lyons ET, Forde TS, Shekhani MT, Adarichev VA, Splitter GA. 2013. Osteoarticular tissue infection and development of skeletal pathology in murine brucellosis. *Disease Models & Mechanisms* **6**: 811-818. doi: 10.1242/dmm.011056.

Mays SA. 2007. Lysis at the anterior vertebral body margin: Evidence for Brucellar spondylitis? *International Journal of Osteoarchaeology* **17**(2): 107-118. doi: 10.1002/oa.903.

Mays SA, Taylor GM, Legge AJ, Young DB, Turner-Walker G. 2001. A Palaepathological and biomolecular study of tuberculosis in a medieval skeletal collection from England. *American Journal of Physical Anthropology* **114**: 298-311.

McElroy A. 1990. Biocultural models in studies of human health and adaptation. *Medical Anthropology Quarterly* **4**: 243-265.

Mehanic S, Baljic R, Mulabdic V, Huric-Jusufi I, Pinjo F, Topalovic-Cetkovic J, Gadziosmanovic V. 2012. Osteoarticular Manifestations of Brucellosis. *Medicinski Arhiv* **66**(3, suppl 1): 24-26.

Merrett DC. 2002. Is pastoralism a pain in the . . . ? Palaepathology in Early Neolithic Iran. American Association of Physical Anthropologists, Annual Meeting, Buffalo, New York. *American Journal of Physical Anthropology* S34: 112-113.

Merrett DC 2004, Bioarchaeology in Early Neolithic Iran: assessment of health status and subsistence strategy. Unpublished PhD thesis dissertation; Winnipeg, Canada: University of Manitoba.

Moreno E. 2014. Retrospective and prospective perspectives on zoonotic brucellosis. *Frontiers in Microbiology* **5**:213.

Murphy EM, Chistov YK, Hopkins R, Rutland P, Taylor GM 2009. Tuberculosis among Iron Age individuals from Tyva, south Siberia: palaeopathological and biomolecular findings. *Journal of Archaeological Science* **36**: 2029-2038.

Mutolo MJ, Jenny LL, Buszek AR, Fenton TW, Foran DR. 2011. Osteological and molecular identification of *Brucellosis* in ancient Butrint, Albania. *American Journal of Physical Anthropology* **147**: 254-263. doi: 10.1002/ajpa.21643.

Ortner DJ. 2003. *Identification of Pathological Conditions in Human Skeletal Remains. 2<sup>nd</sup> Edition*. Academic Press, New York.

Ouahrani S, Michaux S, Sri Widada J, Bourg G, Tournebize R, Ramuz M, Liautard JP. 1993. Identification and sequence analysis of *IS6501*, an insertion sequence in *Brucella spp*: relationship between genomic structure and the number of *IS6501* copies. *Journal of General Microbiology* **139**: 3265-3273.

Pappas G, Papadimitriou P, Akritidis N, Christou L, Tsianos EV. 2006. The new global map of human brucellosis. *The Lancet Infectious Diseases* **6**: 91-99.

Pisarenko SV, Kovalev DA, Volynkina AS, Ponomarenko DG, Rusanova DV, Zharinova NV, Khachaturova AA, Tokareva LE, Khvoynova IG, Kulichenko AN. 2018. Global evolution and phylogeography of *Brucella melitensis* strains. *BMC Genomics* **19**: 353. <https://doi.org/10.1186/s12864-018-4762-2>

Poester FP, Samartino LE, Santos RL. 2013. Pathogenesis and pathobiology of brucellosis in livestock. *Revue scientifique et technique - Office international des epizooties* **32**: 105-15.

Priest Jr, Low D, Wang C, Bush T. 2008. Brucellosis and sacroiliitis: A common presentation of an uncommon pathogen. *Journal of the American Board of Family Medicine* **21**(2): 158-161. doi: 10.3122/jabfm.2008.02.070270.

Ramadan RO, Hashim NH, Bukhari AAE. 1991. Carpal hygromas in sheep. *World Animal Review* **69**: 64-66.

Rashidi JS, Ortner DJ, Frohlich B, Jonsdottir B. 2001. Brucellosis in Early Bronze Age Jordan and Bahrain: An analysis of possible cases of *Brucella spondylitis*. *American Journal of Physical Anthropology* **S114**: 122-123.

Roberts CA, Buikstra JE. 2019. Ch. 11. Bacterial Infections. In *Ortner's Identification of Pathological Conditions in Human Skeletal Remains*, Third Edition, edited by. Academic Press: London; 321-439.

Schlafer DH, Foster RA. 2016. Female genital system. In: Maxie MG (ed.), Jubb, Kennedy and Palmer's Pathology of Domestic Animals. Sixth edition. Volume 3. St. Louis, MO: Elsevier, 358-464.

Sevilla-López S, Quero Valenzuela F, Piedra Fernandez I. 2011. Bilateral pulmonary nodules due to *Brucellosis*. *Archivos de Bronconeumología* **47**: (6).320-321. doi: 10.1016/j.arbres.2011.02.003. Epub 2011 Apr 5.

Sidstedt M, Jansson L, Nilsson E, Noppa L, Forsman M, Peter Rådström P, Hedman J. 2015. Humic substances cause fluorescence inhibition in real-time polymerase chain reaction. *Analytical Biochemistry* **487**: 30–37.

Sokolov AS, Nedoluzhko AV, Boulygina ES, Tsygankova SV, Sharko FS, Gruzdeva NM, Shishlov AV, Kolpakova AV, Rezepkin AD, Skryabin KG, Prokhortchouk EB. 2016. Six complete mitochondrial genomes from Early Bronze Age humans in the North Caucasus. *Journal of Archaeological Science* **73**:138–144.

Taylor GM. 2014. Ancient DNA (aDNA) and the Fingerprints of Disease: Retrieving Human Pathogen Genomic Sequences from Archaeological Remains Using Real-time Polymerase Chain Reaction (RT-PCR). In, *Molecular Diagnostics: Current Research and Applications*, Huggett JF, O'Grady J (eds.). Caister Academic Press.

Taylor GM, Murphy E, Hopkins R, Rutland P, Chistov Y. 2007. First report of *Mycobacterium bovis* DNA in human remains from the Iron Age. *Microbiology* **153**: 1243–1249.

Turan H, Serefhanoglu K, Karadeli E, Togan T, Arslan H. 2011. Osteoarticular Involvement among 202 brucellosis cases identified in Central Anatolia region of Turkey. *Internal Medicine* **50**: 421-428. doi: 10.2169/internalmedicine.50.4700.

Waldron T. 2009. *Palaeopathology*. Cambridge: Cambridge University Press.

Zeder MA. 2008. Animal domestication in the Zagros: an update and directions for future research. In, *Archaeozoology of the Near East VIII Travaux de la Maison de l'Orient et de la Méditerranée* 49, Vila E, Gourichon L, Choyke AM, Buitenhuis H (eds.), pp. 243–277. Lyon, France: Maison de l'Orient et de la Méditerranée.

## Figure and Table Captions

Figure 1. Understanding and tackling zoonoses: **(A)** the ‘epidemiological triad’ summarising factors influencing infectious disease (characteristics after Johnson-Walker and Kaneene 2018); **(B)** the integrative biocultural model (after McElroy 1990); **(C)** conceptualisation of One Health interventions; **(D)** key recoverable datasets for One Health investigations of past zoonoses.

Table 1. Selected examples of human skeletal evidence interpreted as brucellosis from different contexts.

Table 2. Routes of infection of *Brucella* spp. that result in skeletal lesions in domestic animals

Table 3. Published zooarchaeological remains for which brucellosis is considered in the differential diagnosis or as a possible cause.

Table 4. Archaeological human remains with confirmed biomolecular evidence for *Brucella* species

Figure 1. Understanding and tackling zoonoses: **(A)** the ‘epidemiological triad’ summarising factors influencing infectious disease (characteristics after Johnson-Walker and Kaneene 2018); **(B)** the integrative biocultural model (after McElroy 1990); **(C)** conceptualisation of One Health interventions; **(D)** key recoverable datasets for One Health investigations of past zoonoses.




Table 1. Selected examples of human skeletal evidence interpreted as brucellosis from different contexts.

provenance	discussion	references
Sterkfontein, South Africa dated to 2.8-2.4 MYA	<i>Australopithecus africanus</i> with possible evidence for brucellosis; indicates that zoonotic pathogen transfer is possible without the context of agriculture.	D'Anastasio et al., 2009; D'Anastasio et al., 2011
Early Neolithic Ganj Dareh, Iran, 10,000 CalBP	Tentative identification; peri-domestication context - early herd management and intensive exploitation of goats	Merrett, 2002; Merrett 2004
Bronze Age Bhab-Edh-Dhra, Jordan, 5100-4200 BP	Spinal changes and possible Pedro-Pons sign have been interpreted as evidence of brucellosis; full animal domestication context; development of transhumant pastoralism and secondary products (dairy)	Ortner, 2003; Rashidi et al., 2001
Herculaneum, Italy, dated to the Mt. Vesuvius eruption of 79 AD	Lesions consistent with brucellosis have been observed in 16 of 151 individuals recovered. Diagnosis was based on Pedro-Pons sign of lumbar vertebrae, radiographic evidence of sclerosis below the vertebral body lesions and thickening and increased density of trabeculae. Results are consistent with historic records of the importance of milk and milk products in Roman society and the potential for endemic zoonoses in the past.	Capasso, 1999
Fate Bell Rock Shelter, Texas, dating 4,000-1300 BP	Based on bone macroscopic morphology and CT imaging; hunting/gathering context again indicates that zoonotic pathogen transfer is possible without the context of agriculture	Jones, 2019
Medieval Butrint, Albania, 10 <sup>th</sup> to 13 <sup>th</sup> centuries AD	Both osteological and molecular methods were applied to skeletal remains with macroscopic possible brucellar lesions (see Table 4). In this case the diagnosis is definitive; <i>Brucella</i> spp. aDNA was recovered from the lesions.	Mutolo et al., 2012



Table 2. Routes of infection of *Brucella* spp. that result in skeletal lesions in domestic animals

	Osteological response	Reference
Bacteraemia with haematogenous seeding of bone	Direct localisation in vertebra triggering inflammation and bone modelling/periosteal new bone formation	Schlafer and Foster, 2016
Bacteraemia with haematogenous seeding of articular joints, bursae and tendon sheaths	Inflammation at these sites extends locally to impact adjacent bone causing periosteal new bone formation at specific anatomical sites, e.g. inflammation of supra-atlantal bursa (poll evil) causing osseous lesions on adjacent occipital bone of equine cranium	Craig et al., 2016; Denny, 1973

Table 3. Published zooarchaeological remains for which brucellosis is considered in the differential diagnosis or as a possible cause.

Provenance	Species / element	Brief description and diagnosis	Reference
Early Iron Age Arzhan 1, Tyva	Horse skull	Occipital bone lesions interpreted as foci of inflammation and necrosis following local infection. It is suggested that the pathology represents a case of 'poll-evil', most likely due to a bacterial infection such as <i>Brucella abortus</i> , <i>Actinomyces bovis</i> , or <i>Streptococcus zooepidemicus</i>	Bendrey et al., 2011
Late Iron Age/Early Roman, Viables Farm, UK	Horse skeleton	Proliferative periosteal lesions on the atlas, two thoracic vertebrae (one with lytic damage), the sacrum, four rib fragments, and right pelvis suggestive of systemic infection, most likely due to <i>Trueperella pyogenes</i> , <i>Mycobacterium bovis</i> , <i>Brucella abortus</i> or <i>Aspergillus</i> spp.	Bendrey et al., 2008
Late Iron Age/Early Roman, Downlands Farm, UK	Horse skeleton	Proliferative periosteal lesions on six thoracic and one lumbar vertebrae, and eight rib fragments suggestive of systemic infection, most likely due to <i>Trueperella pyogenes</i> , <i>Mycobacterium bovis</i> , <i>Brucella abortus</i> or <i>Aspergillus</i> spp.	Bendrey et al., 2008
Dragonby, UK	Horse mid-cervical vertebra	'lesion closely resembling modern brucella osteomyelitis'. No further detail.	Baker and Brothwell, 1980, 76

Table 4. Archaeological human remains with confirmed biomolecular evidence for *Brucella* species

Provenance	Age and sex	pathology	reference
Bronze Age Novosvobodnaya (Republic of Adygea, Russia); Kurgan (burial mound) 25 grave 1	Not described	Not described	Sokolov et al., 2016
Iron Age Aymyrlyg, Tyva, south Siberia	25-35 year old female (SkXXXI.34)	Lytic lesions in eight vertebrae (C7, T6-9, L3-L5)	Murphy et al., 2009; Taylor et al., 2007
medieval Butrint, in Southwest Albania, burial 2272 (10-12 <sup>th</sup> centuries AD)	young male individual aged between 17-21 years old	Skeletal lesions included, amongst others, cavitating lytic foci in the thoracic vertebrae (T3-T12) and lumbar vertebrae (L1, L2 and L4), sacrum was affected, as were some ribs, which showed cortical thickening and trabecular expansion with porosity on the parietal surface of some fragments	Mutolo et al., 2011
medieval Butrint, in Southwest Albania, burial 4015 (12 <sup>th</sup> -13 <sup>th</sup> centuries AD)	young male individual aged between 17-21 years old	Skeletal lesions included, amongst others, cavitating lytic lesions in vertebrae (T3-T12, L1, L2), sacrum was affected, as were some ribs, which showed some cortical and porosity on the parietal surfaces.	Mutolo et al., 2011
Medieval Geridu, northwestern Sardinia; second half of the 14 <sup>th</sup> century	50-60 year old male individual (Sk2568)	Diffuse idiopathic skeletal hyperostosis (DISH) – fusions between thoracic vertebrae (T4-T10), and L5 and sacrum; also extraspinal enthesopathies. Thirty-two calcified nodules found in the pelvic girdle.	Kay et al., 2014

## Appendix S1. Background to the genus *Brucella*

Bacteria of the genus *Brucella* are named after Sir David Bruce (1855-1931). In 1886 the then Captain Bruce, Scottish physician, assisted by his microbiologist wife Mary, first isolated and cultured the organism responsible for Malta fever whilst posted to the island. At that time, both local inhabitants and soldiers in the British garrison were experiencing an outbreak of fever of unknown origin. Bruce showed that the organism, later named *Micrococcus melitensis*, obeyed the rules laid down by Robert Koch for causation and transmission of infectious diseases (Koch's postulates). Later in 1905, "Temi" Zammit, Maltese archaeologist, medic and member of the Mediterranean Fever Commission, established that infected goats were a reservoir of disease and were transmitting the fever to humans in their milk. Since then, a number of closely related species have been identified and now comprise the genus *Brucella*. These are: *Brucella abortus*, isolated from an infected cow's udder in 1897; *B. suis* identified in pigs (1914), *B. ovis*, a cause of ovine epididymitis (1956), *B. canis* from dogs (1966) and *B. neotomae* from desert rats in 1957.

*Brucella* is one of eight genera in the Family *Brucellaceae* which belong to the alpha-2 proteobacteria, a class of bacteria in the ancient and diverse phylum of proteobacteria. *Brucella* species are all Gram-negative, facultative, non-motile coccoid or rod-shaped aerobic bacteria which replicate within phagocytic cells of the host reticuloendothelial system. Adaptation to the intracellular environment of diverse hosts has been associated with genome downsizing, as genes no longer required for life in the environment are lost. Current evidence suggests that *Brucella* (typical genome size 3.3Mb) originally evolved from free-living *Ochrobactrum* (4.77 Mb) a related genus of the *Brucellaceae* (Wattam et al., 2014).

Species that are generally pathogenic to humans are *B. melitensis*, *B. abortus* and biovars 1 and 3 of *B. suis*. These are species associated with some of the earliest domesticated animals and it seems probable that past human activities such as farming and animal domestication will have influenced the selection and clonal expansion of individual *Brucella* species and their opportunities to cause human disease. *B. canis* infection of humans has been reported, but is relatively rare. This species appears to have diverged from an ancestral strain of *B. suis* in the last 22,500 years (Foster et al., 2009).

In the last decade or so a number of novel species have been described, such as *B. ceti* affecting whales, *B. pinnipedialis* from marine mammals (dolphins, porpoises and seals), *B. microti* from the common vole, *B. vulpis* from red foxes, *B. papionis* from baboons (Whatmore et al., 2014) and *B. inopinata* isolated from an infected human breast implant (Scholz et al., 2010). The preferred host range of *B. inopinata* is as yet unknown but the species appears to be one of the most diverse lineages of the genus and other novel species closely related to *B. inopinata*, continue to be described from amphibians (Eisenberger et al., 2012) and Australian wild rodents (Tiller et al., 2010). Table S1 summarises the classical and more recently recognised and emerging *Brucella* species and their preferred hosts.

The six classical members of the genus were primarily considered to be separate species but were later seen as biovars of a single species (*B. melitensis*) (Verger et al., 1985). They are best viewed as a complex of host-adapted species causing disease in their preferred hosts. However, host specificity is not absolute and other animal species may become infected when in prolonged close contact. For example, both *B. abortus* and *B. melitensis* have been reported in camels kept in association with sheep and goats (Cooper et al., 1991). Further, a novel species closest to *B. microti* (DNA > 99.8% homologous) has been reported in domestic marsh frog (Jaý et al., 2018).

The *Brucella* genomes of a number of the species and biovars have been sequenced and annotated,

for example *B. melitensis* (DelVecchio *et al.*, 2002), *B. suis* (Paulsen *et al.*, 2002), *B. abortus* (Halling *et al.*, 2005) and *B. pinnipedialis* isolate B2/94 (Audic *et al.*, 2011). All species are highly similar at the nucleotide level. The genome is composed of two circular chromosomes, containing a total of approximately 3.3Mb with 2,124 Mb on chromosome 1 and 1,162 Mb on chromosome 2. The G + C content is 57.2%. A total of approximately 3,200 open reading frames (ORFs) have been predicted on both chromosomes. The Brucella Bioinformatics Portal (<http://www.phidias.us/bbp/>) is a gateway to a variety of genomic resources, bioinformatics tools and literature for researchers in the field.

## References

Audic S, Lescot M, Claverie J-M, Cloeckert A, Zygmunt MS. 2011. The genome sequence of *Brucella pinnipedialis* B2/94 sheds light on the evolutionary history of the genus *Brucella*. *BMC Evolutionary Biology* **11**: 200.

Cooper CW. 1991. The epidemiology of human *Brucellosis* in a well defined urban population in Saudi Arabia. *Journal of Tropical Medicine and Hygiene* **94**: 416–422.

DelVecchio VG, Kapatral V, Redkar RJ, Patra G, Mujer C, Los T, Ivanova N, Anderson I, Bhattacharyya A, Lykidis A, Reznik G, Jablonski L, Larsen N, D'Souza M, Bernal A, Mazur M, Goltsman E, Selkov E, Elzer P, Hagius S, O'Callaghan D, Letesson JJ, Haselkorn R, Kyrpides N, Overbeek R. 2002. The genome sequence of the facultative intracellular pathogen *Brucella melitensis*. *Proceedings of the National Academy of Science USA* **99**: 443–448.

Eisenberg T, Hamann HP, Kaim U, Schlez K, Seeger H, Schauerte N, Melzer F, Tomaso H, Scholz HC, Koylass MS, Whatmore AM, Zschöck M. 2012. Isolation of potentially novel *Brucella* spp. from frogs. *Applied and Environmental Microbiology* **78**: 3753-3755. Doi:10.1128/AEM.07509-11.

Foster JT, Beckstrom-Sternberg SM, Pearson T, Beckstrom-Sternberg JS, Chain PS, Roberto FF, Hnath J, Brettin T, Keim P. 2009. Whole genome based phylogeny of the genus *Brucella*. *Journal of Bacteriology* **191**(8): 2864-2870. doi: 10.1128/JB.01581-08.

Halling SM, Peterson-Burch BD, Bricker BJ, Zuerner RL, Qing Z, Li LL, *et al.* 2005. Completion of the genome sequence of *Brucella abortus* and comparison to the highly similar genomes of *Brucella melitensis* and *Brucella suis*. *Journal of Bacteriology* **187**, 2715–2726. doi: 10.1128/JB.187.8.2715-2726.2005.

Hammerl JA, Ulrich RG, Imholt C, Scholz HC, Jacob J, Kratzmann N, Nöckler K, Al Dahouk S. 2015. Molecular survey on *Brucellosis* in rodents and shrews – natural reservoirs of novel *Brucella* species in Germany? *Transboundary and Emerging Diseases* **64**: 663-671.

Jaÿ M, Girault G, Perrot L, Taunay B, Vuilmet T, Rossignol F, Pitel P-H, Picard E, Ponsart C, Mick V. 2018. Phenotypic and Molecular Characterization of *Brucella microti*-Like Bacteria From a Domestic Marsh Frog (*Pelophylax ridibundus*). *Frontiers in Veterinary Science* **5**: 283. doi: 10.3389/fvets.2018.00283.

Paulsen IT, Seshadri R, Nelson KE, Eisen JA, Heidelberg JF, Read TD, Dodson RJ, Umayam L, Brinkac LM, Beanan MJ, Daugherty SC, Deboy RT, Durkin AS, Kolonay JF, Madupu R, Nelson WC, Ayodeji B, Kraul M, Shetty J, Malek J, Van Aken SE, Riedmuller S, Tettelin H, Gill SR, White O, Salzberg SL, Hoover DL, Lindler LE, Halling SM, Boyle SM, Fraser CM. 2002. The *Brucella suis* genome reveals

fundamental similarities between animal and plant pathogens and symbionts. *Proceedings of the National Academy of Science USA*. **99**: 13148–13153.

Scholz HC, Nöckler K, Göllner C, Bahn P, Vergnaud G, Tomaso H, *et al.* 2010. *Brucella inopinata* sp. nov., isolated from a breast implant infection. *International Journal of Systematic and Evolutionary Microbiology* **60**: 801-808. doi: 10.1099/ijs.0.

Tiller RV, Gee JE, Frace MA, Taylor TK, Setubal JC, Hoffmaster AR, De Barun K. 2010. Characterization of novel *Brucella* strains originating from wild native rodent species in North Queensland, Australia. *Applied and Environmental Microbiology* **76**: 5837-5845.

Verger J-M, Grimont F, Grimont PAD, Grayon M. 1985. *Brucella*, a monospecific genus as shown by deoxyribonucleic acid hybridization. *International Journal of Systematic Bacteriology* **35**: 292–295.

Wattam AR, Foster JT, Mane SP, Beckstrom-Sternberg SM, Beckstrom-Sternberg JM, Dickerman AW, *et al.* 2014. Comparative phylogenomics and evolution of the Brucellae reveal a path to virulence. *Journal of Bacteriology* **196**: 920–930. doi: 10.1128/JB.01091-13.

Whatmore AM, Davison N, Cloeckert A, Al Dahouk S, Zygmunt MS, Brew SD, *et al.* 2014. *Brucella papionis* sp. nov., isolated from baboons (*Papio spp.*). *International Journal of Systematic and Evolutionary Microbiology* **64**: 4120–4128. doi: 10.1099/ijs.0.065482-0.

Table S1. *Brucella* species and their preferred hosts.

<i>Brucella</i> species	Preferred host	Strain	IS711 copies	Reference
<i>B. melitensis</i>	Goats/sheep	Geridu-1 16M	8 8	Kay <i>et al</i> , 2014. Suárez-Esquivel <i>et al</i> , 2017.
<i>B. abortus</i>	Cattle	S19 2308	6 6	Primer BLAST <sup>1</sup> search. Suárez-Esquivel <i>et al</i> , 2017.
<i>B. suis</i>	Pigs (most diverse in host range)	QH05 1330	7 7	BLAST search. BLAST search.
<i>B. canis</i>	Dogs	GB1 ATCC23365	7 7	BLAST search. Suárez-Esquivel <i>et al</i> , 2017.
<i>B. ovis</i>	Sheep	ATC25840 ATC25840	33 38	Suárez-Esquivel <i>et al</i> , 2017. BLAST search.
<i>B. microti</i>	Common vole	CCM4915 <sup>T</sup> BCM4915 CCM4915 <sup>T</sup> “	5-6 14 13 13	Scholz <i>et al</i> , 2008. Suárez-Esquivel <i>et al</i> , 2017. Audic <i>et al</i> , 2009. BLAST search.
<i>B. neotomae</i>	Desert woodrats		?	Stoenner <i>et al</i> , 1957.
<i>B. inopinata</i>	Uncertain	BO-1	4-5	Scholz <i>et al</i> , 2010.
<i>B. pinnipedii</i>	Seals	B2/94 6/566	45-46 31 29	Suárez-Esquivel <i>et al</i> , 2017. Audic <i>et al</i> , 2011. BLAST search.
<i>B. ceti</i>	Dolphins, Porpoises, Whales	Various, TE28753	42-53 14	Suárez-Esquivel <i>et al</i> , 2017. BLAST search.
<i>B. papionis</i>	Baboon	F80/08-60 <sup>T</sup> F80/08-61	High, uncertain	Whatmore <i>et al</i> , 2014.
<i>B. vulpis</i>	Red Foxes  Bank vole	F60 F965	27 27 ?	Hofer <i>et al</i> , 2012. Hofer <i>et al</i> , 2012. Hammerl <i>et al</i> , 2015.
1. Novel motile species, closest to <i>B. inopinata</i>	Amphibians	09RB8471 10RB9215 UK 8/14 Various	Uncertain	Eisenberg <i>et al</i> , 2012. Eisenberg <i>et al</i> , 2012. Whatmore <i>et al</i> , 2015. Scholz <i>et al</i> , 2016.
2. Microti-like	Domestic Marsh frog	17-2122-4144	Uncertain	Jaý <i>et al</i> , 2018.

BLAST search available at: [https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE\\_TYPE=BlastSearch](https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE_TYPE=BlastSearch).

## References

- Audic S, Lescot, Claverie J-M, Scholz HC. 2009. *Brucella microti*: the genome sequence of an emerging pathogen. *BMC Genomics* **10**: 352. doi:10.1186/1471-2164-10-352.
- Audic S, Lescot M, Claverie J-M, Cloeckart A, Zygmunt MS. 2011. The genome sequence of *Brucella pinnipedialis* B2/94 sheds light on the evolutionary history of the genus *Brucella*. *BMC Evolutionary Biology* **11**: 200.
- Eisenberg T, Hamann HP, Kaim U, Schlez K, Seeger H, Schauerte N, Melzer F, Tomaso H, Scholz HC, Koylass MS, Whatmore AM, Zschöck M. 2012. Isolation of potentially novel *Brucella* spp. from frogs. *Applied and Environmental Microbiology* **78**: 3753-3755. Doi:10.1128/AEM.07509-11.
- Hammerl JA, Ulrich RG, Imholt C, Scholz HC, Jacob J, Kratzmann N, Nöckler K, Al Dahouk S. 2015. Molecular survey on *Brucellosis* in rodents and shrews – natural reservoirs of novel *Brucella* species in Germany? *Transboundary and Emerging Diseases* **64**: 663-671.
- Hofer E, Revilla-Fernández S, Al Dahouk S, Riehm JM, Nöckler K, Zygmunt MS, Cloeckart A, Tomaso H, Scholz HC. 2012. A potential novel *Brucella* species isolated from mandibular lymph nodes of red foxes in Austria. *Veterinary Microbiology* **155**: 93-99.
- Jaý M, Girault G, Perrot L, Taunay B, Vuilmet T, Rossignol F, Pitel P-H, Picard E, Ponsart C, Mick V. 2018. Phenotypic and Molecular Characterization of *Brucella microti*-Like Bacteria from a Domestic Marsh Frog (*Pelophylax ridibundus*). *Frontiers in Veterinary Science* **5**: 283. doi: 10.3389/fvets.2018.00283.
- Kay GL, Sergeant MJ, Giuffra V, Bandiera P, Milanese M, Bramanti B, Bianucci R, Pallen MJ. 2014. Recovery of a medieval *Brucella melitensis* genome using shotgun metagenomics. *mBio* **5**: (4). e01337-14. doi:10.1128/mBio.01337-14.
- Scholz HC, Hubálek Z, Sedláček I, Vergnaud G, Tomaso H, Al Dahouk S, Melzer F, Kämpfer P, Neubauer H, Cloeckart A, Maquart M, Zygmunt MS, Whatmore AM, Falsen E, Bahn P, Göllner C, Pfeffer M, Huber B, Busse HJ, Nöckler, K. 2008. *Brucella microti* sp. nov., isolated from the common vole *Microtus arvalis*. *International Journal of Systematic and Evolutionary Microbiology* **58**: 375–382.
- Scholz HC, Nöckler K, Göllner C, Bahn P, Vergnaud G, Tomaso H, et al. 2010. *Brucella inopinata* sp. nov., isolated from a breast implant infection. *International Journal of Systematic and Evolutionary Microbiology* **60**: 801-808. doi: 10.1099/ijs.0.
- Scholz HC, Mühldorfer K, Shilton C, Benedict S, Whatmore AM, Blom J, et al. 2016. The Change of a Medically Important Genus: Worldwide Occurrence of Genetically Diverse Novel *Brucella* Species in Exotic Frogs. *PLoS ONE* **11** (12): e0168872. doi:10.1371/journal.pone.0168872.
- Stoenner HG, Lackman DB. 1957. A new species of *Brucella* isolated from the desert wood rat, *Neotoma lepida* Thomas. *American Journal of Veterinary Research* **18**: 947–951.
- Suárez-Esquivel M, Baker KS, Ruiz-Villalobos N, Hernández-Mora G, Barquero-Calvo E, González-Barrientos R, Castillo-Zeledón A, Jiménez-Rojas C, Chacón-Díaz C, Cloeckart A, Chaves-Olarte E, Thomson NR, Moreno E, Guzmán-Verri C. 2017. *Brucella* genetic variability in wildlife marine mammals populations relates to host Preference and Ocean Distribution. *Genome Biology and*



*Evolution* **9**: 1901–1912 doi:10.1093/gbe/evx137.

Whatmore AM, Davison N, Cloeckert A, Al Dahouk S, Zygmunt MS, Brew SD, et al. 2014. *Brucella papionis* sp. nov., isolated from baboons (*Papio spp.*). *International Journal of Systematic and Evolutionary Microbiology* **64**: 4120–4128. doi: 10.1099/ijs.0.065482-0.

Whatmore AM, Dale E-J, Stubberfield E, Muchowski J, Koylass M, Dawson C, Gopaul KK, Perrett LL, Jones M, Lawrie A. 2015. Isolation of *Brucella* from a White's tree frog (*Litoria caerulea*). *JMM Case Reports* 2: doi: 10.1099/jmmcr.0.000017.

## Appendix S2. Suggestions for aDNA studies

Multi-copy loci within pathogen genomes make ideal initial targets for molecular screening methods, as demonstrated (Mutolo et al., 2011). The insertion element *IS711* (Ouahrani et al., 1993) is an ideal target for detection of all *Brucella* species. Other IS are present within the genus, but not in such high copy number. *IS711* is an 842 bp sequence bounded by 20 bp inverted repeats. It appears to be unique to the *Brucella* genus and is not present in the closely related family of alpha-proteobacteriaceae. In the past, movements and transpositions of this element may well have driven genetic variation between *Brucella* species and biovars. Transposition has been described in two species, *B. ovis* and *B. pinnipedialis* using transposon mutagenesis with the plasmid pGBG1 (Ocampo-Sosa and Garcia-Lobo, 2008). This raises the possibility that *IS711* may still initiate variation in some high-copy strains like those present in marine mammals.

The element does share approximately 53% homology with *IS427* found in *Rhizobium radiobacter*, a rod-shaped Gram-negative bacterium belonging to the *Rhizobacteriaceae*, suggesting a possible common distant ancestry. There is also some minor variation in sequence between individual copies of *IS711*. However, it is straightforward to design primers to conserved regions of the transposase to develop specific PCR methods suitable for aDNA applications.

Table S3 details the primers and FAM labelled probe used in our screening PCR method. The four primers allow for various combinations of single round, nested, or hemi-nested variants of the method to be used as either a conventional or real-time method with a single dual-labelled probe. In general however, the use of a DNA binding fluorescent dye like either SYBR Green 1 or (the less PCR Inhibitory) EVAGreen™ are cost effective and more sensitive, as many fluorescent dye molecules intercalate into each amplicon. The minimum detection limit of this method has been found to be 3.6 fg of *Brucella* DNA, or less than 1 genome equivalent. In contrast, the use of a labelled oligonucleotide probe is more specific and reports only the product rather than primer-dimer, but is less sensitive. The recommendation for future studies would be to use primer combinations amplifying shorter templates e.g. F2/R3 (85bp) for initial screening combined with EVAGreen™, confirmed by probe-based reporters as suggested (Table 6.1; Hinić et al., 2008).

The copy number of *IS711* (where known) in the classical and more recently described species, along with key references are shown in Table S1. The higher copy number of *IS711* in *B. ovis* and the marine isolates, means that these species might be easier to detect under some circumstances. Although isolation of a marine species seems unlikely in human remains, infection with *B. pinnipedialis* has been reported in a few instances in humans in the medical literature, showing these species do have some zoonotic potential (Whatmore et al., 2008; McDonald et al., 2006; Sohn et al., 2003). It is also worth noting that the seal-adapted strain of tuberculosis, *Mycobacterium pinnipedii* was recovered from pre-Colombian human remains in Peru with evidence of skeletal tuberculosis (Bos et al., 2014).

Follow-up PCR tests should be undertaken to confirm the presence of the pathogen. Genetic loci which are conserved across all genus members are useful for this purpose. The region coding for membrane protein *Bcsp31*, has been used by several authors (Mutolo et al., 2011; Mukherjee et al., 2007). Other candidate regions include the *recA* gene, coding for a protein involved in DNA repair and the *per* gene (Bogdanovich et al., 2004; Bounaadja et al., 2009). In the six classical species, *recA* is highly conserved but variation has been reported in more recently described members (Scholz et al., 2010; Scholz et al., 2016). These are useful confirmatory tests for the pathogen and indicate if other, single copy targets for genotyping might be successful or if applying a WGS protocol might be worthwhile.

The fragmented nature of aDNA precludes the use of PCR methods which rely on longer sequences for typing and species identification, like the multiple loci variable nucleotide tandem repeat analysis (MLVA-VNTR type approaches (Le Flèche et al., 2006; Whatmore et al., 2007). Identifying species and biovars can be found by amplification and sequencing of polymorphic genes (Cloekaert et al., 2001; Ficht et al., 1996). Moreover, with information from phylogenetic and WGS studies, there now exists more opportunities for developing specific methods based on species and lineage-specific disruptions such as insertions, deletions and SNPs. Deletions provide the best opportunities for designing specific PCR methods with primers spanning the deletion to generate a short amplicon in the region of about 60-120 bp. For added specificity, a dual-labelled probe can be used to hybridize across the deletion breakpoint and report product formation. If the deletion has not occurred, a PCR can be devised to target the sequence and report product with a differently labelled probe. For strict comparison, the amplicon size should match that of the deletion spanning PCR. Hinić and colleagues (2008) describe 7 separate PCRs which target species-specific *loci* or exploit deletions to differentiate the six classical species, *B. melitensis*, *B. abortus*, *B. suis*, *B. ovis*, *B. canis* and *B. neotomae* (Hinić et al., 2008). The amplicon sizes are all small (63-116 bp) and appropriate for aDNA uses. Kaden and colleagues (2017) have developed a real-time PCR for specific detection of *B. melitensis* isolates, the most important human pathogen of the genus. This exploits a very short deletion of 2bp in the acetyl Co-A acetyl transferase gene.

Other approaches for genotyping include exploiting phylogenetically informative SNPs with conventional PCR and sequencing (Scott et al., 2007; Foster et al., 2012). Methods for bioarchaeological studies need to be optimised for constraints imposed by aDNA fragmentation and evaluated for specificity and sensitivity in the complex milieu, which is co-extracted from archaeological contexts. Even with the advent of high-throughput sequencing techniques and the enormous amount of information which can be obtained from isolates, there remains a place for conventional PCR and sequencing to confirm *loci* with poor coverage.

Negative results can sometimes be the result of PCR inhibitory substances co-extracted along with aDNA. An internal amplification control (IAC) assay can be run up to assess if inhibition is present. This can be achieved by addition of a limited amount of synthetic DNA template with identical binding sites for the primers but with a differing internal sequence. This target is reported with different probe chemistry and generation of amplicons shows inhibition is not problematical (Hoorfar et al., 2004). An alternate approach is to use a universal method for assessing inhibition such as the SPUD assay (Nolan et al., 2006). Primers are used to amplify a unique sequence from the potato genome and the effect of aDNA extracts assessed on amplification profiles (Cq) compared to addition of water (non-template controls).

## References

Bogdanovic T, Skumic M, Lübeck PS, Ahrens P, Hoorfar J. 2004. Validated 5' nuclease PCR assay for rapid identification of the genus *Brucella*. *Journal of Clinical Microbiology* **42**: 2261-2263.

Bos K, Harkins KM, Herbig A, Coscolla M, Weber N, Comas I, Forrest SA, Bryant JM, Harris SR, Schuenemann VJ, Campbell TJ, Majander K, Wilbur AK, Guichon RA, et al. 2014. Pre-columbian mycobacterial genomes reveals seals as a source of New World tuberculosis. *Nature* **514**: 494-497. [doi:10.1038/nature13591](https://doi.org/10.1038/nature13591).

Bounaadja L, Albert D, Chénais B, Hénault S, Zygmunt MS, Poliak S, Garin-Bastuji B. 2009. Real-time PCR for identification of *Brucella* spp.: A comparative study of *IS711*, *bcs31* and *per* target genes. *Veterinary Microbiology* **137**: 156-164.

- Cloekaert A, Verger J-M, Grayon M, Paquet J-Y, Garin-Bastuji B, Foster G, Godfroid J. 2001. Classification of *Brucella* spp. isolated from marine mammals by DNA polymorphism at the omp2 locus. *Microbes and Infection* **3**: 729-738.
- Ficht TA, Hussein HS, Derr J, Bearden SW. 1996. Species-specific sequences at the omp2 locus of *Brucella* type strains. *International Journal of Systematic Bacteriology* **46**: 329-331.
- Foster JT, Price LB, Beckstrom-Sternberg SM, Pearson T, Brown WD, Kiesling DM, Allen CA, Liu CM, Beckstrom-Sternberg J, Frank F Roberto FF, Keim P. 2012. Genotyping of *Brucella* species using clade specific SNPs. *BMC Microbiology* **12**:110
- Hinić V, Brodard I, Thomann A, Cvetnić Z, Makaya PV, Frey J, Abril C. 2008. Novel identification and differentiation of *Brucella melitensis*, *B. abortus*, *B. suis*, *B. ovis*, *B. canis*, and *B. neotomae* suitable for both conventional and real-time PCR systems. *Journal of Microbiological Methods* **75**: 375–378.
- Hoorfar J, Malorny B, Abdulmawjood A, Cook N, Wagner M, Fach P. 2004. Practical considerations in design of internal amplification controls for diagnostic PCR assays. *Journal of Clinical Microbiology* **42**: 1863–1868.
- Kaden R, Ferrari S, Alm E, Wahab T. 2017. A novel real-time PCR assay for specific detection of *Brucella melitensis*. *BMC Infectious Diseases* **17**:230.
- Le Flèche P, Jacques I, Grayon M, Al Dahouk S, Bouchon P, Denoëud F, Nöckler K, Neubauer H, Guilloteau LA, Vergnaud G. 2006. Evaluation and selection of tandem repeat loci for a *Brucella* MLVA typing assay. *BMC Microbiology* **6**: 9 doi:10.1186/1471-2180-6-9
- McDonald WL, Jamaludin R, Mackereth G, Hansen M, Humphrey S, Short P, *et al.* 2006. Characterization of a *Brucella* sp. strain as a marine-mammal type despite isolation from a patient with spinal osteomyelitis in New Zealand. *Journal of Clinical Microbiology* **44**: 4363–4370.
- Mukherjee F, Jain J, Patel V, Nair M. 2007. Multiple genus-specific markers in PCR assays improve the specificity and sensitivity of diagnosis of brucellosis in field animals. *Journal of Medical Microbiology* **56**:1309-16.
- Mutolo MJ, Jenny LL, Buszek AR, Fenton TW, Foran, DR 2011. Osteological and molecular identification of *Brucellosis* in ancient Butrint, Albania. *American Journal of Physical Anthropology* **147**: 254-263. doi: 10.1002/ajpa.21643
- Nolan T, Hands RE, Ogunkolade W, Bustin SA. 2006. SPUD. A quantitative PCR assay for the detection of inhibitors in nucleic acid preparations. *Analytical Biochemistry* **351**: 308-310.
- Ocampo-Sosa AA, Garcia-Lobo JM. 2008. Demonstration of *IS711* transposition in *Brucella ovis* and *Brucella pinnipedialis*. *BMC Microbiology* **8**:17 doi:10.1186/1471-2180-8-17.
- Ouahrani S, Michaux S, Sri Widada J, Bourg G, Tournebize R, Ramuz M, Liautard JP, 1993. Identification and sequence analysis of *IS6501*, an insertion sequence in *Brucella* spp: relationship between genomic structure and the number of *IS6501* copies. *Journal of General Microbiology* **139**: 3265-3273.
- Scholz HC, Nöckler K, Göllner C, Bahn P, Vergnaud G, Tomaso H, *et al.* 2010. *Brucella inopinata* sp.

*nov.*, isolated from a breast implant infection. *International Journal of Systematic and Evolutionary Microbiology* **60**: 801-808. doi: 10.1099/ijs.0.

Scholz HC, Mühldorfer K, Shilton C, Benedict S, Whatmore AM, Blom J, et al. 2016. The Change of a Medically Important Genus: Worldwide Occurrence of Genetically Diverse Novel *Brucella* Species in Exotic Frogs. *PLoS ONE* **11** (12): e0168872. doi:10.1371/journal.pone.0168872.

Scott JC, Koylass MS, Stubberfield MR, Whatmore AM. 2007. Multiplex assay based on single-nucleotide polymorphisms for rapid identification of *Brucella* Isolates at the species level. *Applied and Environmental Microbiology* **73**: 7331–7337. Doi: 10.1128/AEM.00976-07.

Sohn AH, Probert WS, Glaser CA, Gupta N, Bollen AW, Wong JD, Grace EM, McDonald WC. 2003. Human neurobrucellosis with intracerebral granuloma caused by a marine mammal *Brucella* spp. *Emerging Infectious Diseases* **9**: 485-488.

Whatmore AM, Perrett LL, MacMillan AP. 2007. Characterisation of the genetic diversity of *Brucella* by multilocus sequencing. *BMC Microbiology* **7**:34 doi:10.1186/1471-2180-7-34.

Whatmore AM, Dawson C, Groussaud P, Koylass MS, King A, Shankster SJ, et al. 2008. Marine mammal *Brucella* genotype associated with zoonotic infection. *Emerging Infectious Diseases* **14**: 517-518. <https://dx.doi.org/10.3201/eid1403.070829>.

### **Appendix S3. Historical perspectives and potential**

The study of history has evolved considerably across the centuries, and the energies of successive generations of historians have been channelled into an increasingly diverse array of sub-disciplinary studies in response to changes in historiographical emphasis. In general terms, these disciplinary shifts have taken historians from a traditional focus upon great men and great deeds, to a much wider exploration of economic, social, and cultural history. In recent decades this had broadened out much further, with historical studies of particular social groups or themes. As the foci of historical studies have shifted, so too have the methodologies employed by historians in order to examine a much broader range of documentary and other evidence, the value of which might have been overlooked in the context of previous historiographical debates. It is for this reason that source materials and previous historical studies should be returned to in the context of new interdisciplinary questions, to interrogate whether the historical record might yet have more to yield in relation to previously unposed questions. In particular, it is hoped that drawing together materials from across previously separated fields of study and temporal periods of investigation may contribute different perspectives to our understanding of this disease in the historical past and to its potential visibility in the historical record.

Each modern day country and region has its own historiographical journey, shaped in part by the trends observed in the broader studies in history across the international stage, but also in part by the documentation and evidence that has survived within each geopolitical region as it has evolved. It is beyond the scope of this appendix to provide a detailed examination of all potential historical sources for the study of brucellosis in the past. However, Table S4 below points to the directions in which the sub-disciplines of history might be explored in the context of English history, as an example, to contribute to the interdisciplinary investigation of brucellosis in the past (Figure 1). This includes some examples of the types of sources that may be utilised from across different periods of English history. Historians are exceptionally lucky in the rich and fertile sources that have been preserved for the study of English history, and also in the phenomenal steps that have already been taken to bring elements of such evidence together in the form of authoritative volumes on specific topics. These seminal works are logical starting points for any such study, pointing the way to other sources and evidence of relevance, and again examples of such works are given in the table below. Both the primary and secondary evidence cited below serve as illustrative examples only, and the exclusion of other materials should not be taken as an oversight.

Table S1. *Brucella* species and their preferred hosts.

<i>Brucella</i> species	Preferred host	Strain	IS711 copies	Reference
<i>B. melitensis</i>	Goats/sheep	Geridu-1 16M	8 8	Kay <i>et al</i> , 2014. Suárez-Esquivel <i>et al</i> , 2017.
<i>B. abortus</i>	Cattle	S19 2308	6 6	Primer BLAST <sup>1</sup> search. Suárez-Esquivel <i>et al</i> , 2017.
<i>B. suis</i>	Pigs (most diverse in host range)	QH05 1330	7 7	BLAST search. BLAST search.
<i>B. canis</i>	Dogs	GB1 ATCC23365	7 7	BLAST search. Suárez-Esquivel <i>et al</i> , 2017.
<i>B. ovis</i>	Sheep	ATC25840 ATC25840	33 38	Suárez-Esquivel <i>et al</i> , 2017. BLAST search.
<i>B. microti</i>	Common vole	CCM4915 <sup>T</sup> BCM4915 CCM4915 <sup>T</sup> “	5-6 14 13 13	Scholz <i>et al</i> , 2008. Suárez-Esquivel <i>et al</i> , 2017. Audic <i>et al</i> , 2009. BLAST search.
<i>B. neotomae</i>	Desert woodrats		?	Stoenner <i>et al</i> , 1957.
<i>B. inopinata</i>	Uncertain	BO-1	4-5	Scholz <i>et al</i> , 2010.
<i>B. pinnipedii</i>	Seals	B2/94 6/566	45-46 31 29	Suárez-Esquivel <i>et al</i> , 2017. Audic <i>et al</i> , 2011. BLAST search.
<i>B. ceti</i>	Dolphins, Porpoises, Whales	Various, TE28753	42-53 14	Suárez-Esquivel <i>et al</i> , 2017. BLAST search.
<i>B. papionis</i>	Baboon	F80/08-60 <sup>T</sup> F80/08-61	High, uncertain	Whatmore <i>et al</i> , 2014.
<i>B. vulpis</i>	Red Foxes  Bank vole	F60 F965	27 27 ?	Hofer <i>et al</i> , 2012. Hofer <i>et al</i> , 2012. Hammerl <i>et al</i> , 2015.
1. Novel motile species, closest to <i>B. inopinata</i>	Amphibians	09RB8471 10RB9215 UK 8/14 Various	Uncertain	Eisenberg <i>et al</i> , 2012. Eisenberg <i>et al</i> , 2012. Whatmore <i>et al</i> , 2015. Scholz <i>et al</i> , 2016.
2. Microti-like	Domestic Marsh frog	17-2122-4144	Uncertain	Jaý <i>et al</i> , 2018.

BLAST search available at: [https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE\\_TYPE=BlastSearch](https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE_TYPE=BlastSearch).

## References

- Audic S, Lescot, Claverie J-M, Scholz HC. 2009. *Brucella microti*: the genome sequence of an emerging pathogen. *BMC Genomics* **10**: 352. doi:10.1186/1471-2164-10-352.
- Audic S, Lescot M, Claverie J-M, Cloeckeaert A, Zygmunt MS. 2011. The genome sequence of *Brucella pinnipedialis* B2/94 sheds light on the evolutionary history of the genus *Brucella*. *BMC Evolutionary Biology* **11**: 200.
- Eisenberg T, Hamann HP, Kaim U, Schlez K, Seeger H, Schauerte N, Melzer F, Tomaso H, Scholz HC, Koylass MS, Whatmore AM, Zschöck M. 2012. Isolation of potentially novel *Brucella* spp. from frogs. *Applied and Environmental Microbiology* **78**: 3753-3755. Doi:10.1128/AEM.07509-11.
- Hammerl JA, Ulrich RG, Imholt C, Scholz HC, Jacob J, Kratzmann N, Nöckler K, Al Dahouk S. 2015. Molecular survey on *Brucellosis* in rodents and shrews – natural reservoirs of novel *Brucella* species in Germany? *Transboundary and Emerging Diseases* **64**: 663-671.
- Hofer E, Revilla-Fernández S, Al Dahouk S, Riehm JM, Nöckler K, Zygmunt MS, Cloeckeaert A, Tomaso H, Scholz HC. 2012. A potential novel *Brucella* species isolated from mandibular lymph nodes of red foxes in Austria. *Veterinary Microbiology* **155**: 93-99.
- Jaý M, Girault G, Perrot L, Taunay B, Vuilmet T, Rossignol F, Pitel P-H, Picard E, Ponsart C, Mick V. 2018. Phenotypic and Molecular Characterization of *Brucella microti*-Like Bacteria from a Domestic Marsh Frog (*Pelophylax ridibundus*). *Frontiers in Veterinary Science* **5**: 283. doi: 10.3389/fvets.2018.00283.
- Kay GL, Sergeant MJ, Giuffra V, Bandiera P, Milanese M, Bramanti B, Bianucci R, Pallen MJ. 2014. Recovery of a medieval *Brucella melitensis* genome using shotgun metagenomics. *mBio* **5**: (4). e01337-14. doi:10.1128/mBio.01337-14.
- Scholz HC, Hubálek Z, Sedláček I, Vergnaud G, Tomaso H, Al Dahouk S, Melzer F, Kämpfer P, Neubauer H, Cloeckeaert A, Maquart M, Zygmunt MS, Whatmore AM, Falsen E, Bahn P, Göllner C, Pfeffer M, Huber B, Busse HJ, Nöckler, K. 2008. *Brucella microti* sp. nov., isolated from the common vole *Microtus arvalis*. *International Journal of Systematic and Evolutionary Microbiology* **58**: 375–382.
- Scholz HC, Nöckler K, Göllner C, Bahn P, Vergnaud G, Tomaso H, et al. 2010. *Brucella inopinata* sp. nov., isolated from a breast implant infection. *International Journal of Systematic and Evolutionary Microbiology* **60**: 801-808. doi: 10.1099/ijs.0.
- Scholz HC, Mühldorfer K, Shilton C, Benedict S, Whatmore AM, Blom J, et al. 2016. The Change of a Medically Important Genus: Worldwide Occurrence of Genetically Diverse Novel *Brucella* Species in Exotic Frogs. *PLoS ONE* **11** (12): e0168872. doi:10.1371/journal.pone.0168872.
- Stoenner HG, Lackman DB. 1957. A new species of *Brucella* isolated from the desert wood rat, *Neotoma lepida* Thomas. *American Journal of Veterinary Research* **18**: 947–951.
- Suárez-Esquivel M, Baker KS, Ruiz-Villalobos N, Hernández-Mora G, Barquero-Calvo E, González-Barrientos R, Castillo-Zeledón A, Jiménez-Rojas C, Chacón-Díaz C, Cloeckeaert A, Chaves-Olarte E, Thomson NR, Moreno E, Guzmán-Verri C. 2017. *Brucella* genetic variability in wildlife marine mammals populations relates to host Preference and Ocean Distribution. *Genome Biology and*



*Evolution* **9**: 1901–1912 doi:10.1093/gbe/evx137.

Whatmore AM, Davison N, Cloeckert A, Al Dahouk S, Zygmunt MS, Brew SD, et al. 2014. *Brucella papionis* sp. nov., isolated from baboons (*Papio spp.*). *International Journal of Systematic and Evolutionary Microbiology* **64**: 4120–4128. doi: 10.1099/ijs.0.065482-0.

Whatmore AM, Dale E-J, Stubberfield E, Muchowski J, Koylass M, Dawson C, Gopaul KK, Perrett LL, Jones M, Lawrie A. 2015. Isolation of *Brucella* from a White's tree frog (*Litoria caerulea*). *JMM Case Reports* 2: doi: 10.1099/jmmcr.0.000017.

Table S2. Selected common infectious causes of abortion in domestic cattle, goats, sheep and pigs, summarised from Givens and Marley (2008).

		Embryonic death	Early foetal death	Mid-gestational foetal death	Late foetal death
<b>Cattle</b>	Bacterial pathogens	<i>Campylobacter fetus</i>		<i>Campylobacter fetus</i> , <i>Trueperella pyogenes</i> ,	<i>Campylobacter fetus</i> , <i>Histophilus somni</i> , <i>Ureaplasma</i> spp., <i>Brucella abortus</i> , <i>Leptospira</i> spp., <i>Listeria monocytogenes</i> , <i>Trueperella pyogenes</i> , <i>Chlamydophila</i> spp., <i>Coxiella burnetti</i> , <i>Salmonella</i> spp.
	Fungal pathogens			<i>Aspergillus fumigatus</i> , <i>Mucor</i> spp., <i>Mortierella</i> <i>wolfii</i>	<i>Aspergillus fumigatus</i> , <i>Mucor</i> spp., <i>Mortierella wolfii</i>
	Protozoan pathogens	<i>Tritrichomonas fetus</i>	<i>Tritrichomonas fetus</i>	<i>Neospora caninum</i>	<i>Anaplasma marginale</i>
	Viral pathogens	Bovine herpesvirus 1, Bovine viral diarrhoea virus	Bovine viral diarrhoea virus, Bluetongue virus	Bovine herpesvirus 1, Bovine viral diarrhoea virus	Bovine herpesvirus 1, Epizootic bovine abortion
<b>goat</b>	Bacterial pathogens				<i>Listeria monocytogenes</i> , <i>Chlamydophila</i> <i>abortus</i> , <i>Brucella melitensis</i> , <i>Coxiella</i> <i>burnetti</i> , <i>Mycoplasma</i> spp., <i>Leptospira</i> spp., <i>Salmonella abortus-ovis</i> , <i>Campylobacter</i> spp.
	Protozoan pathogens			<i>Sarcocystis</i> spp.	<i>Toxoplasma gondii</i>
	Viral pathogens	Border disease virus			Caprine herpesvirus
<b>sheep</b>	Bacterial pathogens				<i>Campylobacter</i> spp., <i>Listeria</i> <i>monocytogenes</i> , <i>Brucella</i> spp., <i>Chlamydophila abortus</i> , <i>Coxiella burnetti</i> <i>Salmonella</i> spp., <i>Leptospira</i> spp.
	Protozoan pathogens	<i>Toxoplasma gondii</i>			<i>Toxoplasma gondii</i>
	Viral pathogens	Cache valley virus	Border disease virus	Border disease virus, Bluetongue	Border disease virus
<b>pig</b>	Bacterial pathogens	<i>Brucella suis</i>	<i>Brucella suis</i>	<i>Brucella suis</i> , <i>Leptospira</i> <i>pomona</i> , <i>Streptococcus suis</i> , <i>Mycoplasma suis</i>	<i>Brucella suis</i> , <i>Leptospira pomona</i> , <i>Chlamydia</i> spp.

Protozoan pathogens	<i>Toxoplasma gondii</i>			
Viral pathogens	Porcine parvovirus, Porcine enterovirus and teschovirus, Pseudorabies virus (Suid herpesvirus-1), Classical swine fever	Porcine parvovirus, Porcine enterovirus and teschovirus, Pseudorabies virus (Suid herpesvirus-1), Classical swine fever virus, Japanese encephalitis virus	Porcine parvovirus, Pseudorabies virus (Suid herpesvirus-1), Classical swine fever virus, Japanese encephalitis virus	Pseudorabies virus (Suid herpesvirus-1), Classical swine fever virus, Porcine circovirus type 2,

### References

Givens D M and Marley M S D 2008, Infectious causes of embryonic and fetal mortality. *Theriogenology* **70**: 270-285.

Table S3. Primers and FAM labelled probe used in our *Brucella* screening PCR method.

Primer	Sequence	Anneal °C	Amplicon bp (primers)
F1	5'-AGCCTTTCCCATACACCGGCGT-3'	60	141 (F1/R1)
R1	5'-TAACCGATTATTTGTCGACGCCATCTT-3'		
F2	5'-TACACCGGCGTGCGACCAGCGC-3'	60	108 (F2/R2)
R2	5'-ATCTTGTGGATGGCTGCCAATGCAGC-3'		
R3	5'-CGCACTGGCGCGATCTGCCT-3'	58	85 (F2/R3)
Probe	5'-(6FAM)-GCATGAACCGCTGTCCATTTGCCGAAGGTCGC-BHQ1-3'	Acquire at 74	

Table S4. Examples of historical approaches and evidence for potential case study examination of brucellosis in English history

<b>Historical sub-disciplines of relevance</b>	<b>Questions/topics of interest</b>	<b>Examples of relevant historical events, topics or source materials</b>
Political History	Contextual evidence of changing practices relating to agricultural production and public health	Impact of national level policies such as labour legislation, taxation, enclosures, food supply strategies, milk pasteurisation etc. (e.g. Steere-Williams, 2015)
Economic History	Economic value of livestock on national and local levels Demographic profiles of sheep flocks	Seminal work on the Agrarian History of England and Wales, including analysis of regional variations and the practices of larger and smaller scale agricultural holdings (eight volumes covering the period 43 A.D. to 1939, e.g. Thirsk, 1985).
Agrarian History	Animal husbandry techniques and farming practices Contribution of small scale animal husbandry to individual households	
Social History	Proximity of people to animals Personal living standards and diet	Probate inventories and records of debt Studies of agrarian communities and peasant society (e.g. Dyer, 1989)
Historical Demography	Health and demographic status of local populations	Examination of life expectancy and other indicators of human health and longevity (e.g. Wrigley et al., 1997)
Medical History	Past understanding of medicine and of disease risks Public health measures to prevent disease risk Diagnosis of disease in the past	Studies of medical knowledge and its application in the past (e.g. Porter, 1997)
Local History	Exploration of detailed case study examples of changing agricultural practices in specific areas and the potential impacts upon local populations	Studies of specific estates or farming communities, such as those utilising the pipe rolls of the medieval Bishops of Winchester (e.g. Stone, 2017)

## References

Dyer C. 1989. *Standards of living in the later middle ages: social change in England, c.1200-1520*. Cambridge University Press: Cambridge.

Porter R. 1997. *The greatest benefit to mankind: a medical history of humanity from antiquity to the present*. Harper Collins: London.

Steere-Williams J. 2015. Milking Science for its Worth: The Reform of the British Milk Trade in the Late Nineteenth Century. *Agricultural History* **89:2**, 263-288. DOI: 10.3098/ah.2015.089.2.263

Stone D. (ed.) 2017. *The Accounts for the Manor of Esher in the Winchester Pipe Rolls, 1235-1376* Surrey Record Society, Vol XLVI. Surrey Record Society: Woking.

Thirsk J. (ed.) 1985. *The Agrarian History of England and Wales, 5: 1640-1750*. 2 pts. Cambridge University Press: Cambridge.

Wrigley EA, Davies RS, Oeppen JE and Schofield RS. 1997. *English population history from family reconstitution, 1580-1837*. Cambridge University Press: Cambridge.