

COUNTING POINTS OF FIXED DEGREE AND BOUNDED HEIGHT

MARTIN WIDMER

ABSTRACT. We consider the set of points in projective n -space that generate an extension of degree e over given number field k , and deduce an asymptotic formula for the number of such points of absolute height at most X , as X tends to infinity. We deduce a similar such formula with instead of the absolute height, a so-called adelic-Lipschitz height.

1. INTRODUCTION

We consider the set of points in projective n -space that generate an extension of degree e over given number field k , and deduce an asymptotic formula for the number of such points of absolute height at most X , as X tends to infinity. We deduce a similar such formula with instead of the absolute height, a so-called adelic-Lipschitz height.

Let k be a number field of degree $m = [k : \mathbb{Q}]$ in a fixed algebraic closure \bar{k} of k and let n be a positive integer. Write $\mathbb{P}^n(\bar{k})$ for the projective space of dimension n over the field \bar{k} and denote by $H(\cdot)$ the non-logarithmic absolute Weil height on $\mathbb{P}^n(\bar{k})$ as defined in [1] p.16. A fundamental property of the height, usually associated with the name of Northcott due to his result Theorem 1 in [11], states that subsets of $\mathbb{P}^n(\bar{k})$ of bounded degree and bounded height are finite. This raises the question of estimating the cardinality of such a set as the height bound gets large. Schanuel proved in [14] that for the counting function

$$Z_H(\mathbb{P}^n(k), X) = |\{P \in \mathbb{P}^n(k); H(P) \leq X\}|$$

one has an asymptotic formula

$$(1.1) \quad Z_H(\mathbb{P}^n(k), X) = S_k(n)X^{m(n+1)} + O(X^{m(n+1)-1} \log X)$$

as X tends to infinity where $S_k(n)$ is a positive constant involving all classical field invariants (see (3.1) for its definition) and the constant implied by the Landau O -symbol depends on k and n . The logarithm can be omitted in all cases except for $n = m = 1$.

Date: May 15, 2009.

2000 Mathematics Subject Classification. Primary 11R04; Secondary 11G50, 11G35.

Key words and phrases. Height, Northcott's Theorem, counting.

A projective point $P = (\alpha_0 : \dots : \alpha_n)$ in $\mathbb{P}^n(\bar{k})$ has a natural degree defined as

$$[k(P) : k]$$

where $k(P)$ denotes the extension we get by adjoining all ratios α_i/α_j ($0 \leq i, j \leq n, \alpha_j \neq 0$) to k . In 1993 Schmidt drew attention to the more general set

$$\mathbb{P}^n(k; e) = \{P \in \mathbb{P}^n(\bar{k}); [k(P) : k] = e\}$$

of points with relative degree e . Clearly $\mathbb{P}^n(k; 1) = \mathbb{P}^n(k)$ and so Schanuel deals with the case $e = 1$. For the counting function

$$Z_H(\mathbb{P}^n(k; e), X) = |\{P \in \mathbb{P}^n(k; e); H(P) \leq X\}|$$

Schmidt [16] proved the following general estimates

$$(1.2) \quad cX^{me(\max\{e, n\}+1)} \leq Z_H(\mathbb{P}^n(k; e), X) \leq CX^{me(e+n)}$$

where $c = c(k, e, n)$ and $C = C(k, e, n)$ are positive constants depending solely on k, e and n . The upper bound holds for $X \geq 0$ and the lower bound holds for $X \geq X_0(k, e, n)$ depending also on k, e and n . Moreover one can choose $C = 2^{me(e+n+3)+e^2+n^2+10e+10n}$.

For $k = \mathbb{Q}$ more is known. Schmidt [17] investigated the quadratic case. Here he provided not only the correct order of magnitude but he found also the precise asymptotics and this in all dimensions n . As X tends to infinity one has

$$(1.3) \quad Z_H(\mathbb{P}^n(\mathbb{Q}; 2), X) = \begin{cases} D_1 X^6 + O(X^4 \log X) & \text{if } n = 1 \\ D_2 X^6 \log X + O(X^6 \sqrt{\log X}) & \text{if } n = 2 \\ D_n X^{2(n+1)} + O(X^{2n+1}) & \text{if } n > 2 \end{cases} .$$

The constant implied by the O -symbol depends only on n . In fact Schmidt's result was more precise since it gave the asymptotics for real and imaginary quadratic points separately. Here $D_1 = \frac{8}{\zeta(3)}$, $D_2 = \frac{8(12+\pi^2)}{\zeta(3)^2}$ and $D_n = D(\mathbb{Q}, 2, n)$ is given by

$$D(\mathbb{Q}, 2, n) = \sum_K S_K(n)$$

where the sum runs over all quadratic fields K . Schmidt proved also an analogue to the above result for a more general kind of height and showed that this leads to asymptotic formulae for the number of decomposable quadratic forms (i.e. product of two linear forms) $f(x_0, \dots, x_n) = \sum_{0 \leq i < j \leq n} a_{ij} x_i x_j$ with coefficients a_{ij} in \mathbb{Z} having $|a_{ij}| \leq X$ and moreover for the number of symmetric $(n+1) \times (n+1)$ matrices with rank ≤ 2 such that $b_{ii} \in \mathbb{Z}$, $|b_{ii}| \leq X$ and $2b_{ij} \in \mathbb{Z}$, $2|b_{ij}| \leq X$ for $i \neq j$. Already way back in 1967

Schmidt [15] introduced more general classes of heights where the maximum norm in (2.1) at the infinite places is replaced by an arbitrary but fixed distance function. More recently Thunder [21] and Roy-Thunder [13] introduced “twisted heights” which allow also modifications at the finite places.

One year after Schmidt’s article on quadratic points Gao [5] made further progress. He proved that if $n > e > 2$ one has

$$(1.4) \quad Z_H(\mathbb{P}^n(\mathbb{Q}; e), X) = D(\mathbb{Q}, e, n)X^{e(n+1)} + O(X^{e(n+1)-1})$$

as X tends to infinity. The constant implied by the O -symbol depends on e and n and the constant $D(\mathbb{Q}, e, n)$ is given by $\sum_K S_K(n)$ where the sum runs over all extensions K of \mathbb{Q} of degree e . For $1 \leq n \leq e$ Gao showed that the correct order of magnitude of $Z_H(\mathbb{P}^n(\mathbb{Q}; e), X)$ is $X^{e(e+1)}$. Here the asymptotics are still unknown, even in the case $e = 3$ and $n = 2$ of cubic points in two dimensions.

Schmidt’s and Gao’s results are restricted to the ground field $k = \mathbb{Q}$. A more recent result of Masser and Vaaler [9] gives the asymptotics for the number of points of fixed degree over an arbitrary fixed number field k , but only in dimension $n = 1$. Masser and Vaaler established the asymptotic formula

$$(1.5) \quad Z_H(\mathbb{P}^1(k; e), X) = eV_{\mathbb{R}}(e)^{r_k}V_{\mathbb{C}}(e)^{s_k}S_k(e)X^{me(e+1)} + O(X^{me(e+1)-e} \log X)$$

as X tends to infinity. The constants $V_{\mathbb{R}}(e), V_{\mathbb{C}}(e)$ have their origins in [3]. Moreover the logarithm can be omitted in all cases except $(m, e) = (1, 1)$ and $(m, e) = (1, 2)$ and the constant implied by the O -symbol depends on k and e . Unfortunately the proof of Masser and Vaaler’s theorem shed no light on the case $n > 1$. Very roughly speaking Masser and Vaaler’s idea was to interpret the height of the root of an irreducible polynomial in $k[x]$ of fixed degree e as a suitable height of the coefficient vector of this polynomial and to proceed by counting minimal polynomials with respect to this modified height. To carry out this plan they had to generalize the class of heights introduced by Schmidt [15], allowing now different distance functions at the infinite places instead of only one for all infinite places as Schmidt did. On the other hand Masser and Vaaler had to impose a technical condition, associated with the name of Lipschitz, on the boundaries of the unit balls given by the respective distance function. They therefore introduced so-called Lipschitz systems, giving what one could call Lipschitz

heights.

In the present article we establish the asymptotics for $\mathbb{P}^n(k; e)$ if n is slightly larger than $5e/2$. Let us write

$$D = D(k, e, n) = \sum_K S_K(n)$$

for the formal sum taken over all extensions of K of k in \bar{k} of degree e . We have the following result.

Theorem 1.1. *Let e, n be positive integers and k a number field of degree m and suppose that $n > 5e/2 + 4 + 2/(me)$. Then the sum defining D converges and as X tends to infinity we have*

$$Z_H(\mathbb{P}^n(k; e), X) = DX^{me(n+1)} + O(X^{me(n+1)-1} \log X).$$

The logarithm can be omitted unless $(me, n) = (1, 1)$ and the constant implied by the O -symbol depends on k, e and n .

If e and n are both larger than one there is a considerable gap between the exponents of the lower and the upper bound in (1.2). Schmidt mentioned that the lower bound is likely to be nearer the truth than the upper bound. Our Theorem 1.1 confirms Schmidt's conjecture at least if n is large enough. We will prove a more general result involving adelic-Lipschitz heights.

Let us give a single new example of our theorem. We take $n = 11$, $k = \mathbb{Q}(i)$, $e = 2$, so that we are counting points in eleven dimensions quadratic over $\mathbb{Q}(i)$. For the number $Z = Z_H(\mathbb{P}^{11}(\mathbb{Q}(i); 2), X)$ of points of height at most X , the Schmidt bounds are $X^{48} \ll Z \ll X^{52}$ for $X \geq X_0$, with absolute implied constants. Our result implies that

$$Z = DX^{48} + O(X^{47})$$

with

$$D = 12 \cdot (2\pi)^{24} \sum_{\substack{K \\ [K:\mathbb{Q}(i)]=2}} \frac{h_K R_K}{w_K \zeta_K(12) |\Delta_K|^6}.$$

Our proof follows the general strategy of Schmidt and Gao. Their audacious idea was to prove a result similar to (1.1) but with $\mathbb{P}^n(K)$ replaced by $\mathbb{P}^n(K/\mathbb{Q})$ the subset of primitive points in $\mathbb{P}^n(K)$; by definition these satisfy $K = \mathbb{Q}(P)$. Now $\mathbb{P}^n(\mathbb{Q}; e)$ is a disjoint union of the sets $\mathbb{P}^n(K/\mathbb{Q})$ where K runs over all number fields of degree e . For each $\mathbb{P}^n(K/\mathbb{Q})$ the main term is the same as that in (1.1) with K instead of k , but for $e = 2$ Schmidt

obtained a more precise error term

$$(1.6) \quad O\left(\frac{\sqrt{h_K R_K \log(3 + h_K R_K)}}{|\Delta_K|^{n/2}} X^{2n+1}\right)$$

where the constant in O depends only on n but is independent of the field K . This is the major step of the proof and involves many new ideas. Now one can sum over all quadratic number fields and the Theorem of Siegel-Brauer ensures that the sum over the main terms $S_K(n)$ as well as over the error terms converges provided $n > 2$. For similar reasons the restriction $n > e$ in Gao's result appears.

We close the introduction with some remarks about the structure of the paper. First we take up the definition of an adelic-Lipschitz system from [23] on a number field and we define a uniform adelic-Lipschitz system on the collection of all extensions of k of degree e . This then gives rise to a class of heights $H_{\mathcal{N}}$ defined on $\mathbb{P}^n(k; e)$. The main result asymptotically estimates the counting function of $\mathbb{P}^n(k; e)$ with respect to the height $H_{\mathcal{N}}$. In Theorem 1.1 we used only the simplest formulation by choosing a special uniform adelic-Lipschitz system with maximum norms at all places so that the corresponding adelic-Lipschitz height $H_{\mathcal{N}}$ becomes just the Weil height H . In Section 3 we state our main theorem for general adelic-Lipschitz heights. It is in [24] and [25] where we see the advantage of working in such generality. In [24] we are concerned with counting points of fixed degree on linear subvarieties of projective space. In [25] we prove the following result: let m, n be positive integers with $n > \max\{6m + 2 + 2/m, m^2 + m\}$. Then as X tends to infinity the number of algebraic numbers α of degree mn such that $\mathbb{Q}(\alpha)$ contains a subfield of degree m and $H(1, \alpha) \leq X$ is asymptotically equal to

$$D'(m, n) X^{mn(n+1)}$$

where $D'(m, n) = \sum_K n V_{\mathbb{R}}(n)^{r_k} V_{\mathbb{C}}(n)^{s_k} S_k(n)$ and the sum runs over all number fields of degree m . Note that the subfield condition reduces the order of magnitude from $X^{mn(mn+1)}$ to $X^{mn(n+1)}$.

In Section 4 we prove the main result Theorem 3.1 which is a generalization of Theorem 1.1 to adelic-Lipschitz heights. Section 5 is devoted to some simple lower and upper bounds for the number of extensions K/k of fixed degree with $\delta(K/k) \leq T$, where $\delta(K/k) = \inf_{\alpha} \{H(1, \alpha); K = k(\alpha)\}$. The invariant $\delta(K/k)$ was already introduced in [23]. Our bounds are essentially by-products of the proof of Theorem 3.1.

ACKNOWLEDGEMENTS

I would like to thank my Ph.D. advisor David Masser for his constant support. His comments led to many significant improvements of this article. I also would like to thank Jeffrey Vaaler for helpful discussions and Gao Xia for showing us his Ph.D. thesis. Finally I would like to thank the referee for carefully reading this manuscript and providing many detailed suggestions to improve the exposition of this article. This work was financially supported by the Swiss National Science Foundation.

2. ADELIC-LIPSCHITZ HEIGHTS REVISITED

The Subsections 2.1, 2.2 and 2.3 of this section are entirely contained in [23]. But adelic-Lipschitz heights are crucial for the entire paper and thus, for convenience of the reader, we introduce this notion here once again. Before we can define adelic-Lipschitz heights we have to fix some basic notation. For a detailed account on heights we refer to [1] and [6].

2.1. Preliminaries. Let K be a finite extension of \mathbb{Q} of degree $d = [K : \mathbb{Q}]$. By a place v of K we mean an equivalence class of non-trivial absolute values on K . The set of all places of K will be denoted by M_K . For each v in M_K we write K_v for the completion of K at the place v and d_v for the local degree defined by $d_v = [K_v : \mathbb{Q}_v]$ where \mathbb{Q}_v is a completion with respect to the place which extends to v . A place v in M_K corresponds either to a non-zero prime ideal \mathfrak{p}_v in the ring of integers \mathcal{O}_K or to an embedding σ of K into \mathbb{C} . If v comes from a prime ideal we call v a finite or non-archimedean place and denote this by $v \nmid \infty$ and if v corresponds to an embedding we say v is an infinite or archimedean place and denote this by $v \mid \infty$. For each place in M_K we choose a representative $|\cdot|_v$, normalized in the following way: if v is finite and $\alpha \neq 0$ we set by convention

$$|\alpha|_v = N\mathfrak{p}_v^{-\frac{\text{ord}_{\mathfrak{p}_v}(\alpha\mathcal{O}_K)}{d_v}}$$

where $N\mathfrak{p}_v$ denotes the norm of \mathfrak{p}_v from K to \mathbb{Q} and $\text{ord}_{\mathfrak{p}_v}(\alpha\mathcal{O}_K)$ is the power of \mathfrak{p}_v in the prime ideal decomposition of the fractional ideal $\alpha\mathcal{O}_K$. Moreover we set

$$|0|_v = 0.$$

And if v is infinite and corresponds to an embedding $\sigma : K \hookrightarrow \mathbb{C}$ we define

$$|\alpha|_v = |\sigma(\alpha)|.$$

If α is in $K^* = K \setminus \{0\}$ then $|\alpha|_v \neq 1$ holds for only a finite number of places v .

Throughout this article n will denote a positive rational integer. The height on K^{n+1} is defined by

$$(2.1) \quad H(\alpha_0, \dots, \alpha_n) = \prod_{M_K} \max\{|\alpha_0|_v, \dots, |\alpha_n|_v\}^{\frac{d_v}{d}}.$$

Due to the remark above this is in fact a finite product. Furthermore this definition is independent of the field K containing the coordinates (see [1] Lemma 1.5.2 or [6] pp.51-52) and therefore defines a height on $\overline{\mathbb{Q}}^{n+1}$ for an algebraic closure $\overline{\mathbb{Q}}$ of \mathbb{Q} . The well-known *product formula* (see [1] Proposition 1.4.4) asserts that

$$\prod_{M_K} |\alpha|_v^{d_v} = 1 \text{ for each } \alpha \text{ in } K^*.$$

This implies in particular that the value of the height in (2.1) does not change if we multiply each coordinate with a fixed element of K^* . Therefore one can define a height on points $P = (\alpha_0 : \dots : \alpha_n)$ in $\mathbb{P}^n(\overline{\mathbb{Q}})$ by

$$(2.2) \quad H(P) = H(\alpha_0, \dots, \alpha_n).$$

Moreover, to evaluate the height, we can assume that one of the coordinates is 1 which shows that $H(\boldsymbol{\alpha}) \geq 1$ for $\boldsymbol{\alpha} \in \overline{\mathbb{Q}}^{n+1} \setminus \{\mathbf{0}\}$. The equations (2.1) and (2.2) define the absolute non-logarithmic projective Weil height or just Weil height.

2.2. Adelic-Lipschitz systems on a number field. Let r be the number of real embeddings and s the number of pairs of complex conjugate embeddings of K so that $d = r + 2s$. Recall that M_K denotes the set of places of K . For every place v we fix a completion K_v of K at v . The value set of v , $\Gamma_v := \{|\alpha|_v; \alpha \in K_v\}$ is equal to $[0, \infty)$ if v is archimedean, and to

$$\{0, (N\mathfrak{p}_v)^0, (N\mathfrak{p}_v)^{\pm 1/d_v}, (N\mathfrak{p}_v)^{\pm 2/d_v}, \dots\}$$

if v is non-archimedean. For $v \mid \infty$ we identify K_v with \mathbb{R} or \mathbb{C} respectively and we identify \mathbb{C} with \mathbb{R}^2 via $\xi \longrightarrow (\Re(\xi), \Im(\xi))$ where we used \Re for the real and \Im for the imaginary part of a complex number.

Before we can introduce adelic-Lipschitz systems we have to give a technical definition. For a vector \mathbf{x} in \mathbb{R}^n we write $|\mathbf{x}|$ for the euclidean length of \mathbf{x} .

Definition 1. Let M and $D > 1$ be positive integers and let L be a non-negative real. We say that a set S is in $Lip(D, M, L)$ if S is a subset of \mathbb{R}^D ,

and if there are M maps $\phi_1, \dots, \phi_M : [0, 1]^{D-1} \longrightarrow \mathbb{R}^D$ satisfying a Lipschitz condition

$$(2.3) \quad |\phi_i(\mathbf{x}) - \phi_i(\mathbf{y})| \leq L|\mathbf{x} - \mathbf{y}| \text{ for } \mathbf{x}, \mathbf{y} \in [0, 1]^{D-1}, i = 1, \dots, M$$

such that S is covered by the images of the maps ϕ_i .

We call L a Lipschitz constant for the maps ϕ_i . By definition the empty set lies in $\text{Lip}(D, M, L)$ for any positive integers M and $D > 1$ and any non-negative L .

Definition 2 (Adelic-Lipschitz system). *An adelic-Lipschitz system (ALS) \mathcal{N}_K or simply \mathcal{N} on K (of dimension n) is a set of continuous maps*

$$(2.4) \quad N_v : K_v^{n+1} \rightarrow \Gamma_v \quad v \in M_K$$

such that for $v \in M_K$ we have

- (i) $N_v(\mathbf{z}) = 0$ if and only if $\mathbf{z} = \mathbf{0}$,
- (ii) $N_v(\omega\mathbf{z}) = |\omega|_v N_v(\mathbf{z})$ for all ω in K_v and all \mathbf{z} in K_v^{n+1} ,
- (iii) if $v \mid \infty$: $\{\mathbf{z}; N_v(\mathbf{z}) = 1\}$ is in $\text{Lip}(d_v(n+1), M_v, L_v)$ for some M_v, L_v ,
- (iv) if $v \nmid \infty$: $N_v(\mathbf{z}_1 + \mathbf{z}_2) \leq \max\{N_v(\mathbf{z}_1), N_v(\mathbf{z}_2)\}$ for all $\mathbf{z}_1, \mathbf{z}_2$ in K_v^{n+1} .

Moreover we assume that

$$(2.5) \quad N_v(\mathbf{z}) = \max\{|z_0|_v, \dots, |z_n|_v\}$$

for all but a finite number of $v \in M_K$. If we consider only the functions N_v for $v \mid \infty$ then we get an (r, s) -Lipschitz system (of dimension n) in the sense of Masser and Vaaler [9]. With M_v and L_v from (iii) we define

$$M_{\mathcal{N}} = \max_{v \mid \infty} M_v,$$

$$L_{\mathcal{N}} = \max_{v \mid \infty} L_v.$$

We say that \mathcal{N} is an ALS with associated constants $M_{\mathcal{N}}, L_{\mathcal{N}}$. The set defined in (iii) is the boundary of the set $\mathbf{B}_v = \{\mathbf{z}; N_v(\mathbf{z}) < 1\}$ and therefore \mathbf{B}_v is a bounded symmetric open star-body in \mathbb{R}^{n+1} or \mathbb{C}^{n+1} (see also [9] p.431). In particular \mathbf{B}_v has a finite volume V_v .

Let us consider the system where N_v is as in (2.5) for all places v . If v is an infinite place then \mathbf{B}_v is a cube for $d_v = 1$ and the complex analogue if $d_v = 2$. Their boundaries are clearly in $\text{Lip}(d_v(n+1), M_v, L_v)$ most naturally with $M_v = 2n+2$ maps and $L_v = 2$ if $d_v = 1$ and with $M_v = n+1$ maps and for example $L_v = 2\pi\sqrt{2n+1}$ if $d_v = 2$. This system is called the standard adelic-Lipschitz system.

We return to arbitrary adelic-Lipschitz systems. We claim that for any $v \in M_K$ there is a c_v in the value group $\Gamma_v^* = \Gamma_v \setminus \{0\}$ with

$$(2.6) \quad N_v(\mathbf{z}) \geq c_v \max\{|z_0|_v, \dots, |z_n|_v\}$$

for all $\mathbf{z} = (z_0, \dots, z_n)$ in K_v^{n+1} . For if v is archimedean then \mathbf{B}_v is bounded open and it contains the origin. Since Γ_v^* contains arbitrary small positive numbers the claim follows by (ii). Now for v non-archimedean N_v and $\max\{|z_0|_v, \dots, |z_n|_v\}$ define norms on the vector space K_v^{n+1} over the complete field K_v . But on a finite dimensional vector space over a complete field all norms are equivalent ([2] Corollary 5. p.93) hence (2.6) remains true for a suitable choice of c_v .

So let \mathcal{N} be an ALS on K of dimension n . For every v in M_K let c_v be an element of Γ_v^* , such that $c_v \leq 1$ and (2.6) holds. Due to (2.5) we can assume that $c_v \neq 1$ only for a finite number of places v . We define

$$(2.7) \quad C_{\mathcal{N}}^{fin} = \prod_{v \nmid \infty} c_v^{-\frac{d_v}{d}} \geq 1$$

and

$$(2.8) \quad C_{\mathcal{N}}^{inf} = \max_{v \mid \infty} \{c_v^{-1}\} \geq 1.$$

Multiplying the finite and the infinite part gives rise to another constant

$$(2.9) \quad C_{\mathcal{N}} = C_{\mathcal{N}}^{fin} C_{\mathcal{N}}^{inf}.$$

It will turn out that besides $M_{\mathcal{N}}$ and $L_{\mathcal{N}}$ this is another important quantity for an ALS. So we say that \mathcal{N} is an ALS with associated constants $C_{\mathcal{N}}, M_{\mathcal{N}}, L_{\mathcal{N}}$.

Remark 1. Let v be an infinite place. Suppose $N_v : K_v^{n+1} \rightarrow [0, \infty)$ defines a norm, so that $N_v(\mathbf{z}_1 + \mathbf{z}_2) \leq N_v(\mathbf{z}_1) + N_v(\mathbf{z}_2)$. Then \mathbf{B}_v is convex and (2.6) combined with (2.7), (2.8) and (2.9) shows that \mathbf{B}_v lies in $B_0(C_{\mathcal{N}}\sqrt{n+1})$. This implies (see Theorem A.1 in [22]) that $\partial\mathbf{B}_v$ lies in $\text{Lip}(d_v(n+1), 1, 8d_v^2(n+1)^{5/2}C_{\mathcal{N}})$.

We denote by $\sigma_1, \dots, \sigma_d$ the embeddings from K to \mathbb{R} or \mathbb{C} respectively, ordered such that $\sigma_{r+s+i} = \bar{\sigma}_{r+i}$ for $1 \leq i \leq s$. We define

$$(2.10) \quad \begin{aligned} \sigma : K &\longrightarrow \mathbb{R}^r \times \mathbb{C}^s \\ \sigma(\alpha) &= (\sigma_1(\alpha), \dots, \sigma_{r+s}(\alpha)). \end{aligned}$$

Sometimes it will be more readable to omit the brackets and simply to write $\sigma\alpha$. We identify \mathbb{C} in the usual way with \mathbb{R}^2 and extend σ componentwise to get a map

$$(2.11) \quad \sigma : K^{n+1} \longrightarrow \mathbb{R}^D$$

where $D = d(n+1)$. On \mathbb{R}^D we use $|\cdot|$ for the usual euclidean norm. For $v \in M_K$ let σ_v be the canonical embedding of K in K_v , again extended componentwise on K^{n+1} .

Definition 3. Let $\mathfrak{D} \neq 0$ be a fractional ideal in K and \mathcal{N} an ALS of dimension n . We define

$$(2.12) \quad \Lambda_{\mathcal{N}}(\mathfrak{D}) = \{\sigma(\boldsymbol{\alpha}); \boldsymbol{\alpha} \in K^{n+1}, N_v(\sigma_v \boldsymbol{\alpha}) \leq |\mathfrak{D}|_v \text{ for all finite } v\}$$

where $|\mathfrak{D}|_v = N\mathfrak{p}_v^{-\frac{\text{ord}_{\mathfrak{p}_v} \mathfrak{D}}{d_v}}$.

It is easy to see that $\Lambda_{\mathcal{N}}(\mathfrak{D})$ is an additive subgroup of \mathbb{R}^D . Now assume $B \geq 1$ and $|\sigma(\boldsymbol{\alpha})| \leq B$; then (2.6) implies $H(\boldsymbol{\alpha})^d \leq (BC_{\mathcal{N}}^{fin})^d N\mathfrak{D}^{-1}$ and by Northcott's Theorem we deduce that $\Lambda_{\mathcal{N}}(\mathfrak{D})$ is discrete. The same argument as for (2.6) yields positive real numbers C_v , one for each non-archimedean place $v \in M_K$, with $N_v(\mathbf{z}) \leq C_v \max\{|z_0|_v, \dots, |z_n|_v\}$ for all $\mathbf{z} = (z_0, \dots, z_n)$ in K_v^{n+1} and $C_v = 1$ for all but finitely many non-archimedean $v \in M_K$. Thus there exists an ideal $\mathfrak{C}_1 \neq 0$ in \mathcal{O}_K with $|\mathfrak{C}_1|_v \leq 1/C_v$ for all non-archimedean places $v \in M_K$. This means that $\sigma(\mathfrak{C}_1\mathfrak{D})^{n+1} \subseteq \Lambda_{\mathcal{N}}(\mathfrak{D})$. It is well-known that the additive group $\sigma(\mathfrak{C}_1\mathfrak{D})^{n+1}$ has maximal rank in \mathbb{R}^D . Therefore $\Lambda_{\mathcal{N}}(\mathfrak{D})$ is a discrete additive subgroup of \mathbb{R}^D of maximal rank. Hence $\Lambda_{\mathcal{N}}(\mathfrak{D})$ is a lattice. Notice that for ε in K^* one has

$$(2.13) \quad \det \Lambda_{\mathcal{N}}((\varepsilon)\mathfrak{D}) = |N_{K/\mathbb{Q}}(\varepsilon)|^{n+1} \det \Lambda_{\mathcal{N}}(\mathfrak{D}).$$

Therefore

$$(2.14) \quad \Delta_{\mathcal{N}}(\mathfrak{D}) = \frac{\det \Lambda_{\mathcal{N}}(\mathfrak{D})}{N\mathfrak{D}^{n+1}}$$

is independent of the choice of the representative \mathfrak{D} but depends only on the ideal class \mathcal{D} of \mathfrak{D} . Let Cl_K denote the ideal class group of K . We define

$$(2.15) \quad V_{\mathcal{N}}^{fin} = 2^{-s(n+1)} |\Delta_K|^{\frac{n+1}{2}} h_K^{-1} \sum_{\mathcal{D} \in Cl_K} \Delta_{\mathcal{N}}(\mathfrak{D})^{-1}$$

for the finite part, where as usual, s denotes the number of pairs of complex conjugate embeddings of K , h_K the class number of K and Δ_K the discriminant of K . The infinite part is defined by

$$V_{\mathcal{N}}^{inf} = \prod_{v|\infty} V_v.$$

By virtue of (2.6) we observe that

$$(2.16) \quad V_{\mathcal{N}}^{inf} = \prod_{v|\infty} V_v \leq \prod_{v|\infty} (2C_{\mathcal{N}}^{inf})^{d_v(n+1)} = (2C_{\mathcal{N}}^{inf})^{d(n+1)}.$$

We multiply the finite and the infinite part to get a global volume

$$(2.17) \quad V_{\mathcal{N}} = V_{\mathcal{N}}^{inf} V_{\mathcal{N}}^{fin}.$$

2.3. Adelic-Lipschitz heights on $\mathbb{P}^n(K)$. Let \mathcal{N} be an *ALS* on K of dimension n . Then the height $H_{\mathcal{N}}$ on K^{n+1} is defined by

$$H_{\mathcal{N}}(\boldsymbol{\alpha}) = \prod_{v \in M_K} N_v(\sigma_v(\boldsymbol{\alpha}))^{\frac{d_v}{d}}.$$

Thanks to the product formula and (ii) from Subsection 2.2 $H_{\mathcal{N}}(\boldsymbol{\alpha})$ does not change if we multiply each coordinate of $\boldsymbol{\alpha}$ with a fixed element of K^* . Therefore $H_{\mathcal{N}}$ is well-defined on $\mathbb{P}^n(K)$ by setting

$$H_{\mathcal{N}}(P) = H_{\mathcal{N}}(\boldsymbol{\alpha})$$

where $P = (\alpha_0 : \dots : \alpha_n) \in \mathbb{P}^n(K)$ and $\boldsymbol{\alpha} = (\alpha_0, \dots, \alpha_n) \in K^{n+1}$.

Remark 2. *Multiplying (2.6) over all places with suitable multiplicities yields*

$$(2.18) \quad H_{\mathcal{N}}(P) \geq C_{\mathcal{N}}^{-1} H(P)$$

for $P \in \mathbb{P}^n(K)$. Thanks to Northcott's Theorem it follows that $\{P \in \mathbb{P}^n(K); H_{\mathcal{N}}(P) \leq X\}$ is a finite set for each X in $[0, \infty)$.

2.4. Adelic-Lipschitz systems on a collection of number fields. Recall that k is a number field of degree m and \bar{k} is an algebraic closure of k . We fix k and \bar{k} throughout and assume finite extensions of k to lie in \bar{k} . Let \mathcal{C} be a collection of finite extensions of k . We are especially interested in the set of all extensions of fixed relative degree. We denote it by

$$\mathcal{C}_e = \mathcal{C}_e(k) = \{K \subseteq \bar{k}; [K : k] = e\}.$$

Let \mathcal{N} be a collection of adelic-Lipschitz systems \mathcal{N}_K of dimension n - one for each K of \mathcal{C} . Then we call \mathcal{N} an *adelic-Lipschitz system (ALS) on \mathcal{C} of dimension n* . We say \mathcal{N} is a *uniform ALS* on \mathcal{C} of dimension n with associated constants $C_{\mathcal{N}}, M_{\mathcal{N}}, L_{\mathcal{N}}$ in \mathbb{R} if the following holds: for each *ALS* \mathcal{N}_K of the collection \mathcal{N} we can choose associated constants $C_{\mathcal{N}_K}, M_{\mathcal{N}_K}, L_{\mathcal{N}_K}$ satisfying

$$C_{\mathcal{N}_K} \leq C_{\mathcal{N}}, \quad M_{\mathcal{N}_K} \leq M_{\mathcal{N}}, \quad L_{\mathcal{N}_K} \leq L_{\mathcal{N}}.$$

Notice that a uniform *ALS* \mathcal{N} (of dimension n) on the collection consisting only of a single field K with associated constants $C_{\mathcal{N}}, M_{\mathcal{N}}, L_{\mathcal{N}}$ is simply an *ALS* \mathcal{N} (of dimension n) on K with associated constants $C_{\mathcal{N}}, M_{\mathcal{N}}, L_{\mathcal{N}}$ in the sense of Subsection 2.2.

A standard example for a uniform *ALS* on \mathcal{C}_e (of dimension n) is given as follows: for each K in \mathcal{C}_e choose the standard *ALS* on K (of dimension n) so that N_v is as in (2.5) for each v in M_K . For this system we may choose $C_{\mathcal{N}} = 1$, $M_{\mathcal{N}} = 2n + 2$ and $L_{\mathcal{N}} = 2\pi\sqrt{2n + 1}$. Choosing l^2 -norms

at all infinite places and N_v as in (2.5) for all finite places yields another important uniform *ALS*.

2.5. Adelic-Lipschitz heights on $\mathbb{P}^n(k; e)$. Let \mathcal{C} be a collection of finite extensions of k and let \mathcal{N} be an *ALS* of dimension n on \mathcal{C} . Now we can define heights on $\mathbb{P}^n(K/k)$ (the set of points $P \in \mathbb{P}^n(K)$ with $k(P) = K$) for K in \mathcal{C} . Let $P \in \mathbb{P}^n(K/k)$. According to Subsection 2.3 we know that $H_{\mathcal{N}_K}(\cdot)$ defines a projective height on $\mathbb{P}^n(K)$. Now we define

$$(2.19) \quad H_{\mathcal{N}}(P) = H_{\mathcal{N}_K}(P).$$

From Subsection 2.3 we know

$$(2.20) \quad H_{\mathcal{N}_K}(P) = \prod_{v \in M_K} N_v(\sigma_v(\boldsymbol{\alpha}))^{\frac{d_v}{d}}$$

for the functions N_v of \mathcal{N}_K and $[K : \mathbb{Q}] = d$, $[K_v : \mathbb{Q}_v] = d_v$. Starting with $\mathcal{C} = \mathcal{C}_e$ we get a height defined on $\mathbb{P}^n(k; e)$.

3. THE MAIN RESULT

Let \mathcal{N} be an *ALS* on \mathcal{C}_e of dimension n . Then $H_{\mathcal{N}}(\cdot)$ defines a height on $\mathbb{P}^n(k; e)$, the set of points $P = (\alpha_0 : \dots : \alpha_n)$ in $\mathbb{P}^n(\bar{k})$ with $[k(P) : k] = e$ where $k(P) = k(\dots, \alpha_i/\alpha_j, \dots)$ for $0 \leq i, j \leq n$, $\alpha_j \neq 0$. The associated counting function $Z_{\mathcal{N}}(\mathbb{P}^n(k; e), X)$ denotes the number of points P in $\mathbb{P}^n(k; e)$ with $H_{\mathcal{N}}(P) \leq X$. Assume \mathcal{N} is a uniform *ALS* on \mathcal{C}_e (of dimension n). Then due to Northcott and (2.18) $Z_{\mathcal{N}}(\mathbb{P}^n(k; e), X)$ is finite for all X in $[0, \infty)$. The Schanuel constant $S_K(n)$ is defined as follows

$$(3.1) \quad S_K(n) = \frac{h_K R_K}{w_K \zeta_K(n+1)} \left(\frac{2^{r_K} (2\pi)^{s_K}}{\sqrt{|\Delta_K|}} \right)^{n+1} (n+1)^{r_K + s_K - 1}.$$

Here h_K is the class number, R_K the regulator, w_K the number of roots of unity in K , ζ_K the Dedekind zeta-function of K , Δ_K the discriminant, r_K is the number of real embeddings of K and s_K is the number of pairs of distinct complex conjugate embeddings of K . Recall also the definition of $V_{\mathcal{N}_K}$ (see (2.17)). Now we define the sum

$$(3.2) \quad D_{\mathcal{N}} = D_{\mathcal{N}}(k, e, n) = \sum_K 2^{-r_K(n+1)} \pi^{-s_K(n+1)} V_{\mathcal{N}_K} S_K(n)$$

where the sum runs over all extensions of k with relative degree e . We will prove that the sum in (3.2) converges if n is large enough compared to e . It will be convenient to use Landau's O -notation. For non-negative real functions $f(X), g(X), h(X)$ we say that $f(X) = g(X) + O(h(X))$ as $X > X_0$ tends to infinity if there is a constant C_0 such that $|f(X) - g(X)| \leq C_0 h(X)$ for each $X > X_0$.

After all this we are ready to state the main result.

Theorem 3.1. *Let e, n be positive integers and k a number field of degree m . Suppose \mathcal{N} is a uniform adelic-Lipschitz system of dimension n on \mathcal{C}_e , the collection of all finite extensions of k of relative degree e , with associated constants $C_{\mathcal{N}}, M_{\mathcal{N}}$ and $L_{\mathcal{N}}$. Write*

$$(3.3) \quad A_{\mathcal{N}} = M_{\mathcal{N}}^{me} (C_{\mathcal{N}}(L_{\mathcal{N}} + 1))^{me(n+1)-1}.$$

Suppose that either $e = 1$ or

$$(3.4) \quad n > 5e/2 + 4 + 2/(me).$$

Then the sum in (3.2) converges and as $X > 0$ tends to infinity we have

$$Z_{\mathcal{N}}(\mathbb{P}^n(k; e), X) = D_{\mathcal{N}} X^{me(n+1)} + O(A_{\mathcal{N}} X^{me(n+1)-1} \mathfrak{L}),$$

where $\mathfrak{L} = \log \max\{2, 2C_{\mathcal{N}}X\}$ if $(me, n) = (1, 1)$ and $\mathfrak{L} = 1$ otherwise. The constant in O depends only on k, e and n .

In subsequent papers [24] and [25] we will explore some applications of Theorem 3.1. Here we are content with some immediate consequences. For $e = 1$ we recover a version of the Proposition in [9], which allows more general norms at the finite places (this generalization will be essential to deduce the results of [24]). Now choose the standard uniform *ALS* as described at the end of Subsection 2.4 so that $H_{\mathcal{N}}$ becomes the Weil height. Schanuel's Theorem implies $S_K(n) = D_{\mathcal{N}}(K, 1, n) = 2^{-r_K(n+1)} \pi^{-s_K(n+1)} V_{\mathcal{N}_K} S_K(n)$. We can verify

$$(3.5) \quad V_{\mathcal{N}_K} = 2^{r_K(n+1)} \pi^{s_K(n+1)}$$

directly by noting that $\Lambda_{\mathcal{N}}(\mathfrak{D}) = (\sigma\mathfrak{D})^{n+1}$ in (2.12), so that $\det \Lambda_{\mathcal{N}}(\mathfrak{D}) = (2^{-s_K} N \mathfrak{D} \sqrt{|\Delta_K|})^{n+1}$ (see [9] Lemma 5). Inserting the latter in definition (2.15) we get $V_{\mathcal{N}_K}^{fin} = 1$ and it is clear that $V_{\mathcal{N}_K}^{inf} = 2^{r_K(n+1)} \pi^{s_K(n+1)}$. Then (3.5) follows from $V_{\mathcal{N}_K} = V_{\mathcal{N}_K}^{inf} V_{\mathcal{N}_K}^{fin}$ and so we find Theorem 1.1 from the introduction. For $k = \mathbb{Q}$ and $e = 2$ we recover essentially Schmidt's theorem (1.3) but only for $n > 10$ while Schmidt does it for all $n \geq 3$ and even (in a modified form) for $n = 1, 2$. For $k = \mathbb{Q}$ and $e > 2$ we find Gao's result (1.4) but again with the stronger restriction $n > 5e/2 + 4 + 2/(me)$ instead of Gao's $n > e$.

It is likely that Theorem 3.1 is valid for $n > e$ instead of (3.4). Gao showed, at least for his definition of height (see also [22] Appendix B), that for $k = \mathbb{Q}$ the bound $n > e$ suffices. On the other hand Schmidt's lower bound in (1.2) implies that Theorem 3.1 cannot hold for $e > 1$ and $n < e$. However, there is a good possibility of obtaining the asymptotics for $e > 1$ and $n = 1$ using a kind of generalized Mahler measure.

4. PROOF OF THE MAIN RESULT

The major part of the work was already done in [23] where we proved estimates for $Z_{\mathcal{N}_K}(\mathbb{P}^n(K/k), X)$. These estimates will be essential to deduce Theorem 3.1.

4.1. Preliminaries. Let K be in \mathcal{C}_e . Then by definition $H_{\mathcal{N}}(P) = H_{\mathcal{N}_K}(P)$ for all P in $\mathbb{P}^n(K/k)$. Since

$$(4.1) \quad \mathbb{P}^n(k; e) = \bigcup_{K \in \mathcal{C}_e} \mathbb{P}^n(K/k)$$

where the right hand side is a disjoint union, we get

$$(4.2) \quad Z_{\mathcal{N}}(\mathbb{P}^n(k; e), X) = \sum_{K \in \mathcal{C}_e} Z_{\mathcal{N}_K}(\mathbb{P}^n(K/k), X).$$

To state the estimates for $Z_{\mathcal{N}_K}(\mathbb{P}^n(K/k), X)$ from [23] we are forced to introduce some more notation. For fields k, K with $k \subseteq K$ and $[K : k] = e$ we define

$$G(K/k) = \{[K_0 : k]; K_0 \text{ is a field with } k \subseteq K_0 \subsetneq K\}$$

if $k \neq K$, and we define

$$G(K/k) = \{1\}$$

if $k = K$. Clearly $|G(K/k)| \leq e$. Then for an integer $g \in G(K/k)$ we define

$$(4.3) \quad \delta_g(K/k) = \inf_{\alpha, \beta} \{H(1, \alpha, \beta); k(\alpha, \beta) = K, [k(\alpha) : k] = g\}$$

(which is ≥ 1) and

$$(4.4) \quad \mu_g = m(e - g)(n + 1) - 1.$$

In [23] the author proved the following result.

Theorem 4.1. *Let k, K be number fields with $k \subseteq K$ and $[K : k] = e$, $[k : \mathbb{Q}] = m$, $[K : \mathbb{Q}] = d$. Let \mathcal{N}_K be an adelic-Lipschitz system of dimension n on K with associated constants $C_{\mathcal{N}_K}, L_{\mathcal{N}_K}, M_{\mathcal{N}_K}$. Write*

$$(4.5) \quad A_{\mathcal{N}_K} = M_{\mathcal{N}_K}^d (C_{\mathcal{N}_K} (L_{\mathcal{N}_K} + 1))^{d(n+1)-1},$$

$$(4.6) \quad D_{\mathcal{N}_K} = 2^{-r_K(n+1)} \pi^{-s_K(n+1)} V_{\mathcal{N}_K} S_K(n),$$

$$(4.7) \quad B_{\mathcal{N}_K} = A_{\mathcal{N}_K} R_K h_K \sum_{g \in G(K/k)} \delta_g(K/k)^{-\mu_g}.$$

Then as $X > 0$ tends to infinity we have

$$Z_{\mathcal{N}_K}(\mathbb{P}^n(K/k), X) = D_{\mathcal{N}_K} X^{d(n+1)} + O(B_{\mathcal{N}_K} X^{d(n+1)-1} \mathfrak{L}),$$

where

$$\mathfrak{L} = \log \max\{2, 2C_{\mathcal{N}_K} X\} \text{ if } (n, d) = (1, 1) \text{ and } \mathfrak{L} = 1 \text{ otherwise}$$

and the implied constant in O depends only on n and d .

Thanks to (4.2) and Theorem 4.1 it suffices to show that $\sum D_{\mathcal{N}_K}$ and $\sum B_{\mathcal{N}_K}$ are convergent (here the sum runs over the same fields as in (3.2) and (4.2)).

We will also deal with $\delta(\cdot)$, a simplified version of $\delta_g(\cdot)$

$$\delta(K/k) = \inf_{\alpha} \{H(1, \alpha); K = k(\alpha)\}.$$

The quantity $\delta(K/\mathbb{Q})$ was already introduced by Roy and Thunder [12].

It will be extremely convenient to use Vinogradov's \ll, \gg -notation. The constants involved in \ll and \gg will depend only on k, n, e unless we indicate the dependence on additional parameters by an index.

The case $e = 1$ of Theorem 3.1 is already covered by Theorem 4.1 by choosing $K = k$. For the rest of this article we assume

$$e > 1.$$

For a non-zero ideal \mathfrak{A} in K let $D_{K/k}(\mathfrak{A})$ be the discriminant-ideal of \mathfrak{A} relative to k (for definitions see [10] p.212 or [7]) and write $D_{K/k}$ for $D_{K/k}(\mathcal{O}_K)$ where \mathcal{O}_K denotes the ring of integers in K . By assumption we have $\mathbb{Q} \subseteq k \subseteq K$ and hence by [10] (2.10) Korollar p.213

$$(4.8) \quad |\Delta_{K/\mathbb{Q}}| = |\Delta_{k/\mathbb{Q}}|^{[K:k]} N_{k/\mathbb{Q}}(D_{K/k})$$

where $N_{k/\mathbb{Q}}(a)$ denotes the absolute norm of an ideal $a \neq 0$ of the ring of integers \mathcal{O}_k , i.e. $N_{k/\mathbb{Q}}(a) = |\mathcal{O}_k/a|$. Let P be in $\mathbb{P}^n(K/k)$, so $K = k(P)$. We use a theorem of Silverman ([20] Theorem 2) with Silverman's S_F (for $F = k$) as the set of archimedean absolute values. Then Silverman's $L_F(\cdot)$ is simply the usual norm $N_{k/\mathbb{Q}}(\cdot)$. Hence we deduce

$$(4.9) \quad H(P)^m \geq \exp\left(-\frac{\delta_k \log e}{2(e-1)}\right) N_{k/\mathbb{Q}}(D_{K/k})^{\frac{1}{2e(e-1)}}$$

where δ_k is the number of archimedean places in M_k . Since Silverman uses not an absolute height but rather an "absolute height relative to k ", we had to take the m -th power on the left hand side of (4.9). Combining (4.8) and (4.9) yields

$$(4.10) \quad \begin{aligned} H(P) &\geq \exp\left(-\frac{\delta_k \log e}{2(e-1)m}\right) |\Delta_k|^{-\frac{1}{2(e-1)m}} |\Delta_K|^{\frac{1}{2e(e-1)m}} \\ &\gg |\Delta_K|^{\frac{1}{2e(e-1)m}}. \end{aligned}$$

Recalling the definitions of δ , δ_g and $G(K/k)$ and taking $P = (1 : \alpha_1 : \alpha_2)$ in $\mathbb{P}^2(K/k)$ we get

$$(4.11) \quad \delta_g(K/k) \gg |\Delta_K|^{\frac{1}{2e(e-1)m}}$$

for any $g \in G(K/k)$; and similarly

$$(4.12) \quad \delta(K/k) \gg |\Delta_K|^{\frac{1}{2e(e-1)m}}.$$

Here it might be worthwhile to point out that (4.10) can be used to prove a version of Theorem 4.1 where $B_{\mathcal{N}_K}$ is redefined in terms of the discriminants; namely

$$(4.13) \quad B_{\mathcal{N}_K} = A_{\mathcal{N}_K} R_K h_K \sum_{g \in G(K/k)} (|\Delta_k|^{-e} |\Delta_K|)^{-\frac{\mu g}{2e(e-1)m}}.$$

At a first glance this error term looks more appropriate due to the unavoidable appearance of Δ_K in the main term. But as it turns out, the summation over Δ_K instead of over $\delta_g(K/k)$ leads to a result weaker than Theorem 3.1, in which we have to assume that n exceeds some quadratic function of e instead of (3.4). The reason for this is, that we have estimates for the number of number fields K with $\delta_g(K/k) \leq T$ which are more accurate than the best available estimates for the number of number fields with $|\Delta_K| \leq T$, see Section 5 for a discussion on this. Thanks to the well-known Theorem of Siegel-Brauer ([7] p.328 Corollary or [19] p.67 Satz 1 for a more precise version) we can use the inequalities (4.11) and (4.12) to bound the product of regulator and class number. More precisely we have

$$(4.14) \quad R_K h_K \ll_{\beta} \delta_g(K/k)^{\beta}$$

and

$$(4.15) \quad R_K h_K \ll_{\beta} \delta(K/k)^{\beta}.$$

for any $\beta > e(e-1)m$ and any $g \in G(K/k)$.

4.2. Three preparatory lemmas. We start with a very simple argument, known as dyadic summation. Since it will be frequently used we state it as a lemma.

Lemma 4.1 (Dyadic summation). *Let \mathcal{C} be a non-empty subset of \mathcal{C}_e and let ι be a map $\iota : \mathcal{C} \rightarrow [1, \infty)$. Write $N_{\iota}(T) = |\{K \in \mathcal{C}; \iota(K) \leq T\}|$ and suppose there are nonnegative real numbers b, c (independent of T) with*

$$N_{\iota}(T) \leq cT^b$$

for every $T > 0$. Let \mathcal{C}' be a non-empty subset of \mathcal{C} . Set $\mathfrak{M} = \lceil \log_2 \max_{\mathcal{C}'} \iota(K) \rceil + 1$ if \mathcal{C}' is finite and $\mathfrak{M} = \infty$ otherwise. Moreover suppose α is a real number such that $\sum_{i=1}^{\mathfrak{M}} 2^{i(\alpha+b)}$ converges. Then we have

$$\sum_{K \in \mathcal{C}'} \iota(K)^{\alpha} \leq c2^{|\alpha|} \sum_{i=1}^{\mathfrak{M}} 2^{i(\alpha+b)}.$$

Proof. From the definition of \mathfrak{M} and since $\mathcal{C}' \subseteq \mathcal{C}$ we have

$$\sum_{K \in \mathcal{C}'} \iota(K)^\alpha = \sum_{i=1}^{\mathfrak{M}} \sum_{\substack{K \in \mathcal{C}' \\ 2^{i-1} \leq \iota(K) < 2^i}} \iota(K)^\alpha \leq \sum_{i=1}^{\mathfrak{M}} \sum_{\substack{K \in \mathcal{C} \\ 2^{i-1} \leq \iota(K) < 2^i}} \iota(K)^\alpha.$$

First suppose $\alpha < 0$. Then the latter is

$$\leq \sum_{i=1}^{\mathfrak{M}} 2^{(i-1)\alpha} N_\iota(2^i) \leq c 2^{-\alpha} \sum_{i=1}^{\mathfrak{M}} 2^{i(\alpha+b)}.$$

If $\alpha \geq 0$ then we even get

$$\sum_{K \in \mathcal{C}'} \iota(K)^\alpha \leq c \sum_{i=1}^{\mathfrak{M}} 2^{i(\alpha+b)}.$$

This proves the lemma. \square

Recall the definition of $G(K/k)$ from Subsection 4.1. In our applications ι will be δ_g and \mathcal{C} will be

$$\mathcal{C}_e^{(g)} = \{K \in \mathcal{C}_e; g \in G(K/k)\}$$

the set of extensions K of k of relative degree e containing an intermediate field $K_0 \subsetneq K$ with $[K_0 : k] = g$. Let G_u be the union of all sets $G(K/k)$

$$G_u = \bigcup_{K \in \mathcal{C}_e} G(K/k),$$

so that $\mathcal{C}_e^{(g)}$ is non-empty if and only if $g \in G_u$. In fact G_u is simply the set of positive, proper divisors of e but we need only

$$\{1\} \subseteq G_u \subseteq \{1, \dots, [e/2]\}.$$

To apply the dyadic summation lemma we need information about the growth rate of $N_{\delta_g}(T)$. In accordance with the notation in Lemma 4.1 we define for an integer $g \in G_u$ and real positive T

$$N_{\delta_g}(T) = |\{K \in \mathcal{C}_e^{(g)}; \delta_g(K/k) \leq T\}|.$$

The set on the right-hand side is finite. More precisely we have the following lemma.

Lemma 4.2. *Set $\gamma_g = m(g^2 + g + e^2/g + e)$. Then for real positive T and g in G_u we have*

$$N_{\delta_g}(T) \ll T^{\gamma_g}.$$

Proof. Since $H(1, \alpha_1, \alpha_2) \geq \max\{H(1, \alpha_1), H(1, \alpha_2)\}$ it suffices to show that the number of tuples $(\alpha_1, \alpha_2) \in \bar{k}^2$ with

$$(4.16) \quad [k(\alpha_1) : k] = g$$

$$(4.17) \quad [k(\alpha_1, \alpha_2) : k(\alpha_1)] = e/g$$

$$(4.18) \quad H(1, \alpha_1), H(1, \alpha_2) \leq T$$

is $\ll T^{\gamma g}$. The number of projective points in $\mathbb{P}(k; g)$ with height not exceeding T is an upper bound for the number of α_1 in \bar{k} of relative degree g with $H(1, \alpha_1) \leq T$. Thus by (1.2) we get the upper bound

$$(4.19) \quad \ll T^{mg(g+1)}$$

for the number of α_1 . Next for each α_1 we count the number of α_2 . By (4.17) we have $[k(\alpha_1, \alpha_2) : k(\alpha_1)] = e/g$ and moreover $H(1, \alpha_2) \leq T$. Applying (1.2) (note that the constant $C(k, e, n)$ in (1.2) depends only on $[k : \mathbb{Q}]$, e, n) once more yields the upper bound

$$(4.20) \quad \ll T^{[k(\alpha_1) : \mathbb{Q}](e/g)(e/g+1)} = T^{me(e/g+1)}$$

for the number of α_2 provided α_1 is fixed. Multiplying the bound (4.19) for the number of α_1 and (4.20) gives the upper bound

$$\ll T^{m(g^2+g+e^2/g+e)}$$

for the number of tuples (α_1, α_2) and thereby proves the lemma. \square

Recall that $\delta_1 = \delta$ and that $N_\delta(T)$ denotes the number of number fields K in \bar{k} of relative degree e with $\delta(K/k) \leq T$. So Lemma 4.2 with $g = 1$ yields an upper bound for the growth rate of $N_\delta(T)$ but applying (1.2) directly gives a slightly better result.

Lemma 4.3. *Set $\gamma = me(e+1)$ and let $C_\delta = C(k, e, 1)$ be as in (1.2). Then for $T > 0$ we have*

$$(4.21) \quad N_\delta(T) \leq C_\delta T^\gamma.$$

Proof. The number of points in $\mathbb{P}(k; e)$ with height not larger than T is clearly an upper bound for $N_\delta(T)$. Thus the lemma follows from the upper bound in (1.2). \square

In fact Lemma 4.2 would suffice to prove the full Theorem 3.1, so one could omit Lemma 4.3. We did not because γ looks nicer than γ_1 and the proof above is essentially simply a reference.

4.3. Proof of Theorem 3.1. Recall the definition of $B_{\mathcal{N}_K}$ and $D_{\mathcal{N}_K}$ from (4.7) and (4.6). We have seen that it suffices to show that $\sum B_{\mathcal{N}_K}$ and $\sum D_{\mathcal{N}_K}$ are convergent where the sums run over all fields in \mathcal{C}_e .

Since \mathcal{N} is a uniform ALS on \mathcal{C}_e with associated constants $C_{\mathcal{N}}, M_{\mathcal{N}}$ and $L_{\mathcal{N}}$ we can assume that

$$(4.22) \quad C_{\mathcal{N}_K} \leq C_{\mathcal{N}},$$

$$(4.23) \quad M_{\mathcal{N}_K} \leq M_{\mathcal{N}},$$

$$(4.24) \quad L_{\mathcal{N}_K} \leq L_{\mathcal{N}}.$$

Hence by definition (3.3) and (4.5)

$$(4.25) \quad A_{\mathcal{N}_K} \leq A_{\mathcal{N}}.$$

Let us now prove that $\sum_K B_{\mathcal{N}_K}$ converges. We set $\beta = e(e-1)m + 1/8$. Using (4.14) and (4.25) we get

$$\sum_{K \in \mathcal{C}_e} B_{\mathcal{N}_K} \ll A_{\mathcal{N}} \sum_{K \in \mathcal{C}_e} \sum_{g \in G(K/k)} \delta_g(K/k)^{\beta - \mu_g}.$$

Recall that $G_u = \bigcup_{\mathcal{C}_e} G(K/k)$. So the term on the right-hand side above is

$$(4.26) \quad \begin{aligned} & A_{\mathcal{N}} \sum_{g \in G_u} \sum_{\substack{K \in \mathcal{C}_e \\ g \in G(K/k)}} \delta_g(K/k)^{\beta - \mu_g} \\ &= A_{\mathcal{N}} \sum_{g \in G_u} \sum_{K \in \mathcal{C}_e^{(g)}} \delta_g(K/k)^{\beta - \mu_g} \end{aligned}$$

provided the sum converges. This will be verified in a moment (see (4.27)). Applying the dyadic summation lemma with $\iota = \delta_g$ and $b = \gamma_g$ from Lemma 4.2 we see that the latter is

$$\ll A_{\mathcal{N}} \sum_{g \in G_u} \sum_{i=1}^{\infty} 2^{i(\gamma_g + \beta - \mu_g)}.$$

The next lemma will tell us that $\gamma_g + \beta - \mu_g \leq -1/8$. Assuming this for a moment we see that the inner sum above is $\ll 1$. Thus we derive the upper bound

$$(4.27) \quad \ll A_{\mathcal{N}} \sum_{g \in G_u} 1 \ll A_{\mathcal{N}},$$

confirming that the whole sum in (4.26) converges. This verifies the convergence of $\sum_K B_{\mathcal{N}_K}$ under the hypothesis $\gamma_g + \beta - \mu_g \leq -1/8$ for all $g \in G_u$. The following lemma shows that this hypothesis holds true. Recall that we assume $e > 1$ and therefore by our assumption in Theorem 3.1 $n > 5e/2 + 4 + 2/(me)$.

Lemma 4.4. *Let g be in G_u . Then*

$$(4.28) \quad \gamma_g + \beta - \mu_g \leq -\frac{1}{8}.$$

Proof. Recall that $G_u \subseteq \{1, \dots, [e/2]\}$ and $\mu_g = m(e-g)(n+1) - 1$. Write

$$F(g) = \frac{1}{m(e-g)}(\gamma_g + \beta + 1).$$

So (4.28) claims that $m(e-g)(F(g) - (n+1)) \leq -1/8$ for all $g \in G_u$. Hence it suffices to show that

$$F(g) - (n+1) \leq -\frac{1}{4me}$$

for $1 \leq g \leq e/2$. By definition

$$F(g) = \frac{g^2 + g + e^2/g + e}{e - g} + \frac{e(e - 1)}{e - g} + \frac{1 + 1/8}{m(e - g)}.$$

We claim that the second derivative $F''(g)$ is positive for $1 \leq g \leq e/2$. One finds

$$\begin{aligned} F''(g) &= \frac{2(e^2/g^3 + 1)(e - g) + 2(2g + 1 - e^2/g^2)}{(e - g)^2} + \frac{2e(e - 1)}{(e - g)^3} \\ &\quad + \frac{2(g^2 + g + e^2/g + e)}{(e - g)^3} + \frac{2(1 + 1/8)}{m(e - g)^3}. \end{aligned}$$

For $1 \leq g \leq e/2$ the last three fractions are certainly positive and so we may focus on the numerator of the first fraction. Now if $2g + 1 - e^2/g^2 \geq 0$ the claim follows at once. If $2g + 1 - e^2/g^2 < 0$ it suffices to show that

$$(e^2/g^3 + 1)(e - g) \geq e^2/g^2 - 2g - 1.$$

With $u = e/g$ the latter is equivalent to $u^3 - u^2 + e - g \geq u^2 - 2g - 1$ and this is equivalent to $u^2(u - 2) + e + g + 1 \geq 0$, which is certainly true since $1 \leq g \leq e/2$ and therefore $2 \leq u \leq e$.

Thus we have shown that $F''(g) > 0$ for $1 \leq g \leq e/2$ so that F is here concave. It suffices to prove $F(g) - (n + 1) \leq -1/(4me)$ for $g = 1$, $g = e/2$. First we use a simple arithmetic argument. Since n is an integer and $n > E = 5e/2 + 4 + 2/(me)$ with denominator dividing $2me$ we see that

$$(4.29) \quad n + 1 \geq E + 1 + 1/(2me).$$

Now $F(e/2) = 5e/2 + 5 + 2/(me) + 1/(4me) = E + 1 + 1/(4me)$ and thus

$$F(e/2) - (n + 1) \leq 1/(4me) - 1/(2me) = -1/(4me).$$

Finally

$$F(1) = 2e + 2 + 4/(e - 1) + 9/(8m(e - 1)).$$

Using (4.29) again yields

$$(4.30) \quad F(1) - (n + 1) \leq \frac{4}{e - 1} + \frac{9}{8m(e - 1)} - \frac{e}{2} - 3 - \frac{2}{me} - \frac{1}{2me}.$$

First suppose $e = 2$. Then (4.30) says $F(1) - (n + 1) \leq -1/(8m) = -1/(4em)$. Next suppose $e > 2$. Then the right-hand side in (4.30) is $\leq 4/2 + 9/(16m) - e/2 - 3 - 5/(2me) < -5/(2me) < -1/(4em)$. This completes the proof of the lemma. \square

To show convergence for $\sum_K D_{\mathcal{N}_K}$ we may use similar arguments but here we use only $\delta = \delta_1$ instead of δ_g . Let $d = me$ so that $[K : \mathbb{Q}] = d$. To estimate $V_{\mathcal{N}_K}$ in (4.6) recall that $V_{\mathcal{N}_K} = V_{\mathcal{N}_K}^{inf} V_{\mathcal{N}_K}^{fin}$. By (2.16) we have

$$V_{\mathcal{N}_K}^{inf} \ll (C_{\mathcal{N}_K}^{inf})^{d(n+1)}.$$

To estimate $V_{\mathcal{N}_K}^{fin}$ we define the non-zero ideal \mathfrak{C}_0 by

$$(4.31) \quad \mathfrak{C}_0 = \prod_{v \nmid \infty} \mathfrak{p}_v^{-\frac{d_v \log c_v}{\log N \mathfrak{p}_v}}$$

with c_v as in (2.7). Thus $|\mathfrak{C}_0|_v = c_v$ and

$$(4.32) \quad N\mathfrak{C}_0 = (C_{\mathcal{N}_K}^{fin})^d.$$

Let $\mathfrak{D} \neq 0$ be a fractional ideal. Clearly $|\alpha|_v \leq |\mathfrak{C}_0^{-1}\mathfrak{D}|_v$ for all non-archimedean v is equivalent to $\alpha \in \mathfrak{C}_0^{-1}\mathfrak{D}$. By (2.6) we conclude

$$(4.33) \quad \Lambda_{\mathcal{N}_K}(\mathfrak{D}) \subseteq \sigma(\mathfrak{C}_0^{-1}\mathfrak{D})^{n+1}$$

(where σ is given by (2.11)) and therefore

$$\det \Lambda_{\mathcal{N}_K}(\mathfrak{D}) \geq \det \sigma(\mathfrak{C}_0^{-1}\mathfrak{D})^{n+1}.$$

It is well-known (see [10] p.33 (5.2) Satz) that

$$\det \sigma(\mathfrak{C}_0^{-1}\mathfrak{D}) = 2^{-s_K} \sqrt{|\Delta_K|} N(\mathfrak{D}) N(\mathfrak{C}_0)^{-1},$$

where s_K is the number of pairs of complex conjugate embeddings of K . Combining the latter with (4.32) we see that

$$\det \sigma(\mathfrak{C}_0^{-1}\mathfrak{D})^{n+1} = 2^{-s_K(n+1)} |\Delta_K|^{(n+1)/2} N\mathfrak{D}^{n+1} (C_{\mathcal{N}_K}^{fin})^{-d(n+1)}.$$

Inserting the latter in definition (2.15) yields

$$V_{\mathcal{N}_K}^{fin} \ll (C_{\mathcal{N}_K}^{fin})^{d(n+1)}.$$

Now on recalling that $C_{\mathcal{N}_K} = C_{\mathcal{N}_K}^{inf} C_{\mathcal{N}_K}^{fin}$ and using (4.22) we conclude

$$V_{\mathcal{N}_K} \ll C_{\mathcal{N}_K}^{d(n+1)} \leq C_{\mathcal{N}}^{d(n+1)}.$$

The number of roots of unity w_K in (3.1) is at least 2. Furthermore $\zeta_K(n+1) > 1$. Hence $S_K(n) \ll R_K h_K |\Delta_K|^{-\frac{n+1}{2}}$. This together with the above estimate for $V_{\mathcal{N}_K}$ implies $D_{\mathcal{N}_K} \ll C_{\mathcal{N}}^{d(n+1)} R_K h_K |\Delta_K|^{-\frac{n+1}{2}}$ and since by Siegel-Brauer $R_K h_K \ll_\epsilon |\Delta_K|^{\frac{1}{2}+\epsilon}$ for any positive ϵ we get

$$(4.34) \quad D_{\mathcal{N}_K} \ll_\epsilon C_{\mathcal{N}}^{d(n+1)} |\Delta_K|^{-\frac{n}{2}+\epsilon}.$$

Remark 3. Let $N_\Delta(T) = |\{K \in \mathcal{C}_e; |\Delta_K| \leq T\}|$. Schmidt [18] showed

$$(4.35) \quad N_\Delta(T) \ll T^{\frac{e+2}{4}}.$$

Thus we could apply the dyadic summation lemma with $\iota = |\Delta_K|$ and $b = (e+2)/4$ to see that $\sum_K D_{\mathcal{N}_K}$ converges for $n > e/2 + 1$.

Instead of using Schmidt's bound (4.35) we will prove a lower bound for $|\Delta_K|$ in terms of $\delta(K/k)$ which might be of interest for its own sake. Then we can apply Lemma 4.3 instead of (4.35).

Lemma 4.5. *We have*

$$(4.36) \quad \delta(K/k) \leq \delta(K/\mathbb{Q}) \ll |\Delta_K|^{\frac{1}{d}}.$$

Proof. The lemma is trivially true for $K = k = \mathbb{Q}$. However we have by assumption $e \geq 2$ and so $[K : \mathbb{Q}] = em \geq 2$. The first inequality follows immediately from the definition. Let σ be as in (2.10) and suppose α is a non-zero integer of K . One gets

$$\begin{aligned}
H(1, \alpha) &= \prod_{i=1}^d \max\{1, |\sigma_i(\alpha)|\}^{1/d} \\
&\leq \max\{1, \max_{1 \leq i \leq d} |\sigma_i(\alpha)|\} \\
(4.37) \quad &\leq |\sigma(\alpha)|
\end{aligned}$$

because $\prod_{1 \leq i \leq d} |\sigma_i(\alpha)| \geq 1$. Let $v_1 = \sigma(\alpha_1), \dots, v_d = \sigma(\alpha_d)$ be linearly independent vectors of the lattice $\sigma\mathcal{O}_K$ with $|v_i| = \lambda_i$ for the successive minima λ_i ($i = 1, \dots, [K : \mathbb{Q}] = d$). Let us temporarily denote by b the maximum of the degrees of the proper subfields of K . Therefore $K = \mathbb{Q}(\alpha_1, \dots, \alpha_{b+1})$. Next we need to construct a primitive element in \mathcal{O}_K with small height. A standard argument (see [8] p.244 or Lemma 3.3 in [23]) yields a primitive $\alpha = \sum_{j=1}^{b+1} m_j \alpha_j$ with rational integers $0 \leq m_j < e$. Hence by (4.37) we get

$$H(1, \alpha) \leq |\sigma(\sum_{j=1}^{b+1} m_j \alpha_j)| \leq \sum_{j=1}^{b+1} m_j |\sigma(\alpha_j)| \ll \lambda_{b+1}.$$

We shall estimate λ_{b+1} :

$$\begin{aligned}
\lambda_{b+1} &= \left(\frac{\lambda_1 \dots \lambda_b \lambda_{b+1}^{d-b}}{\lambda_1 \dots \lambda_b} \right)^{\frac{1}{d-b}} \\
&\leq \left(\frac{\lambda_1 \dots \lambda_d}{\lambda_1 \dots \lambda_b} \right)^{\frac{1}{d-b}} \\
&\ll \left(\frac{\det(\sigma\mathcal{O}_K)}{\lambda_1 \dots \lambda_b} \right)^{\frac{1}{d-b}} \\
&= \left(\frac{|\Delta_K|^{\frac{1}{2}}}{2^{s_K} \lambda_1 \dots \lambda_b} \right)^{\frac{1}{d-b}} \\
&\ll |\Delta_K|^{\frac{1}{2(d-b)}}
\end{aligned}$$

where we used that $\lambda_1 = |\sigma(\alpha_1)| \geq H(1, \alpha_1) \geq 1$. So all this together implies

$$(4.38) \quad \delta(K/\mathbb{Q}) \ll |\Delta_K|^{\frac{1}{2(d-b)}}.$$

Now b is the degree of a proper subfield. Thus $b \leq d/2$ and we get (4.36). \square

Using Lemma 4.5 and (4.34) with ϵ replaced by ϵ/d we deduce

$$(4.39) \quad D_{\mathcal{N}_K} \ll_{\epsilon} C_{\mathcal{N}}^{d(n+1)} \delta(K/k)^{-\frac{dn}{2} + \epsilon}$$

for any positive ϵ . Choosing $\epsilon = 1/2$ we get

$$D_{\mathcal{N}_K} \ll C_{\mathcal{N}}^{d(n+1)} \delta(K/k)^{-\frac{dn}{2} + \frac{1}{2}}.$$

Applying the dyadic summation lemma with $\iota = \delta$ and $b = \gamma$ from Lemma 4.3 we conclude

$$\begin{aligned} \sum_{K \in \mathcal{C}_e} D_{\mathcal{N}_K} &\ll C_{\mathcal{N}}^{d(n+1)} \sum_{K \in \mathcal{C}_e} \delta(K/k)^{-\frac{dn}{2} + \frac{1}{2}} \\ &\ll C_{\mathcal{N}}^{d(n+1)} \sum_{i=1}^{\infty} 2^{(-\frac{dn}{2} + \frac{1}{2} + \gamma)i} \\ &\ll C_{\mathcal{N}}^{d(n+1)} \end{aligned}$$

provided $-\frac{dn}{2} + \frac{1}{2} + \gamma < 0$, which is equivalent to $n > 2e + 2 + 1/d$. But the latter holds since $n > 5e/2 + 4 + 2/(me)$. This completes the proof of Theorem 3.1.

5. COUNTING NUMBER FIELDS

Using results of the previous section we give simple lower bounds for the growth rate of $N_{\delta}(T)$ and $N_{\Delta}(T)$, the number of field extensions K/k of degree e with $\delta(K/k) \leq T$ or $|\Delta_K| \leq T$. The following corollary shows that the estimates for $N_{\delta}(T)$ are more precise than those available for $N_{\Delta}(T)$. Recall that $e > 1$.

Corollary 5.1. *With $c = c(k, e, 1)$, $C = C(k, e, 1)$ and $X_0(k, e, 1)$ from (1.2) set*

$$c_{\delta} = 2^{-5em-22}c, \quad C_{\delta} = C \text{ and } T_0 = X_0(k, e, 1).$$

Then we have

$$c_{\delta} T^{me(e-1)} \leq N_{\delta}(T) \leq C_{\delta} T^{me(e+1)}$$

where the upper bounds holds for $T > 0$ and the lower bound holds for $T \geq T_0$.

Proof. From the definition it is clear that $Z_H(\mathbb{P}(K/k), T) > 0$ if and only if $\delta(K/k) \leq T$. Therefore we have

$$(5.1) \quad N_{\delta}(T) = \sum_{\substack{K \in \mathcal{C}_e \\ \delta(K/k) \leq T}} 1 = \sum_{\substack{K \in \mathcal{C}_e \\ \delta(K/k) \leq T}} \frac{Z_H(\mathbb{P}(K/k), T)}{Z_H(\mathbb{P}(K/k), T)}.$$

Using the equivalence above once again, we see that the term on the far right-hand side of (5.1) is

$$\begin{aligned} &\geq \left(\sup_{K \in \mathcal{C}_e} \{Z_H(\mathbb{P}(K/k), T)\} \right)^{-1} \sum_{\substack{K \in \mathcal{C}_e \\ \delta(K/k) \leq T}} Z_H(\mathbb{P}(K/k), T) \\ &= \left(\sup_{K \in \mathcal{C}_e} \{Z_H(\mathbb{P}(K/k), T)\} \right)^{-1} \sum_{K \in \mathcal{C}_e} Z_H(\mathbb{P}(K/k), T) \\ &= \left(\sup_{K \in \mathcal{C}_e} \{Z_H(\mathbb{P}(K/k), T)\} \right)^{-1} Z_H(\mathbb{P}(k; e), T). \end{aligned}$$

Now $Z_H(\mathbb{P}(K/k), T) \leq Z_H(\mathbb{P}(K; 1), T)$ and by the upper bound in (1.2) and recalling that $[K : \mathbb{Q}] = em$ we get

$$Z_H(\mathbb{P}(K; 1), T) \leq C(K, 1, 1)T^{2me} = 2^{5em+22}T^{2me}.$$

The lower bound in (1.2) with $c = c(k, e, 1)$ yields

$$Z_H(\mathbb{P}(k; e), T) \geq cT^{me(e+1)}$$

for $T \geq X_0(k, e, 1) = T_0$. Hence

$$N_\delta(T) \geq (2^{5em+22}T^{2me})^{-1}cT^{me(e+1)} = c_\delta T^{me(e-1)}$$

for $T \geq T_0$. On the other hand Lemma 4.3 tells us that

$$N_\delta(T) \leq C_\delta T^{me(e+1)}$$

for $T > 0$. □

Corollary 5.1 combined with the lower bound (4.12) for δ in terms of $|\Delta_K|$ yields

Corollary 5.2. *There are positive constants $c_\Delta = c_\Delta(k, e)$ and $T_1 = T_1(k, e)$ depending only on k, e such that*

$$N_\Delta(T) \geq c_\Delta T^{1/2}$$

for $T \geq T_1$.

Proof. From (4.12) we know that there is a positive constant $c_1 = c_1(k, e)$ depending only on $k, e > 1$ such that $\delta(K/k) \geq c_1 |\Delta_K|^{1/(2e(e-1)m)}$. Using Corollary 5.1 and setting $c_\Delta = c_\delta c_1^{me(e-1)}$, $T_1 = (T_0/c_1)^{2e(e-1)m}$ we conclude

$$N_\Delta(T) \geq N_\delta(c_1 T^{1/(2e(e-1)m)}) \geq c_\Delta T^{1/2}$$

provided $T \geq T_1$. □

Ellenberg and Venkatesh's Theorem 1.1 in [4] shows that the exponent $1/2$ in Corollary 5.2 can be replaced by $1/2 + 1/e^2$ and according to Linnik's Conjecture (see [4] p.723) the correct exponent is 1. Although the general Linnik Conjecture is known to be true only for $e \leq 3$ the exponent $1/2$ can always be increased to 1 if e is even or a multiple of 3 (see [4] pp. 723,724).

What about upper bounds for $N_\Delta(T)$? From (4.36) we know that there is a positive constant $c_2 = c_2(d)$ depending only on $d = em$ such that

$$\delta(K/k) \leq c_2 |\Delta_K|^{\frac{1}{em}}.$$

Thus we get

$$N_\Delta(T) \leq N_\delta(c_2 T^{\frac{1}{em}}) \leq C_\delta c_2^{me(e+1)} T^{e+1}$$

for $T > 0$. But Schmidt's bound (4.35) has the much better exponent $(e+2)/4$ on T .

REFERENCES

- [1] E. BOMBIERI AND W. GUBLER, *Heights in Diophantine Geometry*, Cambridge University Press, 2006.
- [2] S. BOSCH, U. GÜNTZER, AND R. REMMERT, *Non-Archimedean Analysis*, Springer, 1984.
- [3] S.-J. CHERN AND J. D. VAALER, *The distribution of values of Mahler's measure*, J. reine angew. Math., 540 (2001), pp. 1–47.
- [4] J. ELLENBERG AND A. VENKATESH, *The number of extensions of a number field with fixed degree and bounded discriminant*, Ann. of Math., 163 (2006), pp. 723–741.
- [5] X. GAO, *On Northcott's Theorem*, Ph.D. Thesis, University of Colorado, (1995).
- [6] S. LANG, *Fundamentals of Diophantine Geometry*, Springer, 1983.
- [7] ———, *Algebraic Number Theory*, Springer, 1994.
- [8] ———, *Algebra*, Springer, 2002.
- [9] D. W. MASSER AND J. D. VAALER, *Counting algebraic numbers with large height II*, Trans. Amer. Math. Soc., 359 (2007), pp. 427–445.
- [10] J. NEUKIRCH, *Algebraische Zahlentheorie*, Springer, 1992.
- [11] D. G. NORTHCOTT, *An inequality in the theory of arithmetic on algebraic varieties*, Proc. Cambridge Phil. Soc., 45 (1949), pp. 502–509 and 510–518.
- [12] D. ROY AND J. L. THUNDER, *A note on Siegel's lemma over number fields*, Monatsh. Math., 120 (1995), pp. 307–318.
- [13] ———, *An absolute Siegel's lemma*, J. reine angew. Math., 476 (1996), pp. 1–26.
- [14] S. H. SCHANUEL, *Heights in number fields*, Bull. Soc. Math. France, 107 (1979), pp. 433–449.
- [15] W. M. SCHMIDT, *On heights of algebraic subspaces and diophantine approximations*, Ann. of Math., 85 (1967), pp. 430–472.
- [16] ———, *Northcott's Theorem on heights I. A general estimate*, Monatsh. Math., 115 (1993), pp. 169–183.
- [17] ———, *Northcott's Theorem on heights II. The quadratic case*, Acta Arith., 70 (1995), pp. 343–375.
- [18] ———, *Number fields of given degree and bounded discriminant*, Astérisque, 228 (1995), pp. 189–195.
- [19] C. L. SIEGEL, *Gesammelte Abhandlungen Band IV*, Springer, 1979.
- [20] J. SILVERMAN, *Lower bounds for height functions*, Duke Math. J., 51 (1984), pp. 395–403.
- [21] J. L. THUNDER, *An adelic Minkowski-Hlawka theorem and an application to Siegel's lemma*, J. reine angew. Math., 475 (1996), pp. 167–185.
- [22] M. WIDMER, *Asymptotically counting points of bounded height*, Ph.D. Thesis, Universität Basel, (2007).
- [23] ———, *Counting primitive points of bounded height*, submitted, (2008).
- [24] ———, *Counting points of fixed degree and bounded height on linear varieties*, submitted, (2009).
- [25] ———, *On number fields with nontrivial subfields*, preprint, (2009).

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS AT AUSTIN, 1 UNIVERSITY STATION C1200, AUSTIN, TEXAS 78712, U.S.A.

E-mail address: `widmer@math.utexas.edu`