

**Title:**

Bohler`s angle and the crucial angle of Gissane in paediatric population

**Authors:**

Bruno Pombo<sup>1</sup>; Ana Cristina Ferreira<sup>2</sup>; Luís Costa<sup>3</sup>

1: MSc, MD, Orthopaedics Department, Centro Hospitalar do Porto – Portugal

2: DPhil, Center of Mathematics, Minho University – Portugal

3 MSc, MD, Consultant, Foot and Ankle Unit, Orthopaedics Department, Centro Hospitalar do Porto – Portugal

**Corresponding author:**

Bruno Pombo

**Running Title:**

Calcaneus angles in children

**Funding:**

This study was not funded

**Conflict of Interest:**

Bruno Pombo, Ana Cristina Ferreira and Luís Costa declare that they have no conflict of interest

**Abstract:**

Bohler`s angle and the crucial angle of Gissane are used on the evaluation of calcaneus fractures. However, few authors have described the variation of the angles when the calcaneus is growing.

In this study, Bohler`s angle and the crucial angle of Gissane in paediatric population were measured using lateral foot radiographs of 429 patients, from 0 to 16 years of age. The control group was composed of 70 adult patients.

The sample had a mean Bohler`s angle of  $35.4 \pm 5.9^\circ$  and a mean crucial angle of Gissane of  $110.5 \pm 7.4^\circ$ . The greater mean difference was identified for Bohler`s angle (eight degrees) in the age group from 5 to 8 years of age.

( $39.6 \pm 5.7^\circ$ ) and for the crucial angle of Gissane (five to six degrees) in the age group from 0 to 4 years of age ( $115.8 \pm 7.3^\circ$ ) ( $p < 0.05$ ).

The influence of the ossification centres on the geometry of the calcaneus across age groups makes Bohler's angle and the crucial angle of Gissane higher in young children.

The increase of Bohler's angle points out the relative development of the posterior facet in young children and the importance of the reconstruction of the posterior facet height in the intra-articular calcaneus fractures.

**Keywords:**

Bohler's angle; Crucial angle of Gissane; Ossification centres; Paediatric population

**Level of Evidence:**

Diagnostic study; Level III

**Introduction:**

The calcaneus fractures in children are the most frequent in tarsal bones. The fracture incidence ranges between 0.05 and 0.15 % and has a peak between 8 and 12 years of age<sup>1</sup>. The aetiology is similar to that of adults.

The extra-articular fractures have an increased incidence in children and the intra-articular fractures in adolescents<sup>2</sup>.

The orthopaedic principles behind the treatment procedures of intra-articular fractures are the same as in adults<sup>4,5</sup>.

The subtalar joint congruence is achieved after the reconstruction of calcaneus angles - Bohler's angle and the crucial angle of Gissane. Specially for bilateral fractures, it is important to recognize its normal range, to determine the degree of deformity<sup>6</sup>, to provide a perfect reduction and evaluate the clinical outcome<sup>7</sup> of displaced, intra-articular calcaneus fractures.

The prenatal and postnatal growth of the calcaneus has been studied in the past<sup>8-15</sup>. The calcaneus is the only tarsal bone that presents two primary and a secondary ossification centres. The process of growth creates geometric variations in the calcaneus. A few studies have described the normal range of calcaneus angles in children and the variation across age groups<sup>16-18</sup>.

The purpose of this study was to characterize the normal range and the variation of Bohler's angle and the crucial angle of Gissane according to the postnatal growth of the calcaneus.

**Methods:**

The study was approved by the appropriate ethics committee in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki and its later amendments. Specific national laws were observed.

An observational and retrospective study was performed through Picture Archiving and Communication System (PACS) consultation at Unidade Local de Saúde do Alto Minho, Portugal. This health centre is composed of two hospitals in the North of Portugal serving a population of more than 200.000 inhabitants. The local population is predominantly Caucasian.

Between 2014 and 2015, 429 patients, from 0 to 16 years of age, were submitted to lateral foot radiographs in outpatient consultation. We defined calcaneal fracture, previous local surgery and inadequate imaging as exclusion criteria. Seventy adult patients, fulfilling the same criteria, were used as a control group.

**Measurement:**

The patients' age, sex and laterality were recorded. Bohler's angle and the crucial angle of Gissane were measured using an on-screen angle tool on PACS system. Bohler's angle is defined by the intersection of two semi-lines at the highest point of the posterior facet, one starting at the highest point of the tuberosity and another at the highest point of the anterior process (Figure 1 B). The crucial angle of Gissane is specified by the intersection of two semi-lines on calcaneus sulcus, one along the posterior facet and another starting on the anterior process (Figure 1 A). All images were evaluated and the angles were measured twice by one junior orthopaedic surgeon. The mean of these two measurements was recorded. All measurements were within a two degree range, indicating acceptable intra-observer measure reliability.

**Statistical analysis:**

The IBM SPSS Statistics for Windows, Version 24.0 Armonk, NY: IBM Corp. program was used for statistical analysis. We defined an error margin of 0.05.

Patients were divided into four groups according to their age (Group A: 0 to 4 years of age; Group B: 5 to 8 years of age; Group C: 9 to 12 years of age; Group D: 13 to 16 years of age). The sample and the control group were defined as Group S or Group E, respectively.

The normality of variables was assessed by graphic analysis and Shapiro Wilk test<sup>23</sup>. Chi-square test was used to evaluate sex and laterality differences between age groups. The independent samples T-test was applied to compare Bohler's angle and the crucial angle of Gissane between the sample and the control group.

The distribution of angles across age was estimated using curve-fitting regression analysis. For Bohler's angle, a cubic function defined by  $y = c_0 + a_1(t) + a_2(t^2) + a_3(t^3)$  was identified. In this model, y represents Bohler's angle in degrees and t the age in years. For the crucial angle of Gissane no model was adequate.

A one-way between groups analysis of variance was used to compare the angle difference between each group. T-testing with Tukey adjustments quantified the mean difference and percentage of variation between groups. With a large sample, even very small differences between groups can become statistically significant. This does not mean that these differences have any practical or theoretical significance. To evaluate this, the strength of relationship between the angles and the age groups was assessed by eta squared ( $\eta^2$ ) which represents the proportion of variance of the angle that is explained by the age groups. Values for  $\eta^2$  can range from 0 to 1. Strength of  $\eta^2$  values were defined as small, moderate and large<sup>24,25</sup>.

## Results:

Bohler's angle and the crucial angle of Gissane were measured in 429 children. The sample was mainly composed of two groups: 9 to 12 years of age (39%) and 13 to 16 years of age (31%). In the sample group, the mean Bohler's angle was  $35.4 \pm 5.9^\circ$  (95% CI: 23.6 – 47.2°) and the mean crucial angle of Gissane was  $110.5 \pm 7.4^\circ$  (95% CI: 95.7 – 125.3°). The difference to the control group regarding Bohler's angle (4°; 10%;  $p < .001$ ) and the crucial angle of Gissane (2°; 2%,  $p = .016$ ) was statistically significant. The distribution of sex ( $p = 0.716$ ) and laterality ( $p = 0.211$ ) by age groups was similar (Table 1).

There was a statistically significant difference in Bohler's angle [ $F(3, 425) = 29.3$ ,  $p < 0.001$ ] and in the crucial angle of Gissane [ $F(3, 425) = 6.8$ ,  $p < 0.001$ ] for the four age groups (Graph 2 and Graph 3). The effect size was large ( $\eta^2 = 0.17$ ) for Bohler's angle and small ( $\eta^2 = 0.05$ ) for the crucial angle of Gissane.

The distribution of Bohler's angle across age follows a cubic function of  $y = 20.43 + 7.11t - 0.93t^2 + 0.033t^3$  [ $F(3; 425) = 39.59$ ,  $R^2: 0.21$ ;  $p < 0.001$ ] (Graph 1). Post-hoc comparisons using the Tukey HSD test indicated that the mean Bohler's angle of Group B ( $39.7 \pm 5.7^\circ$ ) was significantly different from those of Group A ( $33.6 \pm 5.5$ ), Group

C ( $35.1 \pm 5.5^\circ$ ), Group D ( $33.0 \pm 5.0^\circ$ ) and control group ( $31.7 \pm 5.2^\circ$ ). Group C was significantly different from Group D and from the control group.

Regarding the crucial angle of Gissane, Group A ( $115.8 \pm 7.3^\circ$ ) was significantly different from Group B ( $111.1 \pm 7.5^\circ$ ), Group C ( $109.8 \pm 7.2^\circ$ ) and Group D ( $109.8 \pm 7.1^\circ$ ). Only Group C and Group D were significantly different from the control group. There was no difference between the other age groups (Table 2).

### **Discussion:**

Our study showed that the mean Bohler's angle is higher and the crucial angle of Gissane is lower in paediatric age than in adults. The Bohler's angle is more dependent of the age than the crucial angle of Gissane (17%) particularly in transition through 5 to 8 years of age.

Traditionally, calcaneal fractures in children were treated conservatively, but recent publications have drawn attention to the operative treatment of intra-articular fractures<sup>4,5</sup>. Bohler's angle and the crucial angle of Gissane have been used as a method of evaluation of intra-articular fractures of the calcaneus<sup>26</sup>.

In Boyle's et al study<sup>16</sup>, the mean of the crucial angle of Gissane in paediatric age was  $111^\circ$  ( $90 - 147^\circ$ ). There was little angle variation with the exception from 0 to 2 years of age that showed a statistically significant increase in this value compared with other groups. The measurement is unreliable in young children<sup>17</sup> not providing usefulness of this evaluation.

Three studies had measured Bohler's angle in paediatric age<sup>16-18</sup>. The measurement had reliability<sup>17</sup>. The angle rounded  $35^\circ$  ( $14 - 58^\circ$ ). They showed several statistically significant differences between smaller age groups, with interesting age-related variation through 6 to 8 years of age. The variation was justified by the postnatal growth of the foot.

In the first months of life, the primary ossification centre of the calcaneus is oval<sup>13,15</sup>, making the subtalar joint flatter (Figure 2 A). By 1 to 2 years of age, the primary ossification centre elongates and develops the proximal third and the subtalar joint<sup>13</sup>. The primary ossification centre is 62 % ossified at 3 months after birth but continues until 7 years of age<sup>13</sup>.

Between 5 and 7 years of age, one or more secondary ossification centres appear and initiate the development of the calcaneus apophysis<sup>13,15</sup>. They are initially located at the lower third of the calcaneus (Figure 2 B). Then they coalesce and elongate proximally and distally, forming a cap-like structure that surrounds the posterior margin of the

primary ossification centre (Figure 2 C). The ossification process of the posterior apophysis initiates in the middle of the ossification centre and ends on the superior and inferior extremities by the age of 14 to 18 years<sup>13,15</sup> (Figure 2 D, E).

In this study, the initial negative age-related variation of the crucial angle of Gissane in younger children is similar to other descriptions<sup>16</sup>. As we can see in Figures 2 A and 2 B, the development of subtalar joint makes the angle decrease in measure. This is particularly noticeable in the first 8 years of age. After this period, the calcaneus apophysis develops, which has little impact on the subtalar joint and consequently on the angle.

The cubic distribution of Bohler's angle is here described for the first time. We hypothesize that the initial increase, observed in the Graph 1, is justified by the asymmetric development of the posterior facet (Figure 2 B). When we compare Group B and the control group, the mean difference reaches a value of eight degrees (20 %) suggesting that in children with displaced, intra-articular calcaneus fractures, between 5 and 8 years of age, the surgeon should be alert of its upper relative position to reconstruct the posterior facet. Also, particular care should be taken to put the least possible number of screws in the tuberosity, until 7 years of age, to preserve the primary ossification centre that is incompletely ossified<sup>13</sup> and the secondary ossification centres that appear<sup>15</sup>.

Our study had some limitations. Firstly, the retrospective analysis of a sample mainly composed by Caucasian people may not represent the population of all children and can decrease the external validity of the conclusions. Secondly, the measurement by only one rater could bias the study. However, as the reliability of Bohler's angle was accomplished in other studies<sup>17</sup> the impact on the conclusions is residual. In addition, the wide confidence interval for Bohler's angle can create challenges in clinical prediction. It is conceivable that a large sample or the development of a multiple regression model may increase the strength of the conclusion.

### **Conclusion:**

The higher relative position of the posterior facet of the calcaneus makes Bohler's angle higher in children from 5 to 8 years of age and should make the surgeon take particular care in the reconstruction of the congruence of the subtalar joint.

## References:

1. Rammelt S, Godoy-Santos AL, Schneiders W, Fitze G, Zwipp H. Foot and ankle fractures during childhood: review of the literature and scientific evidence for appropriate treatment. *Revista brasileira de ortopedia*. 2016;51(6):630-639.
2. Schmidt TL, Weiner DS. Calcaneal fractures in children. An evaluation of the nature of the injury in 56 children. *Clinical orthopaedics and related research*. 1982(171):150-155.
3. Summers H, Ann Kramer P, Benirschke SK. Pediatric calcaneal fractures. *Orthopedic reviews*. 2009;1(1):e9.
4. Buckingham R, Jackson M, Atkins R. Calcaneal fractures in adolescents: CT classification and results of operative treatment. *Injury*. 2003;34(6):454-459.
5. Petit CJ, Lee BM, Kasser JR, Kocher MS. Operative treatment of intraarticular calcaneal fractures in the pediatric population. *Journal of Pediatric Orthopaedics*. 2007;27(8):856-862.
6. Brunet J. Calcaneal fractures in children: long-term results of treatment. *Bone & Joint Journal*. 2000;82(2):211-216.
7. Pickle A, Benaroch TE, Guy P, Harvey EJ. Clinical outcome of pediatric calcaneal fractures treated with open reduction and internal fixation. *Journal of Pediatric Orthopaedics*. 2004;24(2):178-180.
8. Harding VV. Time schedule for the appearance and fusion of a second accessory center of ossification of the calcaneus. *Child development*. 1952;23(3):180-184.
9. Gardner E, O'Rahilly R, Gray DJ. The Prenatal Development of the Skeleton and Joints of the Human Foot. *JBJS*. 1959;41(5):847-876.
10. Harris EJ. The relationship of the ossification centers of the talus and calcaneus to the developing bone. *Journal of the American Podiatry Association*. 1976;66(2):76-81.
11. Meyer DB, O'Rahilly R. The onset of ossification in the human calcaneus. *Anatomy and embryology*. 1976;150(1):19-33.
12. Agrawal P, Kulkarni DS, Atre PR. The participation of cartilage canals in the ossification of the human fetal calcaneum. *Journal of anatomy*. 1986;147:135-142.

13. Hubbard AM, Meyer JS, Davidson RS, Mahboubi S, Harty MP. Relationship between the ossification center and cartilaginous anlage in the normal hindfoot in children: study with MR imaging. *AJR American journal of roentgenology*. 1993;161(4):849-853.
14. Fritsch H, Brenner E, Debbage P. Ossification in the human calcaneus: a model for spatial bone development and ossification. *Journal of anatomy*. 2001;199(Pt 5):609-616.
15. Rossi I, Rosenberg Z, Zember J. Normal skeletal development and imaging pitfalls of the calcaneal apophysis: MRI features. *Skeletal radiology*. 2016;45(4):483-493.
16. Boyle MJ, Walker CG, Crawford HA. The paediatric Bohler's angle and crucial angle of Gissane: a case series. *Journal of orthopaedic surgery and research*. 2011;6:2.
17. Clint SA, Morris TP, Shaw OM, Oddy MJ, Rudge B, Barry M. The reliability and variation of measurements of the os calcis angles in children. *The Journal of bone and joint surgery British volume*. 2010;92(4):571-575.
18. Parikh SN, Weesner M, Welge J. Postnatal growth of the calcaneus does not simulate growth of the foot. *Journal of pediatric orthopedics*. 2012;32(1):93-99.
19. Seyahi A, Uludag S, Koyuncu LO, Atalar AC, Demirhan M. [The calcaneal angles in the Turkish population]. *Acta orthopaedica et traumatologica turcica*. 2009;43(5):406-411.
20. Sengodan VC, Amruth KH, Karthikeyan. Bohler's and Gissane Angles in the Indian Population. *Journal of clinical imaging science*. 2012;2:77.
21. Khoshhal KI, Ibrahim AF, Al-Nakshabandi NA, Zamzam MM, Al-Boukai AA, Zamzami MM. Bohler's and Gissane's angles of the calcaneus in the Saudi population. *Saudi medical journal*. 2004;25(12):1967-1970.
22. Willmott H, Stanton J, Southgate C. Bohler's angle - What is normal in the uninjured British population? *Foot and ankle surgery : official journal of the European Society of Foot and Ankle Surgeons*. 2012;18(3):187-189.
23. Ghasemi A, Zahediasl S. Normality Tests for Statistical Analysis: A Guide for Non-Statisticians. *International Journal of Endocrinology and Metabolism*. 2012;10(2):486-489.
24. Sink CA, Mvududu NH. Statistical Power, Sampling, and Effect Sizes: Three Keys to Research Relevancy. *Counseling Outcome Research and Evaluation*. 2010;1(2):1-18.
25. Cohen J. Statistical power analysis for the behavioral sciences. 2nd. In: Hillsdale, NJ: Erlbaum; 1988.



26. Isaacs JD, Baba M, Huang P, et al. The diagnostic accuracy of Bohler's angle in fractures of the calcaneus. *The Journal of emergency medicine*. 2013;45(6):879-884.

**Table 1: Mean Bohler`s angle and crucial angle of Gissane across different age groups**

Group	A	B	C	D	S	E
Age, yrs	0-4	5-8	9-12	13-16	0-16	≥ 18
n (%)	31 (7)	95 (22)	170 (39)	133 (31)	429 (100)	70
<b>Sex</b>						
Male, n (%)	17 (55)	43 (45)	83 (49)	58 (44)	201 (47)	30 (43)
Female, n (%)	14 (45)	52 (55)	87 (51)	75 (56)	228 (53)	40 (57)
<b>Side</b>						
Right, n (%)	15 (48)	43 (45)	90 (53)	75 (56)	223 (52)	44 (63)
Left, n (%)	16 (52)	52 (55)	80 (47)	58 (44)	206 (48)	26 (37)
Bohler, dgrs*	33.6 ± 5.5	39.7 ± 5.7	35.1 ± 5.5	33.0 ± 5.0	35.4 ± 5.9	31.7 ± 5.2
Gissane, dgrs*	115.8 ± 7.3	111.1 ± 7.5	109.8 ± 7.2	109.8 ± 7.1	110.5 ± 7.4	112.8 ± 7.4

Group A: ≤ 4 years of age

Group B: 5 to 8 years of age

Group C: 9 to 12 years of age

Group D: 13 to 16 years of age

Group E: ≥ 18 years of age

Group S: 0 to 16 years of age

\* Mean (STD)

**Table 2: Results of Tukey`s post-hoc tests to analyse differences between age groups**

		Bohler`s angle					Crucial angle of Gissane				
		95% Confidence Interval					95% Confidence Interval				
Group 1	Group 2	Mean	%	Std.	Lower	Upper	Mean	%	Std.	Lower	Upper
(I)	(J)	Diff	[(I-J)/I]	Error	Limit	Limit	Diff	[(I-J)/J]	Error	Limit	Limit
		(I-J)					(I-J)				
A	B	-6.02*	-17.92	1.12	-9.08	-2.97	4.78*	4.13	1.50	.67	8.89
	C	-1.46	-4.35	1.05	-4.34	1.43	6.07*	5.24	1.42	2.19	9.96
	D	.61	1.82	1.08	-2.34	3.55	6.04*	5.22	1.45	2.08	10.01
	E	1.92	5.71	1.16	-1.26	5.11	3.04	2.63	1.57	-1.25	7.33
B	C	4.56*	11.49	.69	2.67	6.46	1.29	1.16	.93	-1.25	3.84
	D	6.63*	16.70	.72	4.64	8.61	1.26	1.13	.96	-1.40	3.93
	E	7.95*	20.03	.84	5.62	10.27	-1.74	-1.57	1.14	-4.88	1.39
C	D	2.06*	5.87	.62	.35	3.77	-.03	-0.03	.84	-2.33	2.27
	E	3.38*	9.63	.77	1.28	5.48	-3.04*	-2.77	1.03	-5.86	-.21
D	E	1.32	4.00	.80	-.86	3.49	-3.01*	-2.74	1.07	-5.94	-.07
S <sup>£</sup>	E <sup>£</sup>	3.65*	10.31	.69	2.17	5.13	-2.30*	-2.08	.95	-4.17	-.43

Group A: ≤ 4 years of age

Group B: 5 to 8 years of age

Group C: 9 to 12 years of age

Group D: 13 to 16 years of age

Group E:  $\geq 18$  years of age

Group S: 0 to 16 years of age

Mean Diff (I-J): Mean difference


% [(I-J)/I]: Percentage of variation between Group 1 (I) and Group 2 (J)

Std. Error: Standard error


<sup>£</sup> Independent samples T-test

\*  $p \leq 0.05$

**Graph 1: Regression analysis of distribution of Bohler's angle across age**




**Graph 2: Distribution of Bohler`s angle (mean) across different age groups**



Note: One standard deviation represented by brackets

**Graph 3: Distribution of crucial angle of Gissane (mean) across different age groups**



Note: One standard deviation represented by brackets

**Figure 1: Measurement technique for Bohler's angle and the crucial angle of Gissane**




Figure A: Bohler's angle

Figure B: Crucial angle of Gissane

**Figure 2: Growth of calcaneus across age groups**




Figure A:  $\leq 4$  years of age

Figure B: 5 to 8 years of age

Figure C: 9 to 12 years of age

Figure D: 13 to 16 years of age

Figure E:  $\geq 18$  years of age