




IN BRIEF

Uncovering the Steps Before: Sulfate Induces ABA Biosynthesis and Stomatal Closure^[OPEN]

Plant stomatal aperture regulation via guard cells is an example of how plants dynamically process environmental signals to induce a physiological response. The drought stress hormone abscisic acid (ABA) is a well-characterized signal that induces stomatal closure, preventing water loss. ABA acts via the PYRABACTIN RESISTANCE (PYR)/PYR LIKE-ABA INSENSITIVE1-OPEN STOMATA1 receptor-phosphatase-kinase core signaling pathway. It induces the activity of NADPH oxidases for the production of reactive oxygen species (ROS), important regulators of stomatal closure, in an OPEN STOMATA1-dependent manner. Furthermore, drying of soil induces the root-to-shoot transport of sulfate that, in turn, promotes stomatal closure (Malcheska et al., 2017).

Batool et al. (2018) explored how upstream signaling cascades involving sulfate and Cys biosynthesis lead to ABA accumulation. To study sulfate-induced stomatal closure in maize (*Zea mays*) and *Arabidopsis thaliana*, sulfate was exogenously applied to epidermal peels or administered via petiole feeding, which resulted in stomatal closure. The authors found that sulfate induced ROS formation in guard cells by activating NADPH oxidases. To explore if sulfate-induced ROS formation and stomatal closure act via the ABA signaling pathway, sulfate was applied to *aba3-1*, which is impaired in ABA biosynthesis. Whereas the *aba3-1* mutant exhibited ROS production and stomatal closure in response to ABA treatment, sulfate treatment did not elicit the same response. Therefore, sulfate-induced ROS production depends on de novo ABA biosynthesis via ABA3. Furthermore, exogenous sulfate treatment increased the intracellular ABA concentration in wild-type guard cells, leading

^[OPEN]Articles can be viewed without a subscription. www.plantcell.org/cgi/doi/10.1105/tpc.18.00912



Cys application leads to an increase of intracellular ABA concentration in guard cells to the same extent as ABA application, as indicated by the ABA sensor ABAleon2.1. (Adapted from Batool et al., [2018], Figure 7)

to enhanced transcription of ABA marker genes.

How does sulfur metabolism influence sulfate-induced stomatal closure? The authors used the *sir1-1* mutant, which cannot convert sulfite to sulfide, and the *serat tko* mutant, which is defective in incorporating sulfide into Cys, to examine this question. Neither of these mutants exhibited stomatal closure or ROS formation in response to sulfate treatment. However, exogenous ABA and Cys treatment led to stomatal closure in both mutants, indicating a requirement for the metabolic assimilation of sulfate into Cys by SERAT and SiR for sulfate-induced closure. Furthermore, direct application of Cys in the wild type also elevated the intracellular ABA concentration (see figure 1) and induced ROS formation.

The requirement for the incorporation of sulfide into Cys for sulfate-induced stomatal closure was explored further in the context of ABA biosynthesis. Previous studies showed that ABA3 uses Cys as a sulfur donor to convert abscisic aldehyde to ABA via ABSCISIC ALDEHYDE OXIDASE3 (Bittner et al., 2001; Cao et al., 2014). **Batool et al. (2018)** extended this line of research by demonstrating that petiole feeding of sulfate or Cys induced the transcription of NINE-CIS-EPOXYCAROTENOID DIOXYGENASE3 (*NCED3*), which limits the synthesis of the ABSCISIC ALDEHYDE OXIDASE3 substrate and, consequently, ABA biosynthesis. However, Cys and

sulfate application did not induce stomatal closure in the *aba3-1* and *nced3-2* loss-of-function mutants, indicating that both NCED and ABA3 activity are required for Cys- and sulfate-induced stomatal closure.

Cys synthesis-depleted mutants were found to have enhanced wilting and suffered lower relative water content than wild-type plants under limited water supply. This reflects the need for adequate Cys to promote decreased transpiration in leaves when soil drying occurs, most likely due to the limited ability to synthesize ABA to lead to stomatal closure.

Batool et al. (2018) demonstrate that sulfate promotes stomatal closure by inducing ABA biosynthesis and activating ROS. The incorporation of sulfate into Cys leads to ABA biosynthesis in leaves, highlighting how sulfate can act as a limiting factor to initiate the hormone signaling pathway important for stomatal closure in plants under stress.

Estee E. Tee
Assistant Features Editor
Research School of Biology
Australian National University, Canberra
estee.tee@anu.edu.au
ORCID ID: 0000-0003-2613-5977

REFERENCES

Batool, S., et al. (2018). Sulfate is incorporated into cysteine to trigger ABA

- production and stomata closure. *Plant Cell* **30**: 2973–2987.
- Bittner, F., Oreb, M., and Mendel, R.R.** (2001). ABA3 is a molybdenum cofactor sulfurase required for activation of aldehyde oxidase and xanthine dehydrogenase in *Arabidopsis thaliana*. *J. Biol. Chem.* **276**: 40381–40384.
- Cao, M.J., Wang, Z., Zhao, Q., Mao, J.L., Speiser, A., Wirtz, M., Hell, R., Zhu, J.K., and Xiang, C.B.** (2014). Sulfate availability affects ABA levels and germination response to ABA and salt stress in *Arabidopsis thaliana*. *Plant J.* **77**: 604–615.
- Malcheska, F., et al.** (2017). Drought-enhanced xylem sap sulfate closes stomata by affecting ALMT12 and guard cell ABA synthesis. *Plant Physiol.* **174**: 798–814.

Uncovering the Steps Before: Sulfate Induces ABA Biosynthesis and Stomatal Closure

Estee E. Tee

Plant Cell 2018;30:2894-2895; originally published online December 11, 2018;

DOI 10.1105/tpc.18.00912

This information is current as of June 27, 2019

References	This article cites 4 articles, 3 of which can be accessed free at: /content/30/12/2894.full.html#ref-list-1
Permissions	https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&iissn=1532298X&WT.mc_id=pd_hw1532298X
eTOCs	Sign up for eTOCs at: http://www.plantcell.org/cgi/alerts/ctmain
CiteTrack Alerts	Sign up for CiteTrack Alerts at: http://www.plantcell.org/cgi/alerts/ctmain
Subscription Information	Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: http://www.aspb.org/publications/subscriptions.cfm