Article

CLINICAL REHABILITATION

The role of movement errors in modifying spatiotemporal gait asymmetry post stroke: a randomized controlled trial

Clinical Rehabilitation 2018, Vol. 32(2) 161–172 © The Author(s) 2017 Reprints and permissions: sagepub.co.uk/journalsPermissions.nav DOI: 10.1177/0269215517723056 journals.sagepub.com/home/cre



Michael D Lewek¹, Carty H Braun^{2,3}, Clint Wutzke^{4,5} and Carol Giuliani¹

Abstract

Objective: Current rehabilitation to improve gait symmetry following stroke is based on one of two competing motor learning strategies: minimizing or augmenting symmetry errors. We sought to determine which of those motor learning strategies best improves overground spatiotemporal gait symmetry. **Design:** Randomized controlled trial.

Setting: Rehabilitation research lab.

Subjects: In all, 47 participants $(59 \pm 12 \text{ years old})$ with chronic hemiparesis post stroke and spatiotemporal gait asymmetry were randomized to error augmentation, error minimization, or conventional treadmill training (control) groups.

Interventions: To augment or minimize asymmetry on a step-by-step basis, we developed a responsive, "closed-loop" control system, using a split-belt instrumented treadmill that continuously adjusted the difference in belt speeds to be proportional to the patient's current asymmetry.

Main measures: Overground spatiotemporal asymmetries and gait speeds were collected prior to and following 18 training sessions.

Results: Step length asymmetry reduced after training, but stance time did not. There was no group × time interaction. Gait speed improved after training, but was not affected by type of asymmetry, or group. Of those who trained to modify step length asymmetry, there was a moderately strong linear relationship between the change in step length asymmetry and the change in gait speed.

Conclusion: Augmenting errors was not superior to minimizing errors or providing only verbal feedback during conventional treadmill walking. Therefore, the use of verbal feedback to target spatiotemporal asymmetry, which was common to all participants, appears to be sufficient to reduce step length asymmetry. Alterations in stance time asymmetry were not elicited in any group.

¹Division of Physical Therapy, Department of Allied Health Sciences, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

⁵Department of Rehabilitation Science, George Mason

University, Fairfax, VA, USA

Corresponding author:

Email: mlewek@med.unc.edu

²UNC Health Care System, Chapel Hill, NC, USA ³Children's Hospital of Atlanta, Atlanta, GA, USA ⁴Human Movement Science Program, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

Michael D Lewek, Division of Physical Therapy, Department of Allied Health Sciences, University of North Carolina at Chapel Hill, 3043 Bondurant Hall, CB#7135, Chapel Hill, NC 27599-7135, USA.

Keywords

Gait, stroke, asymmetry, adaptation, learning

Received: 27 March 2017; accepted: 7 July 2017

Introduction

The slow walking speeds commonly observed following stroke are associated with marked temporal and spatial interlimb asymmetries in 48% and 44% of subjects, respectively.¹ Asymmetric gait has been associated with gait energetics,² gait speed,¹ and balance control³ and may lead to an increased risk of falls, lower extremity musculoskeletal injury,⁴ and loss of bone mineral density in the paretic limb.⁵ Because asymmetries are considered movement errors, therapists often use tactile, visual, or verbal cues to *minimize asymmetries* during gait training (error minimization strategy).⁶ Indeed, conventional locomotor training on a treadmill has demonstrated improved gait speed,^{7–9} but no effect on interlimb symmetry.^{7,10,11}

In contrast, the approach of augmenting asymmetry (error augmentation strategy) to modify gait^{12–14} is based on established motor learning principles, including error-based learning¹⁵ and variability of practice.¹⁶ Such an error augmentation strategy may be beneficial for individuals post stroke due to their deficits in recognizing spatiotemporal asymmetries as movement errors.¹⁷ The improved detection of augmented errors by muscle and joint proprioceptors may be the driving signal for adaptation and learning.¹⁵ Walking on a split-belt treadmill with belts operating at different fixed speeds (e.g. 2:1 ratio with short step length limb on fast belt) is a prime example of augmenting error, which will initially exaggerate step length asymmetry.¹⁸ The adaptation over time and resulting after-effect suggest that the neuromuscular system retains the ability to produce nearly symmetric step lengths following stroke.¹⁸ Long-term training (e.g. fourweeks) with this locomotor adaptation approach elicited significant improvement in step length symmetry in more than half of the trained individuals.¹⁹

Rather than using a fixed perturbation, motor learning may be accelerated by transient amplification of movement errors.^{20,21} To test this hypothesis, we designed a closed-loop feedback system coupled with an instrumented split-belt treadmill, in which step-by-step performance of spatiotemporal asymmetry determines the ratio of the belt speeds (i.e. error magnitude).²² The system can either augment or minimize error on a step-by-step basis, by continuously adjusting the difference in belt speed to be proportional to the patient's current asymmetry. For example, if stance time is shorter on the paretic limb, the velocity of the paretic treadmill belt can either be increased (to augment asymmetry) or decreased (to minimize asymmetry) relative to the non-paretic treadmill belt. With our unique system, we can determine which motor learning strategy (error minimization or augmentation) is best for improving overground spatial and temporal symmetry post stroke.

Given that rehabilitation to improve gait symmetry may be based on one of these two competing motor learning strategies (i.e. minimizing or augmenting symmetry errors), the purpose of this randomized controlled trial was to determine which of those motor learning strategies best improves overground spatial and temporal gait symmetry in individuals with chronic hemiparesis post stroke. Based on data from split-belt adaptation work,¹⁸ we hypothesized that training with error augmentation would elicit the largest change in overground gait symmetry because this mode would enhance awareness of movement errors.

Methods

Setting and participants

The study was a randomized controlled trial with three parallel groups. The trial was registered with clinicaltrials.gov (NCT01598675) and all subjects received medical clearance from a physician to participate in training and signed an informed consent form approved by the UNC-Chapel Hill IRB (study #11-1240). We recruited potential subjects with hemiparesis of more than six-month duration after unilateral, supratentorial, ischemic, or hemorrhagic stroke. Participants were recruited from local stroke support groups, email Listservs, and referred by local physicians and physical therapists.

Potential subjects were included if they were able to walk >10 m overground without physical assistance, had an overground comfortable gait speed <1.0 m/s (using assistive devices and bracing below the knee, as needed), were able to walk on a treadmill at >80% of comfortable gait speed using upper extremity support if needed, and exhibited stance time asymmetry and/or step length asymmetry. Asymmetry was defined by a symmetry ratio² (max(paretic, non-paretic)/(paretic+non-paretic)) \geq 0.524 for stance time asymmetry and \geq 0.537 for step length asymmetry. These values represent a doubling of the upper 95% confidence limit of unimpaired individuals.¹ Potential subjects were excluded if they had uncontrolled cardiorespiratory/metabolic disease, other neurologic disorders or orthopedic injury that may affect gait, botulinum toxin to the lower limb in the past six months, or concurrent physical therapy.

Randomization and intervention

Following confirmation of eligibility and a double baseline $(28 \pm 16 \text{ days between baseline sessions})$, participants were randomized into one of three groups (augmentation, minimization, and control) by drawing a card from one of four envelopes that were stratified based on initial overground gait speed (≤ 0.5 or > 0.5 m/s, to help ensure similar gait speeds among groups)²³ and the type of asymmetry (step length or stance time). To avoid excluding subjects who had both stance time asymmetry and step length asymmetry, subjects were assigned, and trained accordingly, based on the larger of these two symmetry ratios. Assessors assigned to perform all outcome measures were initially blinded to group allocation; however, due to staffing constraints, the final four subjects were assessed by an unblinded assessor. All participants refrained from attending their usual physical therapy, if applicable, during the course of the study.

Subjects completed 18 sessions of walking for up to 20 minutes on a split-belt, instrumented treadmill (Bertec Corp, Worthington, OH, USA) wearing a safety harness that did not restrict lower extremity movements. Subjects were permitted to hold onto the handrail, but were discouraged from doing so, if possible. Therapists did not facilitate limb movements. Verbal instructions, using external focus of attention to cue for increased symmetry (e.g. "ride the belt a little longer on your weak side," "reach for a spot near the front of the treadmill") were used for all groups, as necessary.⁶ Heart rate and oxygen saturation were monitored throughout training and blood pressure was measured prior to, at mid-training (~10 minutes) and following each training session to ensure that blood pressure remained below 180/110 mm Hg.24 Participants took rest breaks, as needed.

All three groups began each training session with two minutes of control walking on the treadmill with both belts moving at the same speed, as a warm up. Then, without stopping the treadmill, subjects transitioned to the training phase (asymmetry augmentation, minimization, or control) for up to 18 minutes of additional walking. Each training session on the treadmill was followed by 10–15 minutes of overground gait training to encourage carryover of training to overground surfaces.⁶ During overground training, subjects were permitted to use an assistive device, but typically practiced without, when possible. Similar verbal cues were used during overground and treadmill training.

Treadmill control

The treadmill was controlled by a novel computerdriven control algorithm. Treadmill control began by the therapist setting a base walking speed, v_{base} , which is the average speed of the two treadmill belts. During the first session, participants began walking at $v_{\text{base}} = 80\%$ of comfortable gait speed for overground gait. The difference in speeds between the two belts, v_{diff} was determined by the degree of the patient's asymmetry on a step-by-step basis

For stance time = $v_{diff} = k * \log_2(\text{stance}_{time_{right}} / \text{stance}_{time_{left}}) * v_{base}$

For step length = $v_{diff} = -k * \log_2$ (stance _ length_{right} / stance _ length_{left}) * v_{base}

where *k* is a scale factor (default=3), which can be increased to progress training so that a small amount of asymmetry will yield a larger difference in belt speed. Stance time or step length was calculated on a step-by-step basis from ground reaction force data (treadmill forceplates) at 1080Hz with a Vicon MX40+ system (Vicon/Peak, Los Angeles, CA, USA). Stance time was the time (ms) that the vertical ground reaction force exceeded 10N, and step length was determined from center of pressure data as the anterior/posterior distance between the feet at successive heel strikes.^{25,26} These data were sent to the application computer using VRPN.²⁷ The belt speed difference, v_{diff} , was applied to the treadmill by speeding up one belt and slowing the other

$$v_{right} = \frac{v_{base} - v_{diff}}{2}$$
$$v_{left} = \frac{v_{base} + v_{diff}}{2}$$

Augmenting asymmetry was achieved with a positive "k" value, whereas a negative "k" value was used to minimize asymmetry. Each belt had high and low speed limits applied for safety, such that the belt speeds would not exceed a 2:1 ratio.¹⁸ Training step length or stance time was progressed similarly within and between sessions by altering treadmill speed and the "k" value. The "k" value was adjusted independent of treadmill speed (i.e. v_{base}). Treadmill speed was increased if heart rate remained below 70% of agepredicted maximum²⁸ or the perceived exertion was less than 14 on the Borg scale (for participants receiving medications that blunt heart rate response).²⁴

Outcomes and follow-up

We collected overground gait characteristics from both baseline visits. Because of the tendency for gait speed to increase with repeated testing,²⁹ we used the overground data from the second baseline visit, which occurred one week prior to (pre) training, and the repeated assessments at one week following the final training session (post), and at one month after training (follow-up). At each assessment visit, subjects were instructed to "walk at their normal, comfortable walking speed" as they performed 3 passes across a 14 foot GAITRite mat (CIR Systems, Franklin, NJ, USA) with a 4 foot acceleration and deceleration zone at either end of the mat. GAITRite software was used to determine comfortable gait speed and gait asymmetry, as described above, for both step length and stance time variables.²⁹ Given the automated process of calculating outcome measures, we believe that the lack of blinded assessment for the final four subjects was inconsequential.

Statistical analysis

All data were exported into electronic spreadsheets (Excel 2016, Microsoft Corp) and data analysis was subsequently performed using SPSS (ver 23, IBM). Data residuals were checked for assumptions of normality using the Shapiro-Wilk test, and Mauchly's W was used to test for sphericity with the Huynh-Feldt adjustment, as needed. Primary (gait symmetry ratio) and secondary (comfortable gait speed) variables were analyzed using a two-way, repeated measures analysis of variance (ANOVA; repeated for time) to assess the interaction effect (i.e. effect of time on the three groups: augmentation, minimization, and control). Tukey (between subjects) and Bonferroni corrected paired-samples t-tests (within subjects) were used for post hoc tests. For those with step length asymmetry, the shorter step length can occur on either the paretic or non-paretic limb.^{18,29} We therefore sorted limbs based on which step was longer and shorter for separate comparison across time using a repeated-measures ANOVA. The relationship between changes in spatiotemporal symmetry and change in gait speed from pre- to post-training was assessed using Pearson correlations. Prior to enrollment, we performed a power analysis based on a non-central F-distribution to satisfy concerns about a potentially asymmetric distribution of our primary outcome measure (spatiotemporal asymmetry). Power was calculated using assumptions based on



Figure 1. CONSORT diagram depicting subject allocation through study flow.

our review of reported changes in spatiotemporal asymmetry with locomotor training.^{10,13,30} Estimating an SD of 0.019 and a difference in the change in asymmetry of 0.035 from pre-training to post-training, we expected to have greater than 80% power with sample size of 45.

Results

We recruited 48 subjects to participate (26 in the step length symmetry training and 22 in the stance time symmetry training). As seen in the CONSORT diagram (Figure 1), 37 subjects completed gait training (over a mean (SD) of 48 (10) days) and testing. The three groups did not differ on key baseline characteristics (Table 1). The 11 subjects who dropped out were split between groups (augmentation: N=4; minimization: N=4; control: N=3). Reasons for drop-out varied (transportation

issues=6; exercise intolerance=2; blood pressure/ cardiovascular issues=2; none given=1).

Overall, we observed a significant interaction $(P=0.026; \eta_p^2 = 0.140)$ between time and the selected asymmetry parameter (i.e. step length or stance time asymmetry) indicating that those who were trained to alter step length versus stance time asymmetries responded differently over time. For this reason, all further analyses were separated by the participant's training parameter. No other interaction effects were significant (all P > 0.453).

Step length asymmetry

In all, 26 subjects were assigned to perform training to improve step length asymmetry. Of those subjects, 21 (81%) completed testing and training (N=7: augmentation; N=8: minimization; N=6: control).

	Error augmentation (n=12)	Error minimization (n=14)	Control (n=11)	P values
Age	57.5 (10.5) years	60.7 (13.0) years	57.6 (13.2) years	0.764
Sex	5 F/7 M	5F/9 M	5 F/6 M	0.916
Paretic side	4 L/8 R	9 L/5 R	8 L/3 R	0.181
Time post-stroke	36.3 (21.7) months	54.4 (55.8) months	28.1 (27.9) months	0.228
Overground gait speed (pre)	0.42 (0.21) m/s	0.44 (0.28) m/s	0.36 (0.22) m/s	0.710
Assistive device at pretest	6 canes, 6 none	10 canes, 4 none	9 canes, 2 none	0.275
Ankle bracing at pretest	3 yes, 9 no	7 yes, 7 no	9 yes, 2 no	0.026

Table I. Subject demographics.

Values represent mean (SD).



Figure 2. We observed the presence of a moderate relationship between the change in step length asymmetry and the change in gait speed from pre-training to post-training. Group allocation of individual subjects is shown.

We observed no significant interaction effect (group × time: P=0.472; $\eta_p^2 = 0.086$; Supplementary Figure 2, Table 2) among the subjects who trained to improve step length asymmetry. Likewise, we observed no significant effect for group (P=0.183; $\eta_p^2 = 0.172$). We did, however, note a significant effect of time on step length asymmetry (P=0.005; $\eta_p^2 = 0.311$). Across all groups, step length

asymmetry was significantly higher (i.e. worse) at pretest compared to posttest (P=0.014) and at follow-up (P=0.045). No difference was observed between posttest and follow-up (P=1.000).

To confirm that subjects did not simply improve their symmetry ratio by reducing the longer step length, we compared the step lengths of both limbs (Table 3). Importantly, we observed that subjects

Table 2. Spatiotemporal gait asymmetry.

	Pre-training	Post-training	Follow-up
Step length asymmetry trained	(N=2I)		
Step length asymmetry			
Augmentation	0.598 (0.046)	0.580 (0.039)	0.580 (0.037)
Minimization	0.632 (0.115)	0.571 (0.058)	0.568 (0.052)
Control	0.684 (0.116)	0.626 (0.068)	0.638 (0.084)
Stance time asymmetry			
Augmentation	0.528 (0.013)	0.526 (0.013)	0.525 (0.016)
Minimization	0.536 (0.022)	0.542 (0.028)	0.541 (0.028)
Control	0.539 (0.020)	0.539 (0.017)	0.536 (0.019)
Stance time asymmetry trained	(N=16)		
Stance time asymmetry			
Augmentation	0.544 (0.011)	0.545 (0.008)	0.547 (0.010)
Minimization	0.542 (0.016)	0.535 (0.018)	0.536 (0.016)
Control	0.569 (0.014)	0.565 (0.018)	0.561 (0.014)
Step length asymmetry			
Augmentation	0.527 (0.022)	0.531 (0.015)	0.517 (0.017)
Minimization	0.521 (0.009)	0.517 (0.013)	0.520 (0.014)
Control	0.533 (0.027)	0.539 (0.024)	0.533 (0.029)

Values represent mean (SD).

	Pre-training	Post-training	Follow-up
Shorter steps			
Augmentation	26.8 (11.2)	33.5 (13.7)	33.3 (13.6)
Minimization	26.7 (17.3)	38.8 (16.0)	39.5 (15.1)
Control	20.2 (13.3)	30.5 (13.1)	28.6 (13.1)
Longer steps			
Augmentation	39.1 (12.4)	45.2 (14.6)	45.2 (15.3)
Minimization	41.1 (13.9)	49.9 (15.3)	50.4 (14.1)
Control	38.9 (7.9)	48.3 (8.5)	47.1 (7.4)

Table 3. Step lengths (cm) stratified by shorter and longer steps.

Values represent mean (SD).

significantly increased both their shorter (P < 0.001; $\eta_p^2 = 0.621$) and longer (P < 0.001; $\eta_p^2 = 0.701$) step lengths as a result of training, and maintained those improvements at follow-up (Supplementary Figure 3). Nine subjects (of 21) exhibited changes in step length asymmetry from pre- to post-training that exceeded the minimal detectable change (MDC) of 0.035.²⁹ Of those, one trained with error augmentation, five were in the error minimization group, and three were in the control group (P=0.185). Within this group that trained to improve step length asymmetry, we observed no change in stance time asymmetry (P=0.774; $\eta_p^2 = 0.012$).

Stance time asymmetry

We assigned 22 subjects to train to improve stance time symmetry and 16 (73%) completed training (N=5: augmentation; N=6: minimization; N=5: control). Of those who trained to improve stance time asymmetry, there was no significant effect of time (P=0.302; η_p^2 =0.088) on stance time asymmetry. Likewise, we observed no significant interaction effect (group × time: (P=0.374; $\eta_p^2 = 0.146$; Supplementary Figure 4). These subjects were trained to improve stance time asymmetry, and thus we observed no influence of training on step length asymmetry (P=0.134; $\eta_p^2 = 0.143$).

Gait speed

Because there were no significant interaction effects (all P > 0.342), the post hoc analysis for gait speed was collapsed across all subjects. Gait speed significantly increased over time (P < 0.001; $\eta_p^2 = 0.617$). Specifically, we observed that subjects walked significantly faster after training (mean (SD) gait speed was 0.53 (0.28) m/s) compared to pre-training speeds (0.41 (0.24) m/s; P < 0.001; d = 0.52). Gait speed continued to improve from the posttest to the follow-up session (0.56 (0.29) m/s; P = 0.018; d = 0.08).

Of those who trained to improve their step length asymmetry, we observed a moderate relationship between the change in step length asymmetry from pre- to post-training and the change in gait speed from pre- to post-training (r=-0.45; P=0.040; Figure 2). There was no such relationship between the change in stance time asymmetry and gait speed for those who trained to improve stance time asymmetry (r=-0.153; P=0.571).

We performed one additional comparison because we were particularly concerned about the potential for training with error minimization to increase (i.e. make worse) subject's asymmetry.¹⁸ Indeed, during training, our paradigm has the potential to augment step length symmetry while minimizing stance time symmetry, or vice versa. For example, a participant assigned to augment step length asymmetry during training may also have concomitantly minimized stance time asymmetry during training. For that reason, subjects were recoded as to whether step length asymmetry or stance time asymmetry was minimized or augmented (regardless of group assignment). In fact, we observed that the subjects (N=13) who had their step length asymmetry errors minimized (either intentionally or not) showed a tendency to reduce their step length asymmetry from 0.592 (0.103) at pretest to 0.556 (0.049) at posttest (P=0.097;

d=0.48). Those who had their stance time asymmetry errors minimized (N=13, regardless of intentional or unintentional) did not alter stance time asymmetry from pretest (0.537 (0.014)) to posttest (0.539 (0.029); P=0.682; d=0.12). These data suggest that adaptive minimization of movement errors did not lead to an increase (worsening) in gait asymmetry following training.

Clinical Rehabilitation 32(2)

Discussion

Our hypothesis that gait training with error augmentation would improve spatiotemporal gait asymmetry more than minimizing errors was not supported by these data. Subjects were able to improve step length (spatial) asymmetry, but did not show an appreciable change in stance time (temporal) asymmetry following training. Despite this overall improvement in step length asymmetry after training, we did not observe one method of training to be superior to another. The common training component across all groups was the deliberate focus on spatiotemporal asymmetry and the verbal feedback from therapists during gait training. This attention to specific training variables through verbal feedback during functional practice may, therefore, have been the active ingredient that promoted improvements in step length asymmetry.

We are excited at the prospect that step length asymmetry was significantly improved, and that the improvement was retained for at least a month. Although the training was performed on a treadmill, all testing was performed overground, emphasizing the context-independent nature of the improvements. Although we were initially surprised that the error augmentation algorithm did not produce greater gains, from a clinical perspective this represents a more promising finding. Rather than requiring a splitbelt instrumented treadmill with computer-controlled algorithms, it appears possible to elicit substantial changes to step length symmetry in individuals with chronic stroke during "conventional" gait training, provided that appropriate focus, attention,³¹ and variability in practice are provided.

Based on previous literature,¹⁹ we were surprised that the error augmentation group did not outperform the other two groups. Although some have noted that error augmentation is more beneficial than error minimization during a single session,³² others have reported that error augmentation did not improve the immediate retention of a locomotor balance task.33 In our cohort, the type of error manipulation during training (either minimization or augmentation) did not appear to differentially influence spatial asymmetry changes; comparable changes occurred in all three groups. Importantly, however, we observed that step length symmetry was only improved in the subjects who trained to improve step length symmetry. We can think of two possible reasons for this. First, we could not train subjects to improve stance time asymmetry and expect to see an improvement in step length asymmetry. Despite the known relationship between spatial and temporal asymmetry,¹ the specific attention to step length asymmetry during training appeared important for eliciting change. Second, by the nature of the inclusion criteria, the subjects who trained to improve step length asymmetry exhibited greater step length asymmetry than those who trained to improve stance time asymmetry. Likewise, Reisman et al.¹⁹ demonstrated that responders appeared to have a larger baseline asymmetry than the non-responders. Nevertheless, there were many individuals in our stance time asymmetry trained group (n=9 of 16 who completed training) who exhibited substantial step length asymmetry, but happened to have stance time asymmetry that was larger in magnitude than their step length asymmetry. Therefore, although the difference in baseline step length asymmetry may have contributed to this finding, we think it is more likely that the change in step length asymmetry after training was due to the attention focused on that particular parameter.

We observed, similar to others,^{7,10,11,13} that stance time asymmetry is particularly resistant to change. As a rapidly adapting parameter,²⁶ stance time asymmetry is highly responsive to the speed of the treadmill belts, and thus did not change in response to training. Stance time is a reactive parameter, which relies heavily on peripheral feedback from hip flexor afferents, limb load receptors, and cutaneous feedback. The fact that after-effects are not observed after manipulation of stance time asymmetry suggests that this is a parameter that may not be modified using error manipulation.³⁴ Interestingly, although stance time asymmetry and step length asymmetry are related to each other,¹ there was no concomitant change in stance time asymmetry in our group who trained to improve step length asymmetry. Thus, changing one parameter does not necessitate a change in another.³⁵

Consistent with current neurorehabilitation strategies, which encourage spatiotemporal symmetry during training,6 our "closed-loop," error minimization system was intended to produce nearly symmetric gait. There has been concern that such training may exacerbate gait asymmetry after training.18 Instead, we observed that those who had their asymmetry errors minimized tended to reduce their asymmetry after 18 training sessions. Although this observation would appear to conflict with previous literature,¹⁸ it is important to note the inherent difference in control schemes between our paradigm and that of others. Previous work has used open-loop systems, which are unresponsive to the step-by-step changes in movement that occur during training.¹⁸ Instead, our "closed-loop" system puts the subject in the loop to ensure that the manipulation of errors is responsive to the subject's stepping. Our subjects in the error minimization and error augmentation groups, therefore, received real-time, proprioceptive feedback of gait asymmetry. Despite the likely presence of deficits in spatiotemporal asymmetry perception,¹⁷ our subjects following stroke did not appear to need the altered proprioceptive feedback from the changing treadmill belt speeds, as subjects in all three groups were able to improve step length asymmetry.

The change in gait speed observed following training was comparable to previously published work.^{7,36} Although gait speed was increased within and across sessions, we focused our subject's attention on spatiotemporal gait symmetry rather than gait speed. Additionally, the gait speed change of our subjects was likely an underestimate compared to previous work. Because gait speed is known to increase substantially without the presence of an intervention,^{29,37} we performed a double baseline and used the second (i.e. faster) session's speed as the pre-training speed. Notably, the change in step length asymmetry was related to the change in comfortable gait speed. Although this finding would seem to suggest that step length asymmetry is a therapeutic target for improving gait speed

following stroke, there is evidence to the contrary. In short, we observed that several subjects experienced sizable increases in gait speed, with small changes in step length symmetry. Likewise, others have noted significant improvements in step length symmetry, without a concomitant change in gait speed.¹⁹ Thus, we must question whether changing step length asymmetry is really necessary to elicit improvements in gait speed.

A limitation of this work is the large number of drop-outs. Getting participants to the laboratory for 18 training sessions proved more difficult than we anticipated given our budget limitations. Ultimately, we had 77% (37 of 48) of our participants complete training, which was consistent with some recently reported gait training trials,⁷ but a smaller percentage than others.³⁸ Despite the somewhat high drop-out rate, there were similar numbers of participants who did not complete training in each group. Additionally, the small sample size produced small cell counts within each stratification, which limited our ability to examine potential differences between individuals with a slow versus a fast walking speed. An additional limitation of this work is that the final four participants were not measured by a blinded assessor. Finally, additional information about specific lesion location might have helped with the interpretation of our data.

The differential response to training suggests that better selection criteria are needed to identify responders to treatment. We observed that nine of the 21 subjects trained to improve step length asymmetry exceeded the MDC.29 Likewise, Reisman et al.¹⁹ demonstrated improvement in 7 of 13 subjects using a more liberal definition of "responder." Although some individuals clearly had the capacity to improve their step length asymmetry, there are also some individuals who did not improve as a result of training. For those individuals who did demonstrate improvement in step length symmetry, only one reached the threshold for having symmetric gait. Presently, thresholds of minimally important clinical changes remain unknown for step length asymmetry. If further work to restore symmetric gait following stroke is to continue, it is important to determine the characteristics that make someone successful at altering their step length

asymmetry. Because the locomotor adaptation literature suggests that many individuals have the neuromotor capacity to produce improved symmetry,¹⁸ perhaps verifying this capacity would improve subject selection for future work.

Clinical Messages

- In patients more than six months after stroke and able to walk, persistent step length asymmetry was improved following training, whereas stance time asymmetry was not.
- A specific emphasis on either minimizing or augmenting asymmetry during treadmill training did not appear to influence these results.
- Verbal instruction and feedback were common across all groups and may have contributed to the noted improvement in step length asymmetry.

Acknowledgements

The authors acknowledge Jeff Feasel, MS, for the treadmill control algorithms; Samantha Hansen, Mary O'Brien, Donna Dean, and Robert Sykes III for assistance with scheduling, recruiting, training, and data management; Prue Plummer for helpful insight including review of the article prior to submission.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the National Institutes of Health (R21—HD068805 to Lewek).

References

- Patterson KK, Gage WH, Brooks D, et al. Evaluation of gait symmetry after stroke: a comparison of current methods and recommendations for standardization. *Gait Posture* 2010; 31: 241–246.
- AwadLN, PalmerJA, PohligRT, et al. Walking speedandstep length asymmetry modify the energy cost of walking after stroke. *Neurorehabil Neural Repair* 2015; 29: 416–423.

- Lewek MD, Bradley CE, Wutzke CJ, et al. The relationship between spatiotemporal gait asymmetry and balance in individuals with chronic stroke. *J Appl Biomech* 2014; 30: 31–36.
- Norvell DC, Czerniecki JM, Reiber GE, et al. The prevalence of knee pain and symptomatic knee osteoarthritis among veteran traumatic amputees and nonamputees. *Arch Phys Med Rehabil* 2005; 86: 487–493.
- Jorgensen L, Crabtree NJ, Reeve J, et al. Ambulatory level and asymmetrical weight bearing after stroke affects bone loss in the upper and lower part of the femoral neck differently: bone adaptation after decreased mechanical loading. *Bone* 2000; 27: 701–707.
- Duncan PW, Sullivan KJ, Behrman AL, et al. Protocol for the locomotor experience applied post-stroke (LEAPS) trial: a randomized controlled trial. *BMC Neurol* 2007; 7: 39.
- Hornby TG, Campbell DD, Kahn JH, et al. Enhanced gait-related improvements after therapist- versus roboticassisted locomotor training in subjects with chronic stroke. A randomized controlled study. *Stroke* 2008; 39: 1786–1792.
- Sullivan KJ, Brown DA, Klassen T, et al. Effects of taskspecific locomotor and strength training in adults who were ambulatory after stroke: results of the STEPS randomized clinical trial. *Phys Ther* 2007; 87: 1580–602; discussion 603–607.
- Duncan PW, Sullivan KJ, Behrman AL, et al. Bodyweight-supported treadmill rehabilitation after stroke. N Engl J Med 2011; 364: 2026–2036.
- Patterson SL, Rodgers MM, Macko RF, et al. Effect of treadmill exercise training on spatial and temporal gait parameters in subjects with chronic stroke: a preliminary report. J Rehabil Res Dev 2008; 45: 221–228.
- Silver KH, Macko RF, Forrester LW, et al. Effects of aerobic treadmill training on gait velocity, cadence, and gait symmetry in chronic hemiparetic stroke: a preliminary report. *Neurorehabil Neural Repair* 2000; 14: 65–71.
- Savin DN, Tseng SC, Whitall J, et al. Poststroke hemiparesis impairs the rate but not magnitude of adaptation of spatial and temporal locomotor features. *Neurorehabil Neural Repair* 2013; 27: 24–34.
- Kahn JH and Hornby TG. Rapid and long-term adaptations in gait symmetry following unilateral step training in people with hemiparesis. *Phys Ther* 2009; 89: 474–483.
- Lam T, Luttmann K, Houldin A, et al. Treadmill-based locomotor training with leg weights to enhance functional ambulation in people with chronic stroke: a pilot study. J Neurol Phys Ther 2009; 33: 129–135.
- Kawato M. Feedback-error-learning neural network for supervised learning. In: Eckmiller R (ed.) Advanced neural computers. Amsterdam: North-Holland, 1990, pp.365–372.
- Schmidt RA. A schema theory of discrete motor skill learning. *Psychol Rev* 1975; 82: 225–260.
- Wutzke CJ, Faldowski RA and Lewek MD. Individuals poststroke do not perceive their spatiotemporal gait asymmetries as abnormal. *Phys Ther* 2015; 95: 1244–1253.

- Reisman DS, Wityk R, Silver K, et al. Locomotor adaptation on a split-belt treadmill can improve walking symmetry post-stroke. *Brain* 2007; 130: 1861–1872.
- Reisman DS, McLean H, Keller J, et al. Repeated split-belt treadmill training improves poststroke step length asymmetry. *Neurorehabil Neural Repair* 2013; 27: 460–468.
- Emken JL and Reinkensmeyer DJ. Robot-enhanced motor learning: accelerating internal model formation during locomotion by transient dynamic amplification. *IEEE Trans Neural Syst Rehabil Eng* 2005; 13: 33–39.
- 21. Gordon KE and Ferris DP. Learning to walk with a robotic ankle exoskeleton. *J Biomech* 2007; 40: 2636–2644.
- Feasel J, Whitton MC, Kassler L, et al. The integrated virtual environment rehabilitation treadmill system. *IEEE Trans Neural Syst Rehabil Eng* 2011; 19: 290–297.
- Plummer P, Behrman AL, Duncan PW, et al. Effects of stroke severity and training duration on locomotor recovery after stroke: a pilot study. *Neurorehabil Neural Repair* 2007; 21: 137–151.
- ACSM. ACSM's guidelines for exercise testing and prescription. 7th ed. Philadelphia, PA: Lippincott Williams & Wilkins, 2005.
- Noble JW and Prentice SD. Adaptation to unilateral change in lower limb mechanical properties during human walking. *Exp Brain Res* 2006; 169: 482–495.
- Reisman DS, Block HJ and Bastian AJ. Interlimb coordination during locomotion: what can be adapted and stored? *J Neurophysiol* 2005; 94: 2403–2415.
- Taylor RM, II, Hudson TC, Seeger A, et al. VRPN: a device-independent, network-transparent VR peripheral system. In: *Proceedings of the ACM symposium on virtual reality software and technology*, 15–17 November 2001, Banff, AB, Canada. New York: ACM.
- 28. Gordon NF, Gulanick M, Costa F, et al. Physical activity and exercise recommendations for stroke survivors: an American Heart Association scientific statement from the Council on Clinical Cardiology, Subcommittee on Exercise, Cardiac Rehabilitation, and Prevention; the Council on Cardiovascular Nursing; the Council on Nutrition, Physical Activity, and Metabolism; and the Stroke Council. *Circulation* 2004; 109: 2031–2041.
- 29. Lewek MD and Randall EP. Reliability of spatiotemporal asymmetry during overground walking for individuals following chronic stroke. *J Neurol Phys Ther* 2011; 35: 116–121.
- Lewek MD, Cruz TH, Moore JL, et al. Allowing intralimb kinematic variability during locomotor training poststroke improves kinematic consistency: a subgroup analysis from a randomized clinical trial. *Phys Ther* 2009; 89: 829–839.
- van Vliet PM and Wulf G. Extrinsic feedback for motor learning after stroke: what is the evidence? *Disabil Rehabil* 2006; 28: 831–840.
- Kao PC, Srivastava S, Agrawal SK, et al. Effect of robotic performance-based error-augmentation versus errorreduction training on the gait of healthy individuals. *Gait Posture* 2013; 37: 113–120.

- Domingo A and Ferris DP. The effects of error augmentation on learning to walk on a narrow balance beam. *Exp Brain Res* 2010; 206: 359–370.
- Helm EE and Reisman DS. The split-belt walking paradigm: exploring motor learning and spatiotemporal asymmetry poststroke. *Phys Med Rehabil Clin N Am* 2015; 26: 703–713.
- Malone LA, Bastian AJ and Torres-Oviedo G. How does the motor system correct for errors in time and space during locomotor adaptation? *J Neurophysiol* 2012; 108: 672–683.
- Awad LN, Reisman DS, Kesar TM, et al. Targeting paretic propulsion to improve poststroke walking function:

a preliminary study. Arch Phys Med Rehabil 2014; 95: 840-848.

- Flansbjer UB, Holmback AM, Downham D, et al. Reliability of gait performance tests in men and women with hemiparesis after stroke. *J Rehabil Med* 2005; 37: 75–82.
- 38. Awad LN, Reisman DS, Pohlig RT, et al. Reducing the cost of transport and increasing walking distance after stroke: a randomized controlled trial on fast locomotor training combined with functional electrical stimulation. *Neurorehabil Neural Repair* 2016; 30: 661–670.