

Development of barley cultivars for animal forage in Korea

Han Ouk-kyu, Park Tae-il, Song Tae-hwa, Kim Dea-wook, Lee Yu-yong, Ku Ja-hwan, Hwang Jong-jin, Kim Chung-guk, and Kwon Young-up

National Institute of Crop Science, RDA, Suwon 441-857, Rep. Korea . www.nics.go.kr

Contact email: okhan98@korea.kr

Keywords: Winter season, forage barley, production, Korea.

Introduction

In Korea, the domestic consumption of barley as a cereal crop has been decreasing since the 1980s. It has been considered that crop production in the winter-season rice fields could enhance the global competitiveness of domestic livestock industry by providing better quality fodder to livestock and enhancing field use rate. Therefore, the purpose of barley cultivation for cereal food production has been recently replaced by the production of the barley for forage use. Consequently, the area of barley cultivation for forage is markedly increasing in Korea. While any type of barley can be used as forage for feeding cattle, whole crop barley delivers a higher dry matter yield than conventional feed barley. This paper described the present state of forage barley cultivars developed in Korea.

Methods

All of the cultivars were developed by the breeding team at National Institute of Crop Science (NICS), Rural Development Administration (RDA) in 2002 to 2012. The cross combinations for F₁ production were as follows: Youngyang, between cultivar Bunong and an elite pedigree Miryang55; Sunwoo, a germplasm line P71523 and a pedigree Suweon234; Sangwon, a germplasm line 72sel and a breeding line SB86659-B-YB-22; Soman, a breeding line SB79124 and SB77189; WooHo, F₁ [Olbori // F₆(SB77011) / F₅(Bengei // Hagane / Bunong)] / 3 / (Y7213 / SD607 // CM67 / Miryang12)] and F₁ [(1012.2 / IB65 // Olbori) // (Samheung // Suwon18 / Kangbori)]; Yuyeon, a pedigree Suwon311 and SB86648; Dami, a germplasm line BGS60 and cultivar Kangbori; Younghan, F₁(YB3433-3B-5 / YB3135-3B-2-3) and an elite breeding line YB3135-3B-2-3; Youho, a pedigree Suwon339 and Suwon355; Youhan, a pedigree Milyang100 and an elite an breeding line SB951050-B-B-B-77-0-P-3; and Nokyang, cultivar Nakyoungbori and an elite breeding line SB77368-B-145. After preliminary and advance yield testing for 2 years, the promising lines were subsequently evaluated for earliness (heading date) and forage yield during 3 years in several parts of the country, and each of the cultivars finally named were released to farmers of Korea.

Results

The characteristics of barley cultivars required for whole crop silage are not the same as those required for cereal food. For example high biomass of whole plant including leaves, culms, and grain is more important than grain weight compared to the cultivars grown for food (Sakai *et al.* 2003). We have developed forage barley cultivars with cattle's favourite characters (Choi *et al.* 2007a; Karren *et al.* 1994) such as smooth awn, hooded spike, and auricleless types. Cultivar Yuyeon (Choi *et al.* 2007 a), Youho (unpublished) and Youhan (unpublished) have hood type spike. Cultivar WooHo (Kim *et al.* 2007) and Dami (Park *et al.* 2009) have smooth awn and auricleless type plant, respectively. We also developed rough awn type barley cultivars, such as Yongyang (Park *et al.* 2008), Sunwoo (Park *et al.* 2008), Sangwon (Park *et al.* 2008), Soman (Choi *et al.* 2007 b), Younghan (Park *et al.* 2011), and Nokyang (unpublished), showing early-maturing and high-yielding. They yielded about 12 to 10 t/ha of dry matter yield (average 33 t/ha in fresh matter yield, Table 1), and were evaluated for forage quality which showed a higher silage quality (TDN, ADF, NDF, CP, etc.) for whole crop barley use. Based on these results, it is concluded that barley with smooth awn, hooded, and auricleless could be suitable sources in breeding for whole crop forage use.

Conclusion

We have developed 11 barley cultivars with ruminant-palatable plant characters, such as smooth awn, hooded spike and auricleless, and high biomass as a whole crop as compared with cereal barley. These cultivars have advantages in forage or total digestible nutrients (TDN) yield per unit area. Moreover, it is very economic in land utilization because forage barley is produced in a cropping system following rice during the winter season. The forage barley cultivation requires heavy fertilization to obtain higher yield, it is a break crop, but negatively affects rice grain quality due to increasing protein content. Therefore, improved cultivation technologies are needed to overcome the problem of degrading rice grain quality by residual fertilizer after forage barley cultivation in paddy fields. In the near future, we also plan to develop forage barley cultivars with awn-less,

Table 1. Average agronomic characteristics and forage yield of forage barley cultivars in Korea. Data collected from 4 to 7 different locations for 3 years.

Cultivar	Released year	Heading date	Fresh yield (t/ha)	DM ¹ yield (t/ha)	TDN ² (%)	Characteristics
Olbori	-	April 27	-	9.9	63.0	Cereal barley
Youngyang	2002	May 1	37.5	11.7	65.5	Rough awn
Sunwoo	2002	May 8	36.0	11.0	61.8	Rough awn
Sangwon	2004	April 24	36.0	11.8	63.7	Rough awn
Wooho	2005	April 29	36.1	11.0	65.9	Smooth awn
Yuyeon	2006	April 27	33.5	10.8	66.4	Hooded spike
Soman	2006	April 23	31.7	10.6	67.9	Rough awn
Dami	2007	April 30	32.9	12.0	66.4	Auricleless
Youngghan	2008	April 26	33.4	12.0	67.4	Rough awn
Youho	2008	April 24	30.9	11.6	69.2	Hooded spike
Youhan	2012	June 2	36.9	12.6	67.7	Hooded spike
Nokyang	2012	June 4	37.3	11.7	68.5	Rough awn

¹DM = Dry matter, ²TDN = Total digestible nutrients. Sowing and harvest were mid-October and the end of May (early yellow stage), respectively.

high biomass and lysine content, good silage quality such as high sugar content, and solid and brittle stem types.

References

- Choi JS, Kim JG, Baek SB, Park KH, Kwon YU, Park HH, Kang MS, Park TI, Heo HY, Seo JH, Cheong YK, Sung BR, Kim JG, Lee JJ, Kim SJ, Ryu IM, Ju JI, Kim DH, Jung KY, Lee SH (2007 a) A ruminant-palatable hood type barley cultivar 'Yuyeon' for whole-crop-forage use. *Korean Journal Breeding Science* **39**, 242-243.
- Choi JS, Kim JG, Baek SB, Park KH, Kwon YU, Park HH, Kang MS, Park TI, Heo HY, Seo JH, Cheong YK, Sung BR, Kim JG, Lee JJ, Kim SJ, Ryu IM, Ju JI, Kim DH, Jung KY, Lee SH (2007 b) A new early maturing and high yielding barley cultivar 'Soman' for whole crop barley. *Korean Journal Breeding Science* **39**, 240-241.
- Karren, DB, Goonewardene LA, Bradley JA (1994) The effect of feed type on mouth lesions in slaughter cattle. *Canadian Journal of Animal Science* **74**, 571-573.
- Kim JG, Choi JS, Park HH, Baek SB, Kang MS, Park KH, Kwon YU, Heo HY, Seo SJ, Nam JH, Lee JJ, Cheong YK, Sung BR, Kim JG, Ryu IM, Ju JI, Kim DH, Jung KY, Lee SH (2007) A new smooth awn barley cultivar, 'Wooho' suitable for whole crop forage use. *Korean Journal Breeding Science* **39**, 238-239.
- Park TI, Han OK, Seo JH, Choi JS, Park KH, Kim JG (2008) New barley cultivars with improved morphological characteristics for whole crop in Korea. *Journal of Korean Grassland and Forage Science* **28**, 193-202.
- Park, TI, Seo JH, Han OK, Choi JS, Kim JG, Park JC, Kim HS, Heo HY, Baek SB, Kwon YU, Park HH, Kang MS, Park KG, Suh SJ (2009) A new auricleless barley cultivar 'Dami' for whole crop forage. *Korean Journal Breeding Science* **41**, 349-353.
- Park TI, Seo JH, Han OK, Kim KH, Oh YJ, Park KH, Choi JS, Park JC, Park HH, Kim HS, Kim JG, Kim WH, Park NG, Jeung JH, Ju JI, Kim SY, Kim DH (2011) A new whole crop barley cultivar 'Youngghan' with good quality, resistance to lodging and viral disease and high-yielding. *Journal of Korean Grassland and Forage Science* **43**, 184-189.
- Sakai MS, Iida S, Maeda H, Sunohara Y, Nemoto H, Imbe T (2003) New rice varieties for whole crop silage use in Japan. *Breeding Science* **53**, 271-275.