



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

23rd International Grassland Congress

Nitrogen Retention and Microbial Protein Yield of *Desmodium uncinatum*, *Mucuna pruriens* and *Vigna unguiculata* Forage Legumes in Goats

Joseph J. Baloyi
University of Venda, South Africa

Katsande Simbarashe
University of Venda, South Africa

F. V. Nherera-Chokuda
Agriculture Research Centre Irene, South Africa

N. T. Ngongoni
Zimbabwe Open University, Zimbabwe

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/23/2-1-2/2>

The 23rd International Grassland Congress (Sustainable use of Grassland Resources for Forage Production, Biodiversity and Environmental Protection) took place in New Delhi, India from November 20 through November 24, 2015.

Proceedings Editors: M. M. Roy, D. R. Malaviya, V. K. Yadav, Tejveer Singh, R. P. Sah, D. Vijay, and A. Radhakrishna

Published by Range Management Society of India

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Nitrogen retention and microbial protein yield of *Desmodium uncinatum*, *Mucuna pruriens* and *Vigna unguiculata* forage legumes in goatsJoseph J Baloyi^{1*}, S. Katsande¹, F. V. Nherera-Chokuda² and N. T. Ngongoni³¹University of Venda, Thohoyandou, South Africa²Agricultural Research Council, Irene, Pretoria, South Africa³Zimbabwe Open University, Harare, Zimbabwe*Corresponding author e-mail: joseph.baloyi@univen.ac.za**Keywords:** Goats, Legumes, Microbial protein, Nitrogen retention**Introduction**

Forage legumes have high potential degradability, indicating that they might need to stay in the rumen for a longer time (Molina-Alcaide *et al.*, 1996) to increase microbial protein yield. Microbial protein contributes about two thirds of the amino acids absorbed by ruminants (Pathak 2008). Protein supplements which have proper levels of rumen undegradable protein provide growth limiting amino acids like lysine and methionine (Rezai *et al.*, 2012). Nyambati *et al.* (2003) showed that milk production can be increased by supplementing with velvet bean, highlighting the importance of integrating legumes into the low-input, mixed cropping systems in the tropics. The current study was to determine total microbial protein yield from *Vigna unguiculata* (cowpea), *Desmodium uncinatum* (silverleaf desmodium) and *Mucuna pruriens* (velvet bean) legume forages in goats.

Materials and Methods

Four growing indigenous Nguni-type male goats (29±0.5 kg) were used in a 4x4 Latin- square experimental design. The goats were fed veld hay supplemented with legume forage dietary treatments of either 16.3% CP commercial goat feed (CF) *ad libitum* or supplemented with velvet bean (VB) or silverleaf desmodium (SD) or cowpea (CW). Each goat was fed each treatment for 15 days (10 days of adaptation and 5 days of measurement), before being switched to the next treatment. Microbial protein yield was determined estimating allantoin in the urine using the purine derivatives technique as described by Chen and Gomes (1992).

Results and Discussion

Nitrogen retention was negative for goats supplemented with velvet bean and silverleaf desmodium, but positive with cowpea supplement and commercial feed. There were no significant differences ($P > 0.05$) in microbial protein synthesis and animals on the commercial feed diet were as efficient as those supplemented with silverleaf cowpea and velvet bean in microbial protein production (Table 1). Digestible microbial true protein, digestible organic matter in the rumen, microbial true protein, microbial nitrogen yield and allantoin levels were all significantly different ($P < 0.05$).

Table 1: Nitrogen retention (g/d) and excretion of purine derivatives and the microbial protein production in goats

Parameter	Treatment group				SEM
	VB	SD	CW	CF	
N retention	-1.29 ^c	-0.46 ^b	0.84 ^{ab}	2.13 ^a	0.40
Allantoin (mmol.d ⁻¹)	3.63 ^d	4.27 ^b	3.80 ^c	4.67 ^a	0.04
Microbial nitrogen yield (³ MNY) (g.d ⁻¹)	2.64 ^d	3.10 ^b	2.76 ^c	3.40 ^a	0.03
Digestible Microbial true protein (g.d ⁻¹)	11.2 ^d	13.2 ^b	11.7 ^c	14.43 ^a	0.12
Digestible organic matter in the rumen (DOMR)	136 ^c	162 ^{ab}	156 ^{ab}	192 ^a	0.12
¹ E _{mns} (g.kg ⁻¹ DOMR)	2.21 ^a	2.01 ^a	2.06 ^a	1.95 ^a	0.23

^{a,b,c,d}Means in the same row with different superscripts are significantly different ($P < 0.05$), ¹E_{mns} = efficiency of microbial nitrogen synthesis (AFRC, 1993), SEM = standard error of means

Conclusion

Nitrogen retention was negative for goats supplemented with velvet bean and silverleaf, but positive with cowpea supplement and commercial feed. Digestible microbial true protein, digestible organic matter in the rumen, microbial true protein, microbial nitrogen yield and allantoin were all significantly different amongst the dietary groups.

References

Chen, X. B. and M. J. Gomes. 1992. Estimation of microbial protein supply to sheep and cattle based on urinary excretion of purine derivatives - an overview of technical details. Occasional Publication, International Feed Resources Unit. Aberdeen: Rowett Research Institute. 20 pp.

- Molina-Alcaide E., M. R. Weisbjerg and T. Hvelplund. 1996. Degradation characteristics of shrubs and the effect of supplementation with urea or protein on microbial production using continuous culture system. *Journal of Animal Physiology and Animal Nutrition* 75:121- 132.
- Nyambati, E. M., L. E. Sollenberger and W. E. Kuncle. 2003. Feed intake and lactation performance of dairy cows offered napier grass supplemented with legume hay. *Livestock Production Science* 83: 179–189.
- Pathak, A. K. 2008. Various factors affecting microbial protein synthesis in the rumen. *Veterinary World* 1: 186-189.
- Rezai, F., F. Zamani, and M. Vatankhah. 2012. Effect of rumen undegradable protein (RUP) on colostrum quality and growth of Lori Bakhtiari lambs. *Global Veterinaria* 8: 93-100.

Acknowledgement

We greatly thank the University of Venda, South Africa for sponsoring this research work