

*25th International Symposium on Analytical and Environmental Problems***SELF-ASSEMBLY IN WATER OF BULKY OCTAHEDRAL COORDINATION
COMPLEXES BASED ON RH(III) METAL CENTER****Carmen Cretu¹, Maria A. Spirache¹, Adél Len^{2,3}, Zoltán Dudás⁴, Alessandra Crispini⁵,
Elisabeta I. Szerb¹**

¹*“Coriolan Dragulescu” Institute of Chemistry, Romanian Academy, 24 Mihai Viteazu Bvd.
300223 – Timisoara, Romania, e-mail: cretucami@yahoo.com*

²*Nuclear Analysis and Radiography Department/Centre for Energy Research,
Konkoly-Thege 29-33, 1121 Budapest, Hungary*

³*Faculty of Engineering and Information Technology/University of Pécs, Boszorkány út 2,
7624 Pécs, Hungary*

⁴*Neutron Spectroscopy Department/Wigner Research Centre for Physics, Konkoly-Thege 29-
33, 1121 Budapest, Hungary*

⁵*Università della Calabria, Dipartimento di Chimica e Tecnologie Chimiche, 87030
Arcavacata di Rende, Italy*

Abstract

Metal complexes able to self-assembly in water into chromonic-type liquid-crystalline phases are progressively attracting attention because of their unique features consisting in the combination of properties deriving from liquid crystals (self-ordering, ease of alignment, sensitivity to changing conditions and additives) coupled with the optical and electro-optical properties brought by the metal centre. [1,2]

In the last decade, rhodium (III) cyclometallated complexes were researched for their electrocatalytic activity, fluorescent properties and especially for medicinal applications as drug delivery and therapeutic agents. [3(a-d)] Therefore, supramolecular organized aqueous systems built up with these compounds should bring important developments in bio-related fields. Herein we report new Rh(III) cyclometallated complexes that show the ability to self-assemble into chromonic-like phases at relatively low concentration in water. Their supramolecular arrangements in mesophase will be proposed based on optical polarization microscopy (POM), X-ray diffraction studies on single crystals and Small Angle Neutron Scattering (SANS) measurements in the mesophase. The role of different molecular moieties will be discussed in the self-assembling with respect to previous reported structurally analogues Ir(III) complexes [1,2].

Acknowledgements: The Romanian authors acknowledge the support of the “*Coriolan Dragulescu*” Institute of Chemistry, Project 4.1.

References

- [1] Y.J. Yadav, B. Heinrich, G. De Luca, A.M. Talarico, T.F. Mastropietro, M. Ghedini, B. Donnio, E.I. Szerb, *Adv. Optical Mater.* 1 (2013) 844.
[2] L. Ricciardi, T.F. Mastropietro, M. Ghedini, M. La Deda, E.I. Szerb, *J. Organomet. Chem.* 772-773 (2014) 307.
[3] a) J. Kima, E. Rajkumara, S. Kima, Y.M. Parka, Y. Kima, S.-J. Kima, H.J. Leeb, *Catal. Today* 295 (2017) 75; b) A. Nano, A.N. Boynton, J.K. Barton, *J. Am. Chem. Soc.* 139(48) (2017) 17301; c) J. Markham, J. Liang, A. Levina, R. Mak, B. Johannessen, P. Kappen, C.J. Glover, B. Lai, S. Vogt, P.A. Lay, *Eur. J. Inorg. Chem.* 12 (2017) 1812; d) C.-H. Leung, H.-J. Zhong, D.S.-H. Chan, M. Dik-Lung, *Coord. Chem. Rev.* 257 (2013) 1764.