

*25th International Symposium on Analytical and Environmental Problems***DETERMINATION OF HYDROXYL RADICALS USING COUMARIN AND COUMARIN-3-CARBOXYLIC ACID DURING GAMMA RADIOLYSIS AND HETEROGENEOUS PHOTOCATALYSIS****Máté Náfrádi¹, László Wojnárovits², Erzsébet Takács², Tünde Alapi¹**¹*Department of Inorganic and Analytical Chemistry, University of Szeged, H-6720 Szeged, Dóm tér 7, Hungary*²*Institute for Energy Security and Environmental Safety, Centre for Energy Research, Hungarian Academy of Sciences, Budapest, Hungary
e-mail: nafradim@chem.u-szeged.hu***Abstract**

Coumarin and 3-carboxycoumarinic acid, two fluorescent probes commonly used for HO• detection has been used during gamma radiolysis and heterogeneous photocatalysis. The O₂ dependency and the radiation yield of their hydroxylated fluorescent products (7-hydroxycoumarin and 7-hydroxy-3-carboxycoumarinic acid) has been investigated during gamma radiolysis. The radiation yields were found to be 1.2(±0.2) % in O₂-free solutions, while it was 2.9 (±0.06) % in the presence of O₂, proving the importance of peroxy radicals in the formation of these products. The results obtained from radiolysis experiments, were employed during heterogeneous photocatalysis performed with commercial TiO₂ catalyst. The effect of dissolved O₂ was also investigated, as its electron scavenging role during photocatalysis is also important. The formation rate of HO• during photocatalysis was calculated from the formation rate of the fluorescent products, and were found to be 1.8×10⁻⁷ mol dm⁻³ s⁻¹, while the quantum efficiency for its formation is 0.0038.

Introduction

Advanced oxidation processes (AOPs) have been investigated in the last few decades for their possible use as an additional wastewater purification method. During AOPs different reactive species form, the most important one is the hydroxyl radical (HO•), due to its high reaction rate with most organic pollutants. However several methods can be employed, like time-resolved spectroscopy, ESR, the determination of HO• formation rates is a complicated task. Fluorescent probes, like terephthalic acid or coumarines, have also been applied during heterogeneous photocatalysis to evaluate the formation rate of HO•. [1-5]

In this study coumarin (COU) and coumarin-3-carboxylic acid (3-CCA) have been used as a fluorescent probe for determination of the HO• formation rate. They are reported to form highly fluorescent hydroxylated products in their reaction with HO•, 7-hydroxycoumarin (7-HO-COU) and 7-hydroxy-3-carboxycoumarinic acid (7-HO-3-CCA), respectively. The formation rate of both hydroxylated products are reported to be dependant on dissolved O₂. In the presence of O₂ they form via peroxy type radicals, while in the absence of O₂ they form via dismutation, significantly reducing their formation rate. [1-3]

The application of COU and 3-CCA for HO• detection was investigated during two, different AOPs. In the case of gamma radiolysis the formation rate of all reactive species (HO•, e_{aq}⁻, H•) is well determined, since the values of their radiation yields (G value) are well known. Heterogeneous photocatalysis is also a highly researched field of AOPs, but the reaction mechanisms are often not clear. The transformation of organic compounds is mostly related to the reactions with HO•. In addition, the reactions initiated directly by the photogenerated charges (h_{vb}⁺ and e_{cb}⁻) has to be taken into consideration too. The reactions take place on the surface or close to the surface of photocatalyst, and consequently the interactions between the photocatalyst and substrate may have an important role.

The aim of this study is to determine the radiation yields of 7-HO-COU and 7HO-3-CCA, and investigate the effect of dissolved O₂ during gamma radiolysis. Based on these result, the formation rate of HO• and the quantum yield of the HO• formation ($\Phi_{\text{HO}\cdot}$) may be determined during the heterogeneous photocatalysis. By comparing the transformation of the non-adsorbed COU, and the well adsorbed 3-CCA, we may investigate the importance of adsorption on the reactions of these substrates with HO•.

Experimental

In the gamma-radiolysis experiments a ⁶⁰Co gamma source was used in a panoramic type irradiator, the dose rate was 1.48 Gy min⁻¹. The solutions of COU and 3-CCA were irradiated in sealed ampoules, which were saturated with either O₂, N₂O or N₂. All experiments were performed in 10⁻⁴ mol dm⁻³ solutions of COU and 3-CCA in pH = 7.0 (in 0.01 mol dm⁻³ phosphate buffer).

Photocatalysis experiments were performed in a glass reactor. 1.0 g dm⁻³ TiO₂ Aeroxide P25 (Acros Organics) was added to the 250 cm³ solutions, and irradiated using a fluorescent UV light source (GCL303T5/UVA, Lighttech) emitting in the 300-400 nm range. The photon flux of the lamp was 1.20×10⁻⁵ mol_{photon} min⁻¹, determined by ferrioxalate actinometry. Since 3-CCA adsorbed on the photocatalyst surface (≈30 % adsorbed on TiO₂), NaF was added to the samples, for the desorption of the analytes. All samples were centrifuged at 15000 RPM, and filtered using 0.22 μm syringe filters (FilterBio PVDF-L).

The transformation of COU and 3-CCA ($\lambda_{\text{max}}^{\text{COU}} = 277 \text{ nm}$, $\lambda_{\text{max}}^{3\text{-CCA}} = 291 \text{ nm}$, $\epsilon^{\text{COU}} = 10300 \text{ dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$, $\epsilon^{3\text{-CCA}} = 12170 \text{ dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$) has been followed using UV-Vis spectrophotometry (Agilent 8453). The formation of 7-HO-COU and 7-HO-3-CCA were followed using fluorescence spectroscopy (Hitachi F4500) at 455 and 447 nm, respectively. The initial transformation rates of COU and 3-CCA were determined from linear regression fits to the actual concentration versus the duration of irradiation, up to 15 % conversion. The initial formation rates of 7-HO-COU and 7-HO-3-CCA were obtained from the linear regression fits to the actual concentration versus the duration of irradiation.

Results and discussion

First the effect of O₂ on the formation of 7-HO-COU and 7-HO-3-CCA was investigated. In the case of radiolysis mainly HO• and e_{aq}⁻ forms from water. In the presence of O₂ e_{aq}⁻ transforms into O₂^{-•}. Since both O₂^{-•} and its protonated form, HO₂[•] have a low reactivity towards organic compounds, mainly HO• is responsible for the transformation of COU in this case. Moreover, from carbon centered radicals peroxy radicals form immediately. The formation of peroxy radicals opens a new pathway for the formation of hydroxylated products via unimolecular HO₂[•] elimination. In O₂-free solutions both the HO• and the e_{aq}⁻ are able to initiate the transformation of COU. Without O₂, the formation of 7-HO-COU happens via bimolecular dismutation. The formation rate of HO• in the presence of O₂, air or N₂ can be calculated ($r_0^{\text{HO}\cdot} = 6.91 \times 10^{-9} \text{ mol dm}^{-3} \text{ s}^{-1}$). In N₂O saturated solutions e_{aq}⁻ transforms into HO•, and doubling the HO• yield ($r_0^{\text{HO}\cdot} = 1.33 \times 10^{-8} \text{ mol dm}^{-3} \text{ s}^{-1}$).




Figure 1. Concentration of COU (A) and 3-CCA (B) as a function of dose in O₂ (■), N₂O (■) and N₂ (■) saturated solutions, and the formation of 7-HO-COU during gamma radiolysis in O₂ (◇), N₂O (◇) and N₂ (◇) saturated solutions.

The effect of the different gases on the transformation rate of COU and 3-CCA were negligible. Despite the similar transformation rates, the formation rate of the hydroxylated products show significant differences. The lowest formation rate can be observed in O₂-free solution, due to the lack of the possibility of peroxy radical formation, and probably because of the significant contribution of e_{aq}^- to the transformation of COU/3-CCA, which do not result in hydroxylated products. The formation rates in N₂O saturated solutions are greatly increased, due to the increased HO• formation rate. The formation rates are even higher in the presence of O₂, despite the lower HO• formation, proving the importance of peroxy radicals in the formation of the hydroxylated products. (Figure 1. and Table 1)

From the formation rates of the hydroxylated products and the formation rate of HO•, the radiation yield for both fluorescent product can be calculated. In N₂ saturated solutions 0.94 and 1.35 %, in N₂O saturated solutions 1.16 and 1.29 %, while in O₂ saturated solutions 2.99 and 2.86% of HO• produces 7-HO-COU from COU and 7-HO-3-CCA from 3-CCA, respectively.

Table 1. Initial transformation rates of COU and 3-CCA and initial formation rates of 7-HO-COU and 7-HO-3-CCA during gamma radiolysis

	r_0^{COU} ($\times 10^{-9} \text{ mol dm}^{-3} \text{ s}^{-1}$)	$r_0^{7-HO-COU}$ ($\times 10^{-10} \text{ mol dm}^{-3} \text{ s}^{-1}$)	r_0^{3-CCA} ($\times 10^{-9} \text{ mol dm}^{-3} \text{ s}^{-1}$)	$r_0^{7-HO-3-CCA}$ ($\times 10^{-10} \text{ mol dm}^{-3} \text{ s}^{-1}$)
O ₂	3.35	2.06	3.23	1.98
Air	4.33	1.55	2.40	1.71
N ₂	3.38	0.64	2.73	0.93

During heterogeneous photocatalysis the main reactive species is the HO•. The photogenerated $h_{\nu b}^+$ and e_{cb}^- pair may also react with organic compounds, especially when there is special interaction between the substrate and the catalyst surface. COU and 3-CCA have different adsorption properties, as COU do not adsorb on the catalyst surface, as opposed to 3-CCA, due to the strong interaction between its carboxyl groups and Ti=OH surface groups of photocatalyst.




Figure 2. Transformation of COU (A) and 3-CCA (B) as a function of time, in O₂ (■), air (■) and N₂ (■) saturated suspensions, and the formation of 7-HO-COU during heterogeneous photocatalysis in O₂ (◇), air (◇) and N₂ (◇) saturated suspensions.

O₂ plays a crucial role as an electron scavenger, hindering charge recombination, and helping HO• formation via O₂•⁻. In the case of COU there was no difference in the transformation rates determined in O₂ saturated and airted suspensions. At the same time, the formation rate of 7-HO-COU is nearly 1.5 times greater in the case of higher dissolved O₂ concentration. In the case of 3-CCA the transformation rate is also higher with 25% in the case of O₂ saturated suspension, while the formation rate of 7-HO-3-CCA increased with 33%. In N₂ saturated suspensions the transformation of COU and the formation of 7-HO-COU is negligible. This suggests, that direct charge transfer reactions have a low probability, and HO• formation is negligible. The transformation rate of the well adsorbed 3-CCA is 47% of the value measured in O₂ saturated suspension, and there is no 7-HO-3-CCA formation. This suggests that, 3-CCA transformation can happen via direct charge transfer, but this way does not result in hydroxylated products.

Table 2. Initial transformation rates of COU and 3-CCA and initial formation rates of 7-HO-COU and 7-HO-3-CCA during heterogeneous photocatalysis

	$r_0^{\text{COU}} \times 10^{-8}$ (mol dm ⁻³ s ⁻¹)	$r_0^{7\text{-HO-COU}} \times 10^{-9}$ (mol dm ⁻³ s ⁻¹)	$r_0^{3\text{-CCA}} \times 10^{-8}$ (mol dm ⁻³ s ⁻¹)	$r_0^{7\text{-HO-3-CCA}} \times 10^{-9}$ (mol dm ⁻³ s ⁻¹)
O₂	6.15	2.55	8.98	2.77
Air	6.08	1.73	7.10	2.08
N₂	0.29	-	4.23	-

Using the results obtained from gamma radiolysis, the formation rate of HO• in the case of O₂ saturated TiO₂ suspension was found to be 1.78×10^{-7} and 1.86×10^{-7} mol dm⁻³ in the case of COU and 3-CCA respectively. Assuming that, TiO₂ absorbs completely the emitted photons from the light source, the quantum yield for the formation of HO• is 0.0037 and 0.0039, respectively.

Conclusion

- The formation of 7-HO-COU from COU and 7-HO-3-CCA from 3-CCA requires HO•, while dissolved O₂ highly enhances their formation rate.
- The radiation yield of 7-HO-COU from COU and 7-HO-3-CCA from 3-CCA were determined in the case of gamma-radiolysis
- Based on radiation yields of the hydroxylated products, and the photon flux of the light source, the formation rate of HO• and its quantum efficiency has been determined during heterogeneous photocatalysis, using TiO₂ as photocatalyst

Acknowledgements

This work was financially supported by Industrial Research and development Projects of Hungarian-Indian cooperation (TÉT_15_IN-1-2016-0013). This publication was supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences, ÚNKP-19-3-SZTE-207 and UNKP-19-4-SZTE-115, new national excellence programs of the Ministry for Innovation and Technology.

References

- [1] Loutit, G., Foley, S., Cabillic, J., Coffigny, H., Taran, F., Valleix, A., Renault, J.P., Pina, S., *Radiation Physics and Chemistry* 72 (2005) 119–124.
- [2] Manevich, Y., Held, K., Biaglow, J.E., *Radiation Research* 148 (1997) 580–591.
- [3] Yamashita, S., Baldacchino, G., Maeyama, T., Taguchi, M., Muroya, Y., Lin, M., Kimura, A., Murakami, T., Katsumura, Y., *Free Radical Research* 46 (2011) 861–871.
- [4] Zhang, J., Nosaka, J., *Applied Catalysis B: Environmental* 166–167 (2015) 32–36.
- [5] Nosaka Y., Nishikawa, M., Nosaka, A.Y., *Molecules* 19 (2014) 18248–18267.