



Journal of Applied and Natural Science
11(2): 575 - 580 (2019)
ISSN : 0974-9411 (Print), 2231-5209 (Online)
journals.ansfoundation.org

Effect of cadmium on glycogen content in muscle, liver, gill and kidney tissues of freshwater fish *Channa punctatus* (Bloch)

Seema Tewari*

Department of Zoology, I.T.P.G. College, Faizabad Road, Lucknow - 226007 (Uttar Pradesh), India

Sandeep Bajpai

Department of Zoology, B.V.B.G. Degree College, Gontinagar, Lucknow- 226010 (Uttar Pradesh), India

Madhu Tripathi

Department of Zoology, University of Lucknow-226007 (Uttar Pradesh), India

*Corresponding author. E-mail: tewariseema13@gmail.com

Abstract

Aquatic environment gets polluted by heavy metals because of their environmental persistence and ability to bioaccumulate in aquatic organisms. Cadmium is a ubiquitous toxic heavy metal, biologically non-essential element, it is not biodegradable and has a very long biological half-life. The aim of the present study was to assess the glycogen content in muscle, liver, gill and kidney of *Channa punctatus* exposed to sublethal concentrations of cadmium chloride after 4, 7, 15 and 30 days of exposure. The results clearly showed significant decrease in the glycogen levels in the experimental fish *C. punctatus*. Decrease in muscle glycogen was found highly significant ($P < 0.001$) after 30 days in both low concentration (36.823 mg/L) 6.12 ± 0.41 mg/g and in high concentration (73.646 mg/L) 4.04 ± 0.32 mg/g in comparison to control. Decrease in liver glycogen content was found highly significant ($P < 0.001$) after 30 days in high concentration 9.12 ± 0.49 mg/g when compared with control. The decrease in gill glycogen content after 30 days exposure was found highly significant ($P < 0.001$) 1.36 ± 0.13 mg/g in low concentration and in high concentration 0.79 ± 0.25 mg/g in comparison to control. Decrease in kidney glycogen content was found highly significant ($P < 0.001$) at 30 days in low concentration 3.92 ± 0.05 mg/g and in high concentration 2.81 ± 0.20 mg/g in comparison to control. The influence of toxicant cadmium chloride in selected tissues of fish was taken into account in evaluating fish response against stressor. Hence, we can use glycogen content as biomarker of cadmium stress in fish.

Keywords: Cadmium, *Channa punctatus*, Fish, Glycogen content, Heavy metal

Article Info

DOI: [10.31018/jans.v11i2.2115](https://doi.org/10.31018/jans.v11i2.2115)

Received: May 5, 2019

Revised: June 7, 2019

Accepted: June 10, 2019

How to Cite

Tewari, S. *et al.* (2019). Effect of cadmium on glycogen content in muscle, liver, gill and kidney tissues of freshwater fish *Channa punctatus* (Bloch.). *Journal of Applied and Natural Science*, 11(2): 575 - 580 <https://doi.org/10.31018/jans.v11i2.2115>

INTRODUCTION

Pollution of water is an important dimension of environmental degradation. The contamination of aquatic ecosystem by toxic materials is closely connected with increased concentration of different types of pollutants. Intense activities in industrial and agricultural sectors has led to increased in the levels of heavy metals in natural waters (Gumgum *et al.*, 1994; Nimmo *et al.*, 1998; Jordao *et al.*, 2002; Rauf *et al.*, 2009; Javed and Usmani, 2013). The contamination of water bodies with heavy metals has become a source of great concern not only because of their threat to aquatic life especially fishes (Opaluwa *et al.*, 2012; Bawuro, *et al.*, 2018; Ali *et al.*, 2019) but also due to the public health implications of such contaminations (Sim *et al.*, 2016; Baharom and Ishak, 2015).

Non degradable heavy metals are regarded as hazardous to aquatic ecosystem because of their

environmental persistence and their tendency for bioaccumulation (Das *et al.*, 2001; Agrahari and Gopal, 2007). Cadmium (Cd) is considered as a nonessential element (Viarengo, 1985) that has gained great importance from the toxicological (USDHHS-ATSDR, 1993; Waisberg *et al.*, 2003; USEPA, 2016) and ecotoxicological (WHO, 1992) point of view. It has been listed in the "Black list" of European community (Mason, 1996). Cadmium occurs naturally in ores together with zinc, lead and copper. It is widely used in mining, metallurgical operations, electroplating industries and manufacturing vinyl plastics. Effluents from such activities are a source of cadmium into aquatic environment.

Fish, as they come into intimate contact with large amounts of polluted water can be used as early warning biological indicators of polluted environment (Viana and Lucena Frédo, 2014; Plessl *et al.*, 2017).

To understand the mode of action of toxicants on the aquatic animals biochemical investigations are done. Stress condition cause alterations in metabolic cycles, therefore it becomes necessary to understand the significance of these variations in the organic content of tissues. The present study was planned to investigate the effect of sublethal concentrations of Cd on glycogen content in muscle, liver, gill and kidney of *C. punctatus* after 4, 7, 15 and 30 days of Cd exposure.

MATERIALS AND METHODS

Live and healthy specimens of *C. punctatus*, a fresh water murrel, were procured from local resources having 15-18 cms length and 25±2gm weight and brought to the laboratory. Before introducing into the aquarium, fishes were treated with 0.1% KMnO₄ solution to avoid any cutaneous infection. Fishes were acclimated to laboratory conditions for a period of two weeks. During acclimation water was changed daily and fish were fed ad libitum. *Channa punctatus* was selected as the model organism for this study because of its availability, survival for longer period under laboratory conditions, easy handling etc. Fish were maintained following standard maintenance procedure (APHA, 2012) in glass aquaria.

Experimental design: A total of 108 fishes were used for the experiment. Fish were maintained in static renewal condition. They were divided into three groups of thirty six fishes each, which served as replicates. Each group was further subdivided into three subgroups having twelve fishes in each aquaria. The fish in the subgroup I and subgroup II were treated with 1/5th LC₅₀ (73.646mg/L) and 1/10th (LC₅₀ 36.823mg/L) respectively. Subgroup III served as control. Control group was not treated with Cd. The aquarium of both control and exposed groups were cleansed on every two days one hour after feeding period to reduce contamination with food remains. All experimental waters were completely renewed on every two days.

Three fishes from each subgroup were taken after 4, 7, 15 and 30 days of metal exposure and were sacrificed and tissues like liver, muscle, kidney and gill were excised rapidly and processed for biochemical estimations. The glycogen content in the tissues was estimated by the method of Nicholas *et al.* (1956). Values were compared using student 't' test.

RESULTS

After exposure to sublethal concentrations of Cd, the level of total glycogen content was altered in muscle, liver, gill and kidney of the fish, *C. punctatus*.

Muscle glycogen: Total glycogen content in muscle of control fish and fishes exposed to lower and higher concentrations were recorded as 7.84mg/g, 7.50mg/g and 7.15mg/g respectively after 4 days.

At the end of 7 days of exposure, glycogen was noticed 7.98mg/g in control fish, 7.35mg/g in lower and 6.98mg/g in higher concentrations exposed fishes. Similarly at the end of 15 days of exposure, it was found to be 8.16mg/g in control, 7.10mg/g in lower concentration and 6.25mg/g in higher concentration exposed fishes. Whereas at the end of 30 days of exposure, it was noticed 8.32mg/g in control, 6.12mg/g in lower concentration and 4.04mg/g in fishes exposed to higher concentration. Decrease in muscle glycogen was found highly significant (P<0.001) in comparison to control after 15 days in high concentration and 30 days in both low (36.823 mg/L) and high concentrations (73.646 mg/L) (Table 1). These observations revealed that the decline in total glycogen content in muscle was directly proportional to the concentration of Cd and duration of exposure.

Liver glycogen: Total glycogen content in liver of control fish and fishes exposed to lower and higher concentrations were recorded as 12.05mg/g, 12.02mg/g and 11.52mg/g respectively after 4 days. At the end of 7 days of exposure it was noticed 12.20mg/g in control fish, 12.0mg/g in lower and 11.36mg/g in higher concentrations exposed fishes. Similarly at the end of 15 days of exposure, it was found to be 12.42mg/g in control, 11.15mg/g in lower concentration and 10.26mg/g in higher concentration exposed fishes. Whereas at the end of 30 days of exposure, it was noticed 12.56mg/g in control, 10.08mg/g in lower concentration and 9.12mg/g in fishes exposed to higher concentration.

The decrease in glycogen content was found slight significant (P<0.01) in low concentration (36.823 mg/L) exposed fishes at 7 days. At 15 days in low concentration it was highly significant (P<0.001). The decrease was slight significant (P<0.01) in low concentration (36.823 mg/L) and highly significant (P<0.001) in high concentration (73.646 mg/L) at 30 days of exposure (Table 2). The observations revealed that like muscle, decline in total glycogen content in liver was directly proportional to the concentration of Cd and duration of exposure.

Gill glycogen: Total glycogen content in gills of control fish and fishes exposed to lower and higher concentrations were recorded as 2.30mg/g, 2.23mg/g and 2.10mg/g respectively after 4 days. At the end of 7 days of exposure it was noticed 2.52mg/g in control fish, 2.40mg/g in lower and 2.14mg/g in higher concentrations exposed fishes. Similarly at the end of 15 days of exposure, it was found to be 2.56mg/g in control, 2.12mg/g in lower concentration and 1.82mg/g in higher concentration exposed fishes. Whereas at the end of 30 days of exposure, it was noticed 2.65mg/g in control, 1.36mg/g in lower concentration and 0.79mg/g in fishes exposed to higher concentration.

The decrease in glycogen content was found

slight significant ($P < 0.01$) at 15 days in high concentration exposed fishes. After 30 days the decline was found highly significant ($P < 0.001$) in fishes exposed to lower (36.823 mg/L) and higher concentration (73.646 mg/L) of Cd in comparison to control (Table 3). The observations revealed that the decline in total glycogen content in gill was directly proportional to the concentration of Cd and duration of exposure, though it was not so significant in the short duration exposures.

Kidney glycogen: Total glycogen content in kidney of control group and fishes exposed to lower and higher concentrations were recorded as 4.80mg/g, 4.68mg/g and 4.51mg/g respectively after 4 days. At the end of 7 days of exposure it was noticed 4.83mg/g in control fish, 4.50mg/g in lower and 4.26mg/g in higher concentrations exposed fishes. Similarly at the end of 15 days of exposure, it was found to be 4.91mg/g in control, 4.42mg/g in lower concentration and 3.84mg/g in higher concentration exposed fishes. Whereas at the end of 30 days of exposure, it was noticed 4.98mg/g in control, 3.92mg/g in lower concentration and 2.81mg/g in fishes exposed to higher concentration (Table 4).

The decrease in glycogen content was found slightly significant ($P < 0.01$) at 15 days in high concentration (73.646 mg/L) exposed fishes. At 30 days in low (36.823 mg/L) and high concentration (73.646 mg/L) exposed fishes the decline was found highly significant ($P < 0.001$). It is clear from the results that fishes exposed to lower and higher concentration of Cd showed decrease in glycogen content of kidney from the control values after 4, 7, 15 and 30 days of exposure, with maximum decrease after 30 days.

DISCUSSION

In the present study, glycogen content was depleted significantly in muscle, liver, gill and kidney tissues of *C. punctatus* after Cd exposure. These findings are well supported by the observations of earlier workers who have exposed various experimental models for Cd for different durations (Sastry and Sunita, 1982; Reddy et al., 1989; Kamaraju and Ramasamy, 2011; Prabhakar et al., 2012; Veeraiyah et al., 2013; Goswami et al., 2016). The various experimental models used were *C. punctatus*, fresh water field crab *Barytelphusa guerini*, fish *Hypophthalmichthys molitrix*, *Cirrhinus mrigala* and *Tilapia mossibica*.

Sastry and Subhadra (1982) reported decrease in glycogen content of muscle and liver in *Heteropneustes fossilis* after 15 and 30 days Cd exposure however they found increased glycogen content after 60 days exposure in both tissues. Similar observations have been reported by Malik et al. (1998) after zinc exposure in *C. punctatus*, Jagadeesan (1990) after mercuric chloride exposure in *Anabas testudineus* fingerlings and Karuppasamy (2000) after phenyl mercuric acetate exposure in *C. punctatus*.

Same findings were reported by Cicik and Engin (2005) in *Cyprinus carpio*, Sobha et al. (2007) in *Catla catla* and Kamaraju and Ramasamy (2011) in *H. molitrix* in various tissues after exposure to Cd for different durations.

Sujata (2015) reported significant decrease in protein and glycogen levels in reproductive organs of freshwater fish *C. punctatus* exposed to sub lethal concentrations of Cd for 30 days.

Metal intoxication in fishes usually results in glyco-

Table 1. Alteration in total glycogen content in muscles (mg/g) of *C. punctatus* after exposure to different concentrations of Cd.

Duration of exposure (Days)	Control group Not treated with Cd	Experimental groups	
		Low concentration (36.823 mg/L)	High concentration (73.646 mg/L)
4	7.84±0.21	7.50±0.38 ^{NS}	7.15±0.31 ^{NS}
7	7.98±0.28	7.35±0.34 ^{NS}	6.98±0.23*
15	8.16±0.36	7.10±0.20*	6.25±0.16***
30	8.32±0.15	6.12±0.41***	4.04±0.32***

Values are Mean ± S.E., N=6; N= Number of observations for each value; * $P < 0.05$ and *** $P < 0.001$ (in comparison to control); NS=non significant

Table 2. Alteration in total glycogen content in liver (mg/g) of *C. punctatus* after exposure to different concentrations of Cd.

Duration of exposure (Days)	Control group Not treated with Cd	Experimental groups	
		Low concentration (36.823 mg/L)	High concentration (73.646 mg/L)
4	12.05±0.10	12.02±0.26 ^{NS}	11.52±0.38 ^{NS}
7	12.20±0.34	12.0±0.28**	11.36±0.45 ^{NS}
15	12.42±1.10	11.15±0.21***	10.26±0.20 ^{NS}
30	12.56±0.23	10.08±0.60**	9.12±0.49***

Values are Mean ± S.E., N=6; N= Number of observations for each value; ** $P < 0.01$ and *** $P < 0.001$ (in comparison to control); NS=non significant

Table 3. Alteration in total glycogen content in gill (mg/g) of *C. punctatus* after exposure to different concentrations of Cd.

Duration of exposure (Days)	Control group Not treated with Cd	Experimental groups	
		Low concentration (36.823 mg/L)	High concentration (73.646 mg/L)
4	2.30±0.14	2.23±0.12 ^{NS}	2.10±0.35 ^{NS}
7	2.52±0.13	2.40±0.21 ^{NS}	2.14±0.24 ^{NS}
15	2.56±0.11	2.12±0.18 ^{NS}	1.82±0.19 ^{**}
30	2.65±0.20	1.36±0.13 ^{***}	0.79±0.25 ^{***}

Values are Mean ± S.E., N=6; N= Number of observations for each value; **P<0.01 and ***P<0.001 (in comparison to control); NS=non significant

Table 4. Alteration in total glycogen content in kidney (mg/g) of *C. punctatus* after exposure to different concentrations of Cd.

Duration of exposure (Days)	Control group Not treated with Cd	Experimental groups	
		Low concentration (36.823 mg/L)	High concentration (73.646 mg/L)
4	4.80±0.13	4.68±0.04 ^{NS}	4.51±0.12 ^{NS}
7	4.83±0.18	4.50±0.02 ^{NS}	4.26±0.31 ^{NS}
15	4.91±0.24	4.42±0.29 ^{NS}	3.84±0.14 ^{**}
30	4.98±0.12	3.92±0.05 ^{***}	2.81±0.20 ^{***}

Values are Mean ± S.E., N=6; N= Number of observations for each value; **P<0.01 and ***P<0.001 (in comparison to control); NS=non significant

gen depletion and is reported in several species of fishes such as *H. fossilis* (Qayyum and Shaffi, 1977), *Sarotherodon mossambicus* (Akhilendra Naidu, 1982), *C. punctatus* (Sastry and Sunita, 1983), *Labeo rohita* (Bengery and Patil, 1986), *L. rohita* (Radhakrishnaih et al., 1992), *Perca flavescens* (Levesque et al., 2002), *C. mrigala* (Bhilave et al., 2008).

It is considered that protein and carbohydrate stores are mobilized to a varying degree as a compensatory mechanism in response to energy stress during acute Cd exposure (Xuan et al., 2011). Most of the investigators have found that heavy metals cause glycogen depletion but the glycogenolytic response by different species of fish varies.

Like all other vertebrates, fish also store glucose in the form of glycogen in the liver, skeletal muscle, myocardium and brain (Leibson and Plietskaya, 1968). When required, the glycogen from these stores is broken down (Glycogenolysis) and transported to the muscle as glucose. On reaching the muscle, the glucose may be used at once or reconverted into glycogen.

A fall in the glycogen level clearly indicates its rapid utilization to meet the enhanced energy demands in fish exposed to toxicant. Some workers have also suggested that heavy metals could decrease the glycogen reserve in fish (Levesque et al., 2002) by affecting the activities of enzymes that play a role in carbohydrate metabolism. Sastry and Subhadra (1982) reported decrease in glycogen reserve of *H. fossilis* by stimulating the glycolytic enzymes like lactate dehydrogenase, pyruvate dehydrogenase and succinate dehydrogenase.

Decrease in carbohydrates is probably due to glycogenolysis and utilization of glucose to meet increased metabolic cost as suggested by Viswaraajan and Muthukrishnan (1988) in *Oreochromis mossambicus* under the stress of tannic acid. Several other reasons have been suggested for the decreased glycogen level in fishes after exposure to metals such as acute hypoxia (Heath and Pritchard, 1965) and neuroendocrine stimulation of fish under stress of metal exposure which in turn causes disturbances in carbohydrate metabolism (Mazeand et al., 1977). The duration taken in the present study of 4, 7, 15 and 30 days of Cd exposure to determine the glycogen content of *C. punctatus* makes this work different from others.

Conclusion

In the present investigation, the depletion of glycogen in muscle, liver, gill and kidney tissues of *C. punctatus* was directly proportional to the concentration of Cd and duration of exposure which clearly indicates its rapid utilization to meet the enhanced energy demands in fish exposed to Cd. Glycogen content can be used as a biomarker of Cd stress in fish. There is need to focus on harmful influences of heavy metals on biochemical activities of aquatic organisms and on the environment at large.

REFERENCES

1. Agrahari, S. and Gopal, K. (2007). Fate and toxicity of Cadmium and lead accumulation in different tissues (gills, liver, kidney, brain) of fresh water fish, *Channa punctatus*. *J. Ecophysiol. Occup. Hlth.* 7(3): 151-155
2. Akhilendra Naidu, K. (1982). Physiological studies on freshwater teleost *Sarotherodon mossambicus*

- about mercury toxicity, Ph.D. Thesis, S.V. University, Tirupathi India.
3. Ali, H., Khan, E. and Ilahi, I. (2019). Environmental Chemistry and Ecotoxicology of Hazardous Heavy Metals: Environmental Persistence, Toxicity, and Bioaccumulation. *Journal of Chemistry*. Volume 2019, Article ID 6730305, <https://doi.org/10.1155/2019/6730305>
 4. APHA, AWWA, WEF. (2012). Standard Methods for examination of water and wastewater. 22nd ed. Washington: American Public Health Association; 1360 pp. ISBN 978-087553-013-0. <http://www.standardmethods.org/>
 5. Baharoma, Z.S. and Ishak, M.Y. (2015). Determination of heavy metal accumulation in fish species in Galas River, Kelantan and Beranang mining pool, Selangor. *Procedia Environmental Sciences* . 30: 320-325.
 6. Bawuro, A. A., Voegborlo, R. B. and Adimado, A. A. (2018). Bioaccumulation of Heavy Metals in Some Tissues of Fish in Lake Geriyo, Adamawa State, Nigeria. *Journal of Environmental and Public Health*. Volume 2018, Article ID 1854892, <https://doi.org/10.1155/2018/1854892>
 7. Bengery, K.V. and Patil, H.S. (1986). Respiration, liver glycogen and bioaccumulation in *Labeo rohita* exposed to Zinc. *Indian J. Com. Animal Physiol.*, 4: 79-84.
 8. Bhilave, M. P., Muley, D. V., and Deshpande, V. Y., (2008). Biochemical changes in the fish *Cirrhinus mrigala* after acute and chronic exposure of heavy metal. *Nature Environment and Pollution Technology*, 7(1): 65-71.
 9. Cicik, B. and Engin, K. (2005). The effects of cadmium on levels of glucose in serum and glycogen reserves in the liver and muscle tissues of *Cyprinus carpio* (L., 1758). *Turk J Vet. Anim. Sci.*, 29: 113-117.
 10. Das, S., Patro, S.K. and Sahu, B.K. (2001). Biochemical changes induced by mercury in the liver of penaeid prawns *Penaeus indicus* and *P.monodon* (Crustacea: Penaeidae) from Rushikulya estuary, east coast of India. *Indian J. Marine Sci.*, 30(4): 246-252.
 11. Goswami, P., Kaushik, U., Damor, S., Sharma, P., Sharma, N., (2016). Effect of Cadmium Chloride on Biochemical Profile and Enzyme Activity in *Tilapia mossibica*. *Int. J. Pharma. Res. Health Sci.* 4 (6): 1462-1465. DOI:10.21276/ijprhs.2016.06.05
 12. Gumgum, B. and Unlu E. and Tez Z. Gulsun Z. (1994). Heavy metal pollution in water, sediment and fish from Tigris River in Turkey. *Chemosphere*, 29: 111-116.
 13. Heath, A.G. and Pritchard, A.W. (1965). Effects of severe hypoxia on carbohydrate energy, stores and metabolism in two species of freshwater fish. *Physiol. Zool.* 38: 325-334. <https://www.jstor.org/stable/30152409>
 14. Jagadeesan, G. (1990). Studies on the effect of mercuric chloride on biomodel biochemical analysis and vertebral deformation on freshwater fish *Anabas testudineus* (Bloch) M.Phil Thesis, Annamalai University.
 15. Javed, M. and Usmani, N. (2013). Assessment of heavy metal (Cu, Ni, Fe, Co, Mn, Cr, Zn) pollution in effluent dominated rivulet water and their effect on glycogen metabolism and histology of *Mastacembelus armatus*. *Springer Plus* 2:390 <http://www.springerplus.com/content/2/1/390>.
 16. Jordao, C.P., Pereira, M.G., Bellato, C.R., Pereira, J.L. and Matos, A.T. (2002). Assesment of water systems for contaminants from domestic and industrial sewages. *Environ. Monit. Asses.*, 79(1): 75-100.
 17. Kamaraju, S. and Ramasamy, K. (2011). Effect of cadmium chloride on glycogen content in gill, liver and kidney of edible exotic fish *Hypophthalmichthys molitrix*. *Int. J. Curr. Res.*, 3(5): 53-57.
 18. Karuppasamy, R. (2000). Effect of phenyl mercuric acetate on carbohydrate content of *Channa punctata* Uttar Pradesh. *J. Zool.*, 20(3): 219-225.
 19. Kumari, B. and Ahsan, J. (2011). Study of muscle glycogen content in both sexes of an Indian teleost *Clarius batrachus* (Linn.) exposed to different concentrations of arsenic. *Fish Physiol. Biochem.*, 37: 161-167.
 20. Leibson, L. and Plisetskaya, E.M. (1968). Effect of insulin blood sugar level and glycogen content in organs of some cyclostomes and fish. *Gen. Comp. Endocrinol.*, 11(2): 381-392.
 21. Levesque, H.M., Moon, T.W., Campbell, P.G.C. and Hontela, A. (2002). Seasonal variation in carbohydrate and lipid metabolism of yellow perch (*Perca flavescens*) chronically exposed to metals in the field. *Aquat. Toxicol.*, 60: 257-267.
 22. Malik, D.S., Sastry, K.V. and Hamilton, D.P. (1998). Effects of zinc toxicity on biochemical composition of muscle and liver of Murrel (*Channa punctatus*). *Environ. Int.*, 24(4): 433-438.
 23. Mary Chandravathy, V. and Reddy, S.L.N. (1996). Lead nitrate exposure changes in carbohydrate of freshwater fish, *Anabas scandens*. *J. Environ. Biol.*, 17: 75-79.
 24. Mason, C.F. (1996). Biology of freshwater pollution, 3rd edn., Longman, U.K. 1-4.
 25. Mazeand, M.N., Mazeand, F. and Donaldson, E.M. (1977). Primary and secondary effects of stress in fish some new data with a general view. *Trans. Ame. Fish Soc.*, 106: 201-212.
 26. Nicholas, V.C., Longley, R.W. and Roe, J.H. (1956) Determination of glycogen in liver and muscle by use of Anthrone reagent. *J. Biol. Chem.*, 220(2): 583-593.
 27. Nimmo, D.R. and Willox, M.J., Lafrancois, T.D., Chapman, P.L., Brinkman, S.F. and Greene, J.C. (1998). Effects of metal mining and milling on boundary waters of Yellow Stone National Park, U.S.A. *Environ. Manage.* 22(6): 913-926.
 28. Opaluwa, O.D., Aremu, M.O., Ogbo, L.O., Magaji, J.I., Odiba, I.E. and Ekpo, E.R. (2012). Assessment of Heavy Metals in Water, Fish and Sediments from UKE Stream, Nasarawa State, Nigeria. *Current World Environment*, 7(2): 213-220.
 29. Plessl, C., Otachi, E. O., Körner, W., Avenant-Oldewage, A. and Jirsa, F. (2017). Fish as bioindicators for trace element pollution from two contrasting lakes in the Eastern Rift Valley, Kenya: spatial and temporal aspects. *Environ. Sci. Pollut. Res.*, 24: 19767-19776. DOI 10.1007/s11356-017-9518-z
 30. Prabhakar, C., Saleshrani, K., Tharmaraj, K., and Vellaiyan, M., (2012). Studies on the Effect of Cadmium Compound on the Biochemical Parameters of Fresh Water Fish in *Cirrhinus mrigala*. *International Journal of Pharmaceutical & Biological Archives*. 3 (1): 69-73.
 31. Qayyum, M.A. and Shaffi, S.A. (1977). Changes in tissue glycogen on freshwater catfish *Heteropneustes*

- fossilis* due to mercury intoxication. *Curr. Sci.*, 46 (18): 652-653.
32. Radhakrishaniah, K., Venkataramana, P., Suresh, A. and Sivaramakrishna, B. (1992). Effect of lethal and sublethal concentrations of copper on glycolysis in the liver of the freshwater teleost, *Labeo rohita* (Ham). *J. Environ. Biol.*, 13: 63-68.
 33. Rauf, A., Javed, M. and Ubaidullah, M. (2009) Heavy metal levels in three major carps (*Catla catla*, *Labeo rohita*, *Cirrhina mrigala*) from the river Ravi, Pakistan. *Pak Vet J.*, 29(1):24-26.
 34. Reddy, S.L.N., Venugopal, N.B.R.K. and Ramana Rao, J.V. (1989). In vivo effects of cadmium chloride on certain aspects of carbohydrate metabolism in the tissue of freshwater field crab *Barytelphusa guerini*. *Bull. Environ. Contam. Toxicol.*, 42(6): 847-853.
 35. Sastry, K.V. and Sunita, Km. (1983). Enzymological and biochemical changes produced by chronic chromium exposure in a teleost fish *Channa punctatus*. *Toxicol. Lett.*, 16: 9-15.
 36. Sastry, K.V. and Subhadra, Km. (1982). Effect of cadmium on some aspects of carbohydrate metabolism in a freshwater catfish *Heteropneustes fossilis*. *Toxicol. Lett.*, 14: 45-55.
 37. Sastry, K.V. and Sunita, Km. (1982). Effect of cadmium and chromium on the intestinal absorption of glucose in the snakehead fish *Channa punctatus*. *Toxicol. Lett.*, 10: 293-296.
 38. Sim, S.F., Ling, T.Y., Nyanti, L., Gerunsin, N., Ee-Wong, Y. and Kho, L.P. (2016). Assessment of Heavy Metals in Water, Sediment, and Fishes of a Large Tropical Hydroelectric Dam in Sarawak, Malaysia. *Journal of Chemistry*. Volume 2016, Article ID 8923183, <http://dx.doi.org/10.1155/2016/8923183>
 39. Sobha, K., Poornima, A., Harini, P. and Veeraiah, K. (2007). A study on biochemical changes in the freshwater fish, *Catla catla* (Hamilton) exposed to heavy metal toxicant cadmium chloride. *Katmandu University Journal of Science engineering and technology*, 1 (4): 1-11.
 40. Sujata, K., (2015). Impact of Cadmium on the Biochemical Contents in the Reproductive Organs of Freshwater Fish, *Channa punctatus* (Bloch). *International Journal of Science and Research (IJSR)*, 4 (10): 114-118. ISSN (Online): 2319-7064
 41. US DHHS-ATSDR (1993). Toxicological profile for cadmium (TP-92/06).
 42. US EPA (2016). Aquatic life ambient water quality criteria cadmium. EPA-820-R-16-002. Office of water. Office of Science and Technology. Health and Ecological Criteria Division. Washington, DC.
 43. Veeraiah, K., Venkatrao, G., Vivek, Ch., and Hymaranjani, G., (2013). Heavy metal, cadmium chloride induced biochemical changes in the Indian major carp *Cirrhinus mrigala* (Hamilton). *International Journal of Bioassays*, ISSN: 2278-778X. pg1028-1033.
 44. Viana, A.P. and Lucena Frédou, F. (2014). Ichthyofauna as bioindicator of environmental quality in an industrial district in the amazon estuary, Brazil. *Braz. J. Biol.*, 74(2): 315-324 .<http://dx.doi.org/10.1590/1519-6984.16012>
 45. Viarengo, A. (1985). Biochemical effects of trace metals. *Mar. Pollut. Bull.* 16(4): 153-158. [https://doi.org/10.1016/0025-326X\(85\)90006-2](https://doi.org/10.1016/0025-326X(85)90006-2)
 46. Viswarajan, S. and Muthukrishnan, S. (1988). Impact of tannery effluent on phosphatase activity of fishes. *Proc. Inds. Nat. Sci. Acad.* B55: 314-345.
 47. Waisberg, M., Joseph, P., Hale, B. and Beyersmann, D. (2003). Molecular and cellular mechanisms of cadmium carcinogenesis. *Toxicology*. 192: 95-117. [https://doi.org/10.1016/S0300-483X\(03\)00305-6](https://doi.org/10.1016/S0300-483X(03)00305-6)
 48. WHO, (1992). Cadmium. Environmental Aspects. Environmental Health Criteria Series N 135. Geneva.
 49. Xuan, R., Wang, L., Sun, M., Ren, G. and Jiang, M. (2011). Effects of cadmium on carbohydrate and protein metabolisms in the freshwater Crab *Sinopotamon yangtsekiense*. *Comp. Biochem. Physiol., Part C: Toxicology and Pharmacology*, 154(3): 268-274. <https://doi.org/10.1016/j.cbpc.2011.06.005>