



Journal of Applied and Natural Science
12(1): 1 - 8 (2020)
ISSN : 0974-9411 (Print), 2231-5209 (Online)
journals.ansfoundation.org

Effect of intra row spacing, dates of haulm cutting and fertilizer dose on disease free quality seed tuber production of potato (*Solanum tuberosum* L.) under new Alluvial Zone of West Bengal

Manoj Mandal

AICRP on Potato, Directorate of Research, Bidhan Chandra Krishi Viswavidyalaya, Kalyani-741235 Nadia (West Bengal), India

Sanjib Kumar Das*

AICRP on Potato, Directorate of Research, Bidhan Chandra Krishi Viswavidyalaya, Kalyani-741235 Nadia (West Bengal), India

*Corresponding author. E-mail: sanjibag@gmail.com

Article Info

<https://doi.org/10.31018/jans.v12i1.2204>

Received: December 15, 2019

Revised: January 14, 2020

Accepted: January 22, 2020

Abstract

Potato (*Solanum tuberosum* L.) is the most popular tuber crop in West Bengal, India. Seed tuber is the single most important factor in potato cultivation and if the seed is not of good quality, then optimum production could not be achieved. Unavailability of good quality seed, high price and untimely supply of seed at the village level are the main limiting factors in potato production. Field experiments were conducted during *rabi* season of 2017-18 and 2018-19 at Bidhan Chandra Krishi Viswavidyalaya, Kalyani, West Bengal to determine the effects of intra row spacing, dates of haulm cutting and fertilizer dose on disease free quality seed grade tuber production of potato. The experiment was laid out in a split split plot design with three replications having twelve treatment combinations. The results revealed that with the decrease in intra row spacing from 20cm to 15cm seed grade size (< 75g) tuber yield and numbers and total tuber numbers were significantly increased but marketable grade (> 75 g) tuber yield and numbers were significantly reduced. Haulm cutting at 65 DAP increased the seed grade size tuber yield and numbers over 75 DAP. With the decrease in fertility levels from 100% RDF of NPK to 50% RDF of NPK the seed grade tuber production and number were significantly increased at 5% level of significance. Disease incidence and intensity of Phoma leaf spot decreased and early blight increased with decreasing doses of fertilizers. No viral disease was observed. Dehaulming at 65 DAP was found safer so far as infestation and chances of viral disease transmission by the sucking pests were concerned. From the present study it may be concluded that, for quality seed tuber production of potato and to get highest net return under New Alluvial Zone of West Bengal, use of 50cm X 15cm spacing along with haulm cutting at 65 DAP, when planting is done on first week of November and grown with 50% RDF of NPK was found best.

Keywords: Dates of haulm cutting, Fertility levels, Potato, Seed tuber Production, Spacing

How to Cite

Mandal, M. and Das, S.K. (2020). Effect of intra row spacing, dates of haulm cutting and fertilizer dose on disease free quality seed tuber production of potato (*Solanum tuberosum* L.) under new Alluvial Zone of West Bengal. *Journal of Applied and Natural Science*, 12(1): 1 - 8
<https://doi.org/10.31018/jans.v12i1.2204>

INTRODUCTION

West Bengal is the second largest potato growing state in India with a production of 10.0 million tonnes from an area of 409.7 thousand hectares. (Directorate of Agriculture, WB, 2018). The state accounts for one-third of the country's total potato production (Anonymous, 2019). Potato is the most popular crop in West Bengal next to the cereals. The crop is mostly grown during winter season (November-March). Seed tuber is the single most important factor in potato cultivation, which accounts for nearly 40-50% of the total investment for raising the crop, and if the seed is not of good quality, then optimum production could not be achieved (Chakraborty *et al.*, 2013). Unavailability of good quality seed, high price and untimely sup-

ply of seed at the village level are the main limiting factors for increase of potato production. With the expansion of potato cultivation in the Indo-Gangetic plains (IGP), it became evident that seed production in the Indian hills cannot cope with the increasing demand of good quality seed. Moreover, quarantine of seeds produced in Darjeeling area due to wart (Chakraborty *et al.*, 2013) and Nilgiri hills due to cyst nematode infestations seriously restricted opportunities of seed production in the hills (Priyank *et al.*, 2019). The seed produced from the western hills of Himachal Pradesh, Jammu and Kashmir and Uttarakhand also suffered from problems of true dormancy, tedious transportation and rottage due to late blight infection in tubers. Therefore, an alternate seed production technology was urgently required to sustain the

growing potato acreage in the plains. In West Bengal potato seed tuber can be produced if early planting is adopted. Moreover, the critical level of aphid population (20 aphids per 100 compound leaves) generally appears on 2nd week of January onwards. As a result 8-9 weeks of low aphid pressure period is available, and the problem of viral disease infection is much lower during this period. Moreover as potato is a vegetatively propagated crop and requires huge amount of fertilizers thus making the crop susceptible to so many diseases like late blight, early blight, phoma leaf spot etc and to combat these diseases appropriate management strategies are needed to produce disease free quality seed tuber of potato. Presently, the farmers of this state have no other option but to depend upon the homegrown seed or buy it from the cold stores, open market and seeds from other states at higher prices. In most of the cases, poor quality of seed material causes lower yield. The only solution left for the farmers of this state is to produce their own seed by following the 'Seed Plot Technique' (Wurr, 1978). Approximately 30-40% of the seed potato is procured from outside the state like Punjab, U.P etc. the quality of which is not always good enough resulting in rapid degeneration of tubers due to viral disease (Mondal *et al.*, 2015). Not only that, cost of such seed tubers is also very high resulting in huge monetary loss. Therefore, to save the farmers from losses and to make the potato cultivation profitable an effort has been made to produce potato seed in the state itself. Keeping the above facts in view, this experiment was initiated with the objectives to study the effect of intra row spacing, dates of haulm cutting and fertilizer dose on disease free quality seed grade tuber production of potato through 'Seed plot technique' under New Alluvial Zone of West Bengal.

MATERIALS AND METHODS

Field experiments were conducted for two years at C-unit research farm (Kalyani) of Bidhan Chandra Krishi Viswavidyalaya, Nadia, West Bengal, India situated at 22°58' N latitude and 88°3'E longitude with an altitude of 9.75m above mean sea (MSL) during *rabi* 2017-18 and 2018-19. The soil of the experimental field was sandy loam in texture and slightly alkaline in reaction (pH 7.2) having an organic carbon content of 0.57%, 183.26 Kg available N ha⁻¹, 16.8 kg available P₂O₅ ha⁻¹, 132 kg available K₂O ha⁻¹. The experiment was laid out in a split split plot design with three replications having twelve treatment combinations *viz.* two levels of spacing, S1- 50cm X 20cm and S2- 50cm X 15cm were applied in main plots. Two levels of dates of haulm cutting, H1- 65 days after planting (DAP) and H2- 75 days after planting were applied in sub plots and three levels of fertility levels, F1- 100% RDF (Recommended dose of

fertilizer) of NPK, F2- 75% RDF of NPK, F3- 50% RDF of NPK were applied in sub sub plots with a plot size of 5 m X 3 m. Breeder seeds of potato variety Kufri Himalini was planted on 5th November maintaining proper seed plot techniques. Tubers weighing 30–40 g each were planted in the furrows with a depth of planting of 3–4 cm and finally covered with soil. Nitrogen (N), phosphorus (P) and potassium (K) were applied through urea, single super phosphate and muriate of potash respectively. The recommended dose of fertilizer (RDF) was 200, 150, 150 kg N, P₂O₅, K₂O ha⁻¹. Half of nitrogen, full dose of phosphorus and potassium were applied as basal as per treatment details. Rest half N was top dressed at 30 days after planting (DAP) followed by earthing up. Pre-emergence application of Sencor (Metribuzin) @ 0.75 kg a.i. ha⁻¹ was done at 3 DAP followed by 1 hand-weeding at 20 DAP to promote early crop growth. As a prophylactic measure, spraying (twice) with Dithane M-45 (Mancozeb) @ 0.2% at 40 and 60 DAP was done against late blight. Imidacloprid 17.8SL @0.03% was also sprayed (thrice) at 30, 40 and 60 DAP for controlling aphids and other sucking insects. Continuous monitoring and roughing was done. Dehaulming was done as per treatments. For H1 haulm cutting was done on 09.01.18 and 09.01.19, and for H2 haulm cutting was done on 19.01.18 and 19.01.19 and harvesting was done 10 days after haulm cutting, and the crop lines were opened with the help of plough. Potato tubers were dug out from each plot manually. Data on grade wise tuber number and yield and total tuber number and yield were recorded at harvest from each net plot area. Percent disease incidence [(No. of infected plants / Total no. of plants) X100] and intensity [using 1-9 scale for late blight, Malcolimson, 1976 and 1-5 scale for phoma leaf spot and early blight, Kumar *et al.* (2017)] and aphid population build up were studied. Analysis of variance of the data in the experimental design and comparison of means at p≤0.05 were carried out, using MSTAT-C software.

RESULTS AND DISCUSSION

Effect on plant emergence, plant height and no. of shoots per plant: Experimental results revealed that plant emergence of potato was not significantly influenced by intra row spacing, dates of haulm cutting and fertility levels and their interactions (Table 1). Plant height of potato variety Kufri Himalini at 60 DAP crop growth and no. of shoots per plant of potato were not significantly influenced by different effects of spacing and dates of haulm cutting but it was significantly affected by fertility levels. The highest plant height (73.18 cm) was recorded at 100% RDF of NPK. Decrease in fertility levels from 100% RDF of NPK to 50% RDF of NPK significantly reduced the plant

Table 1. Effects of spacing, dates of haulm cutting and fertilizer doses on emergence, plant height and no. of shoots per plant of potato (Pooled data of two years).

Treatment	Potato		
	Emergence (%)	Plant height (cm)	No. of shoots plant ⁻¹
Levels of spacing			
S1	99.66	71.61	3.14
S2	99.77	70.68	3.07
S.E _m (±)	0.15	0.24	0.03
CD(P=0.05)	NS	NS	NS
Levels of dates of haulm cutting			
H1	99.63	71.36	3.10
H2	99.79	70.94	3.12
S.E _m (±)	0.04	0.18	0.04
CD(P=0.05)	NS	NS	NS
Levels of fertility			
F1	99.78	73.18	3.38
F2	99.75	70.94	3.08
F3	99.61	69.32	2.87
S.E _m (±)	0.12	0.32	0.04
CD(P=0.05)	NS	0.95	0.12

Table 2. Effects of spacing, dates of haulm cutting and fertilizer doses on grade wise and total tuber yield of potato (Pooled data of two years).

Treatment	Grade-wise yield of tubers (t ha ⁻¹)					Yield on dry weight basis (t ha ⁻¹)	
	0-25g	25-50g	50-75g	>75g	Total	Tuber	Haulm
Levels of spacing							
S1	3.00	6.34	6.90	13.23	29.47	5.76	3.45
S2	3.78	7.23	8.69	12.96	32.66	5.81	3.49
S.E _m (±)	0.14	0.18	0.08	0.20	0.11	0.08	0.05
CD(P=0.05)	NS	NS	0.48	NS	0.65	NS	NS
Levels of dates of haulm cutting							
H1	3.70	6.83	7.81	12.02	30.36	5.70	3.42
H2	3.08	6.73	7.78	14.17	31.77	5.87	3.52
S.E _m (±)	0.09	0.20	0.13	0.15	0.20	0.06	0.03
CD(P=0.05)	0.35	NS	NS	0.58	0.78	NS	NS
Levels of fertility							
F1	2.88	6.55	6.91	18.95	35.30	6.41	3.85
F2	3.22	6.68	8.11	12.75	30.77	5.68	3.41
F3	4.07	7.12	8.37	7.58	27.14	5.26	3.16
S.E _m (±)	0.14	0.17	0.31	0.33	0.55	0.11	0.07
CD(P=0.05)	0.40	NS	0.92	0.99	1.66	0.34	0.21

height and no. of shoots per plant of potato. Decrease in plant height and no. of shoots per plant of potato with decreasing fertility levels was confirmed by Das *et al.* (2016). In case of effect of fertility levels the highest no of shoots per plant (3.38 no.) was recorded at 100% RDF of NPK. However, the interaction effect of treatments on plant height and no. of shoots per plant of potato were found statistically insignificant at 5% level of significance.

Effect on grade wise and total tuber yield:

Experimental results revealed that, seed grade (<75g) tuber yield and total tuber yield of potato were significantly influenced by intra row spacing, dates of haulm cutting and fertilizer doses. Results showed that 50-75g grade and total tuber production of potato was significantly influenced by spacing (Table. 2). With the decrease in intra row spacing seed grade tuber yield and total tuber

yield were significantly increased but > 75 g grade tuber yield was significantly reduced. Lower spacing also increased the 0-25g grade and 25-50g grade tuber yield, which was desirable to produce more amount of seed grade size tuber of potato. The result corroborated with the findings of Dua *et al.* (2008) and Das *et al.* (2016) for potato crops at Shimla and Kalyani regions of India respectively. Haulm cutting at 65 DAP increased the seed grade size (<75g) tuber yield over haulm cutting at 75 DAP. However, haulm cutting at 75DAP significantly increased the marketable tuber (> 75 g) yield and total tuber production at 5% level of significance. This result corroborated with the findings of Lal and Sahota (1983) and Das *et al.* (2016). With the decrease in fertility levels from 100% RDF of NPK to 50% RDF of NPK the seed grade (< 75 g) tuber yield was significantly increased but marketable grade tuber yield (> 75 g), total tuber yield

Table 3. Effects of spacing, dates of haulm cutting and fertilizer doses on grade wise and total tuber number of Potato (Pooled data of two years).

Treatment	Grade-wise number of tubers (nos. ha ⁻¹)				
	0-25g	25-50g	50-75g	>75g	Total
Levels of spacing					
S1	180093	161574	105093	96759	543518
S2	231019	200463	141204	85185	657870
S.E _m (±)	5792	4428	982	1823	9729
CD(P=0.05)	35731	27321	6059	11245	60024
Levels of dates of haulm cutting					
H1	220833	189352	126389	86111	622685
H2	190278	172685	119907	95833	578704
S.E _m (±)	7768	7371	1669	1464	11871
CD(P=0.05)	30327	NS	NS	5716	NS
Levels of fertility					
F1	171528	161806	102778	109722	545833
F2	206250	182639	127778	93056	609722
F3	238889	198611	138889	70139	646528
S.E _m (±)	6371	5720	5475	1985	11305
CD(P=0.05)	19103	17149	16417	5950	33896

Table 4. Observation on disease incidence in the experimental field of potato (Mean data of two years).

Treatment	Phoma (%)		Early blight (%)		Late Blight (%)		Viral Disease
	Incidence	Intensity	Incidence	Intensity	Incidence	Intensity	
(S1H1F1)	8.35	2.00	3.50	2.25	0.00	0.00	0.00
(S1H1F2)	8.00	1.50	4.00	2.50	0.00	0.00	0.00
(S1H1F3)	7.25	1.50	4.50	2.50	0.00	0.00	0.00
(S1H2F1)	10.30	2.50	5.00	2.50	0.00	0.00	0.00
(S1H2F2)	7.45	2.00	6.00	3.00	0.00	0.00	0.00
(S1H2F3)	6.65	1.50	7.50	3.70	0.00	0.00	0.00
(S2H1F1)	11.50	2.80	5.50	3.00	0.00	0.00	0.00
(S2F1F2)	10.45	2.00	5.75	3.50	0.00	0.00	0.00
(S2H1F3)	10.00	1.50	6.25	3.50	0.00	0.00	0.00
(S2H2F1)	15.00	3.20	6.50	3.00	0.00	0.00	0.00
(S2H2F2)	10.00	3.00	10.00	4.00	0.00	0.00	0.00
(S2H2F3)	9.00	2.35	10.00	4.50	0.00	0.00	0.00

Table 5. Observation on aphid infestation in the experimental field of potato (Mean data of two years).

Treatment	Population of aphid per 100 compound leaves					
	03.01.18	03.01.19	10.01.18	10.01.19	17.01.18	17.01.19
(S1H1F1)	0.00	0.00	0.00	0.00	-	-
(S1H1F2)	0.00	0.00	0.00	0.00	-	-
(S1H1F3)	0.00	0.00	0.00	0.00	-	-
(S1H2F1)	0.00	0.00	0.00	0.00	0.00	2.25
(S1H2F2)	0.00	0.00	0.00	0.00	0.00	2.00
(S1H2F3)	0.00	0.00	0.00	0.00	0.00	1.75
(S2H1F1)	0.00	0.00	0.00	0.00	-	-
(S2F1F2)	0.00	0.00	0.00	0.00	-	-
(S2H1F3)	0.00	0.00	0.00	0.00	-	-
(S2H2F1)	0.00	0.00	0.00	0.00	0.00	2.40
(S2H2F2)	0.00	0.00	0.00	0.00	0.00	2.15
(S2H2F3)	0.00	0.00	0.00	0.00	0.00	1.80

and dry weight yield of tubers were significantly decreased. This was in conformity with the results of Dua *et al.* (2008), Chakraborty *et al.* (2013) and Das *et al.* (2016) for potato crops at Shimla, Hooghly and Kalyani regions of India respectively. The highest total tuber yield (35.30 t ha⁻¹) was recorded with 100% RDF of NPK. However, the interactions were found mostly non significant.

Effect on grade wise tuber numbers and total

tuber numbers: Experimental results revealed that, grade wise tuber numbers and total tuber numbers of potato were significantly influenced by spacing (Table. 3). With the decrease in intra row spacing from 20cm to 15cm seed grade size (< 75g) tuber numbers and total tuber numbers were significantly increased but marketable grade (> 75 g) tuber number was significantly reduced, which is desirable to produce more numbers of seed

Table 6. Effect of spacing, dates of haulm cutting and fertilizer doses on post-harvest soil nutrient status.

Treatment	Post-harvest soil nutrient (kg ha ⁻¹) status		
	Nitrogen	Phosphorus	Potassium
Levels of spacing			
S1	178.89	18.14	130.20
S2	178.24	18.09	129.49
S.E _m (±)	0.72	0.09	0.40
CD(P=0.05)	NS	NS	NS
Levels of dates of haulm cutting			
H1	178.92	18.21	130.20
H2	178.20	18.02	129.50
S.E _m (±)	0.65	0.07	0.32
CD(P=0.05)	NS	NS	NS
Levels of fertility			
F1	176.58	17.82	128.63
F2	179.38	18.26	130.31
F3	179.73	18.28	130.60
S.E _m (±)	0.87	0.13	0.62
CD(P=0.05)	2.60	0.40	1.86

Table 7. Effect of spacing, dates of haulm cutting and fertility level on total nutrient uptake by potato.

Treatment	Nutrient uptake (kg ha ⁻¹) by Potato		
	Nitrogen	Phosphorus	Potassium
Levels of spacing			
S1	133.32	36.78	185.73
S2	134.10	36.99	186.82
S.E _m (±)	0.13	0.04	0.19
CD(P=0.05)	0.72	NS	1.01
Levels of dates of haulm cutting			
H1	131.91	36.39	183.77
H2	135.51	37.38	188.77
S.E _m (±)	0.58	0.20	0.86
CD(P=0.05)	2.40	0.86	3.20
Levels of fertility			
F1	148.21	40.89	206.47
F2	131.22	36.20	182.80
F3	121.71	33.58	169.56
S.E _m (±)	3.10	0.80	4.20
CD(P=0.05)	9.30	2.40	12.60

grade size tuber of potato. The result corroborated with the findings of Dua *et al.* (2008) and Das *et al.* (2016). Haulm cutting at 65 DAP increased the seed grade size (<75g) tuber numbers and total tuber numbers of potato and significantly reduced the marketable grade (> 75 g) tuber numbers. Similar findings were also reported by Mahmud *et al.* (2009) and Garg *et al.* (1999) for potato crops at Bangladesh and Shimla hills of India respectively. However, the effect of dates of haulm cutting on total tuber number was found statistically insignificant. Fertility levels had a significant effect on grade wise and total tuber numbers of potato. With the decrease in fertility levels from 100% RDF of NPK to 50% RDF of NPK the seed grade size (< 75 g) tuber numbers and total tuber numbers were significantly increased but marketable grade (>75g) tuber numbers were significantly decreased. Similar result was also reported by Dua *et al.* (2008) and Das *et al.* (2016) for potato crops at Shimla and Kalyani regions of India respectively. This result is desirable for potato seed

tuber production as large size tubers (>80g) are discarded under potato seed tuber certification process in the state. Benefits of small sized tubers to be treated as seed was also reported by Wurr *et al.* (2001). However, the interaction effects between intra row spacing, dates of haulm cutting and fertilizer dose were found non significant at 5% level of significance.

Disease incidence: In this experiment late blight was not observed at all in both the years of study (Table 4) because in both the years the crop was dehaulmed before appearance of late blight. As far as leaf spot disease is concerned both phoma and early blight was observed. Both spacing and fertilizer dose had a marked effect on disease incidence and severity. Highest phoma leaf spot incidence (15.00%) and intensity (3.20%) was observed when spacing was 50cm X 20cm and 100% RDF of NPK. With same spacing the disease incidence and intensity decreased with decreasing dose of fertilizers i.e. 75% RDF of NPK and 50% RDF of NPK respectively. When the

Table 8. Economics and net returns of different treatments.

Treat- ment	Seed tuber yield (t ha ⁻¹)	Marketable grade (>75g) tuber yield (t ha ⁻¹)	Total tuber Yield (t ha ⁻¹)	Cost of cultivation (Rs. ha ⁻¹)			Pro- duce	Marketa- ble tuber Sale price (Rs.t ⁻¹)	Seed ber price (Rs.t ⁻¹)	Net turns* (Rs. ha ⁻¹)	B:C ratio (Seed+Ma- rketable)	B:C ratio (Only Seed)
				Seed	Fertilizer	Common cost						
	16.19	17.72	33.92	72000	14614	50000	624400	6000	32000	487786	3.57	2.79
S ₁ H ₁ F ₂	16.3	11.50	27.80	72000	10961	50000	590600	6000	32000	457639	3.44	2.92
S ₁ H ₁ F ₃	17.28	7.26	24.54	72000	7307	50000	596520	6000	32000	467213	3.61	3.28
S ₁ H ₂ F ₁	14.55	20.54	35.09	72000	14614	50000	588840	6000	32000	452226	3.31	2.41
S ₁ H ₂ F ₂	16.1	13.92	30.02	72000	10961	50000	598720	6000	32000	465759	3.50	2.87
S ₁ H ₂ F ₃	17.04	8.42	25.46	72000	7307	50000	595800	6000	32000	466493	3.61	3.22
S ₂ H ₁ F ₁	17.65	17.73	35.38	93600	14614	50000	671180	6000	32000	512966	3.24	2.57
S ₂ H ₁ F ₂	20.23	11.50	31.74	93600	10961	50000	716360	6000	32000	561799	3.63	3.19
S ₂ H ₁ F ₃	22.42	6.38	28.79	93600	7307	50000	755720	6000	32000	604813	4.01	3.75
S ₂ H ₂ F ₁	16.99	19.79	36.79	93600	14614	50000	662420	6000	32000	504206	3.19	2.44
S ₂ H ₂ F ₂	19.42	14.09	33.50	93600	10961	50000	705980	6000	32000	551419	3.57	3.02
S ₂ H ₂ F ₃	21.48	8.28	29.76	93600	7307	50000	737040	6000	32000	586133	3.88	3.55

* Cost of produce - Cost of inputs (cultivation)

spacing was increased to 50cm X 20cm from the spacing 50cm X 15cm the disease incidence and intensity started decreasing. Minimum disease incidence (6.65%) and intensity (1.50%) was observed when the spacing was 50cm X 20cm with 50% RDF of NPK. But in case of early blight, the disease incidence and intensity was increased with decreasing fertilizer dose. Barclay *et al.* (1973) also reported that both high nitrogen and low phosphorus treatments significantly reduced the incidence of early blight and the combination of high nitrogen and low phosphorus consistently gave the lowest incidence of the disease. Highest early blight incidence (10.00%) and intensity (4.50%) was observed when spacing was 50cm X 15cm and fertilizer dose was 50% RDF of NPK. Minimum early blight incidence (3.50%) and intensity (2.25%) was recorded at a spacing of 50cm X 20cm and fertilizer dose 100% RDF of NPK. Decrease in early blight incidence with the increase in fertility levels was also confirmed by Mitra *et al.* (2014) for potato crop at new alluvial zone of West Bengal. No viral disease was observed during both the years. This may be due to intensive insecticide application since thirty days after planting of the crop.

Aphid incidence: The data recorded on aphid population clearly showed (Table 5) that no aphid infestation was found during the entire crop growth period in first year. Whatever little population of aphids observed during the second year of experiment on 17th January was much below the critical level of aphid population (20 aphids per 100 compound leaves) and was easily controlled by the prophylactic measures taken and continuous roughing operation. Thus, there was no chance of viral disease transmission through aphids. In West Bengal dehaulming at 65 DAP was found to be safer so far as infestation and chances of viral disease transmission by the sucking pests are concerned as up to 10th January no aphid infestation was noticed in both the years of study. Similar findings were also reported by Das *et al.* (2016) for potato crop under new alluvial zone of West Bengal.

Effect on post-harvest soil nutrient status: The result showed that the nitrogen, phosphorus and potassium availability in post-harvest soil of potato were not significantly influenced by spacing and dates of haulm cutting but these were significantly influenced by fertility levels. It was clearly evident (Table 6) that, In case of effect of level of spacing the highest soil nitrogen status (178.89 kg ha⁻¹), soil phosphorus status (18.14 kg ha⁻¹) and soil potassium status (130.20 kg ha⁻¹) in post harvest soil was recorded with 20 cm intra row spacing which might be due to the fact that higher intra row spacing increased the soil nutrient status due to lesser plant population and uptake. Similar observation was also reported by Dua *et al.* (2008)

for potato crop at Shimla. In case of dates of haulm cutting effect the highest soil nitrogen status ($178.92 \text{ kg ha}^{-1}$), soil phosphorus status (18.21 kg ha^{-1}) and soil potassium status ($130.20 \text{ kg ha}^{-1}$) in post harvest soil was recorded with haulm cutting at 65 days after planting which might be due to the fact that decreased growth duration increased the soil nutrient status of potato due to lesser uptake by the crop. In case of fertility levels highest soil nitrogen status ($179.73 \text{ kg ha}^{-1}$), soil phosphorus status (18.28 kg ha^{-1}) and soil potassium status ($130.60 \text{ kg ha}^{-1}$) in post harvest soil was recorded with 50% RDF of NPK which might be due to lower uptake by the crop. However, the interaction effects were found non significant at 5% level of significance.

Nutrient uptake by potato: The result showed that intra row spacing, dates of haulm cutting and fertilizer doses significantly influenced nitrogen, phosphorus and potassium uptake by potato crop (Table 7). With the decrease in intra row spacing from 20 cm to 15 cm the uptake of nutrients increased due to increased yield. This finding was corroborated with the findings of Shukla *et al.* (1976). In case of dates of haulm cutting, the highest nitrogen uptake ($135.51 \text{ kg ha}^{-1}$), phosphorus uptake (37.38 kg ha^{-1}) and potassium uptake ($188.77 \text{ kg ha}^{-1}$) of potato cultivar Kufri Himalini was recorded with haulm cutting at 75 DAP which might be due to the fact that increased growth duration increased the total tuber yield and nutrient uptake by the crop. In case of fertility levels the highest nitrogen uptake ($148.21 \text{ kg ha}^{-1}$), phosphorus uptake (40.89 kg ha^{-1}) and potassium uptake ($206.47 \text{ kg ha}^{-1}$) of potato cultivar Kufri Himalini was recorded with 100% RDF of NPK which might be due to higher tuber yield. The nutrient uptake decreased with the decrease in fertility levels. This result corroborated with the findings of Dua *et al.* (2008) and Shukla *et al.* (1976) for potato crops at Shimla.

Economics: It was observed that the net return of potato cultivation varied from Rs. 452226 ha^{-1} to Rs. 604813 ha^{-1} (Table 8). The highest net return (604813 ha^{-1}) and B: C ratio (4.01) was recorded with 15 cm intra row spacing, 65 DAP of haulm cutting and 50% of RDF of NPK followed by 15 cm intra row spacing, 75 DAP of haulm cutting and 50% of RDF of NPK. Higher economics in potato seed tuber production with decrease in intra row spacing and fertility level was confirmed by Das *et al.* (2016). The lowest net return (Rs. 452226 ha^{-1}) was recorded with 20 cm intra row spacing, and 75DAP of haulm cutting and 100% RDF of NPK.

Conclusion

Thus, from the present study it can be concluded that, for quality seed grade potato tuber production use of 50cm X 15cm spacing along with

dehauling at 65 days after planting, when planting is done on first week of November and grown with 50% RDF of NPK i.e. 100:75:75 kg N:P₂O₅:K₂O was found best under New Alluvial Zone West Bengal to get higher numbers of quality seed grade sized potato tubers.

ACKNOWLEDGEMENTS

The authors are thankful to the All India Co-ordinated Research Project on Potato, Shimla, H.P. for valuable guidance, encouragement and financial support during the period of this research programme.

REFERENCES

1. Anonymous (2019). Monthly Report Potato (November), Horticulture Statistics Division, Department of Agriculture, Co-operation and Farmers Welfare, Ministry of Agriculture and Farmers Welfare, Govt. of India, New Delhi.
2. Barclay, G.M., Murphy, H.J., Manzer, F.E., Hutchinson, F.E. (1973). Effects of differential rates of nitrogen and phosphorus on early blight in potatoes. *American potato journal*. 50(2): 42-48.
3. Chakraborty, A, Konar, A., Banerjee, H. and Jatav, M.K (2013) In: Manual for 'Quality potato seed production in Hooghly district (New Alluvial Zone) of west Bengal. AICRP on Potato, Directorate of Research, BCKV, West Bengal. Pp 1-7.
4. Das, S.K., Sarkar, A., and Chakraborty, A. (2016). Effect of spacing, dates of haulm cutting and fertility levels on quality seed grade tuber production of potato (*Solanum tuberosum* L.). *Indian Agriculturist*. 60(3 &4): 225-236.
5. Directorate of Agriculture, W.B. (2018). *Economic Review*. Evaluation wing, Directorate of Agriculture, West Bengal, pp. 65.
6. Dua, V.K., Jaiswal, R.C., Khurana, S.C., Kumar, D., Nandekar, D.N., Raghav, M., Rawal, S., Sasani, G.V., Trishan, S.P., Trivedi, S.K., Jatav, M.K., Lal, S.S., Thakur, K.C., Pandey, S.K. and Naik, P.S. (2008). Manipulation of agronomic practices for tuber-size distribution in potato (*Solanum tuberosum*) seed crop. *Indian Journal of Agronomy*. 53(3): 217-223.
7. Garg V.K., Deshraj and Sarjeet, S. (1999). Influence of dates of planting and haulms cutting on the yield of seed sized tubers in Shimla Hills. *Journal of Indian Potato Association*. 26: 7-13.
8. Kumar, V., Singh, G and Tyagi, A. (2017). Evaluation of different fungicides against alternaria leaf blight of tomato. *Int. J. Curr. Microbiol. App. Sci*. 6(5): 2343-2350.
9. Lal, S.S. and Sahota, T.S. (1983). Effect of time of planting and lifting on tuber yield of potato in Shillong hills. *Journal of Indian Potato Association*. 10:129-133.
10. Mahmud, A.A. Akhter, S., Hossain, M.J., Bhuiyan, M.K.R. and Hoque, M.A. (2009). Effect of dehauling on yield of seed potatoes. *Bangladesh J. Agril. Res*. 34(3): 443-448.
11. Malcolmson, J.F. (1976). Assessment of field resistance to late blight (*Phytophthora infestans*) in potatoes. *Transactions of the British Mycological Society*. 67: 321-325.

12. Mitra, K., Naskar, B., Rana, J.P. and Das, S. (2014). Effect of Different Fertilizers Combination on Early Blight of Potato under Diverse Fertility Gradient of Soil. *International Journal of Bioresource Science*. 1 (1): 73-82.
13. Mondal, S.S., Patra, B.C. and Banerjee, H. (2015). *Advances in Potato Cultivation Technology*. Kalyani Publishers New Delhi. Pp 55-66.
14. Priyank, H.M., Divya, K.L., Venkatasalam, E.P., Aarti, B., Sudha, R and Berliner, J. (2019). Potato cyst nematode: A hidden enemy of potato cultivation in hills. *Bhartiya Krishi Anusandhan Patrika*. (34): 50-53. DOI: 10.18805/BKAP151
15. Wurr D.C.F. (1978). Seed tuber production and management. In: the potato Crop (P.M. Haris Ed.), Chapman & Hall, London, pp. 329-354.
16. Wurr, D. C. E., Fellows, J. R., Akehurst, J. M., Hambidge, A. J., & Lynn., J. R. (2001). The effect of cultural and environmental factors on potato seed tuber morphology and subsequent sprout and stem development. *Journal of Agricultural Science, Cambridge*. 136: 55-63.
17. Shukla, D.N., Samarjit, Singh and Singh, S. (1976). Effect of plant densities and fertility levels on the dry matter production and nutrient uptake in potato. *Indian Journal of Agronomy*. 21(4): 408-411.