

Weierstraß-Institut
für Angewandte Analysis und Stochastik
Leibniz-Institut im Forschungsverbund Berlin e. V.

Preprint

ISSN 2198-5855

**Error estimates for elliptic equations
with not exactly periodic coefficients**

Sina Reichelt

submitted: May 10, 2016

Weierstrass Institute
Mohrenstraße 39
10117 Berlin
Germany
E-Mail: sina.reichelt@wias-berlin.de

No. 2260
Berlin 2016



2010 *Mathematics Subject Classification.* 35B10, 35B27, 35B40, 35D30, 35J70.

Key words and phrases. Homogenization, error estimates, periodic unfolding, gradient folding operator.

Acknowledgments. This research was supported by *Deutsche Forschungsgesellschaft* within *Collaborative Research Center 910: Control of self-organizing nonlinear systems: Theoretical methods and concepts of application* via the project *A5: Pattern formation in systems with multiple scales*.

Edited by
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)
Leibniz-Institut im Forschungsverbund Berlin e. V.
Mohrenstraße 39
10117 Berlin
Germany

Fax: +49 30 20372-303
E-Mail: preprint@wias-berlin.de
World Wide Web: <http://www.wias-berlin.de/>

Abstract

This note is devoted to the derivation of quantitative estimates for linear elliptic equations with coefficients that are not exactly ε -periodic and the ellipticity constant may degenerate with order $O(\varepsilon^{2\gamma})$. Here $\varepsilon > 0$ denotes the ratio between the microscopic and the macroscopic length scale. It is shown that for $\gamma = 0$ and $\gamma = 1$ the error between the original solution and the effective solution is of order $O(\varepsilon^{1/2})$. Therefore suitable test functions are constructed via the periodic unfolding method and a gradient folding operator making only minimal additional assumptions on the given data and the effective solution with respect to the macroscopic scale.

1 Introduction

Periodic homogenization is a very powerful tool to derive effective equations for e.g. biological cells [NeJ07, GPS14], binary mixtures [Eck05], modeling of concrete [PeB08, FA*11], and other processes in porous media [HJM94, PtR10, Muv11, Muv13]. All these articles deal with systems of coupled reaction-diffusion equations which contain linear, elliptic equations of the form

$$-\operatorname{div}\left(\varepsilon^{2\gamma}\mathbb{A}\left(x,\frac{x}{\varepsilon}\right)\nabla u_\varepsilon\right)+\mathbb{B}\left(x,\frac{x}{\varepsilon}\right)u_\varepsilon=F_\varepsilon(x)\quad\text{in }\Omega. \quad (1.1)$$

In this text we study equation (1.1) for $\gamma = 0$ and $\gamma = 1$ on a bounded domain $\Omega \subset \mathbb{R}^d$ supplemented with homogeneous boundary conditions. Within the framework of reaction-diffusion systems $\gamma = 1$ models the slow diffusion of order $O(\varepsilon)$ for some species, whereas for $\gamma = 0$ other species diffuse with order $O(1)$. The aim is to quantify the error between the original solution u_ε of (1.1) and the effective solution, which is for $\gamma = 0$ a one-scale function $u(x)$ and for $\gamma = 1$ a two-scale function $U(x, y)$ solving the equations (2.8) and (4.2). The weak and strong two-scale convergence of solutions u_ε to (1.1) is studied in [PeB08] and [Han11] for all $\gamma \in [0, \infty)$. Here we derive quantitative estimates in the two-scale space $\Omega \times Y$, where $Y = [0, 1]^d$, using the periodic unfolding operator $\mathcal{T}_\varepsilon : L^2(\Omega) \rightarrow L^2(\Omega \times Y)$, cf. [CDG02] or (2.1),

$$\gamma = 0 : \quad \|u_\varepsilon - u\|_{L^2(\Omega)} + \|\mathcal{T}_\varepsilon(\nabla u_\varepsilon) - [\nabla u + \nabla_y U]\|_{L^2(\Omega \times Y)} \leq \varepsilon^{1/2}C, \quad (1.2)$$

$$\gamma = 1 : \quad \|\mathcal{T}_\varepsilon u_\varepsilon - U\|_{L^2(\Omega; H^1(Y))} \leq \varepsilon^{1/2}C. \quad (1.3)$$

The constant C depends on the given data as well as on the norms of the effective solution u and U in the space $H^2(\Omega)$ and $H^1(\Omega; H^1(Y))$, respectively. In the case $\gamma = 0$ the corrector U is given by the well-known unit-cell problem (4.3). For the coefficients \mathbb{A} , \mathbb{B} , and \mathbb{F} we assume Y -periodicity with respect to the microscopic scale $y = x/\varepsilon$ and minimal extra regularity with respect to the macroscopic scale $x \in \Omega$, cf. assumption (2.7). Then the right-hand side of (1.1) is defined via the folding or averaging operator, cf. (2.2),

$$F_\varepsilon(x) = \widehat{F}_\varepsilon(x, x/\varepsilon) \quad \text{where} \quad \widehat{F}_\varepsilon(x, y) = \int_{\varepsilon([x/\varepsilon]+Y)} \mathbb{F}(z, y) \, dz, \quad (1.4)$$

for two-scale functions $\mathbb{F} \in H^1(\Omega; L^2(Y))$ which are in general not continuous - neither with respect to $x \in \Omega$ nor $y \in Y$. This becomes relevant for systems of parabolic equations which are coupled via Lipschitz-terms $F_\varepsilon(u_\varepsilon, v_\varepsilon)$, where u_ε and v_ε only belong to the space $H^1(\Omega)$. For such systems the rigorous derivation of effective two-scale equations was proved

in [MRT14] using the notion of strong two-scale convergence via periodic unfolding. In the proceeding article [Rei14] convergence rate of order $O(\varepsilon^{1/4})$ is shown. For elliptic equations with exactly periodic coefficients and $\gamma = 0$ we refer to [Gri04, Gri05, OnV07, Gri14] for unfolding based error estimates of order $O(\varepsilon^{1/2})$ and higher. For coupled reaction-diffusion equations with $\gamma = 0$ only the gradient estimate is of lower order $O(\varepsilon^{1/4})$ in [FMP12]. Estimates of order $O(\varepsilon^{1/2})$ based on asymptotic expansion are derived in [Eck05, Muv13] containing the cases $\gamma = 0$ and $\gamma = 1$. Therein continuity is assumed for the given data and effective solutions, whereas our approach requires no higher regularity with respect to the microscopic scale.

Our main result (Theorem 2.1) is novel for $\gamma = 1$ and for $\gamma = 0$ (Theorem 4.1) it is a slight generalization of the estimates obtained in [Gri04]. There arise two difficulties deriving the quantitative estimates (1.2)–(1.3) which we briefly outline for $\gamma = 1$. The periodicity defect of $\mathcal{T}_\varepsilon u_\varepsilon \in L^2(\Omega; H^1(Y))$ versus $U \in L^2(\Omega; H_{\text{per}}^1(Y))$ is treated as in [Gri05]. Secondly one has to identify a suitable “folding” or “approximating sequence” for the limit $U(x, y)$ since the naive composition $U(x, x/\varepsilon)$ is only well-defined in $H^1(\Omega)$ for functions $U \in C^1(\Omega; H_{\text{per}}^1(Y))$, see [LNW02] for admissible test functions in two-scale convergence. Also U_ε as in (1.4) is not suitable due to insufficient $H^1(\Omega)$ -regularity. Therefore we employ the gradient folding operator $\mathcal{G}_\varepsilon : H^1(\Omega; H_{\text{per}}^1(Y)) \rightarrow H^1(\Omega)$ as in [Han11, MRT14, Rei14] which is defined by solving the degenerating elliptic problem (3.3). In Theorem 3.4 we control the *folding mismatch* of U_ε and $\mathcal{G}_\varepsilon U$ with respect to the $H^1(\Omega)$ -norm as it is indicated in [Rei14].

This text is structured as follows. In Subsection 2.1 we introduce basic definitions and notations and in Subsection 2.2 our assumptions and Theorem 2.1 for $\gamma = 1$ are stated. The derivation of quantitative estimates is done in Section 3, whereby Subsection 3.1 and 3.2 contain estimates for the approximation errors and the folding mismatch, respectively. The proof of Theorem 2.1 is given in Subsection 3.3. Finally we show in Section 4 how our approach applies to the case $\gamma = 0$ (Subsection 4.1) as well as to parabolic equations (Subsection 4.3). In Subsection 4.2 we compare our estimates with [Gri04].

2 Notations and definitions

2.1 Periodic unfolding

Let $\Omega \subset \mathbb{R}^d$ be a bounded domain with Lipschitz boundary $\partial\Omega$. Following [CDG02], we denote with $Y := [0, 1]^d$ the unit cube and $\mathcal{Y} := \mathbb{R}^d / \mathbb{Z}^d$ is the torus which we obtain by identifying opposite faces of \bar{Y} . Using the mappings $[\cdot] : \mathbb{R}^d \rightarrow \mathbb{Z}^d$ and $\{\cdot\} : \mathbb{R}^d \rightarrow \mathcal{Y}$, any point $x \in \mathbb{R}^d$ admits the additive decomposition $x = [x] + \{x\}$. Here $[x] := ([x_1], \dots, [x_d])$ is the componentwise Gauss bracket and $\{x\} := x - [x]$ is the remainder. With this we define for $\varepsilon > 0$ the cells $\mathcal{C}_\varepsilon(x) := \varepsilon([x/\varepsilon] + Y)$ where the macroscopic part $\varepsilon[x/\varepsilon] \in \varepsilon\mathbb{Z}^d$ denotes the node associated to $\mathcal{C}_\varepsilon(x)$ and $y \in \mathcal{Y}$ respective $y \in Y$ is the microscopic part. Since the domain $\Omega \subset \mathbb{R}^d$ is bounded, not all cells $\mathcal{C}_\varepsilon(x)$ are fully contained in Ω . To handle cells intersecting the boundary $\partial\Omega$ we define the following sets

$$\Omega_\varepsilon^- := \text{int}(\{x \in \Omega \mid \mathcal{C}_\varepsilon(x) \subset \Omega\}) \quad \text{and} \quad \Omega_\varepsilon^+ := \text{int}(\{x \in \Omega \mid \overline{\mathcal{C}_\varepsilon(x)} \cap \bar{\Omega}\})$$

so that $\Omega_\varepsilon^- \subset \Omega \subset \Omega_\varepsilon^+$ as depicted in Figure 2.1. The Lipschitz property of Ω guarantees $\text{vol}(\Omega_\varepsilon^+ \setminus \Omega_\varepsilon^-) \leq \varepsilon 2\sqrt{d} \text{meas}(\partial\Omega)$ with finite surface measure $\text{meas}(\partial\Omega)$.




Figure 1: Covering of the domain Ω with microscopic cells.

The *periodic unfolding operator* $\mathcal{T}_\varepsilon : L^2(\Omega) \rightarrow L^2(\Omega \times \mathcal{Y})$ is defined as in [CDG02] via

$$(\mathcal{T}_\varepsilon u)(x, y) := u\left(\varepsilon\left[\frac{x}{\varepsilon}\right] + \varepsilon y\right) \quad (2.1)$$

if $x \in \Omega_\varepsilon^-$ and $(\mathcal{T}_\varepsilon u)(x, y) := 0$ otherwise. Accordingly we define the *folding operator* (also averaging operator) $\mathcal{F}_\varepsilon : L^2(\Omega \times \mathcal{Y}) \rightarrow L^2(\Omega)$ via

$$(\mathcal{F}_\varepsilon U)(x) := \int_{\mathcal{C}_\varepsilon(x)} U\left(z, \left\{\frac{x}{\varepsilon}\right\}\right) dz \quad (2.2)$$

if $(x, y) \in \Omega_\varepsilon^- \times \mathcal{Y}$ and $(\mathcal{F}_\varepsilon U)(x) := 0$ otherwise, where $\int_A u(z) dz := \text{vol}(A)^{-1} \int_A u(z) dz$ denotes the average. Both operators are linear and bounded, i.e.

$$\|\mathcal{T}_\varepsilon u\|_{L^2(\Omega \times \mathcal{Y})} \leq \|u\|_{L^2(\Omega)} \quad \text{and} \quad \|\mathcal{F}_\varepsilon U\|_{L^2(\Omega)} \leq \|U\|_{L^2(\Omega \times \mathcal{Y})} \quad (2.3)$$

by Jensen's inequality. We emphasize that the periodic unfolding operator \mathcal{T}_ε in (2.1) does *not* satisfy the integral identity $\int_{\Omega \times \mathcal{Y}} \mathcal{T}_\varepsilon u dx dy = \int_\Omega u dx$ and we refer to [MiT07] for a unfolding definition which does do so. However we only use the duality $\mathcal{F}_\varepsilon = \mathcal{T}_\varepsilon^*$, cf. [CDG08, Sect. 2.2], meaning for all $u \in L^2(\Omega)$ and $V \in L^2(\Omega \times \mathcal{Y})$

$$\int_{\Omega \times \mathcal{Y}} (\mathcal{T}_\varepsilon u) V dx dy = \int_\Omega u (\mathcal{F}_\varepsilon V).$$

Furthermore the unfolding operator satisfies $\mathcal{T}_\varepsilon u \in L^2(\Omega; H^1(Y))$ for all $u \in H^1(\Omega)$ and it is $\mathcal{T}_\varepsilon(\varepsilon \nabla u) = \nabla_y(\mathcal{T}_\varepsilon u)$, cf. [CDG08, Eq. (3.1)]. We point out that the spaces $L^2(\Omega \times \mathcal{Y})$ and $L^2(\Omega \times Y)$ can be identified, whereas $L^2(\Omega; H^1(\mathcal{Y}))$ is a closed subspace of $L^2(\Omega; H^1(Y))$ with $H^1(\mathcal{Y}) = H_{\text{per}}^1(Y) \subsetneq H^1(Y)$.

Using periodic unfolding via \mathcal{T}_ε , *weak (respective strong) two-scale convergence* of bounded sequences $(u_\varepsilon)_\varepsilon \subset L^2(\Omega)$, as introduced in [Ngu89], is equivalent to weak (respective strong) convergence of the unfolded sequence $(\mathcal{T}_\varepsilon u_\varepsilon)_\varepsilon$ in the two-scale space $L^2(\Omega \times \mathcal{Y})$. It was first proved in [All92, Prop. 1.14] that sequences $(u_\varepsilon)_\varepsilon \subset H^1(\Omega)$ satisfying $\sup_{\varepsilon > 0} \{\|u_\varepsilon\|_{L^2(\Omega)} + \|\varepsilon \nabla u_\varepsilon\|_{L^2(\Omega)}\} < \infty$ weakly two-scale converge (up to subsequence) to a function $U \in L^2(\Omega; H^1(\mathcal{Y}))$, i.e. $\mathcal{T}_\varepsilon u_\varepsilon \rightharpoonup U$ and $\mathcal{T}_\varepsilon(\varepsilon \nabla u_\varepsilon) \rightharpoonup \nabla_y U$ in $L^2(\Omega \times \mathcal{Y})$. We emphasize that the unfolded function $\mathcal{T}_\varepsilon u_\varepsilon$ is in general not Y -periodic, while its two-scale limit U indeed is, and we call this observation *periodicity defect* as in [Gri04].

2.2 Assumptions and statement of the main result for $\gamma = 1$

We study the following elliptic equation, which is degenerating as $\varepsilon \rightarrow 0$,

$$-\operatorname{div}(\varepsilon^2 A_\varepsilon \nabla u_\varepsilon) + B_\varepsilon u_\varepsilon = F_\varepsilon \quad \text{in } \Omega \quad (2.4)$$

supplemented with homogeneous Neumann boundary conditions, i.e. $(\varepsilon^2 A_\varepsilon \nabla u_\varepsilon) \cdot \nu = 0$ on $\partial\Omega$. Here ν denotes the unit outer normal vector of Ω . The coefficients depend on $x \in \Omega$ and $\varepsilon > 0$ via

$$A_\varepsilon(x) := \mathbb{A}\left(x, \frac{x}{\varepsilon}\right), \quad B_\varepsilon(x) := \mathbb{B}\left(x, \frac{x}{\varepsilon}\right), \quad \text{and} \quad F_\varepsilon(x) := (\mathcal{F}_\varepsilon \mathbb{F})(x). \quad (2.5)$$

Throughout this text we impose the following assumptions on the given data. Let $\mathbb{A}(x, y) \in \mathbb{R}_{\text{sym}}^{d \times d}$ be a *symmetric* matrix and let \mathbb{A} and \mathbb{B} be *positive definite*, i.e. there exist constants $0 < \alpha \leq \beta < \infty$ such that we have uniformly for all $(x, y) \in \Omega \times \mathcal{Y}$ and

$$\alpha |\xi|^2 \leq \mathbb{A}(x, y) \xi \cdot \xi \leq \beta |\xi|^2 \quad \text{and} \quad \alpha \leq \mathbb{B}(x, y) \leq \beta \quad \forall \xi \in \mathbb{R}^d. \quad (2.6)$$

For the derivation of quantitative estimates we need additional regularity with respect to the macroscopic scale $x \in \Omega$, namely let

$$\mathbb{A} \in W^{1, \infty}(\Omega; L^\infty(\mathcal{Y})), \quad \mathbb{B} \in W^{1, \infty}(\Omega; L^\infty(\mathcal{Y})), \quad \text{and} \quad \mathbb{F} \in H^1(\Omega; L^2(\mathcal{Y})). \quad (2.7)$$

Thanks to the continuity of \mathbb{A} and \mathbb{B} with respect to $x \in \Omega$, the definitions of A_ε and B_ε in (2.5) are indeed well-defined. Passing to the limit $\varepsilon \rightarrow 0$ in (2.4) gives the two-scale convergence of u_ε to a limit function $U \in L^2(\Omega; H^1(\mathcal{Y}))$ solving the effective equation (cf. [PeB08, Han11, MRT14])

$$-\operatorname{div}_y(\mathbb{A} \nabla_y U) + \mathbb{B}U = \mathbb{F} \quad \text{in } \Omega \times \mathcal{Y} \quad (2.8)$$

with periodic boundary conditions on the torus \mathcal{Y} and no boundary conditions on Ω . By the Lax–Milgram Theorem and the regularity (2.7) of the given data, the solution U of (2.8) even belongs to the better space $H^1(\Omega; H^1(\mathcal{Y}))$. In particular all solutions are uniformly bounded in the sense

$$\sup_{\varepsilon > 0} \{ \|u_\varepsilon\|_{L^2(\Omega)} + \|\varepsilon \nabla u_\varepsilon\|_{L^2(\Omega)} \} + \|U\|_{H^1(\Omega; H^1(\mathcal{Y}))} \leq \operatorname{Const.}(\mathbb{A}, \mathbb{B}, \mathbb{F}). \quad (2.9)$$

Our main result is in the case $\gamma = 1$ as follows.

Theorem 2.1. *Let the given data satisfy the assumptions (2.5)–(2.7) and let u_ε and U solve the elliptic equation (2.4) and (2.8), respectively. Then there exists a constant $C > 0$ depending on the norm in (2.9) such that*

$$\| \mathcal{T}_\varepsilon u_\varepsilon - U \|_{L^2(\Omega; H^1(\mathcal{Y}))} \leq \varepsilon^{1/2} C.$$

The proof (see Subsection 3.3) combines ideas of [Rei14, Thm. 3.2] and [Gri04, Prop. 4.3], and it is shown that $\mathcal{G}_\varepsilon U$ is an approximate solution of the original equation (2.4). Therefore we proceed in three steps:

1. The choice of an admissible test function for the effective equation (2.8) leads to the periodicity defect.
2. We reformulate by using the duality $\mathcal{F}_\varepsilon = \mathcal{T}_\varepsilon^*$ and control the approximation errors.
3. Choosing a suitable test function for the original equation is equivalent to finding the correct “recovery sequence” (in terms of Γ -convergence) which is $\mathcal{G}_\varepsilon U$ here.

3 Error estimates

3.1 Estimating the periodicity defect and approximation errors

In this subsection we provide preparatory estimates to control the approximation error for the folding operator and the periodicity defect. To handle the error for microscopic cells \mathcal{C}_ε which are not fully contained in the domain Ω , we exploit for functions $U \in H^1(\Omega; L^2(\mathcal{Y}))$ the boundary estimate

$$\|U\|_{L^2((\Omega \setminus \Omega_\varepsilon^-) \times \mathcal{Y})} \leq \varepsilon^{1/2} C \|U\|_{H^1(\Omega; L^2(\mathcal{Y}))}, \quad (3.1)$$

where the constant $C > 0$ only depends on the Lipschitz property of Ω . For one-scale functions, estimate (3.1) can be found in e.g. [Gri04, Eq. (4.6)] or [Gri05, Eq. (2.4)] and for a full proof we refer to [Rei15, Lem. 2.3.5].

Lemma 3.1. *For every $U \in H^1(\Omega; L^2(\mathcal{Y}))$ it holds*

$$\|U - \mathcal{T}_\varepsilon \mathcal{F}_\varepsilon U\|_{L^2(\Omega \times \mathcal{Y})} \leq (\varepsilon + \varepsilon^{1/2}) C \|U\|_{H^1(\Omega; L^2(\mathcal{Y}))},$$

where the constant $C > 0$ only depends on the domains Ω and Y .

Proof. Using the paving property $\Omega_\varepsilon^- = \cup_{\lambda_i} \mathcal{C}_\varepsilon(x)$, where $\lambda_i = [x/\varepsilon] \in \mathbb{Z}^d$ for $x \in \Omega_\varepsilon^-$, and applying the Poincaré–Wirtinger inequality yields

$$\begin{aligned} \|U - \mathcal{T}_\varepsilon \mathcal{F}_\varepsilon U\|_{L^2(\Omega_\varepsilon^- \times \mathcal{Y})}^2 &= \sum_{\lambda_i} \int_{\varepsilon(\lambda_i + Y)} \int_{\mathcal{Y}} \left| U(x, y) - \int_{\mathcal{C}_\varepsilon(x)} U(z, y) \, dz \right|^2 \, dx \, dy \\ &\leq \sum_{\lambda_i} (\text{diam}(\mathcal{C}_\varepsilon(x)))^2 \|\nabla_x U\|_{L^2(\varepsilon(\lambda_i + Y))}^2 \leq \varepsilon^2 C \|\nabla_x U\|_{L^2(\Omega \times \mathcal{Y})}^2. \end{aligned}$$

Here we used that the Poincaré–Wirtinger constant is bounded by the diameter of the convex set $\mathcal{C}_\varepsilon(x)$. Since $\|U - \mathcal{T}_\varepsilon \mathcal{F}_\varepsilon U\|_{L^2(\Omega \times \mathcal{Y})} = \|U - \mathcal{T}_\varepsilon \mathcal{F}_\varepsilon U\|_{L^2(\Omega_\varepsilon^- \times \mathcal{Y})} + \|U\|_{L^2((\Omega \setminus \Omega_\varepsilon^-) \times \mathcal{Y})}$ the boundary estimate (3.1) gives the desired result. \square

Introducing for $\varepsilon > 0$ and $\varphi \in H^1(\Omega)$ the norm

$$\|\varphi\|_\varepsilon := \|\varphi\|_{L^2(\Omega)} + \|\varepsilon \nabla \varphi\|_{L^2(\Omega)} \quad (3.2)$$

we can control the periodicity defect as follows.

Theorem 3.2 ([Gri05, Thm. 2.2]). *For every $\varphi \in H^1(\Omega)$, there exists a \mathcal{Y} -periodic function $\Phi_\varepsilon \in L^2(\Omega; H^1(\mathcal{Y}))$ such that*

$$\|\Phi_\varepsilon\|_{H^1(Y; L^2(\Omega))} \leq C \|\varphi\|_\varepsilon \quad \text{and} \quad \|\mathcal{T}_\varepsilon \varphi - \Phi_\varepsilon\|_{H^1(Y; H^1(\Omega)^*)} \leq (\varepsilon + \varepsilon^{1/2}) C \|\varphi\|_\varepsilon,$$

where the constant $C > 0$ only depends in the domains Ω and Y .

3.2 Gradient folding operator and folding mismatch

By the Sobolev extension Theorem, there exists a linear and continuous operator, cf. e.g. [Neč67, Thm. 3.9], $\mathcal{E} : \mathbf{H}^1(\Omega; \mathbf{L}^2(\mathcal{Y})) \rightarrow \mathbf{H}^1(\mathbb{R}^d; \mathbf{L}^2(\mathcal{Y}))$ such that $(\mathcal{E}U)|_{\Omega \times \mathcal{Y}} = U$ and $\|\mathcal{E}U\|_{\mathbf{H}^1(\mathbb{R}^d; \mathbf{L}^2(\mathcal{Y}))} \leq C\|U\|_{\mathbf{H}^1(\Omega; \mathbf{L}^2(\mathcal{Y}))}$. Moreover, let \mathcal{E} be such that its restriction to one-scale functions $u \in \mathbf{H}^1(\Omega)$ satisfies $(\mathcal{E}u)|_{\Omega} = u$ and $\|\mathcal{E}u\|_{\mathbf{H}^1(\mathbb{R}^d)} \leq C\|u\|_{\mathbf{H}^1(\Omega)}$. Taking care of cells \mathcal{C}_ε intersecting the boundary of Ω , we define the extended folding operator $\mathcal{F}_\varepsilon^+ : \mathbf{H}^1(\Omega; \mathbf{L}^2(\mathcal{Y})) \rightarrow \mathbf{L}^2(\mathbb{R}^d)$ via

$$(\mathcal{F}_\varepsilon^+ U)(x, y) := \int_{\mathcal{C}_\varepsilon(x)} (\mathcal{E}U)(z, \{\frac{x}{\varepsilon}\}) \, dz.$$

The *gradient folding operator* $\mathcal{G}_\varepsilon : \mathbf{H}^1(\Omega; \mathbf{H}^1(\mathcal{Y})) \rightarrow \mathbf{H}^1(\Omega_\varepsilon^+)$ is defined as in [MRT14] following [MiT07, Han11]: for $U \in \mathbf{H}^1(\Omega; \mathbf{H}^1(\mathcal{Y}))$ given, $\mathcal{G}_\varepsilon U := \widehat{u}_\varepsilon \in \mathbf{H}^1(\Omega_\varepsilon^+)$ is the unique solution of the elliptic problem

$$\int_{\Omega_\varepsilon^+} (\widehat{u}_\varepsilon - \mathcal{F}_\varepsilon^+ U)\varphi + (\varepsilon \nabla \widehat{u}_\varepsilon - \mathcal{F}_\varepsilon^+(\nabla_y U)) \cdot \varepsilon \nabla \varphi \, dx = 0 \quad \text{for all } \varphi \in \mathbf{H}^1(\Omega_\varepsilon^+). \quad (3.3)$$

The Lax–Milgram Theorem implies that $\mathcal{G}_\varepsilon U$ is well-defined and bounded in the sense $\|\mathcal{G}_\varepsilon U\|_\varepsilon \leq \|\mathcal{E}U\|_{\mathbf{H}^1(\mathbb{R}^d; \mathbf{H}^1(\mathcal{Y}))}$. Throughout the following calculations the operator \mathcal{E} is omitted in notation, however all generic constants C will also depend on its norm.

Having defined two different folding operators \mathcal{F}_ε and \mathcal{G}_ε , we want to control their difference which we call *folding mismatch*. Therefore we introduce the so-called scale-splitting operator $\mathcal{Q}_\varepsilon : \mathbf{H}^1(\Omega) \rightarrow \mathbf{W}^{1,\infty}(\mathbb{R}^d)$ following [CDG08, Def. 4.1]: for $x \in \mathcal{C}_\varepsilon(x)$ and every $\kappa = (\kappa_1, \dots, \kappa_d) \in \{0, 1\}^d$, we set

$$\bar{x}_l^{(\kappa_l)} := \begin{cases} \frac{x_l - \varepsilon \lfloor x/\varepsilon \rfloor_l}{\varepsilon} & \text{if } \kappa_l = 1 \\ 1 - \frac{x_l - \varepsilon \lfloor x/\varepsilon \rfloor_l}{\varepsilon} & \text{if } \kappa_l = 0 \end{cases}$$

and

$$(\mathcal{Q}_\varepsilon w)(x) := \sum_{\kappa \in \{0,1\}^d} (\mathcal{F}_\varepsilon^+ w)(\varepsilon \lfloor \frac{x}{\varepsilon} \rfloor + \varepsilon \kappa) \cdot \bar{x}_1^{(\kappa_1)} \dots \bar{x}_d^{(\kappa_d)}. \quad (3.4)$$

The function $\mathcal{Q}_\varepsilon w$ interpolates the values of $\mathcal{F}_\varepsilon^+ w$ at the nodes $\varepsilon \lfloor x/\varepsilon \rfloor$ via \mathcal{Q}_1 -Lagrange elements as customary in the finite elements methods. Indeed it is $\mathcal{F}_\varepsilon^+ w \in \mathbf{L}^\infty(\Omega_\varepsilon^+)$ such that for $z \in \mathbf{L}^2(\mathcal{Y})$ the products

$$x \mapsto (\mathcal{F}_\varepsilon^+ w)(x)z(\frac{x}{\varepsilon}) \quad \text{and} \quad x \mapsto (\mathcal{Q}_\varepsilon w)(x)z(\frac{x}{\varepsilon}) \quad (3.5)$$

belong to the space $\mathbf{L}^2(\Omega_\varepsilon^+)$, see e.g. [LNW02, Thm.4]. According to [CDG08, Prop. 4.5], there exists a constant $C > 0$ only depending on Ω and Y such that

$$\|\mathcal{Q}_\varepsilon w\|_{\mathbf{H}^1(\Omega)} \leq C\|w\|_{\mathbf{H}^1(\Omega)} \quad \text{for all } w \in \mathbf{H}^1(\Omega). \quad (3.6)$$

The following auxiliary estimate is proved in [Rei15, Lem. 3.6] or [Rei14, Lem. 2.3.9] based on ideas from [Gri04, Prop. 3.2].

Lemma 3.3. *For $w \in \mathbf{H}^1(\Omega)$ and $z \in \mathbf{L}^2(\mathcal{Y})$, it holds*

$$\|(\mathcal{F}_\varepsilon^+ w - \mathcal{Q}_\varepsilon w)z(\frac{\cdot}{\varepsilon})\|_{\mathbf{L}^2(\Omega_\varepsilon^+)} \leq \varepsilon C\|w\|_{\mathbf{H}^1(\Omega)}\|z\|_{\mathbf{L}^2(\mathcal{Y})},$$

where the constant $C > 0$ only depends on the dimension d .

Finally we have collected all ingredients to control the folding mismatch.

Theorem 3.4. *For every $U \in H^1(\Omega; H^1(\mathcal{Y}))$, the folding mismatch is*

$$\|\mathcal{G}_\varepsilon U - \mathcal{F}_\varepsilon U\|_{L^2(\Omega)} + \|\varepsilon \nabla(\mathcal{G}_\varepsilon U) - \mathcal{F}_\varepsilon(\nabla_y U)\|_{L^2(\Omega)} \leq \varepsilon^{1/2} C \|U\|_{H^1(\Omega; H^1(\mathcal{Y}))}, \quad (3.7)$$

$$\|\mathcal{G}_\varepsilon U - \mathcal{F}_\varepsilon^+ U\|_{L^2(\Omega_\varepsilon^+)} + \|\varepsilon \nabla(\mathcal{G}_\varepsilon U) - \mathcal{F}_\varepsilon^+(\nabla_y U)\|_{L^2(\Omega_\varepsilon^+)} \leq \varepsilon C \|U\|_{H^1(\Omega; H^1(\mathcal{Y}))}, \quad (3.8)$$

where the constant $C > 0$ only depends on the domains Ω and Y .

Proof. For brevity we set $L_\varepsilon^2 := L^2(\Omega_\varepsilon^+)$ throughout this proof.

Step 1: Treatment of the boundary. The triangle inequality, the identity $(\mathcal{F}_\varepsilon U)|_{\Omega_\varepsilon^-} = (\mathcal{F}_\varepsilon^+ U)|_{\Omega_\varepsilon^-}$, Jensen's inequality, and the boundary estimate (3.1) imply

$$\begin{aligned} \|\mathcal{G}_\varepsilon U - \mathcal{F}_\varepsilon U\|_{L^2(\Omega)} &\leq \|\mathcal{G}_\varepsilon U - \mathcal{F}_\varepsilon^+ U\|_{L^2(\Omega)} + \|\mathcal{F}_\varepsilon^+ U\|_{L^2(\Omega \setminus \Omega_\varepsilon^-)} \\ &\leq \|\mathcal{G}_\varepsilon U - \mathcal{F}_\varepsilon^+ U\|_{L_\varepsilon^2} + (\varepsilon + \varepsilon^{1/2}) C \|U\|_{H^1(\Omega; L^2(\mathcal{Y}))}. \end{aligned}$$

An analog estimate holds for the gradient term so that it remains to prove (3.8) and (3.7) follows immediately.

Step 2: Proof of estimate (3.8) for products $U(x, y) = w(x)z(y)$. Let U satisfy the decomposition $U(x, y) = w(x)z(y)$ with $w \in H^1(\Omega)$ and $z \in H^1(\mathcal{Y})$. We introduce the function $\vartheta_\varepsilon(x) := (\mathcal{Q}_\varepsilon w)(x)z(x/\varepsilon)$ which belongs to the space $H^1(\Omega_\varepsilon^+)$ by construction. The definition of $\mathcal{G}_\varepsilon U$ in (3.3) is equivalent to

$$\begin{aligned} \int_{\Omega_\varepsilon^+} (\mathcal{G}_\varepsilon U - \vartheta_\varepsilon) \varphi + \varepsilon \nabla(\mathcal{G}_\varepsilon U - \vartheta_\varepsilon) \cdot \varepsilon \nabla \varphi \, dx \\ = \int_{\Omega_\varepsilon^+} (\mathcal{F}_\varepsilon^+ U - \vartheta_\varepsilon) \varphi + (\mathcal{F}_\varepsilon^+(\nabla_y U) - \varepsilon \nabla \vartheta_\varepsilon) \cdot \varepsilon \nabla \varphi \, dx \quad \text{for all } \varphi \in H^1(\Omega_\varepsilon^+). \end{aligned}$$

Choosing $\varphi = \mathcal{G}_\varepsilon U - \vartheta_\varepsilon$ as well as applying Hölder's inequality, Lemma 3.3, and the boundedness of \mathcal{Q}_ε in (3.6) gives

$$\begin{aligned} \|\mathcal{G}_\varepsilon U - \vartheta_\varepsilon\|_\varepsilon &\leq \|\mathcal{F}_\varepsilon^+ U - \vartheta_\varepsilon\|_{L_\varepsilon^2} + \|\mathcal{F}_\varepsilon^+(\nabla_y U) - \varepsilon \nabla \vartheta_\varepsilon\|_{L_\varepsilon^2} \\ &= \|(\mathcal{F}_\varepsilon^+ w - \mathcal{Q}_\varepsilon w)z(\cdot/\varepsilon)\|_{L_\varepsilon^2} + \|(\mathcal{F}_\varepsilon^+ w - \mathcal{Q}_\varepsilon w) \nabla_y z(\cdot/\varepsilon)\|_{L_\varepsilon^2} + \|\varepsilon \nabla_x (\mathcal{Q}_\varepsilon w)z(\cdot/\varepsilon)\|_{L_\varepsilon^2} \\ &\leq \varepsilon C \|w\|_{H^1(\Omega)} \|z\|_{H^1(\mathcal{Y})}. \end{aligned}$$

Finally the triangle inequality yields

$$\begin{aligned} \|\mathcal{G}_\varepsilon U - \mathcal{F}_\varepsilon U\|_{L_\varepsilon^2} + \|\varepsilon \nabla(\mathcal{G}_\varepsilon U) - \mathcal{F}_\varepsilon(\nabla_y U)\|_{L_\varepsilon^2} \\ \leq \|\mathcal{G}_\varepsilon U - \vartheta_\varepsilon\|_\varepsilon + \|\mathcal{F}_\varepsilon U - \vartheta_\varepsilon\|_{L_\varepsilon^2} + \|\mathcal{F}_\varepsilon(\nabla_y U) - \varepsilon \nabla \vartheta_\varepsilon\|_{L_\varepsilon^2} \\ \leq 2\varepsilon C \|w\|_{H^1(\Omega)} \|z\|_{H^1(\mathcal{Y})} \end{aligned} \quad (3.9)$$

and the desired estimate (3.8) follows for functions U of product form.

Step 3: Proof of estimate (3.8) for general functions $U(x, y)$. Let $\{\Phi_i\}_{i=1}^\infty$ be an orthonormal basis in $H^1(\mathcal{Y})$ which is also orthogonal in $L^2(\mathcal{Y})$. Then we can express $U \in H^1(\Omega_\varepsilon^+; H^1(\mathcal{Y}))$ (extended by \mathcal{E}) via the linear combination

$$U(x, y) = \sum_{i=1}^{\infty} u_i(x) \Phi_i(y) \quad \text{where} \quad u_i(x) := \int_{\mathcal{Y}} U(x, y) \Phi_i(y) \, dy. \quad (3.10)$$

By construction it is $u_i \in H^1(\Omega_\varepsilon^+)$ and we set $U_i(x, y) := u_i(x)\Phi_i(y)$.

The assumptions on Φ_i imply $\Phi_i(\cdot/\varepsilon) \perp \Phi_j(\cdot/\varepsilon)$ and $\nabla_y \Phi_i(\cdot/\varepsilon) \perp \nabla_y \Phi_j(\cdot/\varepsilon)$ in L_ε^2 for all $i \neq j$, i.e. $\{\Phi_i(\cdot/\varepsilon)\}_i$ is an orthogonal system in $H^1(\Omega_\varepsilon^+)$ for each $\varepsilon > 0$. Indeed using the paving property $\Omega_\varepsilon^+ = \cup_{\lambda_i} \mathcal{C}_\varepsilon(x)$, where $\lambda_i = [x/\varepsilon] \in \mathbb{Z}^d$ for $x \in \Omega_\varepsilon^+$, a substitution of variables yields

$$\begin{aligned} \int_{\Omega_\varepsilon^+} \Phi_i\left(\frac{x}{\varepsilon}\right)\Phi_j\left(\frac{x}{\varepsilon}\right) dx &= \sum_{\lambda_i} \int_{\mathcal{C}_\varepsilon(x)} \Phi_i\left(\frac{x}{\varepsilon}\right)\Phi_j\left(\frac{x}{\varepsilon}\right) dx \\ &= \sum_{\lambda_i} \frac{1}{\varepsilon^d} \int_{[\frac{x}{\varepsilon}] + Y} \Phi_i(y)\Phi_j(y) dy = 0 \end{aligned} \quad (3.11)$$

and analogously for $\nabla_y \Phi_i$. Applying the folding operator $\mathcal{F}_\varepsilon^+$ to U_i gives

$$(\mathcal{F}_\varepsilon^+ U_i)(x) = (\mathcal{F}_\varepsilon^+ u_i)(x)\Phi_i\left(\frac{x}{\varepsilon}\right) \quad \text{and} \quad (\mathcal{F}_\varepsilon^+ [\nabla_y U_i])(x) = (\mathcal{F}_\varepsilon^+ u_i)(x)\nabla_y \Phi_i\left(\frac{x}{\varepsilon}\right).$$

Since $(\mathcal{F}_\varepsilon^+ u_i)(x) = \int_{\mathcal{C}_\varepsilon(x)} u_i(z) dz$ is constant on each cell $\mathcal{C}_\varepsilon(x)$, we have as well $\mathcal{F}_\varepsilon^+ U_i \perp \mathcal{F}_\varepsilon^+ U_j$ and $\mathcal{F}_\varepsilon^+ (\nabla_y U_i) \perp \mathcal{F}_\varepsilon^+ (\nabla_y U_j)$ in L_ε^2 for all $i \neq j$. Therefore it suffices to consider the basis functions Φ_i . By the definition of \mathcal{G}_ε we have for $v_i^\varepsilon := \mathcal{G}_\varepsilon \Phi_i$

$$\int_{\Omega_\varepsilon^+} (v_i^\varepsilon - \Phi_i(\frac{\cdot}{\varepsilon})) \varphi + (\varepsilon \nabla v_i^\varepsilon - \nabla_y \Phi_i(\frac{\cdot}{\varepsilon})) \cdot \varepsilon \nabla \varphi dx = 0 \quad \text{for all } \varphi \in H^1(\Omega_\varepsilon^+). \quad (3.12)$$

Inserting the test function $\varphi(x) = \Phi_j(x/\varepsilon)$ in (3.12) yields with (3.11) for all $i \neq j$

$$\int_{\Omega_\varepsilon^+} v_i^\varepsilon \Phi_j\left(\frac{\cdot}{\varepsilon}\right) + \varepsilon \nabla v_i^\varepsilon \cdot \nabla_y \Phi_j\left(\frac{\cdot}{\varepsilon}\right) dx = 0.$$

Furthermore choosing $\varphi(x) = v_j^\varepsilon(x)$ in (3.12) yields with the latter equality

$$\int_{\Omega_\varepsilon^+} v_i^\varepsilon v_j^\varepsilon + \varepsilon \nabla v_i^\varepsilon \cdot \varepsilon \nabla v_j^\varepsilon dx = 0.$$

Hence it is $v_i^\varepsilon \perp v_j^\varepsilon$ in $H^1(\Omega_\varepsilon^+)$ for all $i \neq j$. We continue to estimate the folding mismatch of $\mathcal{F}_\varepsilon^+$ and \mathcal{G}_ε and set $u_i^\varepsilon := \mathcal{G}_\varepsilon U_i$. Replacing $\mathcal{F}_\varepsilon^+ \Phi_i$ and $\mathcal{F}_\varepsilon^+ (\nabla_y \Phi_i)$ with $\mathcal{F}_\varepsilon^+ U_i$ and $\mathcal{F}_\varepsilon^+ (\nabla_y U_i)$ in (3.12), respectively, and exploiting that $\mathcal{F}_\varepsilon^+ u_i$ is constant on each cell $\mathcal{C}_\varepsilon(x)$ implies $u_i^\varepsilon \perp u_j^\varepsilon$ in $H^1(\Omega_\varepsilon^+)$ for all $i \neq j$.

Finally we apply the result of Step 2 with $w = u_i$ and $z = \Phi_i$. Since $\mathcal{F}_\varepsilon^+$ and \mathcal{G}_ε are linear and continuous operators, we have in particular $u^\varepsilon = \sum_{i=1}^\infty u_i^\varepsilon$. Therefore we square estimate (3.9) so that the mixed product terms vanish for $i \neq j$, namely

$$\begin{aligned} &\|u^\varepsilon - \mathcal{F}_\varepsilon^+ U\|_{L_\varepsilon^2}^2 + \|\varepsilon \nabla u^\varepsilon - \mathcal{F}_\varepsilon^+ (\nabla_y U)\|_{L_\varepsilon^2}^2 \\ &= \|\sum_{i=1}^\infty (u_i^\varepsilon - \mathcal{F}_\varepsilon^+ U_i)\|_{L_\varepsilon^2}^2 + \|\sum_{i=1}^\infty (\varepsilon \nabla u_i^\varepsilon - \mathcal{F}_\varepsilon^+ (\nabla_y U_i))\|_{L_\varepsilon^2}^2 \\ &= \sum_{i=1}^\infty \|u_i^\varepsilon - \mathcal{F}_\varepsilon^+ U_i\|_{L_\varepsilon^2}^2 + \sum_{i=1}^\infty \|\varepsilon \nabla u_i^\varepsilon - \mathcal{F}_\varepsilon^+ (\nabla_y U_i)\|_{L_\varepsilon^2}^2 \\ &\leq \sum_{i=1}^\infty \varepsilon^2 C \|u_i\|_{H^1(\Omega)}^2 \|z\|_{H^1(\mathcal{Y})}^2 = \varepsilon^2 C \|U\|_{H^1(\Omega; H^1(\mathcal{Y}))}^2. \end{aligned}$$

The last equality follows by Parseval's identity $\|U\|_{H^1(\Omega_\varepsilon^+; H^1(\mathcal{Y}))}^2 = \sum_{i=1}^\infty \|u_i\|_{H^1(\Omega_\varepsilon^+)}^2$. \square

3.3 Proof of Theorem 2.1

Proof of Theorem 2.1. *Step 1: Periodicity defect.* The weak formulation of the effective equation (2.8) reads

$$\int_{\Omega \times \mathcal{Y}} \mathbb{A} \nabla_y U \cdot \nabla_y \Phi + \mathbb{B} U \Phi - \mathbb{F} \Phi \, dx \, dy = 0 \quad \text{for all } \Phi \in L^2(\Omega; H^1(\mathcal{Y})). \quad (3.13)$$

Thanks to Theorem 3.2 we can choose $\Phi_\varepsilon \in L^2(\Omega; H^1(\mathcal{Y}))$ to estimate the periodicity defect of $\mathcal{T}_\varepsilon \varphi \in L^2(\Omega; H^1(\mathcal{Y}))$. Indeed exploiting the higher x -regularity (2.7) of the given data and the limit $U \in H^1(\Omega; H^1(\mathcal{Y}))$ yields

$$\begin{aligned} & \left| \int_{\Omega \times \mathcal{Y}} \mathbb{A} \nabla_y U \cdot [\nabla_y (\mathcal{T}_\varepsilon \varphi) - \nabla_y \Phi_\varepsilon] + (\mathbb{B} U - \mathbb{F}) [\mathcal{T}_\varepsilon \varphi - \Phi_\varepsilon] \, dx \, dy \right| \\ & \leq \|\mathbb{A} \nabla_y U\|_{L^2(\mathcal{Y}; H^1(\Omega))} \|\nabla_y (\mathcal{T}_\varepsilon \varphi) - \nabla_y \Phi_\varepsilon\|_{L^2(\mathcal{Y}; H^1(\Omega)^*)} \\ & \quad + (\|\mathbb{B} U\| + \|\mathbb{F}\|)_{L^2(\mathcal{Y}; H^1(\Omega))} \|\mathcal{T}_\varepsilon \varphi - \Phi_\varepsilon\|_{L^2(\mathcal{Y}; H^1(\Omega)^*)} \\ & \leq \text{Const.}(\mathbb{A}, \mathbb{B}, \mathbb{F}, U) (\varepsilon + \varepsilon^{1/2}) C \|\varphi\|_\varepsilon. \end{aligned}$$

Note that the identification of the spaces $(L^2(\mathcal{Y}; H^1(\Omega)))^*$ and $L^2(\mathcal{Y}; H^1(\Omega)^*)$ as well as $L^2(\mathcal{Y}; H^1(\Omega))$ and $H^1(\Omega; L^2(\mathcal{Y}))$ holds due to the underlying tensor product structure of the two-scale space $L^2(\mathcal{Y}; H^1(\Omega))$. Overall testing (3.13) with Φ_ε gives

$$\left| \int_{\Omega \times \mathcal{Y}} \mathbb{A} \nabla_y U \cdot \nabla_y (\mathcal{T}_\varepsilon \varphi) + \mathbb{B} U \mathcal{T}_\varepsilon \varphi - \mathbb{F} \mathcal{T}_\varepsilon \varphi \, dx \, dy \right| \leq \varepsilon^{1/2} C \|\varphi\|_\varepsilon.$$

Step 2: Approximation errors. Using $\mathcal{F}_\varepsilon = \mathcal{T}_\varepsilon^*$ and $\nabla_y (\mathcal{T}_\varepsilon \varphi) = \mathcal{T}_\varepsilon (\varepsilon \nabla \varphi)$ gives

$$\left| \int_{\Omega} \mathcal{F}_\varepsilon (\mathbb{A} \nabla_y U) \cdot \varepsilon \nabla \varphi + \mathcal{F}_\varepsilon (\mathbb{B} U) \varphi - F_\varepsilon \varphi \, dx \right| \leq \varepsilon^{1/2} C \|\varphi\|_\varepsilon. \quad (3.14)$$

Thanks to the Lipschitz continuity of \mathbb{A} and \mathbb{B} with respect to $x \in \Omega$ and the boundedness (2.3) of \mathcal{F}_ε , we obtain

$$\begin{aligned} & \|\mathcal{F}_\varepsilon (\mathbb{A} \nabla_y U) - A_\varepsilon \mathcal{F}_\varepsilon (\nabla_y U)\|_{L^2(\Omega)}^2 \\ & = \int_{\Omega_\varepsilon^-} \left| \int_{\mathcal{C}_\varepsilon(x)} (\mathbb{A} (z, \frac{x}{\varepsilon}) - \mathbb{A} (x, \frac{x}{\varepsilon})) \nabla_y U (z, \frac{x}{\varepsilon}) \, dz \right|^2 \, dx \\ & \leq \varepsilon^2 \|\nabla_x \mathbb{A}\|_{L^\infty(\Omega \times \mathcal{Y})} \|\nabla_y U\|_{L^2(\Omega \times \mathcal{Y})} \end{aligned}$$

and analogously $\|\mathcal{F}_\varepsilon (\mathbb{B} U) - B_\varepsilon \mathcal{F}_\varepsilon (U)\|_{L^2(\Omega)}^2 \leq \varepsilon^2 \|\nabla_x \mathbb{B}\|_{L^\infty(\Omega \times \mathcal{Y})} \|U\|_{L^2(\Omega \times \mathcal{Y})}$. Inserting these two estimates into (3.14) yields

$$\left| \int_{\Omega} A_\varepsilon \mathcal{F}_\varepsilon (\nabla_y U) \cdot \varepsilon \nabla \varphi + B_\varepsilon \mathcal{F}_\varepsilon U \varphi - F_\varepsilon \varphi \, dx \right| \leq \varepsilon^{1/2} C \|\varphi\|_\varepsilon.$$

Step 3: Folding mismatch. Thanks to Theorem 3.4 we can replace the folded functions $\mathcal{F}_\varepsilon U$ and $\mathcal{F}_\varepsilon (\nabla_y U)$ with the gradient folding $\mathcal{G}_\varepsilon U$ and $\varepsilon \nabla (\mathcal{G}_\varepsilon U)$, respectively, such that

$$\left| \int_{\Omega} A_\varepsilon \varepsilon \nabla (\mathcal{G}_\varepsilon U) \cdot \varepsilon \nabla \varphi + B_\varepsilon (\mathcal{G}_\varepsilon U) \varphi - F_\varepsilon \varphi \, dx \right| \leq \varepsilon^{1/2} C \|\varphi\|_\varepsilon. \quad (3.15)$$

The choice $\varphi_\varepsilon = u_\varepsilon - \mathcal{G}_\varepsilon U \in H^1(\Omega)$ is indeed admissible in (3.15) and φ_ε is also a test function for the original equation (2.4), i.e.

$$\int_{\Omega} A_\varepsilon \varepsilon \nabla u_\varepsilon \cdot \varepsilon \nabla \varphi_\varepsilon + B_\varepsilon u_\varepsilon \varphi_\varepsilon - F_\varepsilon \varphi_\varepsilon \, dx = 0. \quad (3.16)$$

Subtracting (3.15) from (3.16) and exploiting the positive definiteness (2.6) gives

$$\alpha \|u_\varepsilon - \mathcal{G}_\varepsilon U\|_\varepsilon^2 \leq \varepsilon^{1/2} C \|u_\varepsilon - \mathcal{G}_\varepsilon U\|_\varepsilon. \quad (3.17)$$

Finally applying the triangle inequality, the boundedness (2.3) of \mathcal{T}_ε , estimate (3.7) for the folding mismatch, and the approximation error yields

$$\begin{aligned} & \| \mathcal{T}_\varepsilon u_\varepsilon - U \|_{L^2(\Omega; H^1(\mathcal{Y}))} \\ & \leq \| u_\varepsilon - \mathcal{G}_\varepsilon U \|_\varepsilon + \| \mathcal{G}_\varepsilon U - \mathcal{F}_\varepsilon U \|_{L^2(\Omega)} + \| \varepsilon \nabla (\mathcal{G}_\varepsilon U) - \mathcal{F}_\varepsilon (\nabla_y U) \|_{L^2(\Omega)} \\ & \quad + \| \mathcal{T}_\varepsilon \mathcal{F}_\varepsilon U - U \|_{L^2(\Omega \times \mathcal{Y})} + \| \mathcal{T}_\varepsilon \mathcal{F}_\varepsilon (\nabla_y U) - \nabla_y U \|_{L^2(\Omega \times \mathcal{Y})} \leq \varepsilon^{1/2} C \end{aligned}$$

which finishes the proof. \square

4 Further error estimates and discussion

4.1 The non-degenerating case for $\gamma = 0$

We consider the non-degenerating elliptic equation

$$- \operatorname{div}(A_\varepsilon \nabla u_\varepsilon) + B_\varepsilon u = F_\varepsilon \quad \text{in } \Omega \quad (4.1)$$

with homogeneous Neumann boundary conditions and given data as in (2.5)–(2.7). It is well-known that $u_\varepsilon \rightharpoonup u$ weakly in $H^1(\Omega)$ and u solves the effective equation

$$- \operatorname{div}(A_{\text{eff}} \nabla u) + B_{\text{eff}} u = F_{\text{eff}} \quad \text{in } \Omega \quad (4.2)$$

with the same boundary conditions. The effective matrix A_{eff} is given via the standard unit-cell problem (for arbitrary vectors $\xi \in \mathbb{R}^d$)

$$\xi \cdot A_{\text{eff}}(x) \xi = \min_{\Phi \in H_{\text{av}}^1(\mathcal{Y})} \int_{\mathcal{Y}} (\xi + \nabla_y \Phi) \cdot \mathbb{A}(x, y) (\xi + \nabla_y \Phi) \, dy, \quad (4.3)$$

where $H_{\text{av}}^1(\mathcal{Y}) = \{\Phi \in H^1(\mathcal{Y}) \mid \int_{\mathcal{Y}} \Phi \, dy = 0\}$. The effective data B_{eff} and F_{eff} are the usual averages

$$B_{\text{eff}}(x) = \int_{\mathcal{Y}} \mathbb{B}(x, y) \, dy \quad \text{and} \quad F_{\text{eff}}(x) = \int_{\mathcal{Y}} \mathbb{F}(x, y) \, dy. \quad (4.4)$$

The regularity of \mathbb{A} in (2.7) implies $A_{\text{eff}} \in W^{1,\infty}(\Omega; \mathbb{R}_{\text{sym}}^{d \times d})$. If the domain Ω is additionally convex or of class \mathcal{C}^2 , we obtain higher regularity of u , namely $u \in H^2(\Omega)$, see e.g. [Gri85, Lóp13]. This higher regularity of the limit solution u is needed to derive quantitative estimates and it is not nearly as trivial as in the degenerating case $\gamma = 1$. The corrector U is the unique minimizer in (4.3) corresponding to $\xi = \nabla u(x)$ and it satisfies $U \in H^1(\Omega; H^1(\mathcal{Y}))$ thanks to the higher regularity of \mathbb{A} and u with respect to $x \in \Omega$. Hence we can state our main result for $\gamma = 0$.

Theorem 4.1. *Let the assumptions (2.5)–(2.7) hold and let u_ε and u solves the elliptic equation (4.1) and (4.2), respectively. If u belongs to $H^2(\Omega)$, then there exists a constant $C > 0$ such that*

$$\|u_\varepsilon - u\|_{L^2(\Omega)} + \|\mathcal{T}_\varepsilon(\nabla u_\varepsilon) - [\nabla u + \nabla_y U]\|_{L^2(\Omega \times \mathcal{Y})} \leq \varepsilon^{1/2} C.$$

Proof. The proof is analog to the one of Theorem 2.1 for $\gamma = 1$ and one shows that $u + \varepsilon \mathcal{G}_\varepsilon U$ is an approximate solution of the original equation (4.1). To control the periodicity defect in Step 1 we apply [Gri05, Thm. 2.3], i.e. for every $\varphi \in H^1(\Omega)$, there exists a function $\Phi_\varepsilon \in L^2(\Omega; H^1(\mathcal{Y}))$ and a constant $C > 0$ such that $\|\Phi_\varepsilon\|_{H^1(\mathcal{Y}; L^2(\Omega))} \leq C\|\varphi\|_{H^1(\Omega)}$ and $\|\mathcal{T}_\varepsilon(\nabla \varphi) - [\nabla \varphi + \nabla_y \Phi_\varepsilon]\|_{L^2(\mathcal{Y}; H^1(\Omega)^*)} \leq (\varepsilon + \varepsilon^{1/2})C\|\varphi\|_{H^1(\Omega)}$. \square

Remark 4.2. *For different boundary conditions the error estimates are functioning as before, however one has to modify the approximating (resp. recovery) sequence so that the new boundary conditions are satisfied. In the case of homogeneous Dirichlet boundary conditions, a suitable candidate is $u + \varepsilon \mathcal{G}_\varepsilon(\rho_\varepsilon U)$, where ρ_ε denotes a cut-off function to guarantee the nullity at the boundary $\partial\Omega$.*

4.2 Comparison with [Gri04]

The corrector U admits the decomposition (see e.g. [LNW02, Sec. 4])

$$U(x, y) = \sum_{i=1}^d \frac{\partial u}{\partial x_i}(x) z_i(x, y), \quad (4.5)$$

where $z_i \in W^{1,\infty}(\Omega; H_{\text{av}}^1(\mathcal{Y}))$ is the unique minimizer corresponding to the standard basis vector e_i in \mathbb{R}^d . If A_ε is exactly periodic as in [Gri04], i.e. $A_\varepsilon(x) = \mathbb{A}(x/\varepsilon)$, then $z_i \in H_{\text{av}}^1(\mathcal{Y})$ is it as well. In this case we have the following error estimates for solutions u_ε and u of (4.1) and (4.2), respectively, with $\mathbb{B} \equiv 0$ and either homogeneous Neumann or Dirichlet boundary conditions.

Theorem 4.3 ([Gri04, Prop. 4.3]). *If $u \in H^2(\Omega)$, then there exists $C > 0$ such that*

$$\|u_\varepsilon - u\|_{L^2(\Omega)} + \|\nabla u_\varepsilon - [\nabla u + \sum_{i=1}^d \mathcal{Q}_\varepsilon\left(\frac{\partial u}{\partial x_i}\right) z_i(\cdot/\varepsilon)]\|_{L^2(\Omega)} \leq \varepsilon^{1/2} C.$$

In any case – exactly periodic or not – the question is how to “fold” $U(x, y)$? We remark that $x \mapsto U(x, x/\varepsilon) \in L^2(\Omega)$ is in general not satisfied for $U \in L^\infty(\Omega; L^2(\mathcal{Y}))$, whereas for products $x \mapsto w(x)z(x/\varepsilon) \in L^2(\Omega)$ holds true for $w \in L^\infty(\Omega)$ and $z \in L^2(\mathcal{Y})$ according to [LNW02, Thm.4]. Hence the construction $\sum_{i=1}^d \mathcal{Q}_\varepsilon\left(\frac{\partial u}{\partial x_i}\right)(x) z_i(x/\varepsilon)$ is admissible in $H^1(\Omega)$ for exactly periodic coefficients. However in the non-periodic situation, the choice $\mathcal{Q}_\varepsilon\left(\frac{\partial u}{\partial x_i}\right)(x) z_i(x, x/\varepsilon)$ is not admissible. This technicality is circumvented in our approach by using the gradient folding operator and choosing the recovery sequence $u + \varepsilon \mathcal{G}_\varepsilon U$ respective $u + \varepsilon \mathcal{G}_\varepsilon(\rho_\varepsilon U)$. Nevertheless for more regular data such as $\mathbb{A} \in C^1(\overline{\Omega}; L^\infty(\mathcal{Y}))$ and hence $z_i \in C^1(\overline{\Omega}; H_{\text{av}}^1(\mathcal{Y}))$, the naive folding $z_i(x, x/\varepsilon)$ is again well-defined in $H^1(\Omega)$ and the proof of [Gri04, Prop. 4.3] seems to be valid, too.

The approach of [Gri04] applies as well to systems of reaction-diffusion equations, cf. [FMP12]. And our approach is also generalizable to semilinear parabolic equations with globally Lipschitz continuous right-hand sides, see Subsection 4.3.

4.3 Error estimates for parabolic equations

Our approach can be generalized to semilinear parabolic equations. Following e.g. [Eck05, FMP12, Muv13, Rei15] one aims to apply Gronwall's Lemma and therefore estimates

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|u_\varepsilon - \mathcal{F}_\varepsilon U\|_{L^2(\Omega)}^2 &= \int_{\Omega} (\dot{u}_\varepsilon - \mathcal{F}_\varepsilon \dot{U})(u_\varepsilon - \mathcal{F}_\varepsilon U) \, dx \\ &= \int_{\Omega} (\dot{u}_\varepsilon - \mathcal{F}_\varepsilon \dot{U}) \varphi_\varepsilon \, dx + \int_{\Omega} (\dot{u}_\varepsilon - \mathcal{F}_\varepsilon \dot{U})(\mathcal{G}_\varepsilon U - \mathcal{F}_\varepsilon U) \, dx. \end{aligned}$$

Term 1 Term 2

Choosing the test function $\varphi_\varepsilon = u_\varepsilon - \mathcal{G}_\varepsilon U$ and inserting the reformulated equations, Term 1 can be treated as in the elliptic case. To control Term 2, in particular to obtain the convergence rate $\varepsilon^{1/2}$, we need the uniform boundedness $\sup_{\varepsilon>0} \|\dot{u}_\varepsilon\|_{L^2(\Omega)} < \infty$ and estimate (3.8) of order $O(\varepsilon)$ for the folding mismatch, i.e. $|\text{Term 2}| \leq \varepsilon C$. This can be achieved by assuming either Ω is polyhedral or by extending all solutions, given data, and operators from Ω to Ω_ε^+ such that the duality $\mathcal{F}_\varepsilon^+ = (\mathcal{T}_\varepsilon^+)^*$ is applicable.

References

- [All92] G. ALLAIRE. Homogenization and two-scale convergence. *SIAM J. Math. Anal.*, 23(1), 1482–1518, 1992.
- [CDG02] D. CIORANESCU, A. DAMLAMIAN, and G. GRISO. Periodic unfolding and homogenization. *C. R. Math. Acad. Sci. Paris*, 335(1), 99–104, 2002.
- [CDG08] D. CIORANESCU, A. DAMLAMIAN, and G. GRISO. The periodic unfolding method in homogenization. *SIAM J. Math. Anal.*, 40(4), 1585–1620, 2008.
- [Eck05] C. ECK. Homogenization of a phase field model for binary mixtures. *Multiscale Model. Simul.*, 3(1), 1–27 (electronic), 2004/05.
- [FA*11] T. FATIMA, N. ARAB, E. P. ZEMSKOV, and A. MUNTEAN. Homogenization of a reaction-diffusion system modeling sulfate corrosion of concrete in locally periodic perforated domains. *J. Engrg. Math.*, 69(2-3), 261–276, 2011.
- [FMP12] T. FATIMA, A. MUNTEAN, and M. PTASHNYK. Unfolding-based corrector estimates for a reaction-diffusion system predicting concrete corrosion. *Appl. Anal.*, 91(6), 1129–1154, 2012.
- [GPS14] I. GRAF, M. A. PETER, and J. SNEYD. Homogenization of a nonlinear multiscale model of calcium dynamics in biological cells. *J. Math. Anal. Appl.*, 419(1), 28–47, 2014.
- [Gri85] P. GRISVARD. *Elliptic Problems in Nonsmooth Domains*, volume 24 of *Monographs and Studies in Mathematics*. Pitman (Advanced Publishing Program), Boston, MA, 1985.
- [Gri04] G. GRISO. Error estimate and unfolding for periodic homogenization. *Asymptot. Anal.*, 40(3-4), 269–286, 2004.
- [Gri05] G. GRISO. Interior error estimate for periodic homogenization. *C. R. Math. Acad. Sci. Paris*, 340(3), 251–254, 2005.
- [Gri14] G. GRISO. Error estimates in periodic homogenization with a non-homogeneous Dirichlet condition. *Asymptot. Anal.*, 87(1-2), 91–121, 2014.
- [Han11] H. HANKE. Homogenization in gradient plasticity. *Math. Models Methods Appl. Sci.*, 21, 1651–1684, 2011.
- [HJM94] U. HORNUNG, W. JÄGER, and A. MIKELIĆ. Reactive transport through an array of cells with semi-permeable membranes. *RAIRO Modél. Math. Anal. Numér.*, 31, 1257–1285, 1994.
- [LNW02] D. LUKKASSEN, G. NGUETSENG, and P. WALL. Two-scale convergence. *Int. J. Pure Appl. Math.*, 2, 35–86, 2002.
- [Lóp13] J. LÓPEZ-GÓMEZ. *Linear second order elliptic operators*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2013.

- [MiT07] A. MIELKE and A. TIMOFTE. Two-scale homogenization for evolutionary variational inequalities via the energetic formulation. *SIAM J. Math. Anal.*, 39(2), 642–668, 2007.
- [MRT14] A. MIELKE, S. REICHELDT, and M. THOMAS. Two-scale homogenization of nonlinear reaction-diffusion systems with slow diffusion. *Netw. Heterog. Media*, 9(2), 353–382, 2014.
- [Muv11] A. MUNTEAN and T. L. VAN NOORDEN. Homogenisation of a locally periodic medium with areas of low and high diffusivity. *European J. Appl. Math.*, 22(5), 493–516, 2011.
- [Muv13] A. MUNTEAN and T. L. VAN NOORDEN. Corrector estimates for the homogenization of a locally periodic medium with areas of low and high diffusivity. *European J. Appl. Math.*, 24(5), 657–677, 2013.
- [Neč67] J. NEČAS. *Les méthodes directes en théorie des équations elliptiques*. Masson et Cie, Éditeurs, Paris; Academia, Éditeurs, Prague, 1967.
- [NeJ07] M. NEUSS-RADU and W. JÄGER. Effective transmission conditions for reaction-diffusion processes in domains separated by an interface. *SIAM J. Math. Anal.*, 39(3), 687–720, 2007.
- [Ngu89] G. NGUETSENG. A general convergence result for a functional related to the theory of homogenization. *SIAM J. Math. Anal.*, 20(3), 608–623, 1989.
- [OnV07] D. ONOFREI and B. VERNESCU. Error estimates for periodic homogenization with non-smooth coefficients. *Asymptot. Anal.*, 54(1-2), 103–123, 2007.
- [PeB08] M. A. PETER and M. BÖHM. Different choices of scaling in homogenization of diffusion and interfacial exchange in a porous medium. *Math. Meth. Appl. Sci.*, 24, 119–136, 2008.
- [PtR10] M. PTASHNYK and T. ROOSE. Derivation of a macroscopic model for transport of strongly sorbed solutes in the soil using homogenization theory. *SIAM J. Appl. Math.*, 70(7), 2097–2118, 2010.
- [Rei14] S. REICHELDT. Error estimates for nonlinear reaction-diffusion systems involving different diffusion length scales. *WIAS Preprint No. 2008*, 2014. Accepted in *J. Phys.: Conf. Ser.*
- [Rei15] S. REICHELDT. *Two-scale homogenization of systems of nonlinear parabolic equations*. PhD thesis, Humboldt-Universität zu Berlin, 2015. <http://edoc.hu-berlin.de/dissertationen/reichelt-sina-2015-11-27/PDF/reichelt.pdf>.