



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Foraging strategies and diet composition of Hadza children

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Crittenden, Alyssa N., Nancy L. Conklin-Brittain, Frank W. Marlowe, Margaret J. Schoeninger, Richard W. Wrangham. 2009. Foraging strategies and diet composition of Hadza children. <i>American Journal of Physical Anthropology</i> 138(S48): 112.
Published Version	doi:10.1002/ajpa.21030
Accessed	February 17, 2015 5:55:10 PM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:2797439
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP

(Article begins on next page)

Foraging strategies and diet composition of Hadza children

Alyssa N. Crittenden, Nancy L. Conklin-Brittain, Frank W. Marlowe, Margaret J. Schoeninger, Richard W. Wrangham

Abstract:

Among the Hadza hunter-gatherers of Tanzania, children are active foragers and collect various types of wild plant foods and hunt small sized prey animals. The collection effort of Hadza children is reported to have a positive effect on a mother's foraging yield (Bulurton-Jones et al. 1994, *Journal of Anthropological Research* 50(3): 217; Hawkes et al. 1995, *Current Anthropology* 36(4): 688), yet few quantitative data are available on the caloric values of children's foods and the ways in which children distribute their own foraging yield. Here, we report on foraging return rates, consumption data, and the compositional values for several of the plant foods collected by children. Due to predator pressure, it is not safe for children to wander far from camp without adult supervision, therefore they typically focus on foods that are close to camp and easy to collect and process. We calculated the caloric content for the following foods: baobab fruit (346 kcal/100g dry matter (DM)), berries (320 kcal/100g DM), legumes (311 kcal/100g DM), drupes (325 kcal/100g DM), and figs (365 kcal/100g DM). Legumes, drupes, and figs have not been previously analyzed; our values for baobab and berries agree with previous analyses (Murray et al. 2001, *Journal of Food Consumption and Analysis* 14: 3). In addition, we calculated foraging return rates and daily consumption values for children. Our results suggest that they collect a significant portion of their daily caloric intake and act as allomothers providing caloric contributions to other children.

Support: National Science Foundation, Regents of the University of California San Diego, Friends of the International Center at the University of California San Diego, and Harvard University.