



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Social Networks and Collateral Health Effects

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Christakis, Nicholas A. 2004. Social networks and collateral health effects. <i>British Medical Journal</i> 329, no.7459: 184-185.
Published Version	doi:10.1136/bmj.329.7459.184
Accessed	February 17, 2015 11:09:57 PM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:4276923
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)



Social networks and collateral health effects

Nicholas A Christakis

BMJ 2004;329:184-185
doi:10.1136/bmj.329.7459.184

Updated information and services can be found at:
<http://bmj.com/cgi/content/full/329/7459/184>

These include:

References

This article cites 9 articles, 5 of which can be accessed free at:
<http://bmj.com/cgi/content/full/329/7459/184#BIBL>

5 online articles that cite this article can be accessed at:
<http://bmj.com/cgi/content/full/329/7459/184#otherarticles>

Rapid responses

One rapid response has been posted to this article, which you can access for free at:
<http://bmj.com/cgi/content/full/329/7459/184#responses>

You can respond to this article at:
<http://bmj.com/cgi/eletter-submit/329/7459/184>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top left of the article

Topic collections

Articles on similar topics can be found in the following collections

[Sociology](#) (342 articles)
[Other communication](#) (298 articles)
[Patient - other](#) (548 articles)

Notes

To order reprints follow the "Request Permissions" link in the navigation box

To subscribe to *BMJ* go to:
<http://resources.bmj.com/bmj/subscribers>

Social networks and collateral health effects

Have been ignored in medical care and clinical trials, but need to be studied

Since a patient or a clinical trial participant is connected to other people through social network ties, medical interventions delivered to a patient, quite apart from their health effects in that person, may have unintended health effects in others to whom he is connected. The cumulative impact of an intervention is therefore the sum of the direct health outcomes in the patient plus the collateral health outcomes in others (figure). These effects, in both the patient and in their social contacts, might be positive or negative. Doctors, trialists, patients, or policy makers might see reason to take them into account when choosing treatment or evaluating benefit.


For example, treating depression in parents may increase their propensity to vaccinate their children, thereby saving children's lives. Replacing a hip or preventing a stroke may mean that a person is better able to care for his spouse, thus improving her health. Delivering a weight loss intervention to one person may trigger substantial weight loss in that person's friends. Giving a patient superior end of life care may decrease the stressfulness of the patient's death and thus decrease his spouse's propensity to die during bereavement.¹

These collateral health consequences that accrue to others are known to social scientists as externalities. They are similar to the increase in value a person's neighbour may see if the person refurbishes his property; the person himself derives no benefit from the neighbour's windfall, even though he has invested resources and created this new value. Here, however, we are considering not monetary externalities but rather specifically health externalities. Moreover, in the healthcare arena patients may derive value from such effects. For example, since 89% of patients feel that a good death involves not burdening family,² patients might prefer hospice care over standard terminal care if they felt it offered health benefits to their loved ones.¹

When the cost-benefit assessment is made by policy makers with a collective viewpoint, all the downstream costs and benefits of health care accruing to a group might be relevant, and the argument in favour of accounting for collateral effects might be even more compelling than that perceived by individual doctors or patients. From a societal perspective the assessment of the cost effectiveness of medical interventions might change substantially if the benefits of an intervention are seen as including the collateral positive effects and the costs as including the collateral negative effects.

Such a concern for collateral effects could, however, also lead to unexpected results. For example, preventing a death from heart attack, which is clearly desirable from the individual's perspective, may mean that we have to forego the motivation that would otherwise have accrued to others to whom the patient is connected to improve their own health habits. Another provocative implication is that policy makers might value socially connected individuals—such as married people—more when it comes to health care since benefits might be multiplicative in such people.³

We have scattered evidence supporting the plausibility and likely magnitude of such collateral health effects. The most well known example is that the death of one spouse increases the risk of death in the other.^{4,5} Moreover, morbidity in one spouse can contribute to morbidity in the other—for example, via caregiver burden.⁶ Breast cancer in one woman may motivate others to whom she is connected to have mammography.⁷ Exercise or smoking cessation in one person may prompt numerous others to behave similarly. Conversely, there may be epidemics of disorders such as obesity, alcoholism, suicide, or depression that might spread in a peer to peer fashion.⁸ Even loose social connections can be conduits for such effects; cancer in a celebrity, for example, may motivate many people not known to the index case to undergo cancer



Collateral health effects of medical care in social networks. In the conventional perspective on medical care, the benefits and costs of health care are judged by the way in which they help to achieve a direct, intended outcome in a patient. However, since a patient is connected to others through social ties, health care delivered to one person, quite apart from its health effects on that person, may have health effects on others. The cumulative impact of the intervention is therefore a sum of the direct outcomes in the patient plus the collateral outcomes in others. These outcomes may be both positive and negative in both the patient and in his or her social contacts (for example, side effects of medication in the individual, herd immunity in social contacts). Attention to, and measurement of, the existence of unintended outcomes arising out of the embeddedness of patients in social networks can prompt a rethinking of the relative value of healthcare interventions or of the conduct of clinical trials.

screening or choose particular treatments.⁹⁻¹⁰ Vaccinating some people in a population may cause others (for example, immunocompromised people) to become sick through the spread of the vaccine virus or, conversely, to remain well through the effect of herd immunity.

The existence of collateral health effects and the fact that each individual may be connected to numerous others, including relatives, friends, neighbours, and co-workers, implies developments in research and policy. To explore such effects, new datasets and methods will be needed. Most prospective cohort studies and randomised controlled trials today include only individuals who are followed to observe outcomes. Some social science and epidemiological cohort studies do ask individuals about the health status of their spouse or some other social contact, but only a few (for example, the US health and retirement survey) actually include the spouse or other social contacts in the study cohort. Developing datasets with such features and measurements is necessary to understand fully collateral health effects. Collecting information about the various contacts of people enrolled in clinical trials or epidemiological studies may represent an extension to study design similar to the extension in the 1990s of including cost effectiveness analyses as a standard feature of clinical trials.

Network phenomena are receiving increased attention in fields as diverse as engineering, biology, and sociology,¹¹⁻¹² but they are also relevant to health and medicine. Networks have emergent properties not explained by the constituent parts and not present in the parts. Understanding such properties requires

seeing whole groups of individuals and their interconnections at once. The existence of social networks means that people and events are interdependent and that health and health care can transcend the individual in ways that patients, doctors, policy makers, and researchers should care about.

Nicholas A Christakis *professor*

Department of Health Care Policy, Harvard Medical School, 180 Longwood Avenue, Boston, MA 02115 (christakis@hcp.med.harvard.edu)

Competing interests: None declared.

- 1 Christakis NA, Iwashyna TJ. The health impact on families of health care: a matched cohort study of hospice use by decedents and mortality outcomes in surviving, widowed spouses. *Soc Sci Med* 2003;57:465-75.
- 2 Steinhilber KE, Christakis NA, Clipp EC, McNeilly M, McIntyre L, Tulsky JA. Factors considered important at the end of life by patients, family, physicians, and other care providers. *JAMA* 2000;284:2476-82.
- 3 D Crane. *The sanctity of social life: physicians' treatment of critically ill patients*. New Brunswick, NJ: Transaction Books, 1990.
- 4 Schaefer C, Quesenberry CP, Soora W. Mortality following conjugal bereavement and the effects of a shared environment. *Am J Epidemiol* 1995;141:1142-52.
- 5 Lillard LA, Waite LJ. 'Til death do us part: marital disruption and mortality. *Am J Sociol* 1995;100:1131-56.
- 6 Schulz R, Beach SR. Caregiving as a risk factor for mortality: the caregiver health effects study. *JAMA*, 1999;282:2215-9.
- 7 Murabito JM, Evans JC, Larson MG, Kreger BE, Splansky GL, Mutalik KM, et al. Family breast cancer history and mammography. Framingham Offspring Study. *Am J Epidemiol* 2001;154:916-23.
- 8 Andrews JA, Tildesley E, Hops H, Li F. The influence of peers on young adult substance use. *Health Psychol* 2002;21:349-57.
- 9 Nattinger AB, Hoffmann RG, Howell-Pelz A, Goodwin JS. Effect of Nancy Regan's mastectomy on choice of surgery for breast cancer by US women. *N Engl J Med* 1998;279:762-6.
- 10 Cram P, Fendrick M, Inadomi J, Cowen ME, Carpenter D, Vijan S. The impact of a celebrity promotional campaign on the use of colon cancer screening: the Katie Couric effect. *Arch Intern Med* 2003;163:1601-5.
- 11 B Wellman, SD Berkowitz. *Social structures: a network approach*. Cambridge: Cambridge University Press, 1988.
- 12 Watts DJ, Strogatz SH. Collective dynamics of "Small World" networks. *Nature* 1998;393:440-2.

Condoms and prevention of HIV

Are essential and effective, but additional methods are also needed

Promotion of condoms has been a mainstay of HIV prevention policy. Over the past few years, however, the value and effectiveness of condoms have increasingly been called into question. The growing "abstinence only" movement in the United States questions the provision of condoms as part of the policy and messaging of the US Agency for International Development (USAID) and claims that condoms have had little to do with the successes achieved in reducing HIV in countries such as Uganda.¹⁻² Senior officials in the Roman Catholic Church also continue to argue about the morality of condom use and dispute its efficacy.³

But what does the evidence tell us? A recent review from the National Institutes for Health says that condoms are protective against HIV infection,⁴ reducing the probability of HIV transmission per sex act by as much as 95% and reducing the annual HIV incidence in serodiscordant couples by 90-95% when used consistently.⁵ However, the impact of inconsistent use of condoms is less substantial: a meta-analysis found that condom use of variable consistency among serodiscordant couples reduced the annual HIV incidence by 69%.⁶ This illustrates how the protection

provided by a condom is dependent both on its efficacy against HIV transmission per sex act and the consistency with which it is used. This is intuitive, yet the consistency of condom use is less commonly factored into scientific and policy debate.

Evidence from around the world highlights the extent to which patterns of condom use are influenced by the form of partnership in which they are being used. Interventions can achieve substantial increases in the use of condoms in commercial and casual sex partnerships. Several studies report high levels of condom use after interventions in commercial sex.⁷⁻⁹ But even in settings where HIV infection is widespread, the use of condoms in primary partnerships remains low—representative surveys of women in 13 African countries found that fewer than 7% report condom use in the last sex act with their regular partner.¹⁰ Surveys of sex workers in Asia generally find that, although many use condoms with their clients, fewer than 40% report using condoms in their last non-commercial sex act. Unless one partner knows they are HIV positive or