



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Woody Biomass and Bioenergy Potentials in Southeast Asia between 1990 and 2020

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Sasaki, Nophea, Wolfgang Knorr, David R. Foster, Hiroko Etoh, Hiroshi Ninomiya, Sengtha Chay, Sophanarith Kim, and Sengxi Sun. 2009. Applied Energy 86(Supp 1): S140-S150.
Published Version	doi:10.1016/j.apenergy.2009.04.015
Accessed	February 18, 2015 7:00:04 PM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:11077568
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

1 **Woody Biomass and Bioenergy Potentials in Southeast Asia between 1990 and 2020**

2

3 Nophea Sasaki^{1,2}, Wolfgang Knorr³, David R. Foster¹, Hiroko Etoh², Hiroshi Ninomiya²,

4 Sengtha Chay², Sophanarith Kim⁴, Sengxi Sun²,

5

6

7 ¹Harvard Forest, Harvard University

8 324 North Main Street, Petersham, MA 01366, USA

9 ²Graduate School of Applied Informatics, University of Hyogo

10 Kobe, Japan

11 ³QUEST, Department of Earth Sciences, University of Bristol

12 Bristol, UK

13 ⁴KIMSAS Services, Phnom Penh, Cambodia

14

15

16

17

18

19

20

21

22

23 **Abstract**

24 Forests in Southeast Asia are important sources of timber and other forest products, of
25 local energy for cooking and heating, and potentially as sources of bioenergy. Many of
26 these forests have experienced deforestation and forest degradation over the last few
27 decades. The potential flow of woody biomass for bioenergy from forests is uncertain and
28 needs to be assessed before policy intervention can be successfully implemented in the
29 context of international negotiations on climate change. Using current data, we developed a
30 forest land use model and projected changes in area of natural forests and forest plantations
31 from 1990 to 2020. We also developed biomass change and harvest models to estimate
32 woody biomass availability in the forests under the current management regime. Due to
33 deforestation and logging (including illegal logging), projected annual woody biomass
34 production in natural forests declined from 815.9 million tons (16.3 EJ) in 1990 to 359.3
35 million tons (7.2 EJ) in 2020. Woody biomass production in forest plantations was
36 estimated at 16.2 million tons yr⁻¹ (0.3 EJ), but was strongly affected by cutting rotation
37 length. Average annual woody biomass production in all forests in Southeast Asia between
38 1990 and 2020 was estimated at 563.4 million tons (11.3 EJ) yr⁻¹ declining about 1.5% yr⁻¹.
39 Without incentives to reduce deforestation and forest degradation, and to promote forest
40 rehabilitation and plantations, woody biomass as well as wood production and carbon
41 stocks will continue to decline, putting sustainable development in the region at risk as the
42 majority of the population depend mostly on forest ecosystem services for daily survival.

43

44 **Keywords:** Woody biomass; wood bioenergy; deforestation; forest degradation; land use
45 change; selective logging; Southeast Asia

46

47

48 **1. Introduction**

49

50 International concerns about global warming caused by excessive emissions of greenhouse
51 gases led to the adoption of the Kyoto Protocol to the United Nations Convention on
52 Climate Change (UNFCCC) in 1997. The protocol commits industrialized countries,
53 known as Annex I countries, to reduce greenhouse gas emissions during the first
54 commitment period between 2008 and 2012. As the first year of the first commitment
55 period ended, discussions for the post-Kyoto climate change agreements were carried out
56 in December 2008 in Poznan, Poland. Several industrialized countries have pledged to
57 reduce carbon emissions by up to 80% [1]. In addition to increasing energy efficiency and
58 increased reliance on renewable energy sources such as wind and solar power, reducing
59 emissions from deforestation and forest degradation (REDD) is likely to be a important
60 mitigation option in the post-Kyoto agreements, because deforestation and forest
61 degradation are responsible for the release of about 1.5 to 2.2 Gt C yr⁻¹ [2, 3] or about up to
62 25% of annual global emissions.

63

64 In addition to increasing carbon emissions, deforestation and forest degradation reduce
65 availability of woody biomass, on which approximately 2.5–2.7 billion people [4, 5]
66 depend for daily cooking fuel. Given the widespread dependency on wood for energy and
67 the importance of forests to mitigate climate change, there is a strong need to assess the
68 future availability while developing a path toward the sustainable use and management of
69 forests. Canadell and Raupach [6] proposed four strategies for managing forests for climate

70 change mitigation. One of the strategies is to expand the use of woody biomass to replace
71 the use of fossil fuels. Smeets *et al.* [7] provided an assessment of wood bioenergy
72 potentials on a global scale, concluding that there is high potential of woody biomass from
73 forests. Kinoshita *et al.* [8] evaluated the utilization of thinned wood as bioenergy in Japan
74 and concluded that bioenergy is increasingly important in substituting for the use of oil.
75 Utilization of woody biomass has a potential role in global warming mitigation because of
76 its low emissions of greenhouse gases compared to the utilization of oil or coal for power
77 generation [7, 8, 9]. To avoid power shortages such as occurred in 2001 in Brazil, the
78 Brazilian government has launched incentive programs to encourage the utilization of
79 biomass (including woody biomass) as bioenergy [10]. All these studies show the
80 importance of woody biomass in climate change mitigation and sustainable development.

81

82 Although the Food and Agricultural Organization of the United Nations' Regional Wood
83 Energy Development Program (referred to as FAO-RWEDP hereafter, [5]) provided an
84 estimate of woodfuels in South and Southeast Asia, their estimate did not incorporate the
85 illegal logging activities and significant logging damages that occur commonly in the
86 region [11, 12, 13]. Their estimate also did not consider local uses of wood, an important
87 consideration given the fact that the availability of woody biomass is directly linked to
88 daily survival in this region. About 30–90% of the population in individual countries in
89 Southeast Asia depends entirely on woody biomass for daily cooking and heating [14].
90 Furthermore, as deforestation and forest degradation continue, the future availability of
91 wood for this region is at risk. Between 1990 and 2005, forest area in Southeast Asia

92 declined approximately 2.6 million ha annually (about 1.2%) to 216.4 million ha in 2005
93 [15]. In addition, forest degradation due to logging (including illegal logging) and related
94 damages causes the gradual loss of forest biomass and carbon stocks [16]. As the
95 population and the demand for woody biomass continue to rise, the current and future
96 availability of woody biomass need to be assessed so that appropriate policies can be
97 introduced.

98

99 The aim of this study is to provide an assessment of the availability of woody biomass and
100 bioenergy in eleven countries in Southeast Asia under current forest management regime,
101 which includes illegal logging and logging damages. The paper is structured as follows: 1)
102 forest land use change models are developed to estimate the rate of deforestation and
103 reforestation through forest plantations; 2) woody biomass and harvesting models are
104 developed to estimate the biomass changes under current management regimes, and
105 potential woody biomass for bioenergy generation is estimated.

106

107 **2. Materials and Methods**

108 2.1. Forests in Southeast Asia

109 Southeast Asian countries in our study include Brunei, Burma, Cambodia, East Timor,
110 Indonesia, Laos, Malaysia, Philippines, Singapore, Thailand, and Vietnam. This region has
111 experienced fast economic development and the gradual loss of forest resources. Changes
112 in areas of natural forests and forest plantations between 1990 and 2005 are given in **Table**
113 **1**. According to FAO [15], natural forests consist of production, multiple-purpose, and

114 unspecified forests, protected forest, conservation forest, and forest for social services. The
 115 first three categories are grouped as production forest (*PdF*), where commercial logging
 116 and land development can take place, while the latter three categories are grouped as
 117 protected forest (*PrF*), where traditional firewood collection and small-scale logging for
 118 housing by local forest communities can take place. There are two types of forest
 119 plantations (FP) in the tropics, namely fast growing species plantation (*FPf*), which
 120 account for 47% of the total plantations and slow growing species plantation (*FPs*), which
 121 account for the rest [17]. For our study the proportion of fast and slow growing plantation
 122 remains unchanged during the modeling period between 1990 and 2020.

123 **Table 1**

124 2.2. Land use models

125 Over the last 15 years, although area of natural forests in Southeast Asia continued to
 126 decrease, area of forest plantations slowly increased as shown in Table 1. It could be
 127 argued that part of the deforested lands was replaced by forest plantations. Therefore, for
 128 our study, it is assumed that deforested lands are partially replaced by forest plantations
 129 (see **Fig.1** for illustration). With this assumption, the change in area of natural forests and
 130 forest plantations can be estimated using models developed by Kim Phat *et al.* [16]:

131

132
$$\frac{dPdF(t)}{dt} = -(k_a + k_b) \cdot PdF(t) \quad (1)$$

133
$$\frac{dPrF(t)}{dt} = 0 \quad (2)$$

134
$$\frac{dFP(t)}{dt} = k_a \cdot PdF(t) \quad (3)$$

135 where $PdF(t)$ is production forest at time t , $PrF(t)$ is protected forest, $FP(t)$ is forest
136 plantation, $-(k_a+k_b)$ is the change of $PdF(t)$, and k_a is the change of $FP(t)$

137

138 Data in Table 2 are used to derive $-(k_a+k_b)$, k_a , and the initial values ($t=0$ in 1990) for areas
139 of PdF and FP using linear regression methods. According to FAO [15], the area of
140 protected forests in the tropics increased by approximately 0.07% from 1990 to 2005.

141 During the modeling period of this study, PrF is considered to remain unchanged.

142

143 **Fig. 1**

144 **Table 2**

145

146 2.3. Woody biomass models

147 Standing biomass refers to all above-ground biomass in tons of dry matter, woody biomass
148 refers to biomass available for bioenergy generation, and bioenergy refers to energy
149 content in woody biomass. Leaves and root biomass are not included.

150

151 2.3.1. Natural forests:

152 A conceptual diagram illustrating the allocation of biomass is given in **Fig. 2**.

153

154 **Fig. 2.**

155

156 To estimate the standing biomass change in Southeast Asia, the following equations
157 modified from Kim Phat *et al.* [16] are used:

158

$$159 \quad \frac{dSB_i(t)}{dt} = MAI_i - H_i(t) - ddB_i(t) \quad (4)$$

$$160 \quad H_i(t) = \frac{f_w \cdot f_T}{1-r} \cdot \frac{SB_i(t)}{CC} \quad (5)$$

$$161 \quad ddB_i(t) = H_i(t) \quad (6)$$

$$162 \quad WAS_i(t) = s \cdot H_i(t) \quad (7)$$

163

164 where $SB_i(t)$ is standing biomass in i forest (*PdF*, *PrF*) (ton ha^{-1}), MAI_i is mean annual
165 biomass increment, $H_i(t)$ is harvested biomass, $ddB_i(t)$ is dead biomass caused by logging,
166 $WAS_i(t)$ is biomass waste due to trimming, felling, skidding and/or transporting, f_w is the
167 fraction of harvested stand biomass, f_T is the fraction of mature-tree stand biomass, CC is
168 the cutting cycle, r is the illegal logging rate, s is the rate of biomass waste. It is unlikely
169 that illegal loggers will harvest immature trees because of no market demand for such trees,
170 and therefore $r \leq 1 - f_w$. In our study the values for MAI , WAS , f_w , f_T , CC , and r (**Table 3**)
171 are based on various country reports [16]. Under conventional logging in East Kalimantan,
172 every one cubic meter of harvested wood resulted in the dying of 0.9–1.2 m^3 of life
173 biomass [18]. In the same region, Sist *et al.* [19] estimated that logging 10 trees caused
174 damage to other 309 trees all with a diameter at breast height over 10 cm, of which 206
175 trees were killed immediately. Therefore, for this study, $ddB_i(t)$ is assumed to be the same

176 as $H_i(t)$ for every time step. An energy content of 20 GJ ton^{-1} of dry woody biomass [20] is
177 used for energy estimates for biomass from natural forests and forest plantations.

178 **Table 3**

179

180 Total woody biomass available for bioenergy (BIE) in natural forests (NF) is estimated as:

181
$$\text{BIE}(t) = \sum_{i=1}^2 [\text{ddB}_i(t) + \text{WAS}_i(t) + \text{iuWAS}_i(t)] \cdot \text{NF}_i(t) \quad (8)$$

182

183 where $\text{iuWAS}_i(t)$ is iuWPI is iuWAS_i is in-use wasted wood due to wood processing at the
184 wood processing factories (see Fig. 2), $\text{NF}_i(t)$ is $\text{PdF}(t)$ and $\text{PrF}(0)$

185

186 Total biomass available for furniture making (BIF) is estimated as:

187
$$\text{BIF}(t) = \sum_{i=1}^2 \text{iuWPI}_i(t) \cdot \text{NF}_i(t) \quad (9)$$

188

189 where iuWPI is in-use wood product (see Fig. 2)

190

191 *2.3.2 Forest plantations:*

192 Unlike natural forests, mean annual increment is faster in forest plantations, where a clear-
193 cut system is applied. For this study, a logistic model is used to estimate biomass in forest
194 plantations:

195

196
$$\frac{dSB_j(t)}{dt} = \alpha_j \cdot SB_j(t) \cdot \left(1 - \frac{SB_j(t)}{SB_{MAX,j}}\right) \quad (10)$$

197

198 where $SB_j(t)$ is standing biomass in j plantations (j is fast-growing plantation, FPf and
 199 slow-growing plantation, FPs) (ton ha^{-1}), α_j is the growth rate of a forest plantation, $B_{MAX,j}$
 200 is the maximum wood biomass that a plantation can reach. Based on Brown [26] in Table
 201 4, average standing biomass increment is 7.7 and 5.9 $\text{ton ha}^{-1} \text{yr}^{-1}$ (see note under Table 4
 202 for calculation) over 10-yr and 40-yr cutting rotation (CR) (**Table 4, Table 5**) for FPf and
 203 FPs , respectively (see note under Table 4 for calculation). In reality, $B_{MAX,j}$ is unknown
 204 because forest plantations are usually harvested before they reach maturity age. For this
 205 study, $B_{MAX,j}$ is assumed at 200 and 300 ton ha^{-1} for FPf and FPs . With these assumptions,
 206 α and $SB_j(0)$ for FPf and FPs are derived at 0.2765 and 0.1337, and 7.7 and 5.9 $\text{ton ha}^{-1} \text{yr}^{-1}$,
 207 respectively. All harvested stem biomass is assumed to be used for pulp production
 208 (PPL_j), and the rest in branches and top logs are summed to be woody biomass for
 209 bioenergy generation (ddb_j) (see **Fig. 2**). Biomass in leaves (1.9% of the total above-
 210 ground biomass [23]) is left behind in the field.

211

212

213

Table 4, Table 5

214 Total standing biomass in forest plantation j , $SBFP_j(t)$ at time t , is

215

216
$$SBFP_j(t_n) = FPA_j(t_0) \times SB_j(t_n) + FPA_j(t_1) \times SB_j(t_{n-1}) + \dots + FPA_j(t_n) \times SB_j(t_0) \quad (11)$$

217

218 where $FPA_j(t)$ is the actual planted area at time t (million ha).

219

220 Total standing biomass in all plantations ($SBFP_{TOTAL}$) is therefore:

221

$$222 \quad SBFP_{TOTAL}(t_n) = \sum_{j=1}^2 SBFP_j(t_n) \quad (12)$$

223

224 Once each forest plantation reaches the CR age ($t=CR$), all biomass is harvested.

225 Plantations established in 1990 (start of the model) will be harvested in 1999 for FPf and

226 in 2029 for FPS . Replanting is assumed to be carried out one year after harvesting.

227

228 Total biomass available for pulp production (BIP) at time $t=n$ in forest plantations is

229

$$230 \quad BIP(t_n) = \sum_{j=1}^2 \frac{SBFP_j(t_n)}{BEF_j} \quad (13)$$

231

232 where BEF_j is a biomass expansion factor (see note under Table 4)

233

234 And woody biomass available for bioenergy (BIE) at time $t=n$ is

$$235 \quad BIE(t_n) = \sum_{j=1}^2 SBFP_j(t_n) - BIP_j(t_n) \quad (14)$$

236

237 **3. Results and Discussions**

238 3.1. Changes in area of forests

239 Over the modeling period, the area of natural forests declines from 245.9 million ha (231.1
240 for the 95% lower bound and 262.3 for the upper bound) in 1990 to 173.7 million ha
241 (165.6–182.6) in 2020, losing annually about 2.0% [$-(k_a+k_b)=-0.0202$]. Mean annual
242 changes in area of natural forests and forest plantation are estimated at 2.8 million ha yr⁻¹
243 between 1990 and 2005, and 2.4 million ha yr⁻¹ between 1990 and 2020 (**Table 6**). The
244 area of forest plantations slowly increases to 16.0 million ha (15.2–16.8) from 10.1 million
245 ha (9.8–10.2) in 1990, increasing about 0.2 million ha yr⁻¹ (**Fig. 3**). Because only about
246 0.09% ($k_a=0.0009$) of deforested forestland is converted to forest plantations, our results
247 suggest that most of the deforested land is converted to other types of land uses.
248 Altogether, Southeast Asia loses about 2.2 million ha yr⁻¹ (2.0–2.4) of forests over the
249 modeling period (**Table 6**). A previous study by Kim Phat *et al.* [16] estimated
250 deforestation in this region at 1.6 million ha yr⁻¹ between 1980 and 2050. This variation
251 may be due to the different modeling timeframe and the data used. Deforestation between
252 1990 and 2005 is estimated at 2.6 million ha yr⁻¹ by our model, which matches very well
253 with that estimated by FAO [15].

254 **Fig. 3**

255 **Table 6**

256 3.2. Standing biomass changes

257 Owing to deforestation and forest degradation, standing biomass in natural forests rapidly
258 declines from 45858.7 million tons (about 957.2 EJ) in 1990 to 26597.4 million tons

259 (531.9 EJ) in 2020, losing about 708.7 million tons yr^{-1} (14.2 EJ) or about 1.5% yr^{-1} .
260 Standing biomass in forest plantations is strongly influenced by cutting rotation, increasing
261 to 1013.8 million tons (20.3 EJ) in 2020 from merely 67.8 million tons (1.3 EJ) in 1990.
262 Altogether, Southeast Asian forests are projected to lose about 677.2 million tons yr^{-1} (13.5
263 EJ) between 1990 and 2020 (**Table 7**).

264 **Table 7**

265

266 3.3. Annual woody biomass and bioenergy production

267 In terms of woody biomass, natural forests produce, an average of 547.2 ± 24.6 million tons
268 yr^{-1} (\pm is standard error) (10.9 EJ) between 1990 and 2020, decreasing from 657.8 ± 23.0
269 million tons yr^{-1} (13.1 EJ) between 1990 and 2005 (**Fig. 4, Table 8**). Forest plantations
270 produce another 16.2 ± 7.5 million tons yr^{-1} (0.3 EJ) between 1990 and 2020. Altogether,
271 total annual production of woody biomass is 563.4 million tons (11.3 EJ) over the same
272 period between 1990 and 2020. Total energy consumption in Southeast Asia was estimated
273 at 6.4 EJ in 1990 and 15.7 EJ in 2006, increasing about 9.0% yr^{-1} [30]. Energy from
274 woodfuels in Southeast Asia (excluding Singapore and Brunei) was estimated at 2.4 EJ in
275 1993 [14] or about 33.1% of the total energy consumption in that year [30]. Energy from
276 woodfuels in this region increased, on average about 2.5% yr^{-1} between 1992 and 1995
277 [14]. Therefore, without effective policy to reducing deforestation and forest degradation,
278 energy shortage is likely to occur in Southeast Asia.

279 **Fig. 4**

280

281 Using carbon coefficients of 25 KgC GJ⁻¹ for coal, 20 KgC GJ⁻¹ for petroleum products,
282 and 15 KgC GJ⁻¹ for natural gas [31], carbon emission reductions associated with using
283 woody biomass instead of fossil fuels for energy generation are estimated at 281.7 TgC yr⁻¹
284 ¹ for replacing coal, 225.3 TgC yr⁻¹ for replacing petroleum products, and 169.0 TgC yr⁻¹
285 for replacing natural gas throughout the modeling period (Table 8).

286 **Table 8**

287

288 3.4. Comparison with previous studies

289 Our models project 92.0±4.1 (52.4 million tons) and 64.8±30.2 million m³ (33.3 million
290 tons), of wood for furniture making and pulpwood production over the modeling period
291 (**Table 8**). Industrial roundwood in Cambodia, Indonesia, Laos, Malaysia, Myanmar,
292 Philippines, Thailand, and Vietnam between 1991 and 2001 was reported at 77.2±5.6
293 million m³ yr⁻¹ [32]. With the addition of roundwood from illegal logging ($r=0.53$), the
294 above figure would have been 164.2 million m³ [=77.2/(1-0.53)], which is equivalent to
295 about 82.2 million m³ (=164.2*0.5, 0.5 is wood processing efficiency) of end-use wood
296 products, about 9.8 million m³ lower than our estimate. This difference may be due to the
297 unreported wood production from illegal logging in some countries in the region.

298

299 Results from previous studies on wood bioenergy using different methods and assumptions
300 are also compared here. Surrounded by uncertainties as identified by Koopmans [5], FAO-
301 RWEDP estimated the potential wood bioenergy from forested land in Southeast Asia at
302 about 6.7 EJ in 1994. If no illegal logging would take place, our model estimates wood

303 bioenergy at 7.0 EJ in 1994 and 5.9 EJ yr⁻¹ between 1990 and 2020 in the same region
304 (Table 9). Smeets & Faaij [7] estimated the loss of wood bioenergy due to tropical
305 deforestation at 13.0 EJ yr⁻¹ between 1998 and 2050. Our estimate of wood bioenergy loss
306 due to deforestation and forest degradation is 18.1 EJ yr⁻¹ between 1990 and 2020. This
307 difference may result from different methods and assumptions (Table 9). Using a global
308 land-use and energy model (GLUE), Yamamoto *et al.* [33] estimated wood bioenergy in all
309 developing countries worldwide at 45.9–85.2 EJ in 2100. Because of the difference in
310 study methods, assumptions, and scales, the results of their study are expected to be higher
311 than our estimate for Southeast Asia only.

312

313

Table 9

314

315 **4. Sensitivity Analysis**

316

317 Illegal logging is strongly affected by the political stability and governance in Southeast
318 Asia. If an illegal logging rate of 73% ($r=0.73$) as reported in Indonesia [37] is used in all
319 natural forests (NF), standing biomass in NF declines from 47858.7 million tons (957.2 EJ)
320 in 1990 to 20652.2 million tons (413.0 EJ) in 2020, a loss of about 1.9% annually. If illegal
321 logging is eliminated ($r=0$), standing biomass declines to 32393.3 million tons (647.9 EJ),
322 losing only about 1.1% as a result of deforestation (**Fig. 5**). In terms of woody biomass
323 production, our models project the mean annual production from all forests at 301.0 (6.0
324 EJ), 563.8 (11.3 EJ), and 831.7 million tons (16.6 EJ) for $r=0$, $r=0.53$ ($r=0.53$ was used in

325 our study), and $r=0.73$, respectively (**Fig. 6**). According to Fig. 6, illegal logging is likely
326 to cause a significant decline in annual woody biomass production. This suggestion is also
327 supported by Meyfroidt and Lambin [41] who found a sharp decline in stand density of
328 natural forests in Vietnam. International policy may influence biomass production. For
329 example, if ongoing discussions lead to the inclusion of the reduced emissions from
330 deforestation and degradation (REDD) in the post-Kyoto climate change agreement period
331 from 2013 to 2020, a large amount of biomass loss as well as carbon emissions could be
332 prevented. Therefore, woody biomass production will also change. Once slow growing
333 plantations become harvestable, woody biomass production is expected to increase as well.

334

335 Another uncertainty of our study relates to the potential increase of woody biomass
336 obtaining from forest rehabilitation as being increasingly implemented in Indonesia [38],
337 Philippines [39], and Vietnam ([40], but see Meyfroidt and Lambin [41]). Forest
338 rehabilitation could bring the deforested land or severely degraded forest back to its pre-
339 harvest level, and therefore would eventually increase woody biomass. Annual or biannual
340 re-assessment may reduce the future uncertainties regarding biomass projection.

341

Fig. 5

342

Fig. 6

343

344 **5. Policy Implications for Woody Biomass Production under REDD**

345 The current climate change agreement discussions include REDD in the post-Kyoto
346 agreements and give hope for tropical forest conservation. The Bali Action [42] and the

347 sustained interest in REDD during the 14th conference of the parties in Poznan in
348 December in 2008 [43] have led to increased attention to REDD [44, 6]. If REDD is finally
349 adopted, well-defined land use and logging planning that addresses the causes of
350 deforestation is required. The causes of deforestation in Southeast Asia could be classified
351 to be 1) the need for land for agricultural cultivation to feed increasing population [45], 2)
352 industrial plantation development [46], and 3) indiscriminate logging [12, 24, 47]. The
353 former is unavoidable because of the need for survival and requires well-assessed planning
354 and policies to encourage sustainable practices. The latter two may be due to policy
355 failures or the lack for incentives for long-term conservation of tropical forests. Economic,
356 social, and ecological assessments of different land use options that take into consideration
357 the financial incentives for protecting natural forests under REDD agreements are
358 necessary so that resource managers—be they government or companies— will have a clear
359 picture in terms of the financial returns and long-term social and ecological consequences
360 of their decisions.

361

362 In order to control indiscriminate logging and its associated forest degradation, incentives
363 are needed to promote reduced impact logging (RIL) which has been proven to reduce
364 damages [12, 24] to residual trees and soil, reduce wood waste (the latter is due to
365 untrained trimming, skidding, and transporting), and increase carbon sinks [47]. The
366 REDD agreements are likely to result in decreases in woody biomass, as overexploitation
367 and illegal logging would be gradually brought under control and the perpetual flow of
368 ecosystem services for sustainable development could be ensured. As forest rehabilitation

369 projects have been increasingly implemented in Indonesia [38], Philippines [39], and
370 Vietnam [40, 41], incentives for further promoting the widespread implementation of such
371 projects in other countries in the region could also lead to increase in woody biomasses as
372 well as wood production. Furthermore, alternative sources of energy such as wind and
373 solar power, and bioenergy through accelerating the development of plantations on
374 deforested lands should be sought. Financial incentives made available through REDD
375 agreements should be used wholly or partially for such alternatives.

376

377 Incentives or investment in plantations of hybrid species which, grow faster and are
378 environmentally adaptable on already deforested lands would lead to the increase of woody
379 biomass and pulpwood production for bioenergy and paper. Plantations could also
380 decrease the pressure on natural forests whose ecosystem services and functioning are vital
381 to sustainable development. Mean annual increment of some hybrid fast growing species
382 of Eucalyptus (such as *E. grandis*) reaches 53–60 m³ h⁻¹ yr⁻¹ (about 39.7–45.0 tons of all
383 above-ground biomass) [48]. If this growth rate could be achieved, future supplies of
384 woody biomass and pulp are likely to come from forest plantations, while natural forests
385 are managed for full ecosystem services.

386

387 **6. Conclusion**

388

389 This study developed models to estimate forest land use changes, standing biomass, and
390 woody biomass (for bioenergy generation) in Southeast Asia between 1990 and 2020. It

391 also discussed the incentives for reducing deforestation and implementing sustainable
392 forest management in the region. Our study methods could be applicable to any country or
393 region where selective logging is practiced.

394

395 The results show that Southeast Asian forests produce about 563.8 million tons yr⁻¹ (11.3
396 EJ) of woody biomass for the period spanning 1990 to 2020. The annual production of
397 woody biomass decreases about 1.5% over the same period. Without appropriate measures
398 to reduce deforestation and bring forests under sustainable management, Southeast Asia is
399 likely to face a shortage of woody biomass. Furthermore, if the current deforestation and
400 forest degradation continue, wood production, woody biomass, climate regulation
401 (including carbon sequestration), watershed protection, and ecosystem functioning will be
402 adversely affected, which, in turn could put sustainable development in the region at risk
403 because a large part of population in this region depend on forests and their ecosystems for
404 daily survival. Countries in the region should take advantages of the international
405 agreements such as the Kyoto Protocol or post-Kyoto agreements, i.e. REDD, to reduce
406 deforestation and forest degradation. At the same time, alternative sources of woody
407 biomass, i.e. from forest rehabilitation and plantations, should be made available, because,
408 currently only 0.08% of the 2.4 million ha deforested land is converted to forest
409 plantations, and the majority of these lands are still available for plantation.

410

411 Our results also suggest that using wood biomass to replace the use of fossil fuels for
412 energy generation could prevent carbon emissions of about 169.0–281.7 TgC yr⁻¹ between
413 1990 and 2020.

414

415 **Acknowledgement**

416 This work is funded through the Harvard Forest's Charles Bullard Fellowship in Forest
417 Research for Advanced Research and Study at Harvard University and a Grant-in-Aid for
418 Scientific Research (No. 18402003) from the Ministry of Education, Culture, Sports,
419 Science, and technology of Japan. The authors gratefully thank Betsy Colburn and
420 Jonathan Thompson of Harvard Forest, Harvard University for commenting and English
421 editing. We also thank the editor and reviewers for their invaluable comments.

422

423 **References**

- [1] UNFCCC. Press Release of the Conference of the Parties on its fourteenth session, held in Poznan from 1 to 12 December 2008. UNFCCC 2008
- [2] Gullison FR, Frumhoff CP, Canadell GJ, Field CB, Nepstad DC, Hayhoe K, Avissar R, Curran LM, Friedlingstein P, Jones CD, Nobre C. Tropical forests and climate policy. *Science* 2007; 136: 985–986.
- [3] Houghton RA. Revised estimates of the annual net flux of carbon to the atmosphere from changes in land use and land management 1850-2000. *Tellus* 55B 2003; 2: 378-390.
- [4] Gaye A. Access to energy and human development. *Fighting climate change: Human solidarity in a divided world. Human Development Report 2007/2008.* UNDP: New York; 2007.

- [5] Koopmans A. Biomass energy demand and supply for South and South-East Asia—assessing the resource base. *Biomass and Bioenergy* 2005; 28: 133–150
- [6] Canadell JG, Raupach MR. Managing forests for climate change mitigation. *Science* 2008; 320: 1456-1457.
- [7] Smeets MW, Faaij APC. Bioenergy potentials from forestry in 2050. An assessment of the drivers that determine the potentials. *Climatic Change* 2007; 81: 353-390.
- [8] Kinoshita T, Inoue K, Iwao K, Kagemoto H, Yamagata Y. A spatial evaluation of forest biomass usage using GIS. *Appl Energy* 2009; 86(1): 1-8.
- [9] Sathre R, Gustavsson L. Using wood products to mitigate climate change: External costs and structural change. *Appl Energy* 2009; 86(2) 251-257.
- [10] Filho PA, Badr O. Biomass resources for energy in North-Eastern Brazil. *Appl Energy* 2004; 77: 51-67.
- [11] Sist P, Saridan A. Stand structure and floristic composition of a primary lowland dipterocarp forest in East Kalimantan. *Journal of Tropical Forest Sci.* 1999; 11(4): 704-722.
- [12] Sist P, Sheil D, Kartawinata K, Priyadi H. Reduced-impact logging in Indonesian Borneo: some results confirming the need for new silvicultural prescriptions. *For Ecol Manage* 2003; 179: 415-427.
- [13] FAO. Financial and economic assessment of timber harvesting operations in Sarawak, Malaysia. *Forest Harvesting Case – Studies 17.* FAO, Rome; 2001.
- [14] FAO. Regional study on wood energy today and tomorrow in Asia. *Field Document No.50.* Bangkok: FAO; 1997.
- [15] FAO. Global forest resources assessment 2005. Progress towards sustainable forest management. *FAO Forestry Paper 147,* Rome; 2005.
- [16] Kim Phat N, Knorr W, Kim S. Appropriate measures for conservation of terrestrial carbon stocks – analysis of trends of forest management in Southeast Asia. *For Ecol Manage* 2004; 191: 283–299.
- [17] FAO. Global forest resource assessments 2000 (main report). *FAO forestry paper 140,* Rome; 2001.

- [18] Iskandar H, Snook LK, Toma T, MacDicken KG, Kanninen M. A comparison of damage due to logging under different forms of resource access in East Kalimantan, Indonesia. For Ecol Manage 2006; 237: 83-93.
- [19] Sist P, Nolan T, Bertault JG, Dykstra DP. Harvesting intensity versus sustainability in Indonesia. For Ecol Manage 1998; 108: 251-260.
- [20] Hall DO. Biomass energy in industrialised countries. A view of the future. For Ecol Manag 1997; 91:17–45.
- [21] Loehnertz PS, Cooz VI, Guerrero J. Sawing hardwoods in five tropical countries. Forest Products Journal 1996; 46: 51-56.
- [22] Brown S. Estimating biomass and biomass change of tropical forests: a Primer. FAO Forestry Paper 134, Rome; 1997.
- [23] Nascimentoa EMH, Laurance FW. Total aboveground biomass in central Amazonian rainforests: a landscape-scale study. For Ecol Manage 2002; 168: 311-321
- [24] Holmes TP, Blate GM, Zweede JC, Pereira R, Barreto P, Boltz F, Bauch R. Financial and ecological indicators of reduced impact logging performance in the eastern Amazon. For Ecol Manage 2002; 163: 93-110.
- [25] Sist P, Saridan A. Description of the primary lowland forest of Berau. In: Bertault JG and Kadir K, editors. Silvicultural research in a lowland mixed dipterocarp forest of East Kalimantan. The contribution of STREK project. CIRAD-FORDA-P.T. INHUTANII; 1998, p. 51–94.
- [26] Brown C. The global outlook for future wood supply from forest plantations. FAO Working Paper, GFPOS/WP/03, 156 pp.; 2000.
- [27] Miranda I, Almeida MH, Pereira H (2007) Influence of provenance, subspecies, and site on wood density in *Eucalyptus globulus* Labill. Wood and Fiber Sci 2007; 33: 9-15.
- [28] Arroja L, Dias AC, Capela I (2006) The role of *Eucalyptus globulus* forest and products in carbon sequestration. Climatic Change 2006; 74: 123-140.
- [29] Varmola M, Del Lungo A. Planted forests database: structure and contents. FAO Planted Forests and Trees Working Papers FP/25. FAO, Rome; 2003.

- [30] Energy Information Administration (2006). International Energy Annual 2006. Online data accessed on 17 March 2009.
<http://www.eia.doe.gov/emeu/international/energyconsumption.html>
- [31] Blasing TJ, Broniak C, Marland G. State-by-state carbon dioxide emissions from fossil-fuel use in the United States 1960–2000. *Mitigation and Adaptation Strategies for Global Change* 2005; 10: 659-674
- [32] FAO. Selected indicators of food and agriculture development in Asia-Pacific region 1992–2002. RAP PUBLICATION : 2003/10, Bangkok: FAO; 2003.
- [33] Yamamoto H, Yamaji K, Fujino J. Evaluation of bioenergy resources with a global land use and energy model formulated with SD technique. *Appl Energy* 1999; 63: 101-113.
- [34] FAO. Asia–Pacific forestry towards 2010. Report of the Asia–Pacific forestry outlook study. Asia-Pacific Forestry Commission, Rome: FAO; 1998a.
- [35] FAO. Global fibre supply model. Rome: FAO; 1998b.
- [36] FAO. Global forest products consumption, production, trade and prices: global forest products model projections to 2010. Rome: FAO; 1998c.
- [37] Kato T. Timber trafficking—Illegal logging in Indonesia, Southeast Asia and international consumption of illegally sourced timber. *Kyoto Review of Southeast Asia* 2002. Online Publication http://kyotoreview.cseas.kyoto-u.ac.jp/issue/issue1/article_151_p.html
- [38] Nawir AA, Murniati, Rumboko L (eds). *Forest rehabilitation in Indonesia: Where to after three decades*. Bogor: CIFOR; 2006
- [39] Pulhin JM, Chokkalingam U, Peras RJJ, Acosta RT, Carandang AP, Natividad MQ, Lasco RD, Razal RA. Chapter 2: historical overview. In: Chokkalingam U., Carandang AP, Pulhin JM, Lasco RD, Peras RJJ, Toma T (eds.), *One century of forest rehabilitation in the Philippines: Approaches, outcomes and lessons*. Bogor: CIFOR; 2006, pp. 6-41.

- [40] de Jong W, Sam DD, Hung TV. Forest rehabilitation in Vietnam: Histories, realities and future. Bogor: CIFOR; 2006.
- [41] Meyfroidt P, Lambin EF. Forest transition in Vietnam and its environmental impacts. *Glob Change Biol* 2008; 14: 1-8.
- [42] UNFCCC. Report of the Conference of the Parties on its thirteenth session, held in Bali from 3 to 15 December 2007. FCCC/CP/2007/6/Add, UNFCCC; 2008.
- [43] International Institute for Sustainable Development. Summary of the fourteenth conference of the parties to the UNFCCC and fourteenth meeting of the Kyoto Protocol: 1-12 December 2008. *Earth Negotiations Bulletin* 2008; 12(395).
- [44] Kindermann G, Obersteiner M, Sohngen B, Sathaye J, Andrasko K, Rametsteiner E, Schlamadinger B, Wunder S, Beach R. Global cost estimates of reducing carbon emissions through avoided deforestation. *Proc Natl Acad Sci USA* 2008; 105: 10302-10307.
- [45] Jha S, Bawa KS. Population Growth, Human Development, and Deforestation in Biodiversity Hotspots. *Conservation Biology* 2006; 20: 906-912.
- [46] Morton DC, DeFries RS, Shimabukuro YE, Anderson LO, Arai E, del Bon Espirito-Santo F, Freitas R, Morissette J. Cropland expansion changes deforestation dynamics in the southern Brazilian Amazon. *Proc Natl Acad Sci USA* 2006; 103: 14637-14641.
- [47] Putz FE, Zuidema PA, Pinard MA, Boot RG, Sayer JA, Sheil D, Sist P; Elias, Vanclay JK. Improved tropical forest management for carbon retention. *PLoS Biol* 2008; 6(7): e166 doi:10.1371/journal.pbio.0060166
- [48] Smith NJH, Serrao EAS, Alvim PT, Falesi IC. Amazonia: Resiliency and Dynamism of the Land and Its People. Tokyo: United Nations University; 1995

424

425

426

427

428

429 **Figures and Captions**

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450





Fig. 1. Illustration of forest land use change model

Note: it is assumed that new plantations are established on deforested land only (i.e. deforested PdF).



459 **Fig. 2.** Conceptual diagram for biomass allocation

460 Note

461 SB_i is standing biomass in natural forest i , $iuWP_i$ is in-use wood product; $iuWAS_i$ is in-use
 462 wasted wood, WAS_i is wasted wood due to felling, skidding, trimming and/or transporting;
 463 ddB_i is dead woody biomass caused by logging


464 SB_j is standing biomass in forest plantation j , ddB_j dead woody biomass in branches and
 465 top logs, PPL_j is biomass in stem for pulp production ($PPL_j = SB_j / BEF_j$, where BEF is
 466 biomass expansion factor. BEF_j values are presented in Table 4).

467

468

469

470



471

472

Fig. 3. Changes in area of forests in Southeast Asia (1990-2020)

473

Note: Confidence intervals for FP are not included because they are very small

474




Fig. 4. Annual wood bioenergy production in Southeast Asia

475


476

477

478 Note

479 Fast growing plantation established in 1990 become harvestable in 1999. Its annual woody
 480 biomass production is strongly affected by cutting rotation. Slow-growing plantation will
 481 become harvestable in 2029, and therefore more woody biomass production is expected
 482 thereafter.

483



484

485 **Fig. 5.** Standing biomass in natural forests under different rates of illegal logging

486

487

488

489


490

491

492

493

494



495

496 **Fig. 6** Woody biomass production from all forests under different rates of illegal logging

497

498 Note

499 Illegal logging leads to more production of woody biomass in the beginning, but it starts
 500 to decline sharply. Additionally, deforestation is also responsible for the gradual loss of
 501 woody biomass as seen in the figure above (green line) when all illegal logging is halted.

502

503 **Tables and Captions**

504

505 Table 1 Changes in area of forests in Southeast Asia 1990-2005

Country	1990 ('000 ha)			Total	2005 ('000 ha)		
	NF	FP	Total		NF	FP	Total
Brunei Darussalam	313.0	0.0	313.0	288.0	278.0	0.0	278.0
Cambodia	12946.0	67.0	13013.0	11613.0	10447.0	59.0	10506.0
Indonesia	116567.0	2209.0	118776.0	100854.0	88495.0	3399.0	91894.0
Laos	17314.0	4.0	17318.0	16631.0	16142.0	224.0	16366.0
Malaysia	22376.0	1956.0	24332.0	23250.0	20890.0	1573.0	22463.0
Myanmar	39219.0	394.0	39613.0	35250.0	32222.0	849.0	33071.0
Philippines	10574.0	1780.0	12354.0	8801.0	7162.0	620.0	7782.0
Singapore	2.0	0.0	2.0	2.0	2.0	0.0	2.0
Thailand	15965.0	2640.0	18605.0	17891.0	14520.0	3099.0	17619.0
Timor-Leste	966.0	29.0	995.0	897.0	798.0	43.0	841.0
Viet Nam	9363.0	967.0	10330.0	13775.0	12931.0	2695.0	15626.0
Total	245605.0	10046.0	255651.0	229252.0	203887.0	12561.0	216448.0
Total (million ha)	245.6	10.0	255.6	229.2	203.9	12.6	216.4

506 Source: FAO [15]

507

508

509

510

511

512

513

514

515

516

517 Table 2 Data used to derive*¹ land use model's initial values and parameters

Year	NF (million ha)			FP (million ha)			Tropical Forests (million ha)
	PdF	PrF	Subtotal	FPf	FPs	Subtotal	
1990	158.4	-	245.6	-	-	10.0	255.7
2000	130.5	-	217.7	-	-	11.6	229.3
2005	116.7	87.2	203.9			12.6	216.4
Initial value	158.7	87.2				10.1	
Parameters	$-(k_a+k_b)=-$ -0.0202					$k_a=0.0009$	

518 **Note**

519 *¹ Least square method was used to derive initial values and parameters

520 NF: Natural forests

521 PdF: Natural production forest

522 PrF: Natural protected forest

523 FP: Forest plantations

524 FPf: Fast growing forest plantation

525 FPs: Slow growing forest plantation

526

527

528

529

530

531

532

533 Table 3 Initial values and parameters for modeling biomass in natural forests

	PdF	PrF	Unit	Remarks and Sources
Stem Volume	200	200	m ³ ha ⁻¹	Taken from Kim Phat <i>et al.</i> [16]
SB(0)* ¹ (stand biomass at t=0)	194.6	194.6	ton ha ⁻¹	dry wood including branches, but without leaves
MAI* ² (mean annual increment)	1.0	1.0	ton ha ⁻¹ yr ⁻¹	dry wood including branches (no leaves, 1.9% of all; converted from [16])
f _W (fraction of harvested stand biomass)	0.3	0.1	%	30% of stand biomass of mature trees ([16] for PdF, 10% is assumed for PrF)
f _T (fraction of mature-tree stand biomass)	0.5	0.5	%	50% mature biomass take from [Kim Phat <i>et al.</i> 16]
CC (cutting cycle)	30	30	yrs	[16]
r (rate of illegal logging)	0.53	0.53	%	[16]
s* ³ (fraction of wasted wood)	0.3	0.3	%	See * ³
a* ⁴ (see Fig. 1) (processing efficiency)	0.5	0.5	%	[21]
WD (wood density)	0.57	0.57	ton m ⁻³	[22]
BEF (biomass expansion factor)	1.74	1.74		[22]
Leaves, l* ⁵	0.019	0.019		[23]
Energy Content	20		GJ per oven dry ton	[20]

534 Note

535 *¹= V*WD*BEF*(1-l), leaves are considered as litters that are left behind as nutrients

536 *²= 1*WD*BEF*(1-l), MAI in stem is 1 m³ ha⁻¹ yr⁻¹ (based on Kim Phat *et al.* [16])

537 *³: based on FAO [13], Homes *et al.* [24], and Sist and Sridan [25]

538 *⁴: Based on Loehnertz *et al.* [21]

539 *⁵: based on Nascimentoa and Laurance [23]

540

541

542

543

544

545

546

547

548

549 Table 4 Mean annual increments and cutting rotations for forest plantations

Species	MAI Range (ha ⁻¹ yr ⁻¹)				Rotation (yrs)*	Countries
	X (m ³)		Y (ton)			
	Min	Max	Min	Max		
<i>Acacia auriculiformis</i>	6.5	10.0	4.8	7.4	15	Myanmar, Philippines, Thailand and Vietnam
<i>Acacia mangium</i>	12.0	19.0	8.8	14.0	8	Indonesia, Malaysia and Papua New Guinea
	8.0	12.5	5.9	9.2		Laos, Philippines, and Vietnam
<i>Eucalyptus</i> species	8.0	12.5	5.9	9.2	5-15	Philippines, Thailand
	6.5	10.0	4.8	7.4		Malaysia
Mean	8.2	12.8	6.0	9.4		
For this study (fast growing species)			7.7		10	
<i>Casuarina</i> species	5.0	7.5	4.9	7.3	15-35	India and Vietnam
	1.5	2.5	1.5	2.4		Angola, Benin, Cuba, Kenya, Madagascar, Mauritius, Mozambique, Senegal, Somalia and Thailand
<i>Dalbergia sissoo</i>	3.0	5.0	2.9	4.9	24	Bangladesh, Bhutan, Burkina Faso, India, Nepal, Nigeria and Pakistan
<i>Swietenia macrophylla</i>	5.0	7.5	4.9	7.3	32	Indonesia and Philippines
<i>Terminalia</i> species	5.0	7.5	4.9	7.3		Bhutan, India and Jamaica
<i>Tectona grandis</i>	8.0	18.0	7.8	17.5	44	Colombia, Costa Rica, Jamaica, Nicaragua, Panama and Trinidad and Tobago
	4.0	6.0	3.9	5.8		Indonesia, Laos, Malaysia, Myanmar, Philippines, Thailand, and Vietnam
Mean	4.5	7.7	4.4	7.5		
For this study (slow growing species)			5.9		40	

550 Source: Brown [26]

551

552 Note:

553 $Y = X \times WD \times BEF \times (1 - 0.019)$ where WD is wood density, WD= 0.5 based on Miranda *et al.*
 554 [27] and Arroja *et al.* [28] for fast growing species and WD=0.57 [22] for slow growing
 555 species; and BEF is biomass expansion factor, BEF=1.50 [26]. (2006) and 1.74 [22] for
 556 fast growing and slow growing species, respectively, 0.019 is 1.9% in leaves [23]

557

558 *: Rotation length was taken as an average of rotation length of major species reported in
 559 Varmola and Del Lungo [29]

560

561

562

563

564

565 Table 5 Parameters for modeling biomass in forest plantations

	FPf	FPS	Unit	Remarks and Source
B _{MAX}	200	300	ton ha ⁻¹	Maximum standing biomass (all aboveground but without leaves)
B(0)	7.7	5.9	ton ha ⁻¹	All aboveground but without leaves
α	0.2765	0.1337		
MAI	7.7	5.9	ton ha ⁻¹ yr ⁻¹	[26]
CC	10	40	yrs	[26]
WD	0.50	0.57		[27] for fast, [22] for slow growing plantation
BEF	1.50	1.74		[28] for fast, [22] for slow growing plantation
Litters	0.019	0.019		[23]
Energy Content	20		GJ per oven dry ton	[20]

566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591

592 Table 6 Mean annual changes in area of natural forests and forest plantations (1990-2020)

Forests	1990-2005		1990-2020	
	(million ha)	(% to 1990)	(million ha)	(% to 1990)
Natural Forests	-2.8	-1.7	-2.4	-1.5
PdF	-2.8	-1.7	-2.4	-1.5
PrF	0	0	0	0
Forest Plantations	0.2	1.7	0.2	2.0
PFf	0.1	0.8	0.1	0.9
PFs	0.1	0.9	0.1	1.0
Total	-2.6	-1.0	-2.2	-0.9

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617 Table 7 Total standing biomass in natural forests and forest plantations (1990-2020)

Forest Type	1990	2005	2020	Annual Change	
				1990-2005	1990-2020
	million tons			million tons yr ⁻¹	
Natural Forests	47858.7	34202.9	26597.4	-910.4	-708.7
PdF	30884.7	17765.9	10611.6	-874.6	-675.8
PrF	16974.1	16436.9	15985.8	-35.8	-32.9
Forest Plantations	67.8	367.4	1013.8	20.0	31.5
PFf* ¹	36.4	150.2	92.5	7.6	1.9
PFs* ²	31.4	217.2	921.3	12.4	29.7
Total	47926.6	34570.3	27611.2	-890.4	-677.2
Total (EJ*³)	958.5	691.4	552.2	-17.8	-13.5
In terms of carbon stock changes (TgC yr ⁻¹)* ⁴					
Natural Forests	23929.4	17101.4	13298.7	455.2	354.4
Forest Plantations	33.9	183.7	506.9	-10.0	-15.8
Total	23963.3	17285.1	13805.6	445.2	338.6

618

619 Note:

620 *¹: Standing biomass is strongly affected by cutting rotation621 *²: Standing biomass will be harvested in 2029, thereafter standing biomass will be
622 reduced.623 *³: EJ is exajoule (1 EJ = 10⁹ GJ)624 *⁴: Multiplying by 0.5 carbon content in dry woody biomass. One Tetragram Carbon
625 (TgC) is one million tons of carbon626 *⁵: Minus sign (-) refers to carbon sinks

627

628

629

630

631

632

633

634

635 Table 8 Mean annual woody biomass and bioenergy production, end-use wood and pulp
 636 production in Southeast Asia

Forests	Year	1990-2005				1990-2020			
		million tons yr ⁻¹		EJ yr ⁻¹		million tons yr ⁻¹		EJ yr ⁻¹	
		Mean	s.e.* ³	Mean	s.e.	Mean	s.e.	Mean	s.e.
Natural Forests									
BIE		657.8	23.0	13.2	0.5	547.2	24.6	10.9	0.5
BIF (million m ³)* ¹		110.6	3.9			92.0	4.1		
PdF									
BIE		533.4	22.7	10.7	0.5	424.5	24.3	8.5	0.5
BIF (million m ³)* ¹		89.7	3.8			71.4	4.1		
PrF									
BIE		124.4	0.3	2.5	0.0	122.6	0.4	2.5	0.0
BIF (million m ³)* ¹		20.9	0.1			20.6	0.1		
Forest Plantations									
BIE		15.7	14.3	0.3	0.3	16.2	7.5	0.3	0.2
BIP (million m ³)* ¹		62.8	57.2			64.8	30.2		
FPf									
BIE		15.7	14.3	0.3	0.3	16.2	7.5	0.3	0.2
BIP (million m ³)* ¹		62.8	57.2			64.8	30.2		
FPs									
BIE		0				0			
BIP (million m ³)* ¹		0				0			
Total									
BIE (million ton)		673.5		13.5		563.4		11.3	
BIF (million m ³)		110.6				92.0			
BIP (million m ³)		62.8				64.8			
In terms of carbon emissions reductions* ² (in TgC yr ⁻¹) by using wood bioenergy to replace:									
Coal				336.7				281.7	
Petroleum products				269.4				225.3	
Natural gas				202.0				169.0	

637 Note

638 *¹: is converted by taking biomass dividing by wood density

639 *²: is derived by multiplying bioenergy (1 EJ = 10⁹ GJ) with carbon coefficients of 25 KgC
 640 GJ⁻¹ for coal, 20 KgC GJ⁻¹ for petroleum products, and 15 KgC GJ⁻¹ for natural gas [31]

641 and dividing by 10⁹ (1 TgC = 10⁹ KgC)

642 *³: s.e. is standard error

643 BIE: woody biomass available for bioenergy
644 BIF: biomass available for furniture making
645 BIP: biomass available for pulp production (*BIP*)
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668

669 Table 9 Previous studies on wood bioenergy

Authors	Methods	Major variables	Scale	Results
This study	Land use change model, biomass stock change model, biomass harvesting model	Natural forests, forest plantation, illegal logging, forest degradation	Regional	Deforestation and forest degradation reduce about 18.1 EJ yr ⁻¹ between 1990-2020. Potential bioenergy is 10.9 EJ yr ⁻¹ between 1990 and 2020. Potential wood bioenergy (no illegal logging) is 7.0 EJ in 1994 and 5.9 EJ yr ⁻¹ between 1990 and 2020.
FAO-Regional Wood Energy Development Program Koopmans [5] (2005)	Extrapolation using data 1990-1995. Biomass growth is assumed to increase 1% every year. Biomass growth of plantation was assumed at 6-10 m ³ ha ⁻¹ yr ⁻¹ . 80% of non-wooded lands also produce woodfuels	Natural forests, forest plantations, non-wooded lands. No illegal logging	Regional	Potential wood bioenergy is 6.7 EJ in 1994 from forested land in Southeast Asia
Smeets & Faaij (2007) [7]	Potential woody biomass in all forests is obtained by multiplying forest area and gross annual increment (GAI) under various scenarios. Data on forest area and GAI were taken from FAO [34], [35], [36]	Natural forests, forest plantations, and tree outside forests. Only GAI is harvested.	Global	Deforestation reduces about 13.0 EJ yr ⁻¹ between 1998 and 2050
Yamamoto <i>et al.</i> (1999) [33]	Global land-use and energy model (GLUE)	Natural forests, forest plantations, arable lands	Global	Potential wood bioenergy is 45.9-85.2 EJ in 2100 in all developing countries worldwide

670

671

672