



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Structural Insights into sGC

The Harvard community has made this article openly available. Please share how this access benefits you. Your story matters.

Citation	Akker, Focco van den, Xiaolei Ma, Faye Martin, Priyaranjan Pattanaik, Pius Padayatti, Matthew Warman, and Annie Beuve. 2009. Structural insights into sGC. BMC Pharmacology 9(Suppl. 1): S42.
Published Version	doi:10.1186/1471-2210-9-S1-S42
Accessed	February 19, 2015 9:54:53 AM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:10318217
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

BMC Pharmacology



Oral presentation Open Access

Structural insights into sGC

Focco van den Akker*¹, Xiaolei Ma², Faye Martin¹, Priyaranjan Pattanaik³, Pius Padayatti⁴, Matthew Warman⁵ and Annie Beuve⁶

Address: ¹Department of Biochemistry/RT500, Case Western Reserve University, Cleveland, OH, 44106, USA, ²Genentech, San Francisco, CA, USA, ³Syngene, Bangalore, India, ⁴Polgenix, Cleveland, OH, USA, ⁵Harvard Medical School, Boston, MA, USA and ⁶UMNDJ, Newark, NJ, USA

Email: Focco van den Akker* - focco.vandenakker@case.edu

* Corresponding author

from 4th International Conference of cGMP Generators, Effectors and Therapeutic Implications Regensburg, Germany. 19–21 June 2009

Published: 11 August 2009

BMC Pharmacology 2009, 9(Suppl 1):S42 doi:10.1186/1471-2210-9-S1-S42

 $This\ abstract\ is\ available\ from:\ http://www.biomedcentral.com/1471-2210/9/S1/S42$

© 2009 Akker et al; licensee BioMed Central Ltd.

The membrane and soluble guanylyl cyclases (sGCs) are key families of enzymes that produce the second messenger cGMP. Although both families recognize different ligands and have therefore different N-terminal input domains, they do have similar C-terminal output domains containing a coiled-coil domain and a guanylyl cyclase catalytic domain. We present here our latest structure-function studies on both the sGC and membrane guanylyl cyclases in an effort to enhance our molecular understanding of these receptors to delineate their similarities and differences. Our studies include a new 2.15 Å crystal structure of part of sGC and mutagenesis studies of these families of guanylyl cyclases thereby providing new insights into the structure, activation, and regulation of these receptors.