



Evaluation of early body shape for epidemiological research in absence of objective measurements

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters.

Citation	Akarolo-Anthony, Sally N, and Clement A Adebamowo. 2012. Evaluation of early body shape for epidemiological research in absence of objective measurements. BMC Proceedings 6(Suppl 3): P70.
Published Version	doi:10.1186/1753-6561-6-S3-P70
Accessed	February 19, 2015 10:46:55 AM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:10497289
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

POSTER PRESENTATION

Open Access

Evaluation of early body shape for epidemiological research in absence of objective measurements

Sally N Akarolo-Anthony^{1*}, Clement A Adebamowo²

From Metabolism, diet and disease
Washington, DC, USA. 29-31 May 2012

Background

Developing countries are undergoing epidemiological transitions with increasing prevalence of non-communicable diseases (NCD). There is increasing appreciation of the role of early life exposure in the etiology of many of these conditions particularly the role of body shape and size for given age. However, many of these environments lack early life records of birth and growth. So it is not possible to correlate early life characteristics with risk of disease in these populations. In this study we, evaluated the use of body images as an estimator of body size and characteristics.

Methods

1058 workers at a government office in Nigeria were enrolled in a study of body size, dietary energy intake and physical activity. The participants answered an interviewed administered questionnaire who also measured their anthropometric characteristics.

Results

The mean age (SD) of participants was 41.5 (9.3) years and mean BMI (SD) was 27.0 (4.9), mean waist (SD) = 75.4 (26.8) cm, mean hip circumference (SD) = 86.7 (33.1) cm, mean waist hip ratio (WHR) = 0.88 (0.10) and mean weight (SD) = 74.6 (14.2) kg. There was strong correlation between the interviewee perception of their current body type and that of the interviewer ($r = 0.73$, p -value < 0.001). Compared with anthropometric characteristics, the body images correlated most strongly with BMI ($r = 0.56$, p -value < 0.001) but less so with

waist circumference ($r = 0.33$), hip circumference ($r = 0.33$), WHR ($r = 0.15$).

Conclusions

Our results suggest that study participants are able to evaluate their current body shape and correlate with it images. These images correlate well with BMI and can be used in epidemiological studies where availability of objective measures are limited. The study is now being extended to evaluate body shape in early life.

Author details

¹Nutrition Department, Harvard School of Public Health, Boston, MA, 02215, USA. ²Epidemiology and Public Health, School of Medicine, University of Maryland, Baltimore, MD, 021201, USA.

Published: 27 June 2012

doi:10.1186/1753-6561-6-S3-P70

Cite this article as: Akarolo-Anthony and Adebamowo: Evaluation of early body shape for epidemiological research in absence of objective measurements. *BMC Proceedings* 2012 **6**(Suppl 3):P70.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Nutrition Department, Harvard School of Public Health, Boston, MA, 02215, USA

Full list of author information is available at the end of the article