



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Epitope Specificity Appears To Be an Important Determinant of In Vivo Killing Ability of Simian Immunodeficiency Virus (SIV)-Specific CD8+ T Cells

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Pozzi, L., A. Carville, V. Varner, J. Sen, D. Knipe, and A. Kaur. 2012. Epitope specificity appears to be an important determinant of in vivo killing ability of simian immunodeficiency virus (SIV)-specific CD8+ T cells. Oral Presentation. <i>Retrovirology</i> 9(Suppl 2): O6.
Published Version	doi:10.1186/1742-4690-9-S2-O6
Accessed	February 19, 2015 10:50:20 AM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:10536951
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)



ORAL PRESENTATION

Open Access

Epitope specificity appears to be an important determinant of in vivo killing ability of simian immunodeficiency virus (SIV)-specific CD8+ T Cells

L Pozzi*, A Carville, V Varner, J Sen, D Knipe, A Kaur

From AIDS Vaccine 2012

Boston, MA, USA. 9-12 September 2012

Background

CD8+ cytotoxic T lymphocytes (CTL) are a critical component of antiviral immunity and play an important role in the control of lentiviral infection.

Methods

We have developed an in vivo CTL assay to directly measure the killing capacity of MHC-restricted SIV-specific CTL in rhesus macaques (RM). In order to evaluate the in vivo efficacy of different epitope-specific CTL, we compared the in vivo killing capacity of Mamu-A*02-restricted Nef YY9 and Gag GY9 CD8+ T lymphocytes in three RM vaccinated with a recombinant HSV prime/DNA boost SIV vaccine regimen.

Results

Tetramer frequencies of Mamu-A*02-restricted Nef YY9-specific CD8+ T-cells were at least 10-fold higher than Mamu-A*02-restricted Gag GY9-specific CD8+ T-cells in individual animals both pre- and post-challenge. Prior to SIV challenge, both CTL populations showed poor and incomplete killing (22-35%) of target cells over 18 hours. Seven weeks post-challenge, there was a marked increase in the CTL killing capacity of both Nef YY9 and Gag GY9-specific CTL with 26-38% killing occurring over the first two hours and up to 100% killing over 18 hours. Surprisingly, Gag GY9-specific CTL consistently showed equivalent or greater in vivo killing when compared to Nef YY9-specific CTL even though the percentages of IFN- γ secreting and degranulating cells upon peptide stimulation were comparable.

Conclusion

These data suggest that epitope specificity rather than tetramer frequency determines the ability of SIV-specific CTL to kill infected cells. Taken together, these data may have important implications for the development of a successful HIV vaccine.

Published: 13 September 2012

doi:10.1186/1742-4690-9-S2-O6

Cite this article as: Pozzi et al.: Epitope specificity appears to be an important determinant of in vivo killing ability of simian immunodeficiency virus (SIV)-specific CD8+ T Cells. *Retrovirology* 2012 9(Suppl 2):O6.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Harvard Medical School, Southborough, MA, USA