



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Comprehensive Genomic Profiling of the Lung Transcriptome in Emphysema and Idiopathic Pulmonary Fibrosis Using RNA-Seq

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters.

Citation	Kusko, Rebecca L., John Brothers, Gang Liu, Lingqi Luo, Brenda Juan Guardela, John Tedrow, Yuriy Aleksyev et al. 2012. Comprehensive genomic profiling of the lung transcriptome in emphysema and idiopathic pulmonary fibrosis using RNA-Seq. BMC Proceedings 6(Suppl 6): P21.
Published Version	doi:10.1186/1753-6561-6-S6-P21
Accessed	February 19, 2015 11:49:53 AM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:10587977
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

POSTER PRESENTATION

Open Access

Comprehensive genomic profiling of the lung transcriptome in emphysema and idiopathic pulmonary fibrosis using RNA-Seq

Rebecca L Kusko^{1*†}, John Brothers II^{2†}, Gang Liu³, Lingqi Luo³, Brenda Juan Guardela⁴, John Tedrow⁴, Yuriy Aleksyev^{1,5}, Ivana V Yang⁶, Mick Correll⁷, Mark Geraci⁸, John Quackenbush⁷, Frank Sciruba⁴, Marc Lenburg^{1,2,3,5}, David A Schwartz⁶, Jennifer Beane³, Naftali Kaminski⁴, Avrum Spira^{1,2,3}

From Beyond the Genome 2012
Boston, MA, USA. 27-29 September 2012

Background

As part of the Lung Genomics Research Consortium (LGRC), we sought to characterize transcriptomic alterations underlying the molecular pathogenesis of emphysema and idiopathic pulmonary fibrosis (IPF) using mRNA-Seq, and comparing to mRNA and microRNA microarray data obtained from the same samples.

Methods

87 LGRC lung tissue samples were sequenced on the Illumina GAIIx, generating 75 nt paired-end reads and approximately 30-40 million reads per sample. Using gapped aligner Tophat, an average of 85% of reads aligned to hg19. Gene expression was quantified using Cufflinks and Ensembl59 known gene annotation ($n = 24,249$ genes). All lung tissue samples used in this study, as well as additional LGRC lung tissue samples, were run on Agilent V2 human whole genome arrays and Agilent V3 human miRNA microarrays.

Results

Using a mixture model, expression of 6,359 genes was detected, 9,538 genes were not detected, and 8,397 genes were variably detected across all lung tissue samples. Using a subset of 58 samples from subjects with IPF ($n = 19$), emphysema ($n = 19$) or control samples ($n = 20$), differential gene expression was determined using a t test to compare each disease state with control. The expression levels of 1,770 genes differed between IPF and control, and

220 genes between emphysema and control ($P < 0.001$). Genes that go up in both emphysema and IPF relative to control were enriched for the p53/hypoxia pathway (KEGG, Biocarta) by GSEA. Gene expression estimates were highly correlated between the mRNA-Seq and array datasets across the same samples; however, we identified additional gene expression changes by RNA-Seq (either not significant by or not assayed by microarray) and validated these by quantitative PCR. Array-based gene expression estimates from additional lung tissue samples not sequenced, together with immunohistochemistry, confirmed the upregulation of the p53/hypoxia pathway in emphysema and IPF. Using reads that aligned across known splice junctions, we identified 5 emphysema-associated and 19 IPF-associated alternative splicing events. These events included the loss of exons and changes to the 3' UTR. Using qPCR and Nanostring, we validated two examples of differential splicing, one of which is shared by both chronic obstructive pulmonary disease (COPD) and IPF and involved with the p53 pathway. Finally, using miRconnX, miRNA microarray data and mRNA-Seq, data were integrated with a prior network of computationally predicted and experimentally validated miRNA-mRNA interactions. miRNAs that co-vary with differentially expressed p53/hypoxia genes in IPF and emphysema were identified and validated using *in vitro* miRNA perturbation studies.

Conclusions

Our data indicate that the lung transcriptome is altered in emphysema and IPF, and suggest that these changes may include alterations in gene expression that are regulated by miRNA as well as disease-associated alternative

† Contributed equally

¹Genetics and Genomics Graduate Program, Boston University School of Medicine, Boston, MA, USA

Full list of author information is available at the end of the article

splicing events that are shared between COPD and IPF. Specifically we observed a shared upregulation of the p53/hypoxia pathway and decreased expression of the miRNAs that may regulate this pathway. Our data also reveal disease-associated changes in known splice junctions that may affect gene regulation or protein function. This unprecedented high-resolution view of the lung transcriptome associated with IPF and COPD may ultimately provide biomarkers of risk and response to therapy as well as potential therapeutic targets.

Author details

¹Genetics and Genomics Graduate Program, Boston University School of Medicine, Boston, MA, USA. ²Bioinformatics Program, Boston University, Boston, MA, USA. ³Division of Computational Biomedicine, Department of Medicine Boston University School of Medicine, Boston, MA, USA. ⁴Simmons Center for Interstitial Lung Disease and Department of Medicine, University of Pittsburgh Medical Center, Pittsburgh, PA, USA. ⁵Department of Pathology and Laboratory Medicine, Boston University School of Medicine, Boston, MA, USA. ⁶Center for Genes, Environment and Health and Department of Medicine, National Jewish Health, Denver, CO, USA. ⁷Dana-Farber Cancer Institute and Harvard School of Public Health, Boston, MA, USA. ⁸Department of Medicine, University of Colorado School of Medicine, Aurora, CO, USA.

Published: 1 October 2012

doi:10.1186/1753-6561-6-S6-P21

Cite this article as: Kusko *et al.*: Comprehensive genomic profiling of the lung transcriptome in emphysema and idiopathic pulmonary fibrosis using RNA-Seq. *BMC Proceedings* 2012 **6**(Suppl 6):P21.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

