



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Functional Multiplex Reporter Assay Using Tagged Gaussia Luciferase

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Van Rijn, Sjoerd, Jonas Nilsson, David P. Noske, W. Peter Vandertop, Bakhos A. Tannous, and Thomas Würdinger. 2013. Functional multiplex reporter assay using tagged Gaussia luciferase. <i>Scientific Reports</i> 3: 1046.
Published Version	doi:10.1038/srep01046
Accessed	February 19, 2015 11:56:58 AM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:10622999
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)



Functional multiplex reporter assay using tagged *Gaussia* luciferase

Sjoerd van Rijn¹, Jonas Nilsson^{1,2}, David P. Noske¹, W. Peter Vandertop¹, Bakhos A. Tannous³ & Thomas Würdinger^{1,3}

¹Neuro-oncology Research Group, Department of Neurosurgery, Cancer Center Amsterdam, VU University Medical Center, Amsterdam, The Netherlands, ²Department of Radiation Sciences, Oncology, Umeå University, Umeå, Sweden, ³Neuroscience Center, Department of Neurology, Massachusetts General Hospital and Neuroscience Program, Harvard Medical School, Boston, MA, USA.

We have developed a multiplex reporter system to monitor multiple biological variables in real-time. The secreted *Gaussia* luciferase was fused to ten different epitope tags (Gluc_{tag}), each expressed in different tumor cells. By immunobinding of the tags followed by Gluc_{tag} detection, this system allowed the independent and real-time monitoring of mixed cell cultures *in vitro* and of mixed subcutaneous and intracranial tumor subpopulations *in vivo*.

To study complex processes in systems biology, tools are needed that give insight into multiple molecular and cellular interactions simultaneously and in real-time^{1,2}. Several cellular reporter systems are in use to shed light on biological processes, many based on luciferase enzymes and their photon production upon luciferin substrate addition, including Firefly luciferase (Fluc)³, *Renilla* luciferase (Rluc)⁴, and *Gaussia* luciferase (Gluc)^{5,6}. These bioluminescent reporters have been used to monitor biological processes such as cellular proliferation and differentiation⁷, transcription factor activity⁸, translational repression by miRNA^{9,10}, and intercellular interactions¹¹, reviewed in¹². Other commonly used reporters are based on the green fluorescent protein (GFP), the red fluorescent protein (RFP) and, more recently, near infra-red fluorescent proteins^{13,14} which can be multiplexed together to monitor several processes simultaneously using spectral unmixing in conjunction with fluorescence molecular tomography¹⁵. These reporters have several disadvantages including low sensitivity due to autofluorescence, complexity, and the requirement of expensive instrumentation. Further, similar to bioluminescence, limited numbers of variables can be monitored simultaneously, limiting their use in multiplex screening assays and cellular heterogeneity studies. Here, we developed a multiplex bioluminescence assay that allows monitoring of many variables in real-time within the same biological system. This assay is based on Gluc since this reporter is highly sensitive, naturally secreted, and can be detected in the conditioned medium of cells in culture as well as in the blood of animals *ex vivo*, allowing real-time monitoring of cellular variables^{5,6}.

Results

We constructed a library of lentiviral vectors consisting of the Gluc cDNA fused to ten different epitope tags at its C-terminus, under the constitutively active cytomegalovirus (CMV) promoter resulting in $\text{Gluc}_{\text{Flag}}$, Gluc_{His} , Gluc_{HA} , $\text{Gluc}_{\text{AcV5}}$, Gluc_{V5} , Gluc_{Glu} , Gluc_{Myc} , Gluc_{K13} , Gluc_{Au1} , Gluc_{E2} , and the control reporter construct without tag, $\text{Gluc}_{\text{Ctrl}}$. These lentiviral vectors also express cerulean fluorescent protein (CFP) separated from Gluc_{tag} by an internal ribosomal entry site (IRES) element as a marker for transduction efficiency (Fig. 1a and Methods section). We employed human U87 glioblastoma cells stably expressing Fluc and the mCherry fluorescent protein⁶ (U87-FM) for validation of the Gluc_{tag} multiplex system. These cells were transduced with each of the lentiviral vectors to produce ten different cell lines, each stably expressing a different Gluc_{tag} and CFP (U87-FM- Gluc_{tag} -CFP). Fluorescent microscopy analysis for mCherry and CFP showed that these cells were nearly 100% transduced with the reporter vectors (Fig. 1b). We first demonstrated that all of the individual U87-FM- Gluc_{tag} -CFP cells expressed the corresponding tags by immunostaining using specific antibodies, except for Gluc_{Au1} that was not detectable by immunostaining attributed to anti-Au1 antibody properties (Fig. 1c). As a control we stained cells expressing $\text{Gluc}_{\text{Ctrl}}$ with the different tag antibodies, which did not show any staining (data not shown). We confirmed Gluc_{tag} expression by Western blot using anti-tag antibodies (Sup. Fig. 1a) and RT-PCR of the tag region (Sup. Fig. 1b and c). Next, we determined the effect of the tags on Gluc by measuring luciferase activity at different time points in an aliquot of conditioned medium from U87-FM cells expressing

SUBJECT AREAS:
MOLECULAR IMAGING
BIOLUMINESCENCE IMAGING
ASSAY SYSTEMS
REPORTER GENES

Received
25 July 2012

Accepted
3 December 2012

Published
10 January 2013

Correspondence and requests for materials should be addressed to T.W. (t.wurdinger@vumc.nl) or J.N. (Jonas.nilsson@onkologi.umu.se).




Figure 1 | Gluc_{tag} multiplex assay development and validation *in vitro*. (a) Schematic of the lentiviral vector constructs encoding the luciferase reporters. (b) Fluorescence microscopy analysis of a representative U87-FM-Gluc_{tag}-CFP cells (using Gluc_{Flag} reporter) showing mCherry levels (in red; marker for tumor cells) and CFP expression (in blue; marker for transduction efficiency). (c) Immunostaining against the various tags in U87-FM cells expressing different Gluc_{tag}. (d) Gluc activity in U87 cells expressing different Gluc_{tag} with respect to cell proliferation over time and cell number. The dashed line represents Gluc_{Ctrl} activity. (e) Gluc_{tag} immunobinding assay versus total Gluc activity using serial dilutions of the conditioned medium from a 1-day culture of U87-FM cell lines expressing the individual Gluc_{tag} reporters. (f-g) Equal mixture of ten cell lines each expressing a different Gluc_{tag} reporter. (f) Immunobinding assay of Gluc_{tag} from a mixed population. (g) Immunostaining for the ten different tags expression in the mixed population of U87-FM cell culture. (h-i) U87-FM cells expressing all ten different Gluc_{tag} reporters. (h) Immunostaining for the ten different tags in one cell line. The insert panels show staining of control parental cells. Percentages of positively stained cells are indicated. (i) Immunobinding assay of all ten Gluc_{tag} reporters in one cell line after TMZ treatment (black bars) compared to untreated control (white bars). (j) CFP fluorescence microscopy of cells in (h-i) with or without 600 μM TMZ. Size bar (in b,c, g and h) = 200 μm. Size bar (i) = 400 μm. N.d. = not detectable.



each Gluc_{tag} reporter construct using the Gluc substrate coelenterazine. We demonstrated that fusion of epitope tags to Gluc does not dramatically affect the expression, secretion or activity of Gluc (Fig. 1d and supplemental Fig. 1a–c). We then used the conditioned media containing the different Gluc_{tag} constructs to optimize antibody-tag immunobinding in a 96-well plate format (Methods).

Plates were first coated with monoclonal antibodies against the different tags and incubated with different amounts of conditioned medium allowing Gluc_{tag} immunobinding. Subsequently, the wells were washed to remove unbound Gluc_{tag} and subjected to the Gluc assay by direct addition of coelenterazine to the wells and measuring photon counts using a plate luminometer. As a control, an aliquot of




Figure 2 | Gluc_{tag} multiplex assay *in vivo*. (a) U87-FM cells expressing different Gluc_{tag} were implanted subcutaneously in nude mice ($n=3$) and tumor growth was monitored overtime. Shown are representative Fluc bioluminescence images of the U87-FM-Gluc_{tag}-CFP tumors expressing the Gluc_{Flag} reporter. (b), Quantitation of the different U87-FM tumors expressing different Gluc_{tag} reporters; Fluc activity (black), calliper measurement (green), total Gluc_{tag} activity in the blood (red), and bound Gluc_{tag} activity after immunobinding (blue). (c–d) U87-FM cells expressing the different Gluc_{tag} reporters were mixed in equal amounts and injected subcutaneously in the same mouse ($n=5$). Tumor growth was monitored using Fluc imaging and total Gluc blood assay as in (a–b). Individual cells subpopulation within the same tumor were monitored using the multiplex Gluc_{tag} immunobinding assay. (e) Immunostaining for the ten different tags expression in the mixed U87-FM-Gluc_{tag}-CFP tumor. Black arrows indicate positive staining for cells expressing the corresponding Gluc_{tag}. Size bar = 200 μm . N.d. = not detectable. (f–g) Mixed population of U87-FM cells expressing all ten different tags were implanted intracranially in the brain of nude mice ($n=5$). Tumor growth was monitored with Fluc imaging and Gluc total blood assay, and different tumor cell subpopulations were monitored using the multiplex Gluc_{tag} assay as in (c–d). Fluc bioluminescence imaging of a representative mouse is shown in (f). Data shown in (b,d, and g) as average relative light units (RLU) \pm standard deviation. Heatscales in (a, c and f) indicate relative photon levels (photons/sec) as measured by the CCD camera.



conditioned medium was assayed directly for total Gluc activity. We determined that six of the ten individual Gluc_{tag} constructs, i.e. Gluc_{Flag}, Gluc_{His}, Gluc_{HA}, Gluc_{AcV5}, Gluc_{V5}, Gluc_{Glu}, reported with high-efficiency after immunobinding in the antibody-coated wells, as compared to total Gluc_{tag} activity (Fig. 1e). The Gluc_{Myo}, Gluc_{Kt3}, Gluc_{Au1}, and Gluc_{E2} reporters showed lower activity upon immunobinding, attributed to suboptimal antibody-tag interactions. In order to demonstrate the use of the Gluc_{tag} system for multiplex applications, we mixed equal numbers of the ten different U87-FM-Gluc_{tag}-CFP cells and plated them in a single well. One, three and five days later, we measured the total Gluc level in aliquots of the conditioned medium. The levels of the individual Gluc_{tag} reporters in the mixed culture medium were analyzed by immunobinding with the different tag specific antibodies. We were able to monitor in real-time the growth of the individual U87-FM-Gluc_{tag}-CFP cell populations in the mixed culture (Fig. 1f). Immunostaining confirmed the expression of each tag in the corresponding cells in the mixed culture, again except for Gluc_{Au1} (Fig. 1g). In order to determine whether we can measure the activity of ten different Gluc_{tag} constructs in the same cell, we transduced U87-FM cells with all ten different lentiviral Gluc_{tag} reporters. Immunostaining of the U87-FM-Gluc_{10x tag}-CFP glioblastoma cells with tag specific antibodies showed that almost all cells expressed all ten different Gluc_{tag} constructs (Fig. 1h). The tag expression as demonstrated by immunostaining is less intense in Fig. 1h as compared to Fig. 1c, which we attribute to the transduction pool of ten different lentivectors used in Fig. 1h. We used these cells to monitor cell viability in the presence and absence of the alkylating agent temozolomide (TMZ), the standard chemotherapy for glioblastoma patients. Gluc_{tag} immunobinding assays were used to determine the expression of all ten reporters in the presence or absence of TMZ. We treated the U87-FM-Gluc_{10x tag}-CFP cells with TMZ and measured Gluc_{tag} activity after six days (Fig. 1i). Precipitation of all ten different Gluc_{tag} reporters showed a ~75% decrease in Gluc bioluminescence signal after TMZ treatment, similar to the total Gluc signal and confirmed by CFP immunofluorescence analysis (Fig. 1j). Altogether, these results demonstrate that we can measure biologically relevant differences using our multiplex Gluc_{tag} system.

In order to determine the *in vivo* applicability of the Gluc_{tag} reporter assay, we injected the ten individual U87-FM cell cultures expressing either one of the Gluc_{tag} or Gluc_{Ctrl} reporters subcutaneously in nude mice for singleplex application. In another set, we injected mice with PBS as a negative control (12 groups; n = 3/group). Tumor growth was monitored over time by calliper measurement, and Fluc bioluminescence imaging (Fig. 2a,b). Blood was collected from mice at different time points and 5 µl of whole blood (optimum amount for Gluc blood assay¹⁶) was assayed for total Gluc activity. At the same time, 25 µl of blood was analyzed for Gluc activity after immunobinding on wells coated with the corresponding tag antibody (Fig. 2b). All of the Gluc_{tag} reporters, except Gluc_{E2}, allowed blood monitoring of tumor growth over time in a singleplex assay. We compared the tumor growth as measured by calliper, Fluc imaging and Gluc blood assay to signals obtained from the ten different Gluc_{tag} reporters after immunobinding (Fig. 2b). The six Gluc_{tag} reporters Gluc_{Flag}, Gluc_{His}, Gluc_{HA}, Gluc_{AcV5}, Gluc_{V5}, Gluc_{Glu} were more sensitive in detecting tumor growth than the others, possibly due to a higher affinity of the antibodies to these tags. To demonstrate the use of the Gluc_{tag} system for multiplex applications, we mixed equal numbers of U87-FM cells expressing the ten different Gluc_{tag} and implanted the heterogeneous cell pool subcutaneously in nude mice (n = 5). We monitored tumor growth by Fluc bioluminescence imaging and total Gluc blood assay (Fig. 2c,d). Importantly, we were able to efficiently monitor the growth of the individual subpopulation of the U87-FM cells expressing the different high-efficiency Gluc_{tag} reporters (Gluc_{Flag}, Gluc_{His}, Gluc_{HA}, Gluc_{AcV5}, Gluc_{V5}, Gluc_{Glu}) in the same tumor over time

(Fig. 2d). We confirmed the expression of all tags (except for Gluc_{Au1}) in the same tumor by immunostaining (Fig. 2e). To determine the applicability of the Gluc_{tag} multiplex system in deep tissues, the mixture of the ten different U87-FM-Gluc_{tag}-CFP cells were implanted orthotopically in the brain of nude mice (n = 5). We monitored tumor growth by Fluc bioluminescence imaging and total Gluc blood assay (Fig. 2f,g). Again, we were able to monitor the intracranial growth of the individual subpopulations of U87-FM cells expressing the high-efficiency Gluc_{tag} reporters within the same brain tumor in real-time, by blood sampling and the application of the Gluc_{tag} multiplex reporter system (Fig. 2g).

Discussion

Here we demonstrated the functionality of the Gluc_{tag} multiplex reporter system by the parallel monitoring of mixed cell cultures *in vitro*, and of subpopulations in mixed subcutaneous and intracranial tumors *ex vivo*. The individual epitope tags could also be employed to localize expression by immunostaining. We used ten different Gluc_{tag} reporters, but this tag library can be readily extended allowing for increased multiplexing. In addition, antibody binding of the low-efficiency tags in the Gluc_{tag} immunobinding assay might be further optimized to achieve higher sensitivity since these tags, except for Gluc_{Au1}, were efficiently detected by immunostaining. Real-time monitoring of individual cells in a heterogeneous mixture may allow, for instance, multiplexed RNAi screening or measurement of drug responses of multiple cell populations in parallel. The Gluc_{tag} multiplex system might be extended to monitor different variables in a single cell type. For instance, engineering each individual Gluc_{tag} under the control of different transcription response elements for multiplex transcription factor activity measurements⁷, constructing microRNA (miRNA)-binding sequences in the 3'-UTR of the Gluc_{tag} constructs for multiplex monitoring of miRNA activity¹⁰ or by inserting different protease cleavage sites into the Gluc_{tag} gene for multiplex protease activity measurement¹⁷. This reporter assay provides a versatile tool to study complex processes with different variables in systems biology.

Methods

Cells. For *in vitro* and *in vivo* validation, we used the glioblastoma cell line U87, stably co-expressing Firefly luciferase and mCherry fluorescent protein. Cell lines were maintained in DMEM high glucose complemented with sodium pyruvate, stable glutamine, 10% FBS and pen/strep (all PAA), incubated under standard cell culture conditions of 37°C and 5% CO₂.

Lentivirus vector construction and transduction. First, we amplified Gluc by PCR using Accuprime Pfx DNA polymerase (Life Technologies) from CSCW-Gluc-IRES-CFP⁶ with specific primers that incorporate an XbaI site downstream of the Gluc cDNA (Forward primer: 5'-GCGTGTACGGTGGGAGGTCT-3', reverse primer: 5'-TCTCGAGTAGAGATCTGTACCACCGCCCTT-3'). The reverse primer also covers the original stop-codon that is thereby removed. The Gluc cDNA, now containing a unique XbaI site, was ligated back into the CSCW-Gluc-IRES-CFP vector with T4 DNA ligase (Life Technologies), after removal of the wild-type Gluc using NheI and XhoI (Life Technologies). Plasmid DNA was transformed in XL-10 Gold ultracompetent cells (Agilent Technologies), cultured overnight in LB agar containing 50 µg/ml Ampicillin. We isolated DNA using a DNA plasmid mini kit (Qiagen) and verified successful transformation by XbaI restriction analysis. The Gluc construct was then digested with XbaI and XhoI to insert the different epitope tags. The epitope tags were designed with an XbaI site upstream, a stop codon and XhoI site downstream. A total of 20 µM of both single strand oligonucleotides of the acquired tag DNA (Life Technologies, see Table S1 for sequences) was annealed in annealing buffer (100 mM Tris-HCl pH 7.5, 1 M NaCl and 10 mM EDTA) by heating to 65°C for 10 minutes and slow cooling to room temperature. The epitope tag was then inserted into the vector using T4 DNA ligase and transformed in XL-10 Gold ultracompetent cells. Bacteria were cultured and DNA was isolated. We verified the Gluc_{tag} constructs by sequencing using BigDye Terminator v3.1 Cycle Sequencing kit (Life Technologies, primer: 5'-GCCAACGTGCGAGTGTTC-3'). The Gluc_{tag} construct was co-transfected with a third generation lentiviral packaging mix (pMDLg/pRRE, pRSV-Rev and pMD2.G, Addgene) in HEK293T cells using Lipofectamine 2000 (Life Technologies). Virus was harvested two and three days after transfection and cell debris was spun down for five minutes at 1,000 × g. U87 cells were transduced overnight with lentivirus using a multiplicity of infection of 100 transducing units per cell in the presence of 8 µg/ml polybrene in standard culture conditions.



Fluorescence microscopy. Successful transduction was verified by visualizing CFP (co-expressed with the Gluc_{tag} construct) and mCherry fluorescent protein (co-expressed with Fluc) using fluorescent microscopy (Leica).

Immunostaining and antibodies. For *in vitro* immunostaining, cells were fixed in 3.7% formaldehyde for 20 minutes, washed with PBS and permeabilized with PBS containing 0.1% Triton X-100. Cells were then washed three times in PBS containing 5% FBS and blocked with PBS containing 5% FBS. Primary mouse-anti-tag monoclonal antibody (10 µg/ml in 100 µl of PBS, Table S1) was added and incubated for one hour at room temperature. Cells were washed again and incubated with 1 : 100 goat-anti-mouse horseradish peroxidase (HRP) conjugate (Dako) for one hour. After washing, cells were stained with DAB+ (Life Technologies). For mouse tissue staining, tumors were removed and embedded in paraffin. Microtome sections of 5 µm on glass slides were deparaffinized in xylene and rehydrated in ethanol series of 100%, 96%, and 70% ethanol. Endogenous peroxidase was blocked with 0.3% H₂O₂ in methanol for 30 minutes. After rinsing with water, antigens were retrieved with citrate buffer (pH 6) with 0.05% Tween 20 using a microwave (Bosch, five minutes 900 watt, 10 minutes 360 watt). After slowly cooling, tissues were washed three times with PBS and incubated with primary antibody (10 µg/ml), for one hour at room temperature. After washing three times again, tissues were incubated with REAL Envision Rabbit anti-mouse (Dako and REAL DAB+ stained (Dako). Tissue sections were dehydrated with ethanol series as before and fixed in xylene. Cells and sections were imaged and photographed by light microscopy (Leica).

Western blot. U87-FM-Gluc_{tag}-CFP cells were lysed in RIPA buffer and mixed with Laemli buffer containing β-mecaptoethanol (80 µl/ml). The samples were separated using a 10% SDS-PAGE gel and the XCell-II blot module (Invitrogen). Primary antibodies (Table S1) were used in a 1 : 3,000 dilutions and goat-anti-mouse HRP immunoglobuline (Dako) secondary antibody was used in a 1 : 2,000 dilution. Incubation times were 1 hour at room temperature.

RT-PCR. U87-FM-Gluc_{tag}-CFP RNA was isolated using Trizol (Invitrogen) according to manufacturer's guidelines. cDNA was generated using Omniscript RT kit (Qiagen) and PCR was performed in triplo using FastStart SYBR Green Master (Rox) mix (Roche). As a forward primer we used 5'-GCCAACGTGCAGTGTTC-3', which is located in the Gluc gene. We used the tag antisense oligonucleotides as tag specific reverse primers (Table S1).

In vitro gluc activity assay. For Gluc_{tag} activity measurement over time, 50,000 cells were plated in a 24-well plate and incubated overnight. A total of 10 µl conditioned medium was harvested from cells and Gluc activity was measured by adding 50 µl (5 µg/ml) coelenterazine (Nanolight Technologies) in PBS and 0.1% Triton X-100. Before addition to the sample, the substrate was incubated at room temperature for 30 minutes for stabilization. Photon counts were determined over 10 seconds in a luminometer (Berthold Technologies). For the measurement of Gluc_{tag} activity with respect to cell number, indicated numbers of cells were plated in a 24-well plate and incubated overnight. Gluc_{tag} activity in medium was determined as described above.

Gluc_{tag} immunobinding assay. White goat-anti-mouse-coated 96-well plates (Thermo Scientific) were washed three times with PBS containing 0.05% Tween 20 and incubated with 50 µl (10 µg/ml) of each of the mouse monoclonal antibody directed against the specific epitope tags (Table S1). Incubation time was two hours while centrifuging at 500 × g at 4°C. Wells were washed and blocked three times with PBS containing 5% FBS and 0.05% Tween 20 on a plate shaker at 65–75 rpm. A total of 30 µl of Gluc_{tag} conditioned medium or Gluc_{tag} mouse blood was added to the well and incubated for two hours at room temperature on a microplate shaker at 65–75 rpm. Wells were washed five times for five minutes on a plate shaker at 65–75 rpm. A total of 50 µl Gluc substrate (5 µg/ml coelenterazine in PBS and 0.1% Triton X-100) was added to the well and photon counts were determined in a microplate luminometer (Tecan) at 0.1 second per well. For the comparison of the bound Gluc to the total Gluc activity, aliquots of conditioned medium or 5 µl mouse blood were transferred to a white 96-wells microplate and assayed using 50 µl (5 µg/ml) coelenterazine in PBS and 0.1% Triton X-100.

In vivo experiments. All *in vivo* experiments were subject to ethical committee approval and are in accordance with VU University Medical Center and national regulations. For subcutaneous tumor xenograft model, six weeks old athymic Nude-Foxn1tm mice (Harlan) were implanted with 5 × 10⁵ U87-FM-Gluc_{tag}-CFP cells in 50 µl DMEM and 50 µl Matrigel (BD Biosciences). For intracranial tumor implantation, a stereotactic frame (Harvard Biosciences) was used to inject cells vertically into the right hemisphere. General anesthesia was induced by subcutaneous injection of buprenorphine hydrochloride (0.1 mg/kg). The mice were further anesthetized with oxygen containing 2.5% isoflurane. After skin incision, lidocaine drops (5 mg/ml in PBS) were administered topically. A small drill was used to drill a hole into the skull. Coordinates used for injection were X = 0.5 mm, Y = 2 mm, Z = -2 mm from the bregma. A total of 2 × 10⁵ cells in 5 µl of DMEM was injected vertically. On indicated days after injection, tumor size was monitored using a calliper

(for subcutaneous) and bioluminescence *in vivo* Fluc imaging, by injecting D-luciferin (100 mg/kg) intraperitoneally. Imaging was performed with an IVIS CCD camera and analyzed with Living Image software (Caliper Life Sciences). We collected 300 µl mouse blood aliquots from the tail vein in Microvette CB300 EDTA capillary tubes (Sarstedt) for *ex vivo* whole Gluc measurements and Gluc immunobinding assays.

- Gross, S. & Piwnica-Worms, D. Spying on cancer: molecular imaging in vivo with genetically encoded reporters. *Cancer cell* **7**, 5–15 (2005).
- Pittet, M. J. & Weissleder, R. Intravital Imaging. *Cell* **147**, 983–991 (2011).
- Contag, C. H. & Ross, B. D. It's not just about anatomy: in vivo bioluminescence imaging as an eyepiece into biology. *Journal of magnetic resonance imaging: JMIR* **16**, 378–87 (2002).
- Bhaumik, S. & Gambhir, S. S. Optical imaging of Renilla luciferase reporter gene expression in living mice. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 377–82 (2002).
- Tannous, B. A., Kim, D.-E., Fernandez, J. L., Weissleder, R. & Breakefield, X. O. Codon-optimized Gaussia luciferase cDNA for mammalian gene expression in culture and in vivo. *Molecular therapy: the journal of the American Society of Gene Therapy* **11**, 435–43 (2005).
- Wurdinger, T. *et al.* A secreted luciferase for *ex vivo* monitoring of in vivo processes. *Nature methods* **5**, 171–3 (2008).
- Bhang, H.-eun C., Gabrielson, K. L., Laterra, J., Fisher, P. B. & Pomper, M. G. Tumor-specific imaging through progression elevated gene-3 promoter-driven gene expression. *Nature medicine* **17**, 123–9 (2011).
- Badr, C. E., Hewett, J. W., Breakefield, X. O. & Tannous, B. A. A highly sensitive assay for monitoring the secretory pathway and ER stress. *PLoS one* **2**, e571 (2007).
- Lee, J. Y. *et al.* Development of a dual-luciferase reporter system for in vivo visualization of MicroRNA biogenesis and posttranscriptional regulation. *Journal of nuclear medicine: official publication, Society of Nuclear Medicine* **49**, 285–94 (2008).
- Tian, W. *et al.* High-Throughput Functional MicroRNAs Profiling by Recombinant AAV-Based MicroRNA Sensor Arrays. *PLoS one* **7**, e29551 (2012).
- McMillin, D. W. *et al.* Tumor cell-specific bioluminescence platform to identify stroma-induced changes to anticancer drug activity. *Nature medicine* **16**, 483–9 (2010).
- Badr, C. E. & Tannous, B. A. Bioluminescence imaging: progress and applications. *Trends in Biotechnology* **29**, 624–33 (2011).
- Shaner, N. C., Steinbach, P. A. & Tsien, R. Y. A guide to choosing fluorescent proteins. *Nature methods* **2**, 905–9 (2005).
- Filonov, G. S. *et al.* Bright and stable near-infrared fluorescent protein for in vivo imaging. *Nature biotechnology* **29**, 757–61 (2011).
- Deliolanis, N. C. *et al.* In vivo tomographic imaging of red-shifted fluorescent proteins. *Biomedical optics express* **2**, 887–900 (2011).
- Tannous, B. A. Gaussia luciferase reporter assay for monitoring biological processes in culture and in vivo. *Nature protocols* **4**, 582–91 (2009).
- Niers, J. M., Kerami, M., Pike, L., Lewandrowski, G. & Tannous, B. A. Multimodal in vivo imaging and blood monitoring of intrinsic and extrinsic apoptosis. *Molecular therapy: the journal of the American Society of Gene Therapy* **19**, 1090–6 (2011).

Acknowledgements

This work was supported partly by grants from NWO-VIDI (TW), and NIH/NINDS P30NS045776 and R01NS064983 (BAT). SVR and JN are supported by CCA/VICI Fellowships. We are thankful to Priscilla Jainandunsing, Laura Jonkman, Stephanie van Hoppe, Tonny Lagerweij and Laurine Wedekind for technical support.

Author contributions

JN and TW conceived this study and SVR, BAT, JN and TW designed the experimental setup, with help of DN and WPV. SVR performed most of the experiments. SVR, JN, and TW wrote the manuscript, which was edited and reviewed by all the authors.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: The authors declare no competing financial interests.

License: This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

How to cite this article: van Rijn, S. *et al.* Functional multiplex reporter assay using tagged Gaussia luciferase. *Sci. Rep.* **3**, 1046; DOI:10.1038/srep01046 (2013).