



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

The roles of long noncoding RNA in epigenetic regulation

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Lee, Jeannie T. 2013. The roles of long noncoding RNA in epigenetic regulation. <i>Epigenetics & Chromatin</i> 6(Suppl 1): O31.
Published Version	doi:10.1186/1756-8935-6-S1-O31
Accessed	February 19, 2015 12:01:09 PM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:10609669
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

ORAL PRESENTATION

Open Access

The roles of long noncoding RNA in epigenetic regulation

Jeannie T Lee^{1,2,3}

From *Epigenetics and Chromatin: Interactions and processes*
Boston, MA, USA. 11-13 March 2013

The X-linked region now known as the 'X-inactivation center' (Xic) was once dominated by protein-coding genes but, with the rise of Eutherian mammals some 150-200 million years ago, became infiltrated by genes that produce long noncoding RNA (lncRNA). Some of the noncoding genes have been shown to play crucial roles during X-chromosome inactivation (XCI), including the targeting of chromatin modifiers to the X. The rapid establishment of lncRNA hints at a possible preference for long transcripts in some aspects of epigenetic regulation. We will consider advantages lncRNA offers in delivering allelic, cis-limited, and locus-specific control. Unlike proteins and small RNAs, long ncRNAs are tethered to the site of transcription, and can therefore tag the allele of origin and mark unique regions in a complex genome. We will discuss general roles of lncRNA, with special consideration of XCI and developmental consequences when Xist RNA is aberrantly expressed. Like small RNAs, lncRNAs may emerge as powerful regulators of the epigenome.

Author details

¹Howard Hughes Medical Institute, USA. ²Department of Molecular Biology, Massachusetts General Hospital, USA. ³Department of Genetics, Harvard Medical School, Boston, MA 02114, USA.

Published: 18 March 2013

doi:10.1186/1756-8935-6-S1-O31

Cite this article as: Lee: The roles of long noncoding RNA in epigenetic regulation. *Epigenetics & Chromatin* 2013 **6**(Suppl 1):O31.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Howard Hughes Medical Institute, USA
Full list of author information is available at the end of the article