



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

S cerevisiae Genome as a Confined Equilibrium Polymer Brush

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Goloborodko, Anton, Jon Matthew Belton, Geoffrey Fudenberg, Maxim Imakaev, Job Dekker, and Leonid Alex Mirny. 2013. S cerevisiae genome as a confined equilibrium polymer brush. Epigenetics & Chromatin 6(Suppl 1): P129.
Published Version	doi:10.1186/1756-8935-6-S1-P129
Accessed	February 19, 2015 12:05:26 PM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:11744410
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

POSTER PRESENTATION

Open Access

S cerevisiae genome as a confined equilibrium polymer brush

Anton Goloborodko^{1*}, Jon Matthew Belton², Geoffrey Fudenberg³, Maxim Imakaev¹, Job Dekker², Leonid Mirny^{1,4}

From *Epigenetics & Chromatin: Interactions and processes*
Boston, MA, USA. 11-13 March 2013

Introduction

A series of recent studies using optical and 3C-based experimental approaches [1,2] have shown that at a global level the yeast genome assumes a Rabl-like conformation with the chromosomal centromeres tethered to the spindle pole body and telomeres anchored on the nuclear periphery. This is strikingly different from human genomic organization, where chromosomes have been shown to assume fractal globule conformations with domains of active and inactive chromatin [3].

In this work, we study Rabl-like chromosomal organization using a computationally efficient polymer lattice model.

Results

We find that our lattice polymer model predicts experimentally observed Hi-C contact maps with high precision ($r \sim 0.87$). This demonstrates that a minimal equilibrium model can reconstitute the majority of the observed interaction patterns. Our model predicts that loci preferentially localize to different regions of the nucleus in 3D, depending on their genomic positions. In turn, this leads to different contact preferences between loci. This effect is most clearly demonstrated by the cross-like patterns of interactions between peri-centromeric regions, as observed in yeast Hi-C data. Centromere colocalization and excluded volume interactions cause chromatin fibers to extend away from the spindle pole body; this can be understood as a 'polymer brush' effect. In turn, the spatial localization of peri-centromeric loci becomes mainly determined by their genomic distances to the centromere; each locus preferentially forms contacts with other loci at similar distances from their respective centromeres. This dependence of spatial localization on genomic coordinates produces

prominent cross-like patterns, as observed in *S. cerevisiae* Hi-C maps. We estimate this polymer brush effect to alter contact preferences at distances up to ~ 200 kbp from each centromere. Finally, our lattice model allows us to quickly predict the relative impact of fiber width, flexibility and linear compaction on chromosomal conformation. We find that the observed Hi-C maps are consistent with a range of fiber parameters.

Author details

¹Department of Physics, Massachusetts Institute of Technology, Cambridge, 02143, MA, USA. ²Department of Biochemistry and Molecular Pharmacology, University of Massachusetts, Worcester, 01605, MA, USA. ³Graduate Program in Biophysics, Harvard University, Cambridge, 02115, MA, USA. ⁴Division of Health Science and Technology, Harvard-MIT, Cambridge, MA, 02139, USA.

Published: 8 April 2013

References

1. Duan Z, Andronescu M, Schutz K, McIlwain S, Kim YJ, Lee C, Shendure J, Fields S, Blau CA, Noble WS: **A three-dimensional model of the yeast genome.** *Nature* 2010, **465**(7296):363-367.
2. Therizols P, Duong T, Dujon B, Zimmer C, Fabre E: **Chromosome arm length and nuclear constraints determine the dynamic relationship of yeast subtelomeres.** *Proc Natl Acad Sci* 2010, **107**(5):2025-2030.
3. Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragoczy T, Telling A, Amit I, Lajoie BR, Sabo PJ, Dorschner MO, Sandstrom R, Bernstein B, Bender MA, Groudine M, Gnirke A, Stamatoyannopoulos J, Mirny LA, Lander ES, Dekker J: **Comprehensive Mapping of Long-Range Interactions Reveals Folding Principles of the Human Genome.** *Science* 2009, **326**:289.

doi:10.1186/1756-8935-6-S1-P129

Cite this article as: Goloborodko et al.: *S cerevisiae* genome as a confined equilibrium polymer brush. *Epigenetics & Chromatin* 2013 **6**(Suppl 1):P129.

¹Department of Physics, Massachusetts Institute of Technology, Cambridge, 02143, MA, USA

Full list of author information is available at the end of the article