



# DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

## Hemoglobin A1c Is Associated With Increased Risk of Incident Coronary Heart Disease Among Apparently Healthy, Nondiabetic Men and Women

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters.

<b>Citation</b>	Pai, Jennifer K., Leah E. Cahill, Frank B. Hu, Kathryn M. Rexrode, JoAnn E. Manson, and Eric B. Rimm. 2013. Hemoglobin a1c is associated with increased risk of incident coronary heart disease among apparently healthy, nondiabetic men and women. <i>Journal of the American Heart Association: Cardiovascular and Cerebrovascular Disease</i> 2(2): e000077.
<b>Published Version</b>	<a href="https://doi.org/10.1161/JAHA.112.000077">doi:10.1161/JAHA.112.000077</a>
<b>Accessed</b>	February 19, 2015 12:07:05 PM EST
<b>Citable Link</b>	<a href="http://nrs.harvard.edu/urn-3:HUL.InstRepos:11363030">http://nrs.harvard.edu/urn-3:HUL.InstRepos:11363030</a>
<b>Terms of Use</b>	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <a href="http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA">http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA</a>

*(Article begins on next page)*

# Hemoglobin A<sub>1c</sub> Is Associated With Increased Risk of Incident Coronary Heart Disease Among Apparently Healthy, Nondiabetic Men and Women

Jennifer K. Pai, ScD, MHS; Leah E. Cahill, PhD; Frank B. Hu, MD, PhD; Kathryn M. Rexrode, MD, MPH; JoAnn E. Manson, MD, DrPH; Eric B. Rimm, ScD

**Background**—Hemoglobin A<sub>1c</sub> (HbA<sub>1c</sub>), a time-integrated marker of glycemic control, predicts risk of coronary heart disease (CHD) among diabetics. Few studies have examined HbA<sub>1c</sub> and risk of CHD among women and men without clinically elevated levels or previously diagnosed diabetes.

**Methods and Results**—We conducted parallel nested case–control studies among women (Nurses' Health Study) and men (Health Professionals Follow-up Study). During 14 and 10 years of follow-up, 468 women and 454 men developed incident nonfatal myocardial infarction (MI) and fatal CHD. Controls were matched 2:1 based on age, smoking, and date of blood draw. For these analyses, participants with a history of diabetes or HbA<sub>1c</sub> levels  $\geq 6.5\%$  at baseline were excluded. Compared with HbA<sub>1c</sub> of 5.0% to  $<5.5\%$ , those with an HbA<sub>1c</sub> of 6.0% to  $<6.5\%$  had a multivariable-adjusted relative risk (RR) of CHD of 1.90 (95% CI 1.11 to 3.25) in women and 1.81 (95% CI 1.09 to 3.03) in men. The pooled RR of CHD was 1.29 (95% CI 1.11 to 1.50) for every 0.5%-increment increase in HbA<sub>1c</sub> levels and 1.67 (95% CI 1.23 to 2.25) for every 1%-increment increase, with the risk plateauing around 5.0%. Furthermore, participants with HbA<sub>1c</sub> levels between 6.0% and  $<6.5\%$  and C-reactive protein levels  $>3.0$  mg/L had a 2.5-fold higher risk of CHD compared with participants in the lowest categories of both biomarkers.

**Conclusions**—Our findings suggest that HbA<sub>1c</sub> is associated with CHD risk among apparently healthy, nondiabetic women and men and may be an important early clinical marker of disease risk. (*J Am Heart Assoc.* 2013;2:e000077 doi: 10.1161/JAHA.112.000077)

**Key Words:** coronary heart disease • C-reactive protein • epidemiology • glycosylated hemoglobin • inflammation

Diabetes is a well-known risk factor for cardiovascular disease (CVD),<sup>1</sup> and although the exact mechanisms are unknown, angiopathy from dysglycemia is likely a primary cause. Glycated hemoglobin, assessed clinically by hemoglobin A<sub>1c</sub> (HbA<sub>1c</sub>), is a time-integrated marker of average blood glucose concentration increasingly used in the screening for

and management of diabetes, and it is more closely related to the risk of complications than are single or episodic measures of glucose levels.<sup>2–4</sup> Despite the clinical utility of HbA<sub>1c</sub>, several recent large randomized trials of intensive HbA<sub>1c</sub> lowering among high-risk diabetics have not demonstrated lower risk of macrovascular events.<sup>5–8</sup>

Currently, the degree to which mild elevations of HbA<sub>1c</sub> not in the diabetic range are associated with risk of coronary heart disease (CHD) is unclear.

The association between dysglycemia and risk of CVD may start at levels that are only modestly elevated, well below the glucose threshold for diabetes.<sup>9,10</sup> The majority of studies conducted among nondiabetics have included participants with HbA<sub>1c</sub>  $\geq 6.5\%$ ,<sup>11,12</sup> the threshold set for diabetes diagnosis by the American Diabetes Association<sup>2</sup> and World Health Organization.<sup>4</sup> In several studies, the association between HbA<sub>1c</sub> and CHD is attenuated and no longer significant after adjustment for well-known cardiovascular risk factors.<sup>11,13</sup> Thus, it remains unclear whether HbA<sub>1c</sub> is a clinically relevant marker of CHD risk in apparently healthy, nondiabetic populations, or if it is a proxy for dyslipidemia, hyperinsulinemia, or other metabolic pathways. In the present

From the Departments of Epidemiology (J.K.P., F.B.H., J.E.M., E.B.R.) and Nutrition (L.E.C., F.B.H., E.B.R.), Harvard School of Public Health, Boston, MA; and Division of Preventive Medicine (K.M.R., J.E.M.) and Channing Division of Network Medicine (J.K.P., F.B.H., E.B.R.), Department of Medicine, Brigham and Women's Hospital, and Harvard Medical School, Boston, MA.

An accompanying Table S1 is available at <http://jaha.ahajournals.org/content/2/2/e000077.full>

**Correspondence to:** Eric B. Rimm, ScD, Channing Division of Network Medicine, Brigham and Women's Hospital and Harvard Medical School, 181 Longwood Ave, 3rd Floor, Boston, MA 02115. E-mail: Eric.Rimm@channing.harvard.edu

Received January 4, 2013; accepted February 6, 2013.

© 2013 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley-Blackwell. This is an Open Access article under the terms of the Creative Commons Attribution Noncommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

study, we examined the association between HbA<sub>1c</sub> and risk of incident CHD in 2 prospective nested case-control studies of health professionals without diabetes or without HbA<sub>1c</sub> concentrations  $\geq 6.5\%$ .

## Methods

### Study Population

The Nurses' Health Study (NHS) and the Health Professionals Follow-Up Study (HPFS) are prospective cohort investigations among 121 700 female US registered nurses aged 30 to 55 years at baseline in 1976 (NHS) and 51 529 male US health professionals aged 40 to 75 years at baseline in 1986 (HPFS). Information about health and disease is assessed biennially, and information about diet is assessed every 4 years via self-administered questionnaires.<sup>14,15</sup> Between 1989 and 1990, a blood sample was requested from all participants in the NHS, and it was returned from 32 826 women. Similarly, between 1993 and 1995, a blood sample was returned from 18 225 men in the HPFS. Participants who provided blood samples were similar to those who did not, albeit somewhat younger. In the NHS, among women without CVD or cancer before 1990, we identified 468 women with incident nonfatal myocardial infarction (MI) or fatal CHD between the date of blood draw and June 2004. In the HPFS, we identified 454 men with incident nonfatal MI or fatal CHD between the date of blood draw and return of the 2004 questionnaire. Using risk-set sampling,<sup>16</sup> controls were randomly selected on a 2:1 basis matched for age, smoking, date of blood draw, and fasting status (women only) from participants free of CVD at the time the case was diagnosed. To assess men and women without diabetes, participants with a history of diabetes or HbA<sub>1c</sub> levels  $\geq 6.5\%$  at baseline (153 women; 123 men) were excluded from further analyses. We used 6.5% as the cutoff because this value has been recommended as a level defining diabetes by the American Diabetes Association<sup>2</sup> and the World Health Organization<sup>4</sup> and is the level at which complications of diabetes such as retinopathy arise.

The study protocol was approved by the Institutional Review Board of the Brigham and Women's Hospital and by the Harvard School of Public Health Human Subjects Committee Review Board.

### Assessment of CHD

MI was confirmed by study physicians blinded to participants' exposure status if it met World Health Organization criteria (symptoms plus either diagnostic electrocardiographic changes or elevated levels of cardiac enzymes).<sup>17,18</sup> Deaths were identified from state vital records and the National

Death Index or reported by the participant's next of kin or the postal system. Fatal CHD was confirmed by use of hospital records or on autopsy or if CHD was listed as the cause of death on the death certificate, if it was the underlying and most plausible cause, and if evidence of previous CHD was available.

### Assessment of Other Factors

Anthropometric data, lifestyle behavior, and diet were derived from the questionnaire administered in 1990 to women and 1994 to men, with missing information substituted from previous questionnaires. Body mass index (BMI) was calculated as body weight divided by height squared. Average nutrient intake was computed using the semiquantitative food frequency questionnaire. Physical activity was expressed as metabolic equivalent-hours. The questionnaires and the validity and reproducibility of measurements have been previously reported.<sup>15,19</sup>

### Measurement of Biochemical Variables

Blood samples were collected in liquid sodium heparin blood tubes for women and EDTA tubes for men, placed on ice packs, stored in polystyrene foam containers, returned to our laboratory via overnight courier, and centrifuged and aliquoted for storage in liquid nitrogen freezers ( $-130^{\circ}\text{C}$  or colder). The first nested case-control set included cases and controls selected from 1990 to 1998 for NHS and 1994 to 2000 for HPFS and were previously documented.<sup>20</sup> These samples were assayed in 2002. Since then, additional nested case-control sets with follow-up through 2004 have been completed for both NHS and HPFS and were sent for assaying using similar methods in 2007. All analyses were performed in a laboratory certified by the Centers for Disease Control and Prevention/National Heart, Lung, and Blood Institute Lipid Standardization Program with commercially available analytic systems.

HbA<sub>1c</sub> was measured by turbidimetric immunoinhibition using packed red cells (Roche Diagnostics), which is a standard approved by the US National Glycohemoglobin Standardization Program and the Food and Drug Administration for clinical use, and traceable to the Diabetes Controls and Complications Trial. The day-to-day variability at HbA<sub>1c</sub> values of 5.5 and 9.1 were 1.9% and 3.0%, respectively. In a substudy of 83 participants measured 3 years apart, HbA<sub>1c</sub> was highly correlated between draws (Spearman=0.88) and had excellent intraclass correlations (0.73). C-reactive protein (CRP) concentrations were determined with an immunoturbidimetric high-sensitivity assay using reagents and calibrators from Denka Seiken with assay day-to-day variability between 1% and 2% and were previously shown to be largely unaffected by transport conditions and to be reproducible

within persons over time.<sup>21,22</sup> Total, high-density lipoprotein (HDL), and directly assayed low-density lipoprotein cholesterol and triglycerides were measured using standard methods with reagents from Roche Diagnostics and Genzyme. Study samples were sent to the laboratory for analysis in randomly ordered batches, and the laboratory was blinded to case-control status. The coefficients of variation for all the assays were <10%.

## Statistical Analyses

We analyzed the 2 cohorts separately. HbA<sub>1c</sub> levels were categorized into quintiles based on the sex-specific distributions among the controls as well as according to categories (<5.0%, 5.0% to <5.5%, 5.5% to <6.0%, and 6.0% to <6.5%). In the categorical analyses, the HbA<sub>1c</sub> 5.0% to <5.5% category was used as the reference because there were more participants than in the <5.0% category. Because of this study design, the odds ratio derived from the logistic regression directly estimates the incidence rate (hazard) ratio and, thus, the RR.<sup>16,23</sup> We analyzed the association between HbA<sub>1c</sub> levels and risk of CHD using both conditional and unconditional logistic regression with adjustment for matching factors and laboratory batch. Because both analyses provided essentially the same results, we present unconditional logistic regression, which parallels the subgroup analyses. In our multivariable model, we further adjusted for parental history of MI before the age of 60 (yes/no), alcohol intake (nondrinker; 0.1 to 4.9, 5.0 to 14.9, 15.0 to 29.9, or  $\geq 30.0$  g/d; or missing), physical activity (quintiles), total cholesterol:HDL cholesterol ratio (quintiles), BMI in categories (<20, 20 to 24.9, 25 to 29.9, 30 to 34.9, and  $\geq 35$  kg/m<sup>2</sup>), and postmenopausal hormone therapy (yes/no, women only). Additional adjustment for hypertension (yes/no) at baseline, triglyceride levels (quintiles), and CRP levels (quintiles) was also conducted. Correlation coefficients were calculated using age-adjusted Spearman partial correlation coefficients. To test for linear trend, we used the median HbA<sub>1c</sub> levels in the control categories as a continuous variable. In addition, we conducted analyses per 0.5%-increment increase in HbA<sub>1c</sub> levels. We examined the possibly nonlinear relation between HbA<sub>1c</sub> and CHD nonparametrically with restricted cubic splines.<sup>24</sup> Tests for nonlinearity used the likelihood ratio test, comparing the model with only the linear term to the model with the linear and the cubic spline terms. To pool the RR estimates for men and women, we used the weighted average of estimates using the DerSimonian and Laird random effects model.<sup>25</sup> Because we are using a single measure of HbA<sub>1c</sub> to predict future disease over long-term follow-up, fluctuations in baseline biological risk factors over time may lead to an underestimation of the true association in the form of “regression dilution bias.”<sup>26,27</sup> To account for potential changes in our baseline

measure over time, we also present RR estimates corrected for potential regression dilution bias using the model described by Clarke et al.<sup>28</sup> In brief, the  $\beta$  coefficients and standard errors for the entire cohort were divided by the regression dilution ratio, based on the within-person correlation coefficient calculated from the subset of participants with repeated measurements, to obtain the corrected risk estimates for the usual relation of HbA<sub>1c</sub> levels and CHD.

All *P* values presented are 2-tailed, and *P* values <0.05 were considered statistically significant. All analyses were performed using SAS version 9 (SAS Institute).

## Results

Cardiovascular risk factors were higher or more frequent among the cases compared with the controls in both men and women (Table 1). As expected, cases were more likely to have parental history of early MI, history of hypercholesterolemia, and history of hypertension. Also, cases were more likely to have higher CRP levels and adverse lipid profiles compared with controls for both women and men. Age-adjusted Spearman partial correlation coefficients are presented among controls for women and men (Table 2). HbA<sub>1c</sub> percentage was positively correlated with CRP levels, triglycerides:HDL cholesterol ratio, triglycerides, and BMI and inversely correlated with HDL cholesterol and alcohol intake.

Among participants with HbA<sub>1c</sub> levels <6.5% and without history of diabetes at baseline, greater HbA<sub>1c</sub> levels were significantly associated with an increased risk of CHD in both women and men. Comparing extreme quintiles of HbA<sub>1c</sub> levels, the multivariable-adjusted RR of CHD was 1.70 (95% CI 1.22 to 2.36) in the pooled analysis, with a *P* for trend of <0.001 (Table 3). Because additional adjustment for history of hypertension, triglycerides, and CRP levels did not significantly attenuate the risk estimates, we refer to model 3 in multivariable-adjusted and pooled analyses in Table 3.

We further examined the association using fixed cut points. Compared with HbA<sub>1c</sub> category 5.0% to <5.5% (reference), the pooled multivariable-adjusted RRs for CHD were 1.14 (95% CI 0.82 to 1.59) for <5.0%, 1.34 (95% CI 1.08 to 1.68) for 5.5% to <6.0%, and 1.85 (95% CI 1.28 to 2.69) for 6.0% to <6.5% (Figure 1). A significantly elevated risk of CHD was observed among nondiabetic women and men with HbA<sub>1c</sub> levels of  $\geq 5.5\%$ . Even in these populations restricted to lower HbA<sub>1c</sub> levels, we found that for every 0.5%-increment increase in HbA<sub>1c</sub> levels, the RR of CHD was 1.32 (95% CI 1.05 to 1.65) in women and 1.27 (95% CI 1.04 to 1.55) in men, and 1.29 (95% CI 1.11 to 1.50) when pooled. This association did not differ by health and lifestyle factors such as age, smoking status, BMI <25 and  $\geq 25$  kg/m<sup>2</sup>, history of hypertension, or parental history of early MI (Table S1). We further examined the joint

**Table 1.** Baseline Characteristics of Subjects With Incident Myocardial Infarction (Cases) and Matched\* Event-Free Controls Among Women (the Nurses' Health Study, 14 Years of Follow-up) and Men (the Health Professionals Follow-Up Study, 10 Years of Follow-up) Without History of Diabetes and Hemoglobin A<sub>1c</sub> <6.5%

Characteristics <sup>†</sup>	Nurses' Health Study			Health Professionals Follow-Up Study		
	Cases	Controls	P Value <sup>‡</sup>	Cases	Controls	P Value <sup>‡</sup>
N	371	832		396	843	
Age, y	59.9±6.6	59.7±6.5	—	64.1±8.8	64.0±8.6	—
Current smokers, %	28.3	26.7	—	9.6	9.1	—
Body mass index, kg/m <sup>2</sup>	25.8±4.7	24.9±4.1	0.004	25.9±3.1	25.6±3.3	0.10
Hormone replacement therapy use among postmenopausal women, %	43.0	45.3	0.76	—	—	—
Parental history of myocardial infarction before age 60, %	24.5	13.9	<0.001	15.9	11.4	0.03
History of hypercholesterolemia, %	52.3	39.2	<0.001	50.0	41.2	0.004
History of hypertension, %	44.7	25.4	<0.001	34.3	28.5	0.04
NSAID use, %	67.7	69.1	0.62	46.7	43.5	0.29
Total caloric intake, kcal	1796±561	1765±510	0.37	2076±645	2044±642	0.41
Alcohol consumption, g/day	1.1 (0.0 to 6.0)	1.8 (0.0 to 8.8)	0.09	4.7 (0.9 to 15.4)	7.0 (0.9 to 18.3)	0.02
Physical activity, MET-h/wk	12.2 (4.3 to 28.0)	12.4 (5.4 to 25.2)	0.69	22.9 (10.2 to 46.4)	26.9 (12.0 to 48.9)	0.08
HbA <sub>1c</sub> , %	5.5 (5.2 to 5.7)	5.4 (5.2 to 5.7)	0.01	5.6 (5.3 to 5.8)	5.5 (5.3 to 5.7)	0.01
CRP, mg/L	2.40 (1.07 to 5.44)	1.82 (0.80 to 3.80)	<0.001	1.33 (0.58 to 2.59)	0.97 (0.48 to 2.09)	<0.001
Total cholesterol, mg/dL	235±40.7	227±40.8	0.001	212±38.3	204±36.2	<0.001
HDL cholesterol, mg/dL	54.0±14.8	60.9±16.7	<0.001	42.9±11.5	46.4±12.7	<0.001
LDL cholesterol, mg/dL	146±37.0	135±37.5	<0.001	135±34.3	127±31.0	<0.001
Triglycerides, mg/dL	115 (82 to 167)	101 (74 to 141)	<0.001	139 (94 to 199)	116 (85 to 167)	<0.001
Total cholesterol:HDL cholesterol ratio	4.66±1.5	4.00±1.3	<0.001	5.19±1.3	4.66±1.3	<0.001

MET-h indicates metabolic equivalent hours; HbA<sub>1c</sub>, hemoglobin A<sub>1c</sub>; CRP, C-reactive protein; HDL, high-density lipoprotein; LDL, low-density lipoprotein.

\*Matching criteria were age, smoking status, and date of blood draw; in women, additional matching criteria included fasting status.

<sup>†</sup>Values are mean±SD for continuous variables and proportions for categorical variables, except HbA<sub>1c</sub>, CRP, triglycerides, alcohol, and physical activity, which are shown as median (interquartile range).

<sup>‡</sup>P for difference between cases and controls (unadjusted), determined by Student *t* test for rows with means, by Wilcoxon's rank sum test for rows with medians, and by  $\chi^2$  test for rows with proportions.

effect of CRP levels based on clinical cut points for CRP and categories of HbA<sub>1c</sub>. Of note, in the lower risk group with CRP levels  $\leq 3.0$  mg/L, a 0.5%-increment increase of HbA<sub>1c</sub> was significantly associated with risk of incident CHD among women (RR 1.42, 95% CI 1.04 to 1.94) and men (RR 1.28, 95% CI 1.03 to 1.60). In pooled analyses, participants in the highest CRP and HbA<sub>1c</sub> categories had an RR of CHD of 2.49 (95% CI 1.32 to 4.71) compared with participants in the lowest of both biomarkers (Figure 2).

A single biomarker at baseline may be a better proxy for exposure over the first several years and possibly become less correlated with exposure over a longer period. When we stratified according to the midpoint of follow-up, the association was strongest among cases and controls in the first 7 years of follow-up. For every 0.5%-increment increase in HbA<sub>1c</sub> levels in the earlier follow-up (<7 years) only, the multivariable-adjusted RR of incident CHD was 1.39 (95% CI

0.97 to 1.98) for women, 1.33 (95% CI 1.02 to 1.73) for men, and 1.35 (95% CI 1.09 to 1.67) when we pooled the estimates. In the later follow-up ( $\geq 7$  years) only, the corresponding RRs were 1.17 (95% CI 0.90 to 1.52) for women, 1.26 (95% CI 0.92 to 1.72) for men, and 1.20 (95% CI 0.98 to 1.47) when pooled. Finally, when we controlled for regression dilution bias to account for variability in the baseline measure, the pooled RR per 0.5% increment was 1.51 (95% CI 1.13 to 2.02) when restricted to the earlier follow-up cycle and 1.42 (95% CI 1.16 to 1.74) for the entire follow-up cycle.

## Discussion

In these prospective nested case-control studies in 2 cohorts of US health professionals, we found that HbA<sub>1c</sub> was predictive of CHD risk in participants without diagnosed

**Table 2.** Age-Adjusted Spearman Partial Correlation Coefficients Between Hemoglobin A<sub>1c</sub> and Cardiovascular Risk Factors Among Controls in Women (the Nurses' Health Study) and Men (the Health Professionals Follow-Up Study)

HbA <sub>1c</sub> %	NHS	HPFS
CRP	0.15 <sup>‡</sup>	0.13 <sup>‡</sup>
Total cholesterol	0.04	0.08*
LDL cholesterol	0.04	0.05
HDL cholesterol	-0.09 <sup>†</sup>	-0.08*
Total cholesterol:HDL cholesterol ratio	0.09 <sup>†</sup>	0.11 <sup>†</sup>
Triglycerides	0.07*	0.12 <sup>‡</sup>
BMI	0.09 <sup>†</sup>	0.10 <sup>†</sup>
Alcohol, g/d	-0.03	-0.08*

HbA<sub>1c</sub> indicates hemoglobin A<sub>1c</sub>; NHS, Nurses' Health Study; HPFS, Health Professionals Follow-Up Study; CRP, C-reactive protein; LDL, low-density lipoprotein; HDL, high-density lipoprotein; BMI, body mass index.  
 \*P<0.05, †P<0.01, ‡P<0.001.

diabetes and with HbA<sub>1c</sub> concentrations <6.5%. Men and women with an HbA<sub>1c</sub> of 6.0% to <6.5% had an 85% higher risk of CHD than those with levels of 5.0% to <5.5%. The positive association persisted when adjusted for traditional cardiovascular risk factors, including hyperlipidemia, anthropometric variables, and CRP. Our results suggest that HbA<sub>1c</sub> is an important marker of risk across a broad range of levels and that the causal etiologic pathway may be complex and include mechanisms other than lipids or inflammation.

The relationship of HbA<sub>1c</sub> with CHD has been studied in several other nondiabetic populations<sup>11,12,29-32</sup> with inconsistent results, possibly due to diverse study protocols, a wide range of length of follow-up, inconsistent exclusion criteria, and differences in cohort demographics and outcome definitions. Two previous studies in particular have examined HbA<sub>1c</sub> in relation to risk of CHD in participants without diabetes or clinically elevated HbA<sub>1c</sub> levels.<sup>13,29</sup> After 7 years of follow-up in a nested case-control analysis from the WHS, Blake et al reported an RR of 2.25 (95% CI 1.59 to 3.19) for CVD events

**Table 3.** Estimated Relative Risks (and 95% CIs) of Myocardial Infarction During Follow-up According to Quintiles of Baseline Percent Hemoglobin A<sub>1c</sub> Among Women (Nurses' Health Study, 14 Years of Follow-up) and Men (the Health Professionals Follow-Up Study, 10 Years of Follow-up)

	Quintile of HbA <sub>1c</sub> *					P Trend <sup>†</sup>
	1	2	3	4	5	
<b>Nurses' Health Study (371 cases/832 controls)</b>						
HbA <sub>1c</sub> , %	<5.1	5.1 to <5.3	5.3 to <5.5	5.5 to <5.7	≥5.7	
Cases/controls	62/166	61/168	75/165	84/167	89/166	
Model 1 <sup>‡</sup>	1.0	1.06 (0.70 to 1.62)	1.43 (0.94 to 2.18)	1.73 (1.12 to 2.66)	2.01 (1.26 to 3.21)	<0.001
Model 2 <sup>§</sup>	1.0	1.06 (0.69 to 1.65)	1.39 (0.89 to 2.16)	1.59 (1.01 to 2.50)	1.78 (1.09 to 2.91)	0.009
Model 3 <sup>¶</sup>	1.0	1.05 (0.68 to 1.63)	1.35 (0.86 to 2.10)	1.55 (0.98 to 2.44)	1.71 (1.04 to 2.80)	0.01
Model 4 <sup>  </sup>	1.0	1.05 (0.67 to 1.65)	1.39 (0.88 to 2.18)	1.52 (0.96 to 2.43)	1.70 (1.03 to 2.81)	0.02
<b>Health Professionals Follow-Up Study (396 cases/843 controls)</b>						
HbA <sub>1c</sub> , %	<5.2	5.2 to <5.4	5.4 to <5.6	5.6 to <5.8	≥5.8	
Cases/controls	64/168	66/169	87/168	73/170	106/168	
Model 1 <sup>‡</sup>	1.0	1.12 (0.73 to 1.70)	1.56 (1.03 to 2.38)	1.35 (0.87 to 2.11)	1.97 (1.28 to 3.03)	0.001
Model 2 <sup>§</sup>	1.0	1.10 (0.71 to 1.68)	1.52 (0.99 to 2.33)	1.19 (0.75 to 1.88)	1.67 (1.07 to 2.59)	0.03
Model 3 <sup>¶</sup>	1.0	1.10 (0.72 to 1.69)	1.55 (1.01 to 2.39)	1.19 (0.75 to 1.88)	1.69 (1.08 to 2.64)	0.02
Model 4 <sup>  </sup>	1.0	1.08 (0.70 to 1.66)	1.57 (1.02 to 2.42)	1.17 (0.74 to 1.86)	1.69 (1.08 to 2.65)	0.02
<b>Pooled</b>						
Cases/controls	126/334	127/337	162/333	157/337	195/334	
Model 3 <sup>¶</sup>	1.0	1.08 (0.79 to 1.46)	1.45 (1.06 to 1.97)	1.36 (0.98 to 1.88)	1.70 (1.22 to 2.36)	<0.001

HbA<sub>1c</sub> indicates hemoglobin A<sub>1c</sub>; NHS, Nurses' Health Study; HDL, high-density lipoprotein; CRP, C-reactive protein.

\*Quintiles of HbA<sub>1c</sub> are based on controls in each sex.


<sup>†</sup>P for trend based on median HbA<sub>1c</sub> levels in quintiles of the control subjects.

<sup>‡</sup>Adjusted for age, smoking status, date of blood drawing, fasting status (in NHS only), and laboratory batch.

<sup>§</sup>Additionally adjusted for family history of myocardial infarction before age of 60 years, alcohol intake, physical activity, nonsteroidal antiinflammatory drug use, total cholesterol:HDL cholesterol ratio, and hormone replacement therapy use among women.

<sup>¶</sup>Additionally adjusted for body mass index.


<sup>||</sup>Further adjusted for hypertension and CRP.



**Figure 1.** Multivariable-adjusted\* relative risk (RR) of incident CHD among women and men according to categories of HbA<sub>1c</sub> levels. \*Adjusted for age, smoking status, date of blood draw, fasting status (in NHS only), laboratory batch, family history of myocardial infarction before age of 60 years, alcohol intake, physical activity, NSAID use, total cholesterol:HDL cholesterol ratio, BMI, and hormone replacement therapy use among women. CHD indicates coronary heart disease; HbA<sub>1c</sub>, hemoglobin A1c; NHS, Nurses' Health Study; NSAID, nonsteroidal antiinflammatory drug; HDL, high-density lipoprotein; BMI, body mass index.

(368 cases of incident MI, stroke, or coronary revascularization) comparing the highest ( $\geq 5.50\%$ ) with the lowest ( $< 5.23\%$ ) quartile of HbA<sub>1c</sub>.<sup>13</sup> However, this association was attenuated (RR 1.00; 95% CI 0.65 to 1.54) after adjustment for BMI, blood pressure, CRP, low-density lipoprotein and HDL cholesterol, and triglycerides.<sup>13</sup> The authors note that the same results were observed when the end point of CHD was examined alone, although with limited power because the CHD analysis was limited to 136 cases. In a recent analysis from the Atherosclerosis Risk in Communities (ARIC) study, the risk factor-adjusted hazard ratios for CHD in participants with HbA<sub>1c</sub> of 5.5% to  $< 6.0\%$  and of 6.0% to  $< 6.5\%$  were 1.23 (95% CI 1.07 to 1.41) and 1.78 (95% CI 1.48 to 2.15), respectively, compared with participants with baseline HbA<sub>1c</sub> concentrations of 5.0% to  $< 5.5\%$ .<sup>29</sup> Our results were similar in magnitude to this study. Furthermore, we showed a significant joint effect of mildly elevated HbA<sub>1c</sub> and CRP levels on the risk of incident CHD among nondiabetic men and women.

Interestingly, we observed a joint association similar to that of Sander et al,<sup>33</sup> who reported that among nondiabetics, the combination of elevated HbA<sub>1c</sub> and CRP was associated with advanced early carotid atherosclerosis progression and a 3-fold increased risk of major vascular events. This association between elevated HbA<sub>1c</sub> and early atherosclerosis progression supports the hypothesis that HbA<sub>1c</sub> may be causally related to CHD even at low levels.<sup>33</sup> Hyperglycemia, in addition to diabetes, can lead to the formation of advanced



**Figure 2.** Multivariable-adjusted\* relative risk of incident CHD among women and men according to categories of HbA<sub>1c</sub> and clinical cut points of CRP levels. \*Adjusted for age, smoking status, date of blood draw, fasting status (in NHS only), laboratory batch, family history of myocardial infarction before age of 60 years, alcohol intake, physical activity, NSAID use, total cholesterol:HDL cholesterol ratio, BMI, and hormone replacement therapy use among women. † $P < 0.05$ . CHD indicates coronary heart disease; HbA<sub>1c</sub>, hemoglobin A1c; CRP, C-reactive protein; NHS, Nurses' Health Study; NSAID, nonsteroidal antiinflammatory drug; HDL, high-density lipoprotein; BMI, body mass index.

glycosylated end products, which have been shown to induce production and secretion of inflammatory cytokines.<sup>34</sup> Indeed, HbA<sub>1c</sub> has been reported to predict CRP in established diabetes.<sup>35</sup> In addition, nitric oxide bioavailability, a marker of intimal reactivity, has been shown to be reduced by both CRP<sup>36</sup> and HbA<sub>1c</sub>.<sup>37</sup> This joint effect was observed at even mildly elevated levels of both biomarkers and may influence CHD risk through mechanisms other than diabetes, atherosclerosis, and inflammation.

A recent meta-analysis<sup>38</sup> examined the association between HbA<sub>1c</sub> and risk of CHD in persons without diabetes, although most studies included participants with HbA<sub>1c</sub> well beyond the established clinical threshold of 6.5%. Using data from 9 cohorts (1639 cases), the meta-analysis reported an RR of 1.20 (95% CI 1.10-1.31) for each 1%-increment increase in levels of HbA<sub>1c</sub>.<sup>38</sup> This risk estimate was not adjusted for established CHD risk factors and was weaker than our own. These differences may be due to the inherent weakness of meta-analyses, where pooling results from many studies may lead to random error from differences in lab methods, study design, and residual confounding. In our nested prospective studies, the pooled RR of CHD for men and women was 1.67 (95% CI 1.23 to 2.25) per 1%-increment increase in HbA<sub>1c</sub>, even after detailed adjustment for other risk factors.

In the present study, even a 0.5%-increment increase of HbA<sub>1c</sub> was significantly associated with risk of incident CHD among both women and men. Although a 0.5%-increment decrease in HbA<sub>1c</sub> in patients undergoing treatment for diabetes<sup>39</sup> and morbid obesity<sup>40</sup> has proven health benefits, there are no published experimental trials that show a similar decrease in HbA<sub>1c</sub> leads to a reduction in CHD or CVD end points in the general population. Nevertheless, the most effective ways of reducing HbA<sub>1c</sub>, namely dietary changes, increased physical activity, and weight loss, have been shown to have beneficial effects on other cardiovascular risk factors. HbA<sub>1c</sub> in nondiabetic individuals may be successfully reduced through nonpharmacologic lifestyle interventions, as demonstrated in the Finnish Diabetes Prevention Study,<sup>41,42</sup> Diabetes Prevention Program<sup>43</sup> and Outcomes Study,<sup>44</sup> and Look AHEAD study.<sup>45</sup> In a small intervention study of nondiabetic obese males, the mean HbA<sub>1c</sub> levels were significantly lower after the weight loss intervention compared with the before-intervention levels (4.9% versus 5.1%),<sup>46</sup> and in a randomized controlled trial of male construction workers, lifestyle intervention aimed at energy balance was significantly associated with improved cardiovascular risk factors, including body weight, blood pressure, and lipid and HbA<sub>1c</sub> levels.<sup>47</sup> Although whether lowering HbA<sub>1c</sub> lowers risk of future CHD among nondiabetic individuals has not been directly tested, working to optimize cardiovascular health in individuals with mildly higher HbA<sub>1c</sub> seems prudent and clinically important.

Strengths of the present analysis include the long duration of follow-up, the prospective nature of the data collection, replication in a second cohort, and comprehensive lifestyle and health factor data used for detailed multivariable adjustment. A few limitations should be addressed. First, the potential exists for measurement error in lifestyle data ascertained from self-reported questionnaires. However, the reproducibility and validity of the self-reported questionnaire have been well-documented in these 2 populations of health professionals.<sup>15,48</sup> Second, our study was conducted in 2 cohorts of middle-aged, predominantly white health professionals and may not be generalizable to other populations. However, the relationship between HbA<sub>1c</sub> and CHD risk was similar in a recent community-based study among whites and African Americans.<sup>29</sup> Third, we only had a single measurement of HbA<sub>1c</sub> at baseline and did not have glucose measurements. However, one advantage of using HbA<sub>1c</sub> is that it is stable in stored samples and reflects glycemic control over an average of 3 months rather than measures such as glucose, which are susceptible to daily fluctuations.<sup>49</sup> In addition, we observed excellent within-person reproducibility in repeated samples measured 3 years apart and conducted additional analyses correcting for potential changes in the baseline measurement over time.

In summary, we found that HbA<sub>1c</sub> is a predictor of CHD risk among apparently healthy women and men without diabetes

and with HbA<sub>1c</sub> <6.5%. Further examination is warranted to determine whether reducing HbA<sub>1c</sub> among otherwise intermediate- and low-risk individuals without diagnosed diabetes will prevent incident CHD events.

## Sources of Funding

This study was supported by grants (HL35464, CA55075, HL34594, CA87969, and CA49449) from the National Institutes of Health. Dr Pai was supported in part by National Institutes of Health grant HL088372. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

## Disclosures

None.

## References

1. Sarwar N, Gao P, Seshasai SR, Gobin R, Kaptoge S, Di Angelantonio E, Ingelsson E, Lawlor DA, Selvin E, Stampfer M, Stehouwer CD, Lewington S, Pennells L, Thompson A, Sattar N, White IR, Ray KK, Danesh J. Diabetes mellitus, fasting blood glucose concentration, and risk of vascular disease: a collaborative meta-analysis of 102 prospective studies. *Lancet*. 2010;375:2215–2222.
2. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2012;35(suppl 1):S64–S71.
3. Gerstein HC. Glycosylated hemoglobin: finally ready for prime time as a cardiovascular risk factor. *Ann Intern Med*. 2004;141:475–476.
4. World Health Organization. Use of Glycated Haemoglobin (HbA1c) in the Diagnosis of Diabetes Mellitus. Who/nmh/chp/cpm/11.1. Geneva, Switzerland: World Health Organization; 2011:25.
5. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). UK Prospective Diabetes Study (UKPDS) group. *Lancet*. 1998;352:837–853.
6. Gerstein HC, Miller ME, Byington RP, Goff DC Jr, Bigger JT, Buse JB, Cushman WC, Genuth S, Ismail-Beigi F, Grimm RH Jr, Probstfield JL, Simons-Morton DG, Friedewald WT. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med*. 2008;358:2545–2559.
7. Patel A, MacMahon S, Chalmers J, Neal B, Billot L, Woodward M, Marre M, Cooper M, Glasziou P, Grobbee D, Hamet P, Harrap S, Heller S, Liu L, Mancia G, Mogensen CE, Pan C, Poulter N, Rodgers A, Williams B, Bompont S, de Galan BE, Joshi R, Travert F. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med*. 2008;358:2560–2572.
8. Duckworth W, Abraira C, Moritz T, Reda D, Emanuele N, Reaven PD, Zieve FJ, Marks J, Davis SN, Hayward R, Warren SR, Goldman S, McCarren M, Vitek ME, Henderson WG, Huang GD. Glucose control and vascular complications in veterans with type 2 diabetes. *N Engl J Med*. 2009;360:129–139.
9. Gerstein HC. More insights on the dysglycaemia-cardiovascular connection. *Lancet*. 2010;375:2195–2196.
10. Gerstein HC, Yusuf S. Dysglycaemia and risk of cardiovascular disease. *Lancet*. 1996;347:949–950.
11. Selvin E, Coresh J, Golden SH, Brancati FL, Folsom AR, Steffes MW. Glycemic control and coronary heart disease risk in persons with and without diabetes: the Atherosclerosis Risk in Communities Study. *Arch Intern Med*. 2005;165:1910–1916.
12. Khaw KT, Wareham N, Bingham S, Luben R, Welch A, Day N. Association of hemoglobin A1c with cardiovascular disease and mortality in adults: the European prospective investigation into cancer in Norfolk. *Ann Intern Med*. 2004;141:413–420.
13. Blake GJ, Pradhan AD, Manson JE, Williams GR, Buring J, Ridker PM, Glynn RJ. Hemoglobin A1c level and future cardiovascular events among women. *Arch Intern Med*. 2004;164:757–761.
14. Colditz GA. The Nurses' Health Study: a cohort of US women followed since 1976. *J Am Med Womens Assoc*. 1995;50:40–44.



15. Rimm EB, Giovannucci EL, Stampfer MJ, Colditz GA, Litin LB, Willett WC. Reproducibility and validity of an expanded self-administered semiquantitative food frequency questionnaire among male health professionals. *Am J Epidemiol*. 1992;135:1114–1126; discussion 1127–1136.
16. Prentice RL, Breslow NE. Retrospective studies and failure time models. *Biometrika*. 1978;65:153–158.
17. Luepker RV, Apple FS, Christenson RH, Crow RS, Fortmann SP, Goff D, Goldberg RJ, Hand MM, Jaffe AS, Julian DG, Levy D, Manolio T, Mendis S, Mensah G, Pajak A, Prineas RJ, Reddy KS, Roger VL, Rosamond WD, Shahar E, Sharrett AR, Sorlie P, Tunstall-Pedoe H. Case definitions for acute coronary heart disease in epidemiology and clinical research studies: a statement from the AHA Council on Epidemiology and Prevention; AHA Statistics Committee; World Heart Federation Council on Epidemiology and Prevention; the European Society of Cardiology Working Group on Epidemiology and Prevention; Centers for Disease Control and Prevention; and the National Heart, Lung, and Blood Institute. *Circulation*. 2003;108:2543–2549.
18. Rose GA, Blackburn H. Cardiovascular Survey Methods. WHO monograph series no. 58. Geneva, Switzerland: World Health Organization; 1982.
19. Willett W, Lenart E. Reproducibility and validity of food-frequency questionnaires. In: Willett W, ed. *Nutritional Epidemiology*. New York, NY: Oxford University Press; 1998:101–147.
20. Pai JK, Pischon T, Ma J, Manson JE, Hankinson SE, Joshipura K, Curhan GC, Rifai N, Cannuscio CC, Stampfer MJ, Rimm EB. Inflammatory markers and the risk of coronary heart disease in men and women. *N Engl J Med*. 2004; 351:2599–2610.
21. Pai JK, Curhan GC, Cannuscio CC, Rifai N, Ridker PM, Rimm EB. Stability of novel plasma markers associated with cardiovascular disease: processing within 36 hours of specimen collection. *Clin Chem*. 2002;48:1781–1784.
22. Pischon T, Hankinson SE, Hotamisligil GS, Rifai N, Willett WC, Rimm EB. Habitual dietary intake of n-3 and n-6 fatty acids in relation to inflammatory markers among US men and women. *Circulation*. 2003;108:155–160.
23. Rothman KJ, Greenland S. Case-control studies. In: Rothman KJ, Greenland S, ed. *Modern Epidemiology*. Philadelphia, Pa: Lippincott-Raven Publishers; 1998:93–114.
24. Durrleman S, Simon R. Flexible regression models with cubic splines. *Stat Med*. 1989;8:551–561.
25. Takkouche B, Cadarso-Suarez C, Spiegelman D. Evaluation of old and new tests of heterogeneity in epidemiologic meta-analysis. *Am J Epidemiol*. 1999; 150:206–215.
26. Lewington S, Whitlock G, Clarke R, Sherliker P, Emberson J, Halsey J, Qizilbash N, Peto R, Collins R. Blood cholesterol and vascular mortality by age, sex, and blood pressure: a meta-analysis of individual data from 61 prospective studies with 55,000 vascular deaths. *Lancet*. 2007;370:1829–1839.
27. Wu JH, Lemaitre RN, King IB, Song X, Sacks FM, Rimm EB, Heckbert SR, Siscovick DS, Mozaffarian D. Association of plasma phospholipid long-chain omega-3 fatty acids with incident atrial fibrillation in older adults: the Cardiovascular Health Study. *Circulation*. 2012;125:1084–1093.
28. Clarke R, Shipley M, Lewington S, Youngman L, Collins R, Marmot M, Peto R. Underestimation of risk associations due to regression dilution in long-term follow-up of prospective studies. *Am J Epidemiol*. 1999;150:341–353.
29. Selvin E, Steffes MW, Zhu H, Matsushita K, Wagenknecht L, Pankow J, Coresh J, Brancati FL. Glycated hemoglobin, diabetes, and cardiovascular risk in nondiabetic adults. *N Engl J Med*. 2010;362:800–811.
30. Pradhan AD, Rifai N, Buring JE, Ridker PM. Hemoglobin A<sub>1c</sub> predicts diabetes but not cardiovascular disease in nondiabetic women. *Am J Med*. 2007; 120:720–727.
31. Adams RJ, Appleton SL, Hill CL, Wilson DH, Taylor AW, Chittleborough CR, Gill TK, Ruffin RE. Independent association of HbA<sub>1c</sub> and incident cardiovascular disease in people without diabetes. *Obesity (Silver Spring)*. 2009;17:559–563.
32. Lawlor DA, Fraser A, Ebrahim S, Smith GD. Independent associations of fasting insulin, glucose, and glycated haemoglobin with stroke and coronary heart disease in older women. *PLoS Med*. 2007;4:e263.
33. Sander D, Schulze-Horn C, Bickel H, Gnahn H, Bartels E, Conrad B. Combined effects of hemoglobin A<sub>1c</sub> and C-reactive protein on the progression of subclinical carotid atherosclerosis: the INVADE study. *Stroke*. 2006;37:351–357.
34. Vlassara H, Brownlee M, Manogue KR, Dinarello CA, Pasagian A. Cachectin/TNF and IL-1 induced by glucose-modified proteins: role in normal tissue remodeling. *Science*. 1988;240:1546–1548.
35. King DE, Mainous AG III, Buchanan TA, Pearson WS. C-reactive protein and glycemic control in adults with diabetes. *Diabetes Care*. 2003;26:1535–1539.
36. Verma S, Wang CH, Li SH, Dumont AS, Fedak PW, Badiwala MV, Dhillon B, Weisel RD, Li RK, Mickle DA, Stewart DJ. A self-fulfilling prophecy: C-reactive protein attenuates nitric oxide production and inhibits angiogenesis. *Circulation*. 2002;106:913–919.
37. Giugliano D, Marfella R, Coppola L, Verrazzo G, Acampora R, Giunta R, Nappo F, Lucarelli C, D'Onofrio F. Vascular effects of acute hyperglycemia in humans are reversed by L-arginine. Evidence for reduced availability of nitric oxide during hyperglycemia. *Circulation*. 1997;95:1783–1790.
38. Sarwar N, Aspelund T, Eiriksdottir G, Gobin R, Seshasai SR, Forouhi NG, Sigurdsson G, Danesh J, Gudnason V. Markers of dysglycaemia and risk of coronary heart disease in people without diabetes: Reykjavik prospective study and systematic review. *PLoS Med*. 2010;7:e1000278.
39. Umpierre D, Ribeiro PA, Kramer CK, Leitao CB, Zucatti AT, Azevedo MJ, Gross JL, Ribeiro JP, Schaun BD. Physical activity advice only or structured exercise training and association with HbA<sub>1c</sub> levels in type 2 diabetes: a systematic review and meta-analysis. *JAMA*. 2011;305:1790–1799.
40. Li Q, Chen L, Yang Z, Ye Z, Huang Y, He M, Zhang S, Feng X, Gong W, Zhang Z, Zhao W, Liu C, Ou S, Hu R. Metabolic effects of bariatric surgery in type 2 diabetic patients with body mass index <35 kg/m<sup>2</sup>. *Diabetes Obes Metab*. 2011;14:262–270.
41. Tuomilehto J, Lindstrom J, Eriksson JG, Valle TT, Hamalainen H, Ilanne-Parikka P, Keinanen-Kiukaanniemi S, Laakso M, Louheranta A, Rastas M, Salminen V, Uusitupa M. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med*. 2001;344: 1343–1350.
42. Lindstrom J, Ilanne-Parikka P, Peltonen M, Aunola S, Eriksson JG, Hemio K, Hamalainen H, Harkonen P, Keinanen-Kiukaanniemi S, Laakso M, Louheranta A, Mannelini M, Paturi M, Sundvall J, Valle TT, Uusitupa M, Tuomilehto J. Sustained reduction in the incidence of type 2 diabetes by lifestyle intervention: follow-up of the Finnish Diabetes Prevention Study. *Lancet*. 2006; 368:1673–1679.
43. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med*. 2002;346:393–403.
44. Perreault L, Pan Q, Mather KJ, Watson KE, Hamman RF, Kahn SE. Effect of regression from prediabetes to normal glucose regulation on long-term reduction in diabetes risk: results from the Diabetes Prevention Program Outcomes Study. *Lancet*. 2012;379:2243–2251.
45. Wing RR, Lang W, Wadden TA, Safford M, Knowler WC, Bertoni AG, Hill JO, Brancati FL, Peters A, Wagenknecht L. Benefits of modest weight loss in improving cardiovascular risk factors in overweight and obese individuals with type 2 diabetes. *Diabetes Care*. 2011;34:1481–1486.
46. Kotani K, Yamada T, Taniguchi N. The association between circulating secreted protein acidic and rich in cysteine (SPARC) and glycosylated haemoglobin (HbA<sub>1c</sub>) during lifestyle-modified weight reduction intervention in obese male subjects. *J Int Med Res*. 2011;39:528–532.
47. Groeneveld IF, Proper KI, van der Beek AJ, van Mechelen W. Sustained body weight reduction by an individual-based lifestyle intervention for workers in the construction industry at risk for cardiovascular disease: results of a randomized controlled trial. *Prev Med*. 2010;51:240–246.
48. Feskanich D, Rimm EB, Giovannucci EL, Colditz GA, Stampfer MJ, Litin LB, Willett WC. Reproducibility and validity of food intake measurements from a semiquantitative food frequency questionnaire. *J Am Diet Assoc*. 1993;93: 790–796.
49. Little RR, Rohlfing CL, Tennill AL, Connolly S, Hanson S. Effects of sample storage conditions on glycated hemoglobin measurement: evaluation of five different high performance liquid chromatography methods. *Diabetes Technol Ther*. 2007;9:36–42.