



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Draft Genome Sequence of Myxococcus xanthus Wild-Type Strain DZ2, a Model Organism for Predation and Development

The Harvard community has made this article openly available.

[Please share](#) how this access benefits you. Your story matters.

Citation	Müller, Susanne, Jonathan W. Willett, Sarah M. Bahr, Cynthia L. Darnell, Katherine R. Hummels, Carolyn K. Dong, Hera C. Vlamakis, and John R. Kirby. 2013. Draft genome sequence of Myxococcus xanthus wild-type strain DZ2, a model organism for predation and development. Genome Announcements 1(3): e00217-13.
Published Version	doi:10.1128/genomeA.00217-13
Accessed	February 19, 2015 12:08:30 PM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:11181054
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

Draft Genome Sequence of *Myxococcus xanthus* Wild-Type Strain DZ2, a Model Organism for Predation and Development

Susanne Müller,^a Jonathan W. Willett,^b Sarah M. Bahr,^a Cynthia L. Darnell,^a Katherine R. Hummels,^a Carolyn K. Dong,^c Hera C. Vlamakis,^d John R. Kirby^a

Department of Microbiology, University of Iowa, Iowa City, Iowa, USA^a; University of Chicago, Department of Biochemistry and Molecular Biology, Chicago, Illinois, USA^b; U.S. Department of State, Bureau of Economic and Business Affairs, Office of Agriculture, Biotechnology, and Textile Trade Affairs, Washington, DC, USA^c; Harvard Medical School, Department of Microbiology and Immunobiology, Boston, Massachusetts, USA^d

S.M., J.W.W., and S.M.B. contributed equally to this work.

***Myxococcus xanthus* is a member of the *Myxococcales* order within the *Deltaproteobacteria* subdivision. The myxobacteria reside in soil, have relatively large genomes, and display complex life cycles. Here, we report the whole-genome shotgun sequence of strain DZ2, which includes unique genes not found previously in strain DK1622.**

Received 25 March 2013 Accepted 27 March 2013 Published 9 May 2013

Citation Müller S, Willett JW, Bahr SM, Darnell CL, Hummels KR, Dong CK, Vlamakis HC, Kirby JR. 2013. Draft genome sequence of *Myxococcus xanthus* wild-type strain DZ2, a model organism for predation and development. *Genome Announc*. 1(3):e00217-13. doi:10.1128/genomeA.00217-13.

Copyright © 2013 Müller et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to John R. Kirby, john-kirby@uiowa.edu.

Myxococcus xanthus is a member of the *Myxococcales* order (fruiting and gliding bacteria) within the *Deltaproteobacteria* subdivision. The myxobacteria typically reside in soil, are characterized by relatively large genomes (up to ~13 Mb), and display complex life cycles. In response to a variety of environmental cues, *M. xanthus* organizes into multicellular fruiting bodies harboring stress-resistant spores. Fruiting body formation is regulated by several mechanisms involving complex signal transduction cascades, production of cytosolic and extracellular signals, and gliding motility (1–34). Previous work determined the sequence of the *M. xanthus* laboratory strain DK1622 (accession number NC_008095.1) (34). While both DK1622 and the closely related strain DZ2 originate from the Roger Y. Stanier collection originally housed at the University of California, Berkeley, strain DK1622 was genetically modified from a parent (35). Moreover, phenotypic differences between DZ2 and DK1622 regarding rates of autolysis, rippling behavior, aggregation, and fruiting body formation have been reported (9, 36, 37).

M. xanthus strain DZ2 was sequenced at the University of Iowa DNA Core Facility using 454 GS-FLX titanium technology. Genomic DNA was prepared by resuspending cell pellets in SET buffer (75 mM NaCl, 10 mM Tris [pH 7.5], 25 mM EDTA, 1% SDS, and 1 mg/ml proteinase K) and then incubating the suspension for 2 h at 55°C. DNA samples were subsequently extracted with chloroform (3×), precipitated with isopropanol, and resuspended in 10 mM Tris (pH 8.0). The sample was processed for 454 sequencing according to established protocols. The resulting sequence represents approximately 20-fold coverage. The genome was assembled *de novo* using Newbler software version 2.7. The sequence comprises 292,633 reads totaling 186 Mb and was assembled into 87 contigs. Using the RAST genome annotation server, we were able to predict a total of 7,709 coding sequences (CDS) within the *M. xanthus* DZ2 genome (38).

The genome of DZ2 is 9.287 Mb, approximately 196 kb larger

than the published genome of *M. xanthus* strain DK1622. A similar size differential was described previously and is presumed to be the result of UV mutagenesis on the DK1622 progenitor strain DK101 (39, 40). The vast majority of the DZ2 genome is identical to the DK1622 genome, with the notable exception of DZ2-specific genes. Many DZ2-specific genes encode hypothetical proteins with high homology to sequences found within other myxobacteria, including *Myxococcus fulvus*, *Stigmatella aurantiaca*, and *Sorangium cellulosum*. Importantly, we identified unique genes likely to encode proteins involved in regulation of transcription or translation, signal transduction, fatty acid modification, and protein transport. The presence of DNA sequences unique to DZ2 has been verified by PCR. We are currently investigating whether the reported phenotypic differences between strains DZ2 and DK1622 might be attributable to production of unique proteins.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [AKYI000000000](https://www.ncbi.nlm.nih.gov/nuccore/AKYI000000000), corresponding to BioProject PRJNA168264. The version described in this paper is the first version, [AKYI020000000](https://www.ncbi.nlm.nih.gov/nuccore/AKYI020000000).

ACKNOWLEDGMENTS

Support for this work was provided by the University of Iowa and NIH NIAID R01 AI59682 and NSF MCB-1244021 to J.R.K. Additional support for J.W.W. was provided by NIH T32 GM077973. Support for C.K.D. was provided by NIH T32 AI007511.

We thank Tom Willett, Tom Bair, and Kevin Knudtson for helpful discussions and data analysis.

The content is the responsibility of the authors and does not represent the official views of the NIH, the NSF, or the University of Iowa.

REFERENCES

1. Willett JW, Kirby JR. 2011. CrdS and CrdA Comprise a two-component system that is cooperatively regulated by the Che3 Chemosensory system

- in *Myxococcus xanthus*. mBio 2:e00110–e00111. <http://dx.doi.org/10.1128/mBio.00110-11>.
2. Shimkets LJ, Gill RE, Kaiser D. 1983. Developmental cell interactions in *Myxococcus xanthus* and the spoC locus. Proc. Natl. Acad. Sci. U. S. A. 80:1406–1410.
 3. Singer M, Kaiser D. 1995. Ectopic production of guanosine penta- and tetraphosphate can initiate early developmental gene expression in *Myxococcus xanthus*. Genes Dev. 9:1633–1644.
 4. Zusman DR, Scott AE, Yang Z, Kirby JR. 2007. Chemosensory pathways, motility and development in *Myxococcus xanthus*. Nat. Rev. Microbiol. 5:862–872.
 5. Campos JM, Zusman DR. 1975. Regulation of development in *Myxococcus xanthus*: effect of 3':5'-cyclic AMP, ADP, and nutrition. Proc. Natl. Acad. Sci. U. S. A. 72:518–522.
 6. Blackhart BD, Zusman DR. 1985. "Frizzy" genes of *Myxococcus xanthus* are involved in control of frequency of reversal of gliding motility. Proc. Natl. Acad. Sci. U. S. A. 82:8767–8770.
 7. O'Connor KA, Zusman DR. 1999. Induction of β -lactamase influences the course of development in *Myxococcus xanthus*. J. Bacteriol. 181:6319–6331.
 8. Mignot T, Merlie JP, Zusman DR. 2005. Regulated Pole-to-pole oscillations of a bacterial gliding motility protein. Science 310:855–857.
 9. Berleman JE, Kirby JR. 2007. Multicellular development in *Myxococcus xanthus* is stimulated by predator-prey interactions. J. Bacteriol. 189:5675–5682.
 10. Kirby JR, Zusman DR. 2003. Chemosensory regulation of developmental gene expression in *Myxococcus xanthus*. Proc. Natl. Acad. Sci. U. S. A. 100:2008–2013.
 11. Willett JW, Kirby JR. 2012. Genetic and biochemical dissection of a HisKA domain identifies residues required exclusively for kinase and phosphatase activities. PLoS Genet. 8:e1003084. <http://dx.doi.org/10.1371/journal.pgen.1003084>.
 12. Zhang Y, Ducret A, Shaevitz J, Mignot T. 2012. From individual cell motility to collective behaviors: insights from a prokaryote, *Myxococcus xanthus*. FEMS Microbiol. Rev. 36:149–164.
 13. Luciano J, Agrebi R, Le Gall AV, Wartel M, Fiegna F, Ducret A, Brochier-Armanet C, Mignot T. 2011. Emergence and modular evolution of a novel motility machinery in bacteria. PLoS Genet. 7:e1002268. <http://dx.doi.org/10.1371/journal.pgen.1002268>.
 14. Mauriello EM, Mignot T, Yang Z, Zusman DR. 2010. Gliding motility revisited: how do the myxobacteria move without flagella? Microbiol. Mol. Biol. Rev. 74:229–249.
 15. Mignot T, Kirby JR. 2008. Genetic circuitry controlling motility behaviors of *Myxococcus xanthus*. Bioessays 30:733–743.
 16. Mignot T, Shaevitz JW, Hartzell PL, Zusman DR. 2007. Evidence that focal adhesion complexes power bacterial gliding motility. Science 315:853–856.
 17. Mignot T. 2007. The elusive engine in *Myxococcus xanthus* gliding motility. Cell. Mol. Life Sci. 64:2733–2745.
 18. Huntley S, Hamann N, Wegener-Feldbrügge S, Treuner-Lange A, Kube M, Reinhardt R, Klages S, Müller R, Ronning CM, Nierman WC, Søgaard-Andersen L. 2011. Comparative genomic analysis of fruiting body formation in Myxococcales. Mol. Biol. Evol. 28:1083–1097.
 19. Kruse T, Lobedanz S, Berthelsen NM, Søgaard-Andersen L. 2001. C-signal: a cell surface-associated morphogen that induces and coordinates multicellular fruiting body morphogenesis and sporulation in *Myxococcus xanthus*. Mol. Microbiol. 40:156–168.
 20. Jelsbak L, Søgaard-Andersen L. 2002. Pattern formation by a cell surface-associated morphogen in *Myxococcus xanthus*. Proc. Natl. Acad. Sci. U. S. A. 99:2032–2037.
 21. Bulyha I, Schmidt C, Lenz P, Jakovljevic V, Höne A, Maier B, Hoppert M, Søgaard-Andersen L. 2009. Regulation of the type IV pili molecular machine by dynamic localization of two motor proteins. Mol. Microbiol. 74:691–706.
 22. Nudleman E, Wall D, Kaiser D. 2005. Cell-to-cell transfer of bacterial outer membrane lipoproteins. Science 309:125–127.
 23. Wei X, Pathak DT, Wall D. 2011. Heterologous protein transfer within structured myxobacteria biofilms. Mol. Microbiol. 81:315–326.
 24. Pathak DT, Wei X, Bucuvalas A, Haft DH, Gerloff DL, Wall D. 2012. Cell contact-dependent outer membrane exchange in myxobacteria: genetic determinants and mechanism. PLoS Genet. 8:e1002626.
 25. Caberoy NB, Welch RD, Jakobsen JS, Slater SC, Garza AG. 2003. Global mutational analysis of NtrC-like activators in *Myxococcus xanthus*: identifying activator mutants defective for motility and fruiting body development. J. Bacteriol. 185:6083–6094.
 26. Giglio KM, Caberoy N, Suen G, Kaiser D, Garza AG. 2011. A cascade of coregulating enhancer binding proteins initiates and propagates a multicellular developmental program. Proc. Natl. Acad. Sci. U. S. A. 108:E431–E439.
 27. Welch R, Kaiser D. 2001. Cell behavior in traveling wave patterns of myxobacteria. Proc. Natl. Acad. Sci. U. S. A. 98:14907–14912.
 28. Jakobsen JS, Jelsbak L, Jelsbak L, Welch RD, Cummings C, Goldman B, Stark E, Slater S, Kaiser D. 2004. Sigma54 enhancer binding proteins and *Myxococcus xanthus* fruiting body development. J. Bacteriol. 186:4361–4368.
 29. Curtis PD, Taylor RG, Welch RD, Shimkets LJ. 2007. Spatial organization of *Myxococcus xanthus* during fruiting body formation. J. Bacteriol. 189:9126–9130.
 30. Kroos L, Hartzell P, Stephens K, Kaiser D. 1988. A link between cell movement and gene expression argues that motility is required for cell-cell signaling during fruiting body development. Genes Dev. 2:1677–1685.
 31. MacNeil SD, Mouzeyan A, Hartzell PL. 1994. Genes required for both gliding motility and development in *Myxococcus xanthus*. Mol. Microbiol. 14:785–795.
 32. Youderian P, Burke N, White DJ, Hartzell PL. 2003. Identification of genes required for adventurous gliding motility in *Myxococcus xanthus* with the transposable element mariner. Mol. Microbiol. 49:555–570.
 33. Youderian P, Hartzell PL. 2006. Transposon insertions of Magellan-4 that impair social gliding motility in *Myxococcus xanthus*. Genetics 172:1397–1410.
 34. Goldman BS, Nierman WC, Kaiser D, Slater SC, Durkin AS, Eisen JA, Ronning CM, Barbazuk WB, Blanchard M, Field C, Halling C, Hinkle G, Iartchuk O, Kim HS, Mackenzie C, Madupu R, Miller N, Shvartsbeyn A, Sullivan SA, Vaudin M, Wiegand R, Kaplan HB. 2006. Evolution of sensory complexity recorded in a myxobacterial genome. Proc. Natl. Acad. Sci. U. S. A. 103:15200–15205.
 35. Wall D, Kolenbrander PE, Kaiser D. 1999. The *Myxococcus xanthus* pilQ (sglA) gene encodes a secretin homolog required for type IV pilus biogenesis, social motility, and development. J. Bacteriol. 181:24–33.
 36. O'Connor KA, Zusman DR. 1988. Reexamination of the role of autolysis in the development of *Myxococcus xanthus*. J. Bacteriol. 170:4103–4112.
 37. Berleman JE, Chumley T, Cheung P, Kirby JR. 2006. Rippling is a predatory behavior in *Myxococcus xanthus*. J. Bacteriol. 188:5888–5895.
 38. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. BMC Genomics 9:75.
 39. Chen HW, Kuspa A, Keseler IM, Shimkets LJ. 1991. Physical map of the *Myxococcus xanthus* chromosome. J. Bacteriol. 173:2109–2115.
 40. Chen H, Keseler IM, Shimkets LJ. 1990. Genome size of *Myxococcus xanthus* determined by pulsed-field gel electrophoresis. J. Bacteriol. 172:4206–4213.