



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Draft Genome of a Type 4 Pilus Defective *Myxococcus xanthus* Strain, DZF1

The Harvard community has made this article openly available.

[Please share](#) how this access benefits you. Your story matters.

Citation	Müller, Susanne, Jonathan W. Willett, Sarah M. Bahr, Jodie C. Scott, Janet M. Wilson, Cynthia L. Darnell, Hera C. Vlamakis, and John R. Kirby. 2013. "Draft Genome of a Type 4 Pilus Defective <i>Myxococcus xanthus</i> Strain, DZF1." <i>Genome Announcements</i> 1 (3): e00392-13. doi:10.1128/genomeA.00392-13. http://dx.doi.org/10.1128/genomeA.00392-13 .
Published Version	doi:10.1128/genomeA.00392-13
Accessed	April 17, 2018 4:35:52 PM EDT
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:11717575
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

Draft Genome of a Type 4 Pilus Defective *Myxococcus xanthus* Strain, DZF1

Susanne Müller,^a Jonathan W. Willett,^b Sarah M. Bahr,^a Jodie C. Scott,^c Janet M. Wilson,^d Cynthia L. Darnell,^a Hera C. Vlamakis,^e John R. Kirby^a

Department of Microbiology, University of Iowa, Iowa City, Iowa, USA^a; University of Chicago, Department of Biochemistry and Molecular Biology, Chicago, Illinois, USA^b; The Forsyth Institute, Department of Microbiology, Cambridge, Massachusetts, USA^c; Division of Select Agents and Toxins, Office of Public Health Preparedness and Response, Centers for Disease Control and Prevention, Atlanta, Georgia, USA^d; Harvard Medical School, Department of Microbiology and Immunobiology, Boston, Massachusetts, USA^e

S.M., J.W.W., and S.M.B. contributed equally to this work.

***Myxococcus xanthus* is a member of the *Myxococcales* order within the deltaproteobacterial subdivision. Here, we report the whole-genome shotgun sequence of the type IV pilus (T4P) defective strain DZF1, which includes many genes found in strain DZ2 but absent from strain DK1622.**

Received 9 May 2013 Accepted 13 May 2013 Published 20 June 2013

Citation Müller S, Willett JW, Bahr SM, Scott JC, Wilson JM, Darnell CL, Vlamakis HC, Kirby JR. 2013. Draft genome of a type 4 pilus defective *Myxococcus xanthus* strain, DZF1. *Genome Announc*. 1(3):e00392-13. doi:10.1128/genomeA.00392-13.

Copyright © 2013 Müller et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to John R. Kirby, john-kirby@uiowa.edu.

Myxococcus xanthus is a soil-dwelling deltaproteobacterium with a genome length of >9.2 Mb. The *Myxococcales* were described in the late 19th century for their capacity to produce macroscopic sporangioles or fruiting bodies (1). Organization into fruiting bodies requires extracellular signaling and coordination of two genetically distinct motility systems (2–29). Previously, the sequence of *M. xanthus* strain DK1622 was determined (NC_008095.1) (30). Recently, we sequenced strain DZ2 (31). These two laboratory strains are noted for behavioral differences (2, 32, 33), and DZ2 has a larger genome (31).

M. xanthus DZF1 is directly descended from the intermediate strain DK101, the progenitor for DK1622, and displays reduced capacity for type IV pilus (T4P)-mediated motility. An earlier study (34) mapped two point mutations in DK101 to *pilQ*, which encodes the T4P secretin (G741S/N762G), accounting for some phenotypic differences between DZ2 or DK1622 and those strains harboring the *pilQ* allele. Because the DZ2 genome contains additional genes relative to DK1622 and because differences in motility have been noted, we sequenced DZF1 to determine if the additional genes were more likely gained by DZ2 or lost from DK1622, relative to the progenitor.

M. xanthus DZF1 was sequenced at the University of Iowa DNA Core Facility using 454 GS-FLX titanium technology. Chromosomal DNA was prepared as described previously (31) and processed for sequencing following established protocols. The resulting sequence comprises 388,477 reads totaling 249 Mb, representing 27-fold coverage. The genome was assembled *de novo* into 75 contigs using Newbler software version 2.7. The resulting genome is approximately 9.28 Mb, similar to that for DZ2 (31), and is approximately 147 kb larger than the DK1622 genome. The RAST annotation server (35) predicts a total of 7,704 coding sequences (CDS) within the DZF1 genome.

The *M. xanthus* DZF1 sequence reveals a single nucleotide

polymorphism (SNP) in the *pilQ* gene, producing a G741S substitution, but lacks the N762G substitution found in DK101 (34). The impact of these SNPs has not been systematically determined but affects the interpretation of several previous studies. Indeed, deletion of *mazF* (encoding RNA interferase as part of a toxin-antitoxin system) is synthetic with the *pilQ* allele in both DZF1 and DK101 to affect cell death (36, 37).

Current analysis is ongoing to determine the role of genes found in DZF1, but not in DK1622, encoding proteins predicted to function in transcription, translation, signal transduction, fatty acid modification, and protein transport. Homologs to these genes are found in DZ2 as well as other myxobacteria, including *Myxococcus fulvus*, *Stigmatella aurantiaca*, and *Sorangium cellulosum*. The presence of sequences unique to both DZF1 and DZ2, while absent from DK1622, has been verified by PCR. Thus, the differences between DK1622 and both DZ2 and DZF1 are attributable to a loss of DNA from the DK1622 genome, likely following UV mutagenesis of DK101, which led to excision of one large prophage (38, 39) and may have induced additional lesions. We are investigating several unique sequences found in DZF1 and DZ2 for their role in *M. xanthus* biology.

Nucleotide sequence accession number. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [AOBT000000000](https://www.ncbi.nlm.nih.gov/nuclink/AB000000000). The version described in this paper is the first version.

ACKNOWLEDGMENTS

Support for this work was provided by the University of Iowa and NSF MCB-1244021 to J.R.K. Additional support for J.W.W. was provided by NIH T32 GM077973.

We thank Tom Willett, Tom Bair, and Kevin Knudtson for helpful discussions and data analysis.

The content is the responsibility of the authors and does not represent the official views of NIH, NSF, or the University of Iowa.

REFERENCES

- Thaxter R. 1892. On the Myxobacteriaceae, a new order of Schizomycetes Bot. Gaz. 17:389–406.
- Berleman JE, Kirby JR. 2007. Multicellular development in *Myxococcus xanthus* is stimulated by predator-prey interactions. J. Bacteriol. 189:5675–5682.
- Blackhart BD, Zusman DR. 1985. “Frizzy” genes of *Myxococcus xanthus* are involved in control of frequency of reversal of gliding motility. Proc. Natl. Acad. Sci. U. S. A. 82:8767–8770.
- Bulyha I, Schmidt C, Lenz P, Jakovljevic V, Höne A, Maier B, Hoppert M, Søgaard-Andersen L. 2009. Regulation of the type IV pili molecular machine by dynamic localization of two motor proteins. Mol. Microbiol. 74:691–706.
- Caberoy NB, Welch RD, Jakobsen JS, Slater SC, Garza AG. 2003. Global mutational analysis of NtrC-like activators in *Myxococcus xanthus*: identifying activator mutants defective for motility and fruiting body development. J. Bacteriol. 185:6083–6094.
- Campos JM, Zusman DR. 1975. Regulation of development in *Myxococcus xanthus*: effect of 3':5'-cyclic AMP, ADP, and nutrition. Proc. Natl. Acad. Sci. U. S. A. 72:518–522.
- Giglio KM, Caberoy N, Suen G, Kaiser D, Garza AG. 2011. A cascade of coregulating enhancer binding proteins initiates and propagates a multicellular developmental program. Proc. Natl. Acad. Sci. U. S. A. 108:E431–E439.
- Huntley S, Hamann N, Wegener-Feldbrügge S, Treuner-Lange A, Kube M, Reinhardt R, Klages S, Müller R, Ronning CM, Nierman WC, Søgaard-Andersen L. 2011. Comparative genomic analysis of fruiting body formation in Myxococcales. Mol. Biol. Evol. 28:1083–1097.
- Jelsbak L, Søgaard-Andersen L. 2002. Pattern formation by a cell surface-associated morphogen in *Myxococcus xanthus*. Proc. Natl. Acad. Sci. U. S. A. 99:2032–2037.
- Kirby JR, Zusman DR. 2003. Chemosensory regulation of developmental gene expression in *Myxococcus xanthus*. Proc. Natl. Acad. Sci. U. S. A. 100:2008–2013.
- Kroos L, Hartzell P, Stephens K, Kaiser D. 1988. A link between cell movement and gene expression argues that motility is required for cell-cell signaling during fruiting body development. Genes Dev. 2:1677–1685.
- Kruse T, Lobedan S, Berthelsen NM, Søgaard-Andersen L. 2001. C-signal: a cell surface-associated morphogen that induces and coordinates multicellular fruiting body morphogenesis and sporulation in *Myxococcus xanthus*. Mol. Microbiol. 40:156–168.
- Luciano J, Agrebi R, Le Gall AV, Wartel M, Fiegna F, Ducret A, Brochier-Armanet C, Mignot T. 2011. Emergence and modular evolution of a novel motility machinery in bacteria. PLoS Genet. 7:e1002268.
- MacNeil SD, Mouzeyan A, Hartzell PL. 1994. Genes required for both gliding motility and development in *Myxococcus xanthus*. Mol. Microbiol. 14:785–795.
- Mauriello EM, Mignot T, Yang Z, Zusman DR. 2010. Gliding motility revisited: how do the myxobacteria move without flagella? Microbiol. Mol. Biol. Rev. 74:229–249.
- Mignot T. 2007. The elusive engine in *Myxococcus xanthus* gliding motility. Cell. Mol. Life Sci. 64:2733–2745.
- Mignot T, Kirby JR. 2008. Genetic circuitry controlling motility behaviors of *Myxococcus xanthus*. BioEssays 30:733–743.
- Mignot T, Merlie JP, Jr, Zusman DR. 2005. Regulated pole-to-pole oscillations of a bacterial gliding motility protein. Science 310:855–857.
- Mignot T, Shaevitz JW, Hartzell PL, Zusman DR. 2007. Evidence that focal adhesion complexes power bacterial gliding motility. Science 315:853–856.
- Nudleman E, Wall D, Kaiser D. 2005. Cell-to-cell transfer of bacterial outer membrane lipoproteins. Science 309:125–127.
- Pathak DT, Wall D. 2012. Identification of the *cglC*, *cglD*, *cglE*, and *cglF* genes and their role in cell contact-dependent gliding motility in *Myxococcus xanthus*. J. Bacteriol. 194:1940–1949.
- Shimkets LJ, Gill RE, Kaiser D. 1983. Developmental cell interactions in *Myxococcus xanthus* and the *spoC* locus. Proc. Natl. Acad. Sci. U. S. A. 80:1406–1410.
- Singer M, Kaiser D. 1995. Ectopic production of guanosine penta- and tetraphosphate can initiate early developmental gene expression in *Myxococcus xanthus*. Genes Dev. 9:1633–1644.
- Wei X, Pathak DT, Wall D. 2011. Heterologous protein transfer within structured myxobacteria biofilms. Mol. Microbiol. 81:315–326.
- Welch R, Kaiser D. 2001. Cell behavior in traveling wave patterns of myxobacteria. Proc. Natl. Acad. Sci. U. S. A. 98:14907–14912.
- Youderian P, Burke N, White DJ, Hartzell PL. 2003. Identification of genes required for adventurous gliding motility in *Myxococcus xanthus* with the transposable element mariner. Mol. Microbiol. 49:555–570.
- Youderian P, Hartzell PL. 2006. Transposon insertions of Magellan-4 that impair social gliding motility in *Myxococcus xanthus*. Genetics 172:1397–1410.
- Zhang Y, Ducret A, Shaevitz J, Mignot T. 2012. From individual cell motility to collective behaviors: insights from a prokaryote, *Myxococcus xanthus*. FEMS Microbiol. Rev. 36:149–164.
- Zusman DR, Scott AE, Yang Z, Kirby JR. 2007. Chemosensory pathways, motility and development in *Myxococcus xanthus*. Nat. Rev. Microbiol. 5:862–872.
- Goldman BS, Nierman WC, Kaiser D, Slater SC, Durkin AS, Eisen JA, Ronning CM, Barbazuk WB, Blanchard M, Field C, Halling C, Hinkle G, Iartchuk O, Kim HS, Mackenzie C, Madupu R, Miller N, Shvartsbeyn A, Sullivan SA, Vaudin M, Wiegand R, Kaplan HB, Kaplan HB. 2006. Evolution of sensory complexity recorded in a myxobacterial genome. Proc. Natl. Acad. Sci. U. S. A. 103:15200–15205.
- Müller S, Willett JW, Bahr SM, Darnell CL, Hummels KR, Dong CK, Vlamakis HC, Kirby JR. 2013. Draft genome sequence of *Myxococcus xanthus* wild-type strain DZ2, a model organism for predation and development. Genome Announc. 1(3):e00217-13. doi:10.1128/genomeA.00217-13.
- Berleman JE, Chumley T, Cheung P, Kirby JR. 2006. Rippling is a predatory behavior in *Myxococcus xanthus*. J. Bacteriol. 188:5888–5895.
- O'Connor KA, Zusman DR. 1988. Reexamination of the role of autolysis in the development of *Myxococcus xanthus*. J. Bacteriol. 170:4103–4112.
- Wall D, Kolenbrander PE, Kaiser D. 1999. The *Myxococcus xanthus* *pilQ* (*sglA*) gene encodes a secretin homolog required for type IV pilus biogenesis, social motility, and development. J. Bacteriol. 181:24–33.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formisano K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. BMC Genomics 9:75.
- Boynton TO, McMurry JL, Shimkets LJ. 2013. Characterization of *Myxococcus xanthus* MazF and implications for a new point of regulation. Mol. Microbiol. 87:1267–1276.
- Lee B, Holkenbrink C, Treuner-Lange A, Higgs PI. 2012. *Myxococcus xanthus* developmental cell fate production: heterogeneous accumulation of developmental regulatory proteins and reexamination of the role of MazF in developmental lysis. J. Bacteriol. 194:3058–3068.
- Chen H, Keseler IM, Shimkets LJ. 1990. Genome size of *Myxococcus xanthus* determined by pulsed-field gel electrophoresis. J. Bacteriol. 172:4206–4213.
- Chen HW, Kuspa A, Keseler IM, Shimkets LJ. 1991. Physical map of the *Myxococcus xanthus* chromosome. J. Bacteriol. 173:2109–2115.