



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Microvesicles and intercellular communication in the context of parasitism

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters.

Citation	Barteneva, Natasha S., Natalia Maltsev, and Ivan A. Vorobjev. 2013. "Microvesicles and intercellular communication in the context of parasitism." <i>Frontiers in Cellular and Infection Microbiology</i> 3 (1): 49. doi:10.3389/fcimb.2013.00049. http://dx.doi.org/10.3389/fcimb.2013.00049 .
Published Version	doi:10.3389/fcimb.2013.00049
Accessed	April 17, 2018 4:38:10 PM EDT
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:11877146
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)



Microvesicles and intercellular communication in the context of parasitism

Natasha S. Barteneva^{1*}, Natalia Maltsev^{2*} and Ivan A. Vorobjev^{3,4}

¹ Program in Cellular and Molecular Medicine, Children's Hospital Boston and Department of Pediatrics, Harvard Medical School, Boston, MA, USA

² Department of Human Genetics, University of Chicago, Chicago, IL, USA

³ A.N. Belozersky Institute of Physico-Chemical Biology, Lomonosov Moscow State University, Moscow, Russia

⁴ Department of Cell Biology and Histology, Faculty of Biology, Lomonosov Moscow State University, Moscow, Russia

Edited by:

Robert Heinzen, National Institutes of Health, USA

Reviewed by:

Stacey Gilk, Indiana University School of Medicine, USA

Antonio Marcilla, Universitat de Valencia, Spain

*Correspondence:

Natasha S. Barteneva, Program in Cellular and Molecular Medicine, 200 Longwood Avenue, Boston, MA, 02115, USA

e-mail: Natasha.Barteneva@childrens.harvard.edu;

Natalia Maltsev, Department of Human Genetics, University of Chicago, E. 58th str., Chicago, IL 60637, USA

e-mail: maltsev@uchicago.edu

There is a rapidly growing body of evidence that production of microvesicles (MVs) is a universal feature of cellular life. MVs can incorporate microRNA (miRNA), mRNA, mtDNA, DNA and retrotransposons, camouflage viruses/viral components from immune surveillance, and transfer cargo between cells. These properties make MVs an essential player in intercellular communication. Increasing evidence supports the notion that MVs can also act as long-distance vehicles for RNA molecules and participate in metabolic synchronization and reprogramming eukaryotic cells including stem and germinal cells. MV ability to carry on DNA and their general distribution makes them attractive candidates for horizontal gene transfer, particularly between multi-cellular organisms and their parasites; this suggests important implications for the co-evolution of parasites and their hosts. In this review, we provide current understanding of the roles played by MVs in intracellular pathogens and parasitic infections. We also discuss the possible role of MVs in co-infection and host shifting.

Keywords: microvesicles, exosomes, miRNA, parasite, metabolism synchronization, horizontal gene transfer, co-infection, *Plasmodium*

INTRODUCTION

Production of membrane-enclosed microvesicles (MVs) is a universal feature of cellular life and has been demonstrated for organisms as diverse as Proteobacteria, Archaea, plants, and animals (Ellis and Kuehn, 2010; Silverman and Reiner, 2012; Deatherage and Cookson, 2012). Several distinct categories of membrane-enclosed MVs exist, including exosomes, ectosomes, and apoptotic bodies (in multi-cellular organisms). MVs are grouped based on their size, density, method of isolation, and markers, and purified MVs usually represent a mixture of aforementioned vesicular fractions.

Secretion of MVs is well-documented for prokaryotic and eukaryotic cells (György et al., 2011; Silverman and Reiner, 2012), and in infected organisms they can contain both host and parasitic antigens. Vesicles from a number of pathogens, such as *Leishmania*, *Cryptococcus*, and *Trypanosoma*, may carry on virulence factors and participate in their delivery to host

cells (Silverman and Reiner, 2011, 2012; Lambertz et al., 2012; Torrecilhas et al., 2012), promoting dissemination of the pathogen.

Though it has become clear that MVs possess immunomodulatory features, little is known about the role of MVs in host-parasite co-existence and co-evolution. We anticipate that recent findings regarding the participation of MVs in the transfer of genetic information will expand the functions attributed to MVs in the host-parasite evolution. We will hypothesize about the role MVs play, as a vehicle for regulatory molecules important for synchronization of host and parasite metabolism, and for delivery of nucleic acids.

MICROVESICLES ARE IMPORTANT INTERCELLULAR COMMUNICATORS

MVs are considered a universal transport vehicle for intercellular communication. MVs incorporate peptides, proteins, lipids, miRNA, and mRNA, all of which can be transferred and become functional in target cells (Ratajczak et al., 2006; Valadi et al., 2007; Skog et al., 2008; Iglesias et al., 2012). MVs bind to cells through receptor-ligand interactions, fuse with target cell membranes, and deliver their cargo to the cytoplasm of the target cell. As has been observed with tumor-derived MVs, MVs can be enriched in specific coding and non-coding RNAs, chromosomal and mitochondrial DNA, retrotransposon RNA, and Alu transposon elements (Ronquist et al., 2009, 2012; Guescini et al., 2010; Balaj et al., 2011; Rak and Guha, 2012; Waldenstroem et al., 2012).

Abbreviations: 7-AAD, 7-aminoactinomycin D; EAE, experimental allergic encephalomyelitis; EGFP, enhanced green fluorescent protein; GFP, green fluorescent protein; HDM, helminth defense molecules; HDP, host defense peptide; HGT, horizontal gene transfer; LDLRAP1 protein, low density lipoprotein receptor adapter protein; MAPK/ERK pathway, mitogen-activated protein kinases/Extracellular signal-regulated kinases pathway; miRNA, microRNA; mtDNA, mitochondrial DNA; MV, microvesicles; NETs, neutrophil extracellular traps; NETosis, *in vivo* NETs are released during a form of pathogen-induced cell death, which was recently named NETosis; ORF, open reading frame; PV, parasitophorous vacuole; PVM, parasitophorous vacuole membrane; ROS, reactive oxygen species.

Transfer of functional genetic information by MVs was initially shown in the experiments with a reporter mRNA encoding GFP (Deregibus et al., 2007), where intact RNA transcripts capable of serving as templates for protein translation were enriched in shed MVs (Li et al., 2012).

MVs are considered the major “miRNA transporter” between cells, since most extracellular miRNAs are found in vesicles (Gallo et al., 2012; Xu et al., 2013). miRNAs have been identified in helminthes and in protozoa possessing *Argonaute/Dicer* genes, while they are absent in protozoa lacking enzymes required for RNAi-based interference, such as *Plasmodium* spp and *Cryptosporidium* (Baum et al., 2009; Manzano-Roman and Siles-Lucas, 2012). MV-mediated export of miRNA is selective (Zhang et al., 2010a; Jaiswal et al., 2012; Vickers and Remaley, 2012). mRNA and miRNA packaged in vesicles appear to be more stable and resistant to RNase digestion in the body fluids, due to the lipid membrane of MVs (Li et al., 2012; Vickers and Remaley, 2012).

During their release, MVs may incorporate components that are originally alien to the cell, such as proteins and nucleic acids that are transiently or constitutively expressed via plasmid or viral vector. Recently, it was shown that exogenous plant miRNA is present in human plasma and animal tissues. These results invoked the idea that miRNAs could regulate gene expression across the kingdoms (Kosaka and Ochiya, 2011; Zhang et al., 2012). We speculate that miRNA derived from the bacterial gut community may be packaged by epithelial cells into MVs and then delivered to different parts of the body, starting with the liver. A recent intriguing finding by two independent groups of the exogenous RNAs of different origin in human plasma samples (Bacteria and Archaea, Fungi, Plants—Wang et al., 2012; microbial RNA sequences—Semenov et al., 2012) supports this assertion.

Overall, MVs as vehicles for miRNA and other regulatory molecules, such as regulatory sequences of mRNA, may play important role in the synchronization of metabolism between the host and its parasites.

MICROVESICLE PROTEOMICS

The production of MVs rises sharply in many parasitic and infectious diseases (Campos et al., 2010; Barteneva et al., 2013; **Table 1**). Proteins identified in these MVs were related to vesicle trafficking, signaling molecules and transmembrane small channels and transporters (Rodrigues et al., 2008; Silverman et al., 2010a). We recently showed that significant percentage of proteins identified in MVs during malaria infection belong to classical and alternative complement pathway, components of cytoskeleton, glycolysis and lipid transport (Mantel et al., 2013). Metabolic enzymes related to glycolysis constitute the largest protein family in excretory/secretory proteome of helminth *E. caproni* (Sotillo et al., 2010). Some glycolytic enzymes in parasitic MVs have separate function that make them important for parasite survival and dissemination (for example, binding of plasminogen for enolase in *Leishmania*) (Chandra et al., 2010). Furthermore, the MVs-production may explain the presence of atypical proteins lacking

classical secretion signal peptides, like enolase, in the parasite secretions.

Parasite-induced MVs also contain constituent host proteins different depending on the species of parasite. For example, while mucin-2 was found in *E. caproni* vesicles, only CD19 and the constant region of the IgA heavy chain were found in *Fasciola hepatica* vesicles (Wilson et al., 2011; Marcilla et al., 2012). Conversely, the same helminth species when develops in several intermediate hosts exhibit host adaptation via differential expression of certain gene families (example: antigen B gene family from *E. granulosa*) in subsequent life cycle stages (Mamuti et al., 2007; Zhang et al., 2010b), however, no parasite proteomes from MVs produced in different intermediate hosts are currently available. MVs production increased during different developmental stages of parasites and proportion of specific antigens may be changed [as shown for RESA-antigen, during ring-stage, trophozoite and shizont stages of *P. falciparum* development (Natakamol et al., 2011)].

In sum, extracellular MVs contain parasite-specific excretory/secretory proteins (Silverman et al., 2010a; Marcilla et al., 2012), often lacking signal sequences (*Leishmania*), and participate in delivery of virulence factors and regulation of parasite virulence (Silverman and Reiner, 2011; Torrecilhas et al., 2012). Majority of proteome studies of parasite-produced MVs identified virulence factors in the MVs proteomes (Geiger et al., 2010; Silverman et al., 2010a; Bayer-Santos et al., 2013). MVs deliver virulence factors such as toxins, proteases, adhesins (Amano et al., 2010; Torrecilhas et al., 2012), *Entamoeba histolytica* rhomboid protease (EhROM1) (Baxt et al., 2008), participate in regulation of gene expression and help to escape immune evasion (Lambertz et al., 2012).

IMMUNOMODULATORY ACTIVITIES OF MICROVESICLES AND MOLECULAR MIMICRY

Parasites have developed many strategies that support their transmission and allow them to survive and reproduce, such as development of novel cellular pathways that enable invasion into different hosts and diverse immune evasion strategies including: alteration of host antigens, establishment of self-tolerance, functional immune inactivation, immunosuppression, molecular mimicry between parasite polypeptides and host antigens, acquisition of sialic acid motifs from host cells and adsorption of host serum sialoglycoconjugates leading to the modulation of NETosis (Hahn et al., 2013), and antigenic variability regulated by parasite methyltransferases (Figueiredo et al., 2008). Production of MVs appears to be involved in many of these processes.

Molecular mimicry as a strategy for host manipulation and evasion of immune response is well-known in viruses because of their ability to acquire host proteins or genetic material during virion assembly (Alcami, 2003; Bernet et al., 2003). There is significant evidence that autoimmune disease can develop after bacterial or parasitic infection, such as Chagas disease, where a cross-reaction between cardiac muscle cells and *T. cruzi* occurs (Acosta and Santos-Buch, 1985; Sepulveda et al., 2000). Recently, molecular mimicry between a family of peptides produced by trematode helminthes, and human defense peptides, including defensins and cathelicidins was found (Robinson et al., 2011). This family of helminth defense molecules (HDMs) is conserved

Table 1 | Microvesicles produced in response to different parasitic pathogens.

Pathogen	Type of microvesicles (according to publication authors)	References
FUNGI		
<i>Cryptococcus neoformans</i>	Exosomes	Yoneda and Doering, 2006; Rodrigues et al., 2008; Nicola et al., 2009; Panepinto et al., 2009; Oliveira et al., 2010; Huang et al., 2012
<i>Malassezia sympodialis</i>	Exosomes	Gehrmann et al., 2011
<i>Paracoccidioides</i>	Conditioned medium(secreted proteins and vesicles)	Weber et al., 2012
<i>Paracoccidioides brasiliensis</i>		Vallejo et al., 2011, 2012a,b
PROTOZOA		
<i>Giardia lamblia</i>	Secretory vesicles	Benchimol, 2004; Gottig et al., 2006
<i>Leishmania</i>	Exosomes from infected macrophages	Silverman et al., 2010a,b; Silverman and Reiner, 2011; Figuera et al., 2012; Hassani and Olivier, 2013
<i>Plasmodium vivax</i>	Plasma-derived MPs	Campos et al., 2010
<i>Plasmodium berghei</i>	Plasma-derived MPs (from infected mice)	Combes et al., 2005; Couper et al., 2010
<i>Plasmodium falciparum</i>	Vesicles(60–100 nm); microvesicles (100–1000 nm)	Trelka et al., 2000; Bhattacharjee et al., 2008; Mantel et al., 2013; Regev-Rudzki et al., 2013
<i>Plasmodium yoelii</i>	Plasma-derived exosomes	Martin-Jaular et al., 2011
<i>Toxoplasma gondii</i>	Exosomes	Bhatnagar et al., 2007
<i>Trypanosoma brucei</i>	Exosomes	Geiger et al., 2010
<i>Trypanosoma cruzi</i>	Outer membrane-derived vesicles, exosomes	Goncalves et al., 1991; Ouassi et al., 1992; Trocoli Torrecilhas et al., 2009; Cestari et al., 2012; Bayer-Santos et al., 2013
MYCOPLASMA		
<i>Mycoplasma</i>	Exosomes	Quah and O'Neill, 2007; Yang et al., 2012
BACTERIA		
<i>Borrelia burgdoferi</i>	Ectosomes (outer membrane vesicles)	Toledo et al., 2012
<i>Brucella abortus</i>	Ectosomes (outer membrane vesicles)	Pollak et al., 2012
<i>Chlamydia trachomatis</i>	Exosomes, outer membrane vesicles	Zhong, 2011; Frohlich et al., 2012
<i>Francisella novacida</i>		Pierson et al., 2011
<i>Legionella pneumophila</i>	Membrane vesicles	Galka et al., 2008
<i>Mycobacterium tuberculosis</i>	Exosomes; shedding microvesicles	Giri et al., 2010; Ramachandra et al., 2010; Singh et al., 2011, 2012; Duarte et al., 2012
<i>Mycobacterium avium</i>	Exosomes	Bhatnagar and Schorey, 2007
<i>Mycobacterium bovis</i>	Exosomes	Giri and Schorey, 2008
<i>Salmonella thyphimurium</i>	Outer membrane-derived vesicles	Yoon et al., 2011
HELMINTHS		
<i>Caenorhabditis elegans</i>	Exosomes	Liegeois et al., 2006
<i>Echinostoma caproni</i>	Exosomes	Andresen et al., 1989; Marcilla et al., 2012
<i>Echinococcus multilocularis</i>	Vesicles derived from metacestodes	Eger et al., 2003; Walker et al., 2004; Huebner et al., 2006; Nono et al., 2012
<i>Fasciola hepatica</i>	Exosomes	Marcilla et al., 2012

throughout trematodes, and these proteins participate in the host immune response modulation and anti-inflammatory action (Robinson et al., 2011).

Infections caused by intracellular pathogens and parasites are often chronic and lead to significant immunomodulation of host immune response by the parasite. MVs produced during protozoan infections were shown to participate in this process (Bhatnagar and Schorey, 2007; Bhatnagar et al., 2007; Barreto et al., 2010; Silverman et al., 2010a; Hassani and Olivier, 2013). For example, during *Plasmodium* infection, there are increased quantities of MVs in plasma, and they contain a significant amount of parasite material. These MVs induce neutrophil activation (Mantel et al., 2013) and strong pro-inflammatory activation of macrophages as measured by CD40 and TNF up-regulation (Couper et al., 2010). Besides, vesiculation, which utilizes host cell machinery, is an important mechanism for parasite egress in the case of *P. falciparum*, the cause of malaria and a member of the phylum *Apicomplexa* (Lew, 2011). Secreted vesicles, which in the case of helminthes, present among other parasite secretion products, have been shown to modulate host immune responses and strongly influence the outcome of infections to the parasite's advantage (Spolski et al., 2000; Allen and MacDonald, 1998; Silverman et al., 2010b).

In our recent publication more than thirty parasite proteins in MVs derived from red blood cells infected with 3D7 or CS2 strains of *P. falciparum* were identified (Mantel et al., 2013). A modified approach, first described by Ludin et al. (2011) was employed, allowing for rigorous analysis of *P. falciparum* proteins that may potentially contribute to the infectious process via molecular mimicry of host molecules. Identified potential candidates include erythrocyte-binding proteins 1, 2, and 3, liver-stage antigen, and others (e.g., Rex2) (manuscript in preparation). **Figures 1A–C** shows the similarity between the *P. falciparum* short (119 amino acids) PEXEL (*Plasmodium* export element)-negative ring-exported protein 2 (Rex2) and the *H. sapiens* Rac1 and Rac2 proteins, providing one example of possible molecular mimicry and parasite-human HGT in parasitic invasion (**Figure 1D**). It was previously demonstrated (Haase et al., 2009) that a short sequence in the N-terminus and transmembrane domain of the Rex2 protein are both required for parasite export. The N-terminus of Rex2 exhibits significant similarity to the human small GTP-ases Rac1 and Rac2. A number of studies have demonstrated that deleterious mutations in Rac2 lead to defective chemotaxis, impaired phagocytosis, and decreased pathogen killing by macrophages and/or neutrophils (Roberts et al., 1999; Koh et al., 2005; Yamauchi et al., 2005; Zhang et al., 2009). Because it has been reported that neutrophils from malaria patients have reduced chemotactic activity (Nielsen et al., 1986; Leoratti et al., 2012), a role for Rex2 in molecular mimicry of Rac2 is likely. It is anticipated that *in silico* analysis of other pathogen-derived MV-associated proteins will be helpful in further understanding how MVs function as intercellular communicators during disease states, and provide insights on what to base future experimental studies on.

Molecular mimicry may be a more prevalent parasitic strategy than was previously thought (Ludin et al., 2011). Acquisition of complete nucleotide sequences or sequence motifs from the host

may happen at different stages of parasite-host co-existence, and MVs may play a significant role in this molecular exchange.

DO MICROVESICLES PARTICIPATE IN CO-INFECTION?

Parasitic and symbiotic associations are ubiquitous and often life-long relationships (Eckburg et al., 2005; Weiss and Aksoy, 2011). Every mammal possesses complex microbial communities that reside on all mucosal surfaces. The human gastrointestinal tract harbors an estimated 1014 species of microbes from over 500 distinct microbial taxa (Eckburg et al., 2005). Although infectious parasite biology research is still dominated by studies of single infections poly-parasitism is very common in nature (Petney and Andrews, 1998; Bordes and Morand, 2009). Infection with one parasitic species can have a large impact on host susceptibility to secondary infection, and this phenomenon is partially dependent on the duration of infection (Telfer et al., 2010).

There is a growing body of evidence that concurrent parasitic infections can confer benefits to their hosts. For example, *Wolbachia*, which is considered a reproductive parasite in arthropods (Werren et al., 2008), can provide metabolic advantages to their hosts during stressful conditions, such as increased haem and riboflavin availability (Brownlie et al., 2009; Hosokawa et al., 2010). There are a number of studies outlining associations between different parasitic co-infections. Thus, helminth infection can prevent or suppress autoimmune and allergic diseases depending on helminth burden (reviewed in Zaccone and Cooke, 2013). A similar finding has been described for malaria infection, where the disease can be asymptomatic or less severe, with concomitant lower parasitaemia, in helminth-infected patients (Adegunica and Kremsner, 2012). The infection of *Schistosoma haematobium* is associated with protection against acute *P. falciparum* infection (Lyke et al., 2005).

The role of MVs as messengers between parasite and host immune cells is well-established (Silverman and Reiner, 2012). We have shown involvement of MVs in cross-communication within a *P. falciparum* population (Mantel et al., 2013). The exchange of MVs derived from different parasites as well as from host cells can participate in the mechanisms of co-infection (**Figure 1E**). However, there are currently no published reports describing the communication of MVs derived from different parasites and the issue deserves in-depth elucidation.

SEARCHING FOR NEW CLUES IN INTERCELLULAR COMMUNICATION BY *IN SILICO* GENOMICS AND PROTEOMICS

Amoebae, as well as other free-living protozoan hosts for bacteria, fungi, giant DNA viruses and virophages are “melting pots” for HGT exchanges (Hotopp et al., 2007; Moliner et al., 2010; Raoult and Boyer, 2010; Lamrabet et al., 2012). Besides, DNA exchange may also occur in reverse from microorganisms to protozoa (Ricard et al., 2006), and to animals (McNulty et al., 2010; Dunning Hotopp, 2011). Examples of gene transfer from the animal host to the ancestor of the apicomplexan parasites include genes encoding proteins involved in cell adhesion, O-linked glycosylation and a major epigenetic regulator histone methyltransferase Set8 (Kishore et al., 2013). In many instances, interdomain HGT involves transfers between endosymbiotic bacteria and their

FIGURE 1 | Continued

cells, and epithelial cell polarization. Rac2 activity also includes regulation of human neutrophil NADPH oxidase and activation of the production of reactive oxygen species (ROS). **(B)** Three-dimensional structure of human Rac2 (critical aminoacids are shown with numbers). **(C)** Sequence alignment between *P. falciparum* Rex2 and Rac2 and Rac1 proteins reveals an exact match in a number of functionally important amino acids positions, including (a) Asp57. RAC2 Asp57Asn mutation has been shown to be associated with severely impaired fMLP- or IL8-induced neutrophil responsiveness, including adhesion, chemotaxis, and superoxide production. The Asp57-mutant Rac2 does not bind GTP and was found to act in a dominant-negative fashion for both Rac1 and Rac2 because of its tight GEF binding; (b) H103, which is involved in Rac1-mediated oxidase activation, and (c) the ubiquitination sites K96 and K123. The alignment also revealed exact matches found exclusively between the Rac1 and Rex2 proteins, including G48, F90 and A151. However, the functional impact of these amino acids is not yet known. **(D)** PSI-Blast

analysis of *Plasmodium falciparum* 3D7 Rex2 protein against NCBI non-redundant database showed weak similarities to hypothetical proteins from parasitic *Apicomplexa* *Theileria annulata*, ciliate *Protozoa Oxytricha trifallax*, *Felis catus* (gi410982116), *Drosophila virilis* (gi195395466), as well as mouse Nrde2 protein (gi19344080). Multiple sequence alignment of these proteins was developed using NCBI Cobalt (Papadopoulos and Agarwala, 2007). Maximum likelihood phylogenetic tree was developed using the iTOL server and default parameters (Letunic and Bork, 2011). As it follows from the tree the Rex2 protein most closely evolutionary relates to a hypothetical protein from algae *Heterosigma akashiwo* and Human Rac2 protein. Evolutionary relations between algae and *Apicomplexa* are well-established (Lemgruber et al., 2013), however, relatedness to Human Rac2 protein suggests HGT from parasite to Human. **(E)** A hypothetical scheme of MVs exchange in parasite-host interaction. Host and multiple parasites produce and exchange microvesicles, which transfer lipids, proteins, nucleic acids such as miRNA, mRNA, DNA, and may camouflage virions.

hosts and from bacteria to asexual animals (Dunning Hotopp, 2011). For example, *Wolbachia*-to-arthropod HGT has been seen in the genomes of the bean beetle (Coleoptera) (Kondo et al., 2002), mosquitoes (Diptera) (Klasson et al., 2009; Woolfit et al., 2009), and other arthropods, as well as HGT in the opposite direction—from arthropod to *Wolbachia* (Duploux et al., 2013). Some *Microsporidia* species acquired a gene from arthropods that encodes purine nucleotide phosphatase, though most HGT to *Microsporidia* identified to date derives from prokaryotes (Selman and Corradi, 2011).

Analysis of *T. cruzi* ribosomal proteins *in silico* identified significant homology not only with members of the animal kingdom (*H. sapiens*, *C. elegans*, *D. melanogaster*), but also with plants and protozoa (Wayengera, 2009). Recent massive bioinformatics analyses of whole genome sequences have shown that many intracellular prokaryotes have the ability to manipulate the eukaryotic ubiquitin system through molecular mimicry of the F-box component of the SCF E3-ubiquitin ligase eukaryotic-like F-box proteins (Price et al., 2009; Price and Kwaik, 2010).

The ability of MVs to serve as vehicles not only for proteins and lipids, but also for nucleic acids (Ronquist et al., 2009, 2012; Guescini et al., 2010; Balaj et al., 2011; Pisetsky et al., 2011; Rak and Guha, 2012), make us hypothesize that MV production and exchange are an important mechanisms in gene information exchange between parasites and their hosts. As described above, this is indirectly confirmed by a number of similarities between parasite protein sequences and host molecules. Recent studies of mammalian (equine) ovarian follicles revealed that intercellular communication in the ovarian follicle may involve transfer of miRNA and other bioactive molecules by MVs between follicular fluid and granulosa cells (da Silveira et al., 2012). In addition, the transfer of chromosomal DNA fragments by prostate-derived MVs to human sperm (vertical transfer) was described by Ronquist et al. (2009). Moreover, MVs camouflage viruses from immune surveillance and facilitate their access to cells (Kadiu et al., 2012).

Thus, we hypothesize that one of the important functions of MVs in parasite-host and parasite-parasite co-evolution is their participation in HGT. This hypothesis is based on the following properties of MVs: (1) the ability to transfer nucleic acids, including DNA; (2) the capability of MVs to protect and deliver

genetic information to different organs, including reproductive organs in the case of multi-cellular organisms; and (3) the ability to transfer cargo to alien cells and tissues (in the case of parasites). Regev-Rudzki et al. (2013) recently provided additional support to this hypothesis confirming that *P. falciparum* derived MVs are capable of delivering genes between parasite populations inside their host.

CONCLUDING REMARKS

MVs are emerging as critical players in HGT including small non-coding RNAs. Major challenges in the field of extracellular vesicle research include (1) development of new comprehensive methods for their isolation and characterization, and (2) isolation of pure populations of specific MVs. Improved understanding of the mechanisms involved in vesicle shuttling of genetic information and proteins is crucial in order for new diagnostic and therapeutic strategies to be designed and implemented.

Dissemination of parasite components with MVs provides (1) a unique advantage in protection against host-mediated immune responses and nucleases from blood and other body fluids, (2) the possibility of reaching distant regions and evading immune attacks due to their small size (<1 μm), and (3) the capability of transferring genetic information long-distance—this may lead to direct participation of pathogenic components in the regulation of gene expression in the different host cells and metabolism synchronization between host and parasite, as well as HGT. This may contribute to further co-adaptation and co-evolution of the parasite and its host. Remarkably, MV exchange may happen similarly in dramatically different animal host cells, as well as in simple multicellular organisms. This can lead to broader host ranges and an increase in the virulence of certain parasites.

Our present knowledge of MVs derived from parasites comes from studies involving limited numbers of parasites and hosts. In the future our understanding of how parasite-host exchange of regulatory molecules and genetic information happens may change, especially when taking into account interactions between multiple parasitic species within the same host organism.

ACKNOWLEDGMENTS

We are thankful to Dr. Luke Jasenosky for critical reading and Aleksandra Gorelova (Harvard University) for editorial

help. Natasha S. Barteneva was supported by Harvard Pilot Grant and Program in Cellular and Molecular Medicine, and Ivan A. Vorobjev was supported by Russian Foundation for Basic Research grants 11-04-01749a and 13-04-40189H. Natalia

Maltsev is grateful to Mr. and Ms. Lawrence Hilibrand and the Boler Family Foundation for their generous support of the project. Space considerations limited us to a selected list of available literature.

REFERENCES

- Acosta, A. M., and Santos-Buch, C. A. (1985). Autoimmune myocarditis induced by *Trypanosoma cruzi*. *Circulation* 71, 1255–1261. doi: 10.1161/01.CIR.71.6.1255
- Adegnica, A. A., and Kremsner, P. G. (2012). Epidemiology of malaria and helminth interaction: a review from 2001 to 2011. *Curr. Opin. HIV AIDS* 7, 221–224. doi: 10.1097/COH.0b013e3283524d90
- Alcami, A. (2003). Viral mimicry of cytokines, chemokines and their receptors. *Nat. Rev. Immunol.* 3, 36–50. doi: 10.1038/nri980
- Allen, J. E., and MacDonald, A. S. (1998). Profound suppression of cellular proliferation mediated by the secretions of nematodes. *Parasite Immunol.* 20, 241–247. doi: 10.1046/j.1365-3024.1998.00151.x
- Amano, A., Takeuchi, H., and Furuta, N. (2010). Outer membrane vesicles function as offensive weapons in host-parasite interactions. *Microbes Infect.* 12, 791–798. doi: 10.1016/j.micinf.2010.05.008
- Andresen, K., Simonsen, P. E., Andersen, B. J., and Birch-Andersen, A. (1989). *Echinostoma caproni* in mice: shedding of antigens from the surface of an intestinal trematode. *Int. J. Parasitol.* 19, 111–118. doi: 10.1016/0020-7519(89)90028-3
- Balaj, L., Lessard, R., Dai, L., Cho, Y.-J., Pomeroy, S. L., Breakfield, X. O., et al. (2011). Tumour microvesicles contain retrotransposon elements and amplified oncogene sequences. *Nat. Commun.* 2, 180. doi: 10.1038/ncomms1180
- Barreto, A., Rodriguez, L. S., Rojas, O. L., Wolf, M., Greenberg, H. B., Franco, M. A., et al. (2010). Membrane vesicles released by intestinal epithelial cells infected with rotavirus inhibit T-cell function. *Viral Immunol.* 23, 595–608. doi: 10.1089/vim.2009.0113
- Barteneva, N. S., Fasler-Kan, E., Bernimoulin, M., Stern, J. N., Ponomarev, E. D., Duckett, L., et al. (2013). Circulating microparticles: square the circle. *BMC Cell Biol.* 14:23. doi: 10.1186/1471-2121-14-23
- Baum, J., Papenfuss, A. T., Mair, G. R., Janse, C. J., Vlachou, D., Waters, A. P., et al. (2009). Molecular genetics and comparative genomics reveal RNAi is not functional in malaria parasites. *Nucleic Acids Res.* 37, 3788–3798. doi: 10.1093/nar/gkp239
- Baxt, L. A., Baker, R. P., Singh, U., and Urban, S. (2008). An *Entamoeba histolytica* rhomboid protease with atypical specificity cleaves a surface lectin involved in phagocytosis and immune evasion. *Genes Dev.* 22, 1636–1646. doi: 10.1101/gad.1667708
- Bayer-Santos, E., Aguilar-Bonavides, C., Rodrigues, S. P., Cordero, E. M., Margues, A. F., Varela-Ramirez, A., et al. (2013). Proteomic analysis of *Trypanosoma cruzi* secretome: characterization of two populations of extracellular vesicles and soluble proteins. *J. Proteome Res.* 12, 883–897. doi: 10.1021/pr300947g
- Benchimol, M. (2004). The release of secretory vesicle in encysting *Giardia lamblia*. *FEMS Microbiol. Lett.* 235, 81–87. doi: 10.1111/j.1574-6968.2004.tb09570.x
- Bernet, J., Mullick, J., Singh, A. K., and Sahu, A. (2003). Viral mimicry of the complement system. *J. Biosci.* 28, 249–264. doi: 10.1007/BF02970145
- Bhatnagar, S., and Schorey, J. S. (2007). Exosomes released from infected macrophages contain *Mycobacterium avium* glycopeptidolipids and are proinflammatory. *J. Biol. Chem.* 282, 25779–25789. doi: 10.1074/jbc.M702277200
- Bhatnagar, S., Shinagawa, K., Castellino, F. J., and Schorey, J. S. (2007). Exosomes released from macrophages infected with intracellular pathogens stimulate a proinflammatory response *in vitro* and *in vivo*. *Blood* 110, 3234–3244. doi: 10.1182/blood-2007-03-079152
- Bhattacharjee, S., van Ooij, C., Balu, B., Adams, J. H., and Haldar, K. (2008). Maurer's clefts of *Plasmodium falciparum* are secretory organelles that concentrate virulence protein reporters for delivery to the host erythrocyte. *Blood* 111, 2418–2426. doi: 10.1182/blood-2007-09-115279
- Bordes, F., and Morand, S. (2009). Coevolution between multiple helminth infestations and basal immune investment in mammals: cumulative effects of polyparasitism. *Parasitol. Res.* 106, 33–37. doi: 10.1007/s00436-009-1623-6
- Brownlie, J. C., Cass, B. N., Riegler, M., Witsenburg, J. J., Iturbe-Ormaetxe, I., McGraw, E. A., et al. (2009). Evidence for metabolic provisioning by a common invertebrate endosymbiont, *Wolbachia pipiens*, during periods of nutritional stress. *PLoS Pathog.* 5:e1000368. doi: 10.1371/journal.ppat.1000368
- Campos, F. M. F., Franklin, B. S., Teixeira-Carvalho, A., Filho, A. L. S., de Paula, S., Fontes, C. J., et al. (2010). Augmented plasma microparticles during acute *Plasmodium vivax* infection. *Malar. J.* 9:327. doi: 10.1186/1475-2875-9-327
- Cestari, I., Ansa-Addo, E., Deolindo, P., Inal, J. M., and Ramirez, M. I. (2012). *Trypanosoma cruzi* immune evasion mediated by host cell-derived microvesicles. *J. Immunol.* 188, 1942–1952. doi: 10.4049/jimmunol.1102053
- Chandra, S., Ruhela, D., Deb, A., and Vishwakarma, R. A. (2010). Glycobiology of the *Leishmania* parasite and emerging targets for anti-leishmanial drug discovery. *Expert Opin. Ther. Targets* 14, 739–757. doi: 10.1517/14728222.2010.495125
- Combes, V., Coltel, N., Alibert, M., van Eck, M., Raymond, C., Juhan-Vague, I., et al. (2005). ABCA1 gene deletion protects against cerebral malaria. Potential pathogenic role of microparticles in neuropathology. *Am. J. Pathol.* 166, 295–302. doi: 10.1016/S0002-9440(10)62253-5
- Couper, K. N., Barnes, T., Hafalla, J. C. R., Combes, V., Ryffel, B., Secher, T., et al. (2010). Parasite-derived plasma microparticles contribute significantly to malaria infection-induced inflammation through potent macrophage stimulation. *PLoS Pathog.* 6:e1000744. doi: 10.1371/journal.ppat.1000744
- da Silveira, J. C., Veeramachaseni, D. N., Winger, Q. A., Carnevale, E. M., and Bouma, G. J. (2012). Cell-secreted vesicles in equine ovarian follicular fluid contain miRNAs and proteins: a possible new form of cell communication within the ovarian follicle. *Biol. Reprod.* 86, 71. doi: 10.1095/biolreprod.111.093252
- Deatherage, B. L., and Cookson, B. T. (2012). Membrane vesicle release in bacteria, eukaryotes, and archaea: a conserved yet underappreciated aspect of microbial life. *Infect. Immun.* 80, 1948–1957. doi: 10.1128/IAI.06014-11
- Deregibus, M. C., Cantaluppi, V., Calogero, R., Lo Iacono, M., Tetta, C., Biancone, L., et al. (2007). Endothelial progenitor cell derived microvesicles activate an angiogenic program in endothelial cells by a horizontal transfer of mRNA. *Blood* 110, 2440–2448. doi: 10.1182/blood-2007-03-078709
- Duarte, T. A., Noronha-Dutra, A. A., Nery, J. S., Ribeiro, S. B., Pitanga, T. N., Lapa E Silva, J. R., et al. (2012). *Mycobacterium tuberculosis*-induced neutrophil ectosomes decrease macrophage activation. *Tuberculosis (Edinb)* 92, 218–225. doi: 10.1016/j.tube.2012.02.007
- Dunning Hotopp, J. C. (2011). Horizontal gene transfer between bacteria and animals. *Trends Genet.* 27, 157–163. doi: 10.1016/j.tig.2011.01.005
- Duploux, A., Iturbe-Ormaetxe, I., Beatson, S. A., Szaubert, J. M., Brownlie, J. C., McMeniman, C. J., et al. (2013). Draft genome sequence of the male-killing *Wolbachia* strain wBoll reveals recent horizontal gene transfer from diverse sources. *BMC Genomics* 14:20. doi: 10.1186/1471-2164-14-20
- Eckburg, P. B., Bik, E. M., Bernstein, C. N., Purdom, E., Dethlefsen, L., Sargent, M., et al. (2005). Diversity of the human intestinal microbial flora. *Science* 308, 1635–1638. doi: 10.1126/science.1110591
- Eger, A., Kirch, A., Manfras, B., Kern, P., Schulz-Key, H., and Soboslay, P. T. (2003). Pro-inflammatory (IL-1 beta, IL-18) cytokines and IL-8 chemokine release by PBMC in response to *Echinococcus multilocularis* metacystode vesicles. *Parasite Immunol.* 25, 103–105. doi: 10.1046/j.1365-3024.2003.00601.x
- Ellis, T. N., and Kuehn, M. J. (2010). Virulence and immunomodulatory roles of bacterial outer membrane vesicles. *Microbiol. Rev.* 74, 81–94. doi: 10.1128/MMBR.00031-09

- Figueiredo, L. M., Janzen, C. J., and Cross, G. A. (2008). A histone methyltransferase modulates antigenic variation in African trypanosomes. *PLoS Biol.* 6:e161. doi: 10.1371/journal.pbio.0060161
- Figuera, L., Acosta, H., Gomez-Arreaza, A., Davila-Vera, D., Balza-Quintero, A., Quinones, W., et al. (2012). Plasminogen binding proteins in secreted membrane vesicles of *Leishmania mexicana*. *Mol. Biochem. Parasitol.* 187, 14–20. doi: 10.1016/j.molbiopara.2012.11.002
- Frohlich, K., Hua, Z., Wang, J., and Shen, L. (2012). Isolation of *Chlamydia trachomatis* and membrane vesicles derived from host and bacteria. *J. Microb. Methods* 91, 222–230. doi: 10.1016/j.mimet.2012.08.012
- Galka, F., Wai, S. N., Kusch, H., Engelmann, S., Hecker, M., Schmeck, B., et al. (2008). Proteomic characterization of the whole secretome of *Legionella pneumophila* and functional analysis of outer membrane vesicles. *Infect. Immun.* 76, 1825–1836. doi: 10.1128/IAI.01396-07
- Gallo, A., Tandon, M., Alevizos, I., and Illei, G. G. (2012). The majority of microRNAs detectable in serum and saliva is concentrated in exosomes. *PLoS ONE* 7:e30679. doi: 10.1371/journal.pone.0030679
- Gehrmann, U., Qazi, K. R., Johansson, C., Hultenby, K., Karlsson, M., Lundeberg, L., et al. (2011). Nanovesicles from *Malassezia sympodialis* and host exosomes induce cytokine responses—novel mechanisms for host-microbe interactions in atopic eczema. *PLoS ONE* 6:e21480. doi: 10.1371/journal.pone.0021480
- Geiger, A., Hirtz, C., Becue, T., Bellard, E., Centeno, D., Gargani, D., et al. (2010). Exocytosis and protein secretion in *Trypanosoma*. *BMC Microbiol.* 10:20. doi: 10.1186/1471-2180-10-20
- Giri, P. K., Kruh, N. A., Dobos, K. M., and Schorey, J. S. (2010). Proteomic analysis identifies highly antigenic proteins in exosomes from *M.tuberculosis*-infected and culture filtrate protein-treated macrophages. *Proteomics* 10, 3190–3202. doi: 10.1002/pmic.200900840
- Giri, P. K., and Schorey, J. S. (2008). Exosomes derived from *M.bovis* BCG infected macrophages activate antigen-specific CD4⁺- and CD8⁺- T-cells *in vitro* and *in vivo*. *PLoS ONE* 3:e2461. doi: 10.1371/journal.pone.0002461
- Goncalves, M. F., Umezawa, E. S., Katzin, A. M., de Souza, W., Alves, M. J. M., Zingales, B., et al. (1991). *Trypanosoma cruzi*: shedding of surface antigens as membrane vesicles. *Exp. Parasitol.* 72, 43–53. doi: 10.1016/0014-4894(91)90119-H
- Gottig, N., Elias, E. V., Quiroga, R., Nores, M. J., Solari, A. J., Touz, M. C., et al. (2006). Active and passive mechanisms drive secretory granule biogenesis during differentiation of the intestinal parasite *Giardia lamblia*. *J. Biol. Chem.* 281, 18156–18166. doi: 10.1074/jbc.M602081200
- Guescini, M., Genedani, S., Stocchi, V., and Agnati, L. F. (2010). Astrocytes and glioblastoma cells release exosomes carrying mtDNA. *J. Neural. Transm.* 117, 1–4. doi: 10.1007/s00702-009-0288-8
- György, B., Szabó, T. G., Pásztói, M., Pál, Z., Misiák, P., Aradi, B., et al. (2011). Membrane vesicles, current state-of-the-art: emerging role of extracellular vesicles. *Cell. Mol. Life Sci.* 68, 2667–2688. doi: 10.1007/s00018-011-0689-3
- Haase, S., Herrmann, S., Gruering, C., Heiber, A., Jansen, P. W., Langer, C., et al. (2009). Sequence requirements for the export of the *Plasmodium falciparum* Maurer's clefts protein REX2. *Mol. Microbiol.* 71, 1003–1017. doi: 10.1111/j.1365-2958.2008.06582.x
- Hahn, S., Giaglis, S., Chowdhury, C. S., Hoesli, I., and Hasler, P. (2013). Modulation of neutrophil NETosis: interplay between infectious agents and underlying host physiology. *Semin. Immunopathol.* 35, 439–453. doi: 10.1007/s00281-013-0380-x
- Hassani, K., and Olivier, M. (2013). Immunomodulatory impact of *Leishmania*-induced macrophage exosomes: a comparative proteomic and functional analysis. *PLoS Negl. Trop. Dis.* 7:e2185. doi: 10.1371/journal.pntd.0002185
- Hosokawa, T., Koga, R., Kikuchi, Y., Meng, X. Y., and Fukatsu, T. (2010). *Wolbachia* as a bacteriocyte-associated nutritional mutualist. *Proc. Natl. Acad. Sci. U.S.A.* 107, 769–774. doi: 10.1073/pnas.0911476107
- Hotopp, J. C., Clark, M. E., Oliveira, D. C., Foster, J. M., Fischer, P., Munoz-Torres, M. C., et al. (2007). Widespread lateral gene transfer from intracellular bacteria to multicellular eukaryotes. *Science* 317, 1753–1756. doi: 10.1126/science.1142490
- Huang, S. H., Wu, C. H., Chang, Y. C., Kwon-Chung, K. J., Brown, R. J., and Jong, A. (2012). *Cryptococcus neoformans*-derived microvesicles enhance the pathogenesis of fungal brain infection. *PLoS ONE* 7:e48570. doi: 10.1371/journal.pone.0048570
- Huebner, M. P., Manfras, B. J., Margos, M. C., Eiffler, D., Hoffmann, W. H., Schulz-Key, H., et al. (2006). *Echinococcus multilocularis* metacystodes modulate cellular and chemokine release by peripheral blood mononuclear cells in alveolar echinococcosis patients. *Clin. Exp. Immunol.* 145, 243–251. doi: 10.1111/j.1365-2249.2006.03142.x
- Iglesias, D. M., El-Kares, R., Taranta, A., Bellomo, F., Emma, F., Besouw, M., et al. (2012). Stem cell microvesicles transfer cystinosin to human cystinotic cells and reduce cystine accumulation *in vitro*. *PLoS ONE* 7:e42840. doi: 10.1371/journal.pone.0042840
- Jaiswal, R., Luk, F., Gong, J., Mathys, J. M., Grau, G. E., and Bebawy, M. (2012). Microparticle conferred microRNA profiles - implications in the transfer and dominance of cancer traits. *Mol. Cancer* 11:37. doi: 10.1186/1476-4598-11-37
- Kadiu, I., Narayanasamy, P., Dash, P. K., Zhang, W., and Gendelman, H. E. (2012). Biochemical and biologic characterization of exosomes and microvesicles as facilitators of HIV-1 infection in macrophages. *J. Immunol.* 189, 744–754. doi: 10.4049/jimmunol.1102244
- Kishore, S. P., Stiller, J. W., and Deitsch, K. W. (2013). Horizontal transfer of epigenetic machinery and evolution of the parasitism in the malaria parasite *Plasmodium falciparum* and other apicomplexans. *BMC Evol. Biol.* 13:37. doi: 10.1186/1471-2148-13-37
- Klasson, L., Kambris, Z., Cook, P. E., Walker, T., and Sinkins, S. P. (2009). Horizontal gene transfer between *Wolbachia* and the mosquito *Aedes aegypti*. *BMC Genomics* 10:33. doi: 10.1186/1471-2164-10-33
- Koh, A. L., Sun, C. X., Zhu, F., and Glogauer, M. (2005). The role of Rac1 and Rac2 in bacterial killing. *Cell. Immunol.* 235, 92–97. doi: 10.1016/j.cellimm.2005.07.005
- Kondo, N., Nikoh, N., Ijichi, N., Shimada, M., and Fukatsu, T. (2002). Genome fragment of *Wolbachia* endosymbiont transferred to X chromosome of host insect. *Proc. Natl. Acad. Sci. U.S.A.* 99, 14280–14285. doi: 10.1073/pnas.222228199
- Kosaka, N., and Ochiya, T. (2011). Unraveling the mystery of cancer by secretory microRNA: horizontal microRNA transfer between living cells. *Front. Genet.* 2:97. doi: 10.3389/fgene.2011.00097
- Lambertz, U., Silverman, J. M., Nandan, D., McMaster, W. R., Clos, J., Foster, L. J., et al. (2012). Secreted virulence factors and immune evasion in visceral leishmaniasis. *J. Leukoc. Biol.* 91, 887–899. doi: 10.1189/jlb.0611326
- Lamrabet, O., Merhej, V., Pontarotti, P., Raouf, D., and Drancourt, M. (2012). The genealogic tree of mycobacteria reveals a long sympatric life into free-living protozoa. *PLoS ONE* 7:e34754. doi: 10.1371/journal.pone.0034754
- Lemgruber, L., Kudryashev, M., Dekiwadia, S., Riglar, D. T., Baum, J., Stahlberg, H., et al. (2013). Cryo-electron tomography reveals four-membrane architecture of the *Plasmodium* apicoplast. *Malar. J.* 12, 25. doi: 10.1186/1475-2875-12-25
- Leoratti, F. M., Trevelin, S. C., Cunha, F. Q., Rocha, B. C., Costa, P. A., Gravina, H. D., et al. (2012). Neutrophil paralysis in *Plasmodium vivax* malaria. *PLoS Negl. Trop. Dis.* 6:e1710. doi: 10.1371/journal.pntd.0001710
- Letunic, I., and Bork, P. (2011). Interactive Tree Of Life v2: online annotation and display of phylogenetic trees made easy. *Nucleic Acids Res.* 39(Web Server issue), W475–W478. doi: 10.1093/nar/gkr201
- Lew, V. L. (2011). Malaria: surprising mechanism of merozoite egress revealed. *Curr. Biol.* 21, R314–R316. doi: 10.1016/j.cub.2011.03.066
- Li, H., Huang, S., Guo, C., Guan, H., and Xiong, C. (2012). Cell-free seminal mRNA and microRNA exist in different forms. *PLoS ONE* 7:e34566. doi: 10.1371/journal.pone.0034566
- Liegeois, S., Benedetto, A., Garnier, J. M., Schwab, Y., and Labouesse, M. (2006). The V0-ATPase mediates apical secretion of exosomes containing Hedgehog-related proteins in *Caenorhabditis elegans*. *J. Cell Biol.* 173, 949–961. doi: 10.1083/jcb.2005.11.072
- Ludin, P., Nilsson, D., and Maeser, P. (2011). Genome-wide identification of molecular mimicry candidates in

- parasites. *PLoS ONE* 6:e17546. doi: 10.1371/journal.pone.0017546
- Lyke, K. E., Dicko, A., Dabo, A., Sangare, L., Kone, A., Coulibaly, D., et al. (2005). Association of *Schistosoma haematobium* infection with protection against acute *Plasmodium falciparum* malaria in Malian children. *Am. J. Trop. Med. Hyg.* 73, 1124–1130.
- Mamuti, W., Sako, Y., Bart, J. M., Nakao, M., Ma, X., Wen, H., et al. (2007). Molecular characterization of a novel gene encoding an 8-kDa-subunit of antigen B from *Echinococcus granulosus* genotypes 1 and 6. *Parasitol. Int.* 56, 313–316. doi: 10.1016/j.parint.2007.06.003
- Mantel, P. Y., Hoang, A. N., Goldowitz, I., Potashnikova, D., Hamza, B., Vorobjev, I., et al. (2013). Malaria microvesicles act as messengers that induce potent responses in immune cells and parasites. *Cell Host Microbe* 13, 521–534. doi: 10.1016/j.chom.2013.04.009
- Manzano-Roman, R., and Siles-Lucas, M. (2012). MicroRNAs in parasitic diseases: potential for diagnosis and targeting. *Mol. Biochem. Parasitol.* 186, 81–86. doi: 10.1016/j.molbiopara.2012.10.001
- Marcilla, A., Trelis, M., Cortes, A., Sotillo, J., Cantalapedra, F., Minguez, M. T., et al. (2012). Extracellular vesicles from parasitic helminths contain specific excretory/secretory proteins and are internalized in intestinal host cells. *PLoS ONE* 7:e45974. doi: 10.1371/journal.pone.0045974
- Martin-Jaular, L., Nakayasu, E. S., Ferrer, M., Almeida, I. C., and del Portillo, H. A. (2011). Exosomes from *Plasmodium yoelii*-infected reticulocytes protect mice from lethal infections. *PLoS ONE* 6:e26588. doi: 10.1371/journal.pone.0026588
- McNulty, S. N., Foster, J. M., Mitreva, M., Dunning Hotopp, J. C., Martin, J., Fischer, K., et al. (2010). Endosymbiont DNA in endobacteria-free filarial nematodes indicates ancient horizontal genetic transfer. *PLoS ONE* 5:e11029. doi: 10.1371/journal.pone.0011029
- Moliner, C., Fournier, P. E., and Raoult, D. (2010). Genome analysis of microorganisms living in amoebae reveals a melting pot of evolution. *FEMS Microbiol. Rev.* 34, 281–294. doi: 10.1111/j.1574-6976.2009.00209.x
- Natakamol, D., Dondorp, A. M., Krudsood, S., Udomsangpetch, R., Pattanapanyasat, K., Combes, V., et al. (2011). Circulating red-cell derived microparticles in human malaria. *J. Infect. Dis.* 203, 700–706. doi: 10.1093/infdis/jiq104
- Nicola, A. M., Frases, S., and Casadevall, A. (2009). Lipophilic dye staining of *Cryptococcus neoformans* extracellular vesicles and capsule. *Eukaryot. Cell* 8, 1373–1380. doi: 10.1128/EC.00044-09
- Nielsen, H., Kharazmi, A., and Theander, T. G. (1986). Suppression of blood monocyte and neutrophil chemotaxis in acute human malaria. *Parasite Immunol.* 8, 541–550. doi: 10.1111/j.1365-3024.1986.tb00868.x
- Nono, J. K., Pletinckx, K., Lutz, M. B., and Brehm, K. (2012). Excretory/secretory-products of *Echinococcus multilocularis* larvae induce apoptosis and tolerogenic properties in dendritic cells *in vitro*. *PLoS Negl. Trop. Dis.* 6:e1516. doi: 10.1371/journal.pntd.0001516
- Oliveira, D. L., Freire-de-Lima, C. G., Nosanchuk, J. D., Casadevall, A., Rodrigues, M. L., and Nimrichter, L. (2010). Extracellular vesicles from *Cryptococcus neoformans* modulate macrophage functions. *Infect. Immun.* 78, 1601–1609. doi: 10.1128/IAI.01171-09
- Ouassi, A., Aguirre, T., Plumas-Marty, B., Piras, M., Schoneck, R., Gras-Masse, H., et al. (1992). Cloning and sequencing of a 24-kDa *Trypanosoma cruzi* specific antigen released in association with membrane vesicles and defined by a monoclonal antibody. *Biol. Cell* 75, 11–17. doi: 10.1016/0248-4900(92)90119-L
- Panepinto, J., Komperda, K., Frases, S., Park, Y.-D., Djordjevic, J. T., Casadevall, A., et al. (2009). Sec6-dependent sorting of fungal extracellular exosomes and laccase of *Cryptococcus neoformans*. *Mol. Microbiol.* 71, 1165–1176. doi: 10.1111/j.1365-2958.2008.06588.x
- Papadopoulos, J. S., and Agarwala, R. (2007). COBALT: constraint-based alignment tool for multiple protein sequences. *Bioinformatics* 23, 1073–1079. doi: 10.1093/bioinformatics/btm076
- Petney, T. N., and Andrews, R. H. (1998). Multiparasite communities in animals and humans: frequency, structure and pathogenic significance. *Int. J. Parasitol.* 28, 377–393. doi: 10.1016/S0020-7519(97)00189-6
- Pierson, T., Matrakas, D., Taylor, Y. U., Manyam, G., Morozov, V. N., Zhou, W., et al. (2011). Proteomic characterization and functional analysis of outer membrane vesicles of *Francisella novicida* suggests possible role in virulence and use as a vaccine. *J. Proteome Res.* 10, 954–967. doi: 10.1021/pr1009756
- Pisetsky, D. S., Gauley, J., and Ullal, A. J. (2011). Microparticles as a source of extracellular DNA. *Immunol. Res.* 49, 227–234. doi: 10.1007/s12026-010-8184-8
- Pollak, C. N., Delpino, M. V., Fossati, C. A., and Baldi, P. C. (2012). Outer membrane vesicles from *Bruceella abortus* promote bacterial internalization by human monocytes and modulate their innate immune response. *PLoS ONE* 7:e50214. doi: 10.1371/journal.pone.0050214
- Price, C. T., Al-Khodori, S., Al-Quanadan, T., Santic, M., Habyarimana, F., Kalia, A., et al. (2009). Molecular mimicry by an F-box effector of *Legionella pneumophila* hijacks a conserved polyubiquitination machinery within macrophages and *Protozoa*. *PLoS Pathog.* 5:e1000704. doi: 10.1371/journal.ppat.1000704
- Price, C. T. D., and Kwai, Y. A. (2010). Exploitation of host polyubiquitination machinery through molecular mimicry by eukaryotic-like bacterial F-box effectors. *Front. Microbiol.* 1:122. doi: 10.3389/fmicb.2010.00122
- Quah, B. J., and O'Neill, H. C. (2007). *Mycoplasma* contaminants present in exosome preparations induce polyclonal B cell responses. *J. Leukoc. Biol.* 82, 1070–1082. doi: 10.1189/jlb.0507277
- Rak, J., and Guha, A. (2012). Extracellular vesicles-vehicles that spread cancer genes. *Bioessays* 34, 489–497. doi: 10.1002/bies.20110169
- Ramachandra, L., Qu, Y., Wang, Y., Lewis, C. J., Cobb, B. A., Takatsu, K., et al. (2010). *Mycobacterium tuberculosis* synergizes with ATP to induce release of microvesicles and exosomes containing major histocompatibility complex class II molecules capable of antigen presentation. *Infect. Immun.* 78, 5116–5125. doi: 10.1128/IAI.01089-09
- Raoult, D., and Boyer, M. (2010). Amoebae as genitors and reservoirs of giant viruses. *Intervirology* 53, 321–329. doi: 10.1159/000312917
- Ratajczak, J., Miekus, K., Kucia, M., Zhang, J., Reza, R., Dvorak, P., et al. (2006). Embryonic stem cell-derived microvesicles reprogram hematopoietic progenitors: evidence for horizontal transfer of mRNA and protein delivery. *Leukemia* 20, 847–856. doi: 10.1038/sj.leu.2404132
- Regev-Rudzki, N., Wilson, D. W., Carvalho, T. G., Sissiquella, X., Coleman, B. M., Rug, M., et al. (2013). Cell-cell communication between malaria-infected red blood cells via exosome-like vesicles. *Cell* 153, 1120–1133. doi: 10.1016/j.cell.2013.04.029
- Ricard, G., McEwan, N. R., Dutilh, B. E., Jouany, J. P., Macheboeuf, D., Mitsumori, M., et al. (2006). Horizontal gene transfer from Bacteria to rumen Ciliates indicates adaptation to their anaerobic, carbohydrates-rich environment. *BMC Genomics* 7:22. doi: 10.1186/1471-2164-7-22
- Roberts, A. W., Kim, C., Zhen, L., Lowe, J. B., Kapur, R., Petryniak, B., et al. (1999). Deficiency of the hematopoietic cell-specific Rho family GTPase Rac2 is characterized by abnormalities in neutrophil function and host defense. *Immunity* 10, 183–196. doi: 10.1016/S1074-7613(00)80019-9
- Robinson, M. W., Donnelly, S., Hutchinson, A. T., To, J., Taylor, N. L., Norton, R. S., et al. (2011). A family of helminth molecules that modulate innate cell responses via molecular mimicry of host antimicrobial peptides. *PLoS Pathog.* 7:e1002042. doi: 10.1371/journal.ppat.1002042
- Rodrigues, M. L., Nakayasu, E. S., Oliveira, D. L., Nimrichter, L., Nosanchuk, J. D., Almeida, E. S., et al. (2008). Extracellular vesicles produced by *Cryptococcus neoformans* contain protein components associated with virulence. *Eukaryot. Cell* 7, 58–67. doi: 10.1128/EC.00370-07
- Ronquist, G. K., Larsson, A., Stavreus-Evers, A., and Ronquist, G. (2012). Prostatomes are heterogeneous regarding size and appearance but affiliated to one DNA-containing exosome-family. *Prostate* 72, 1736–1745. doi: 10.1002/pros.22526
- Ronquist, K. G., Ronquist, G., Carlsson, L., and Larsson, A. (2009). Human prostatomes contain chromosomal DNA. *Prostate* 69, 737–743. doi: 10.1002/pros.20921
- Selman, M., and Corradi, N. (2011). Microsporidia: horizontal gene transfer in the vicious parasites. *Mob. Genet. Elements* 1, 251–255. doi: 10.4161/mge.18611
- Semenov, D. V., Baryakin, D. N., Brenner, E. V., Kurilshikov, A. M., Vasiliev, G. V., Bryzgalov, L. A., et al. (2012). Unbiased

- approach to profile the variety of small non-coding RNA of human blood plasma with massively parallel sequencing technology. *Expert Opin. Biol. Ther.* (Suppl. 1), S43–S51. doi: 10.1517/14712598.2012.679653
- Sepulveda, P., Liegeard, P., Wallukat, G., Levin, M. J., and Hontebeyrie, M. (2000). Modulation of cardiocyte functional activity by antibodies against *Trypanosoma cruzi* Ribosomal P2 protein C terminus. *Infect. Immun.* 68, 5114–5119. doi: 10.1128/IAI.68.9.5114-5119.2000
- Silverman, J. M., Clos, J., deOliveira, C. C., Shirvani, O., Fang, Y., Wang, C., et al. (2010a). An exosome-based secretion pathway is responsible for protein export from *Leishmania* and communication with macrophages. *J. Cell Sci.* 123, 842–852. doi: 10.1242/jcs.056465
- Silverman, J. M., Clos, J., Horakova, E., Wang, A. Y., Wiesigil, M., Kelly, I., et al. (2010b). *Leishmania* exosomes modulate innate and adaptive immune responses through effects on monocytes and dendritic cells. *J. Immunol.* 185, 5011–5022. doi: 10.4049/jimmunol.1000541
- Silverman, J. M., and Reiner, N. E. (2011). Exosomes and other microvesicles in infection biology: organelles with unanticipated phenotypes. *Cell. Microbiol.* 13, 1–9. doi: 10.1111/j.1462-5822.2010.01537.x
- Silverman, J. M., and Reiner, N. E. (2012). *Leishmania* exosomes deliver preemptive strikes to create an environment permissive for early infection. *Front. Cell. Infect. Microbiol.* 1:26. doi: 10.3389/fcimb.2011.00026
- Singh, P. P., LeMaire, C., Tan, J. C., Zeng, E., and Schorey, J. S. (2011). Exosomes released from *M. tuberculosis* infected cells can suppress IFN-gamma mediated activation of naive macrophages. *PLoS ONE* 6:e18564. doi: 10.1371/journal.pone.0018564
- Singh, P. P., Smith, V. L., Karakousis, P. C., and Schorey, J. S. (2012). Exosomes isolated from mycobacteria-infected mice or cultured macrophages can recruit and activate immune cells *in vitro* and *in vivo*. *J. Immunol.* 189, 777–785. doi: 10.4049/jimmunol.1103638
- Skog, J., Wuerdinger, T., van Rijn, S., Meijer, D. H., Gainche, L., Sena-Esteves, M., et al. (2008). Glioblastoma microvesicles transport RNA and proteins that promote tumour growth and provide diagnostic biomarkers. *Nat. Cell Biol.* 10, 1470–1476. doi: 10.1038/ncb1800
- Sotillo, J., Valero, M., Sanchez del Pino, M. M., Fried, B., Esteban, J. G., Marcilla, A., et al. (2010). Excretory/secretory proteome of the adult stage of *Echinostoma caproni*. *Parasitol. Res.* 107, 691–697. doi: 10.1007/s00436-010-1923-x
- Spolski, R. J., Corson, J., Thomas, P. G., and Kuhn, R. E. (2000). Parasite-secreted products regulate the host response to larval *Taenia crassiceps*. *Parasite Immunol.* 22, 297–305. doi: 10.1046/j.1365-3024.2000.00301.x
- Telfer, S., Lambin, X., Birtles, R., Beldomenico, P., Burthe, S., Paterson, S., et al. (2010). Species interactions in a parasite community drive infection risk in a wildlife population. *Science* 330, 243–246. doi: 10.1126/science.1190333
- Toledo, A., Coleman, J. L., Crowley, J. T., and Benach, J. L. (2012). The enolase of *Borrelia burgdorferi* is a plasminogen receptor released in outer membrane vesicles. *Infect. Immun.* 80, 359–368. doi: 10.1128/IAI.05836-11
- Torrecilhas, A. C., Schimacher, R. I., Alves, M. J., and Colli, W. (2012). Vesicles as carriers of virulence factors in parasitic protozoan diseases. *Microb. Infect.* 14, 1465–1474. doi: 10.1016/j.micinf.2012.07.008
- Trelka, D. P., Schneider, T. G., Reeder, J. C., and Tarashi, T. F. (2000). Evidence for vesicle-mediated trafficking of parasite proteins to the host cytosol and erythrocyte surface membrane in *Plasmodium falciparum* infected erythrocytes. *Mol. Biochem. Parasitol.* 196, 131–145. doi: 10.1016/S0166-6851(99)00207-8
- Trocoli Torrecilhas, A. C., Tonelli, R. R., Pavanelli, W. R., da Silva, J. S., Schumacher, R. I., de Souza, W., et al. (2009). *Trypanosoma cruzi*: parasite shed vesicles increase heart parasitism and generate an intense inflammatory response. *Microbes Infect.* 11, 29–39. doi: 10.1016/j.micinf.2008.10.003
- Valadi, H., Ekstrom, K., Bossios, A., Sjostrand, M., Lee, J. J., and Loetvall, J. O. (2007). Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat. Cell Biol.* 9, 654–659. doi: 10.1038/ncb1596
- Vallejo, M. C., Matsuo, A. L., Ganiko, L., Soares Medeiros, L. C., Miranda, K., Silva, L. S., et al. (2011). The pathogenic fungus *Paracoccidioides brasiliensis* exports extracellular vesicles containing highly immunogenic alpha-galactosyl epitopes. *Eukaryot. Cell* 10, 343. doi: 10.1128/EC.00227-10
- Vallejo, M. C., Nakayasu, E. S., Longo, L. V. G., Ganiko, L., Lopes, F. G., Matsuo, A. L., et al. (2012a). Lipidomic analysis of extracellular vesicles from the pathogenic phase of *Paracoccidioides brasiliensis*. *PLoS ONE* 7:e39463. doi: 10.1371/journal.pone.0039463
- Vallejo, M. C., Nakayasu, E. S., Matsuo, A. L., Sobreira, T. J. P., Longo, L. V. G., Ganiko, L., et al. (2012b). Vesicle and vesicle-free extracellular proteome of *Paracoccidioides brasiliensis*: comparative analysis with other pathogenic fungi. *J. Proteome Res.* 11, 1676–1685. doi: 10.1021/pr200872s
- Vickers, K. C., and Remaley, A. T. (2012). Lipid-based carriers of microRNAs and intracellular communication. *Curr. Opin. Lipidol.* 23, 91–97. doi: 10.1097/MOL.0b013e328350a425
- Waldenstroem, A., Gennebaeck, M., Hellman, G., and Ronquist, G. (2012). Cardiomyocytes microvesicles contain DNA/RNA and convey biological messages to target cells. *PLoS ONE* 7:e34653. doi: 10.1371/journal.pone.0034653
- Walker, M., Baz, A., Dematteis, S., Stettler, M., Gottstein, B., Schaller, J., et al. (2004). Isolation and characterization of a secretory component of *Echinococcus multilocularis* metacystodes potentially involved in modulating the host-parasite interface. *Infect. Immun.* 71, 527–536. doi: 10.1128/IAI.72.1.527-536.2004
- Wang, K., Li, H., Yuan, Y., Etheridge, A., Zhou, Y., Huang, D., et al. (2012). The complex exogenous RNA spectra in human plasma: an interface with human gut biota. *PLoS ONE* 7:e51009. doi: 10.1371/journal.pone.0051009
- Wayengera, M. (2009). Searching for new clues about the molecular cause of endomyocardial fibrosis by way of *silico* proteomics and analytical chemistry. *PLoS ONE* 4:e7420. doi: 10.1371/journal.pone.0007420
- Weber, S. S., Parente, A. F. A., Borges, C. L., Parente, J. A., Bailao, A. M., and de Almeida Soares, C. M. (2012). Analysis of the secretomes of *Paracoccidioides mycelia* and yeast cells. *PLoS ONE* 7:e52470. doi: 10.1371/journal.pone.0052470
- Weiss, B., and Aksoy, S. (2011). Microbiome influences on insect host vector competence. *Trends Parasitol.* 27, 514–522. doi: 10.1016/j.pt.2011.05.001
- Werren, J. H., Baldo, L., and Clark, M. E. (2008). *Wolbachia*: master manipulators of invertebrate biology. *Nat. Rev. Microbiol.* 6, 741–751. doi: 10.1038/nrmicro1969
- Wilson, R. A., Wright, J. M., de Castro-Borges, W., Parker-Manuel, S. J., Dowle, A. A., Ashton, P. D., et al. (2011). Exploring the *Fasciola Hepatica* tegument proteome. *Int. J. Parasitol.* 41, 1347–1359. doi: 10.1016/j.ijpara.2011.08.003
- Woolfit, M., Iturbe-Ormaetxe, I., McGraw, E. A., and O'Neill, S. L. (2009). An ancient horizontal gene transfer between mosquito and the endosymbiotic bacterium *Wolbachia pipientis*. *Mol. Biol. Evol.* 26, 367–374. doi: 10.1093/molbev/msn253
- Xu, L., Yang, B. F., and Ai, J. (2013). MicroRNA transport: a new way in cell communication. *J. Cell. Physiol.* 228, 1713–1719. doi: 10.1002/jcp.24344
- Yamauchi, A., Marchal, C. C., Molitoris, J., Pech, N., Knaus, U., Towe, J., et al. (2005). Rac GTPase isoform-specific regulation of NADPH oxidase and chemotaxis in murine neutrophils *in vivo*. Role of C-terminal polybasic domain. *J. Biol. Chem.* 280, 953–964. doi: 10.1074/jbc.M408820200
- Yang, C., Chalasani, G., Ng, Y. H., and Robbins, P. D. (2012). Exosomes released from *Mycoplasma* infected tumor cells activate inhibitory B cells. *PLoS ONE* 7:e36138. doi: 10.1371/journal.pone.0036138
- Yoneda, A., and Doering, T. L. (2006). A eukaryotic capsular polysaccharide is synthesized intracellularly and secreted via exocytosis. *Mol. Biol. Cell* 17, 5131–5140. doi: 10.1091/mbc.E06-08-0701
- Yoon, H., Ansong, C., Adkins, J. N., and Heffron, F. (2011). Discovery of *Salmonella* virulence factors translocated via outer membrane vesicles to murine macrophages. *Infect. Immun.* 79, 2182–2192. doi: 10.1128/IAI.01277-10
- Zaccone, P., and Cooke, A. (2013). Vaccine against autoimmune disease: can helminths or their products provide a therapy. *Curr. Opin. Immunol.* 25, 418–423. doi: 10.1016/j.coi.2013.02.006
- Zhang, H., Sun, C., Glogauer, M., and Bokoch, G. M. (2009). Human

- neutrophils coordinate chemotaxis by differential activation of Rac1 and Rac2. *J. Immunol.* 183, 2718–2728. doi: 10.4049/jimmunol.0900849
- Zhang, L., Hou, D., Chen, X., Li, D., Zhu, L., Zhang, Y., et al. (2012). Exogenous plant mir168a specifically targets mammalian LDLRAP1: evidence of cross-kingdom regulation by microRNA. *Cell Res.* 22, 107–126. doi: 10.1038/cr.2011.158
- Zhang, Y., Liu, D., Chen, X., Li, J., Li, L., Bian, Z., et al. (2010a). Secreted monocytic miR-150 enhances targeted endothelial cell migration. *Mol. Cell* 39, 133–144. doi: 10.1016/j.molcel.2010.06.010
- Zhang, W., Li, J., Jones, M. K., Zhang, Z., Zhao, L., Blair, D., et al. (2010b). The *Echinococcus granulosus* antigen B gene family comprises at least 10 unique genes in five subclasses which are differentially expressed. *PLoS Negl. Trop. Dis.* 4:e784. doi: 10.1371/journal.pntd.0000784
- Zhong, G. (2011). *Chlamydia* secretion of proteases for manipulating host signaling pathways. *Front. Microbiol.* 2:14. doi: 10.3389/fmicb.2011.00014
- Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.
- Received: 17 June 2013; accepted: 20 August 2013; published online: 06 September 2013.
- Citation: Barteneva NS, Maltsev N and Vorobjev IA (2013) Microvesicles and intercellular communication in the context of parasitism. *Front. Cell. Infect. Microbiol.* 3:49. doi: 10.3389/fcimb.2013.00049
- This article was submitted to the journal *Frontiers in Cellular and Infection Microbiology*. Copyright © 2013 Barteneva, Maltsev and Vorobjev. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.