



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Weight loss decreases progressive left ventricular remodeling: The Multi-Ethnic Study of Atherosclerosis

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Abbasi, S. A., R. V. Shah, V. L. Murthy, J. Eng, C. Wu, P. Ouyang, R. Y. Kwong, et al. 2014. "Weight loss decreases progressive left ventricular remodeling: The Multi-Ethnic Study of Atherosclerosis." <i>Journal of Cardiovascular Magnetic Resonance</i> 16 (Suppl 1): O2. doi:10.1186/1532-429X-16-S1-O2. http://dx.doi.org/10.1186/1532-429X-16-S1-O2 .
Published Version	doi:10.1186/1532-429X-16-S1-O2
Accessed	February 16, 2015 10:33:41 AM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:12406590
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

ORAL PRESENTATION

Open Access

Weight loss decreases progressive left ventricular remodeling: The Multi-Ethnic Study of Atherosclerosis

Siddique A Abbasi^{1*}, Ravi V Shah^{2,1}, Venkatesh L Murthy³, John Eng⁴, Colin Wu⁴, Pamela Ouyang⁵, Raymond Y Kwong¹, Allison Goldfine⁶, David Bluemke⁷, Joao A Lima⁵, Michael Jerosch-Herold⁸

From 17th Annual SCMR Scientific Sessions
 New Orleans, LA, USA. 16-19 January 2014

Background

Obesity (body mass index, BMI > 30 kg/m²) is an independent risk factor for incident heart failure (HF). Effects of weight change on cardiac structure have not been extensively investigated in large community-based populations. We hypothesized that weight loss and gain in the Multi-Ethnic Study of Atherosclerosis (MESA) would coincide with changes in left ventricular (LV) mass – key features in the progression to obesity-related HF.


Methods

To investigate the association of longitudinal changes in weight on ventricular remodeling, we investigated 2,351 patients in MESA who underwent two serial cardiac magnetic resonance imaging (CMR) examinations at initial enrollment (2002) and at follow-up (2011) with available obesity status. Canonical parameters of LV structure and function (height-indexed LV mass, LV volumes, and LV ejection fraction) were measured. MESA participants were classified by obesity status (normal weight: 18.5-25 kg/m²; overweight/obese ≥ 25 kg/m²). We constructed splines for linear and logistic models using generalized additive models to assess the form of the continuous relationship between indexed LV mass changes and weight change qualitatively. Next, we constructed multivariable linear models, adjusted for confounders involved in the pathogenesis of LV hypertrophy that could be altered by weight change, including: glycemic status, hypertension, waist-to-hip ratio, body-mass-index, and biomarkers of inflammation. Finally, the multivariable linear model was adjusted

Table 1 Linear regression for percent change in height-indexed LV mass between Exam 1 and Exam 5

Parameter	Estimate	Linear Model	P-value
		95% Confidence Interval	
Age, y	-0.05	(-0.12 to 0.02)	0.18
Female gender	-7.67	(-8.93 to -6.4)	< 0.0001
Race			
White	referent	-	-
Chinese	0.11	(-1.67 to 1.88)	0.9
African-American	2.83	(1.42 to 4.23)	< 0.0001
Hispanic	-0.42	(-2.04 to 1.19)	0.61
Smoking history (ever smoker)	-0.68	(-1.76 to 0.39)	0.21
Income	-0.13	(-0.33 to 0.06)	0.19
Education	-0.04	(-0.31 to 0.23)	0.77
BMI at Exam 1 (per log), kg/m ²	24.51	(17.67 to 31.35)	< 0.0001
Glycemia			
Normal	referent	-	-
IFG	-0.75	(-2.49 to 0.99)	0.4
Untreated DM	-2.56	(-6.5 to 1.38)	0.2
Treated DM	0.92	(-1.29 to 3.12)	0.42
Δ Glycemia	-0.13	(-0.83 to 0.58)	0.72
Hypertension stage	2.73	(2.19 to 3.28)	< 0.0001
Δ Hypertension stage	2.88	(2.43 to 3.33)	< 0.0001
Number of antihypertensive classes at Exam 1	0.2	(-0.43 to 0.83)	0.54
C-reactive protein (per log)	-0.46	(-0.98 to 0.07)	0.09
Waist Circumference at Exam 1 (per cm)	0.07	(-0.02 to 0.15)	0.11
LVMi at Exam 1 (g/m ^{2.7})	-0.8	(-0.88 to -0.73)	< 0.0001
Weight change from Exam 1 to 5 (per 10% decrease)	-2.76	(-3.42 to -2.1)	< 0.0001

¹Section of Non-invasive Cardiovascular Imaging (Department of Internal Medicine/Division of Cardiology and Department of Radiology), Brigham and Women's Hospital/Harvard Medical School, Boston, Massachusetts, USA
 Full list of author information is available at the end of the article



for age, gender, race, income, educational status, smoking, Exam 1 BMI, and height-indexed LV mass at Exam 1.

Results

Of the overall cohort studied, 257 individuals (11%) experienced $\geq 10\%$ weight loss (median 10.2 kilograms) and 194 (8%) had $\geq 10\%$ weight gain (median 10.0 kilograms). After adjustment for hypertension, diabetes, age, race, and other clinical risk factors, every 10% decrease in weight was associated with a fully covariate-adjusted 3% additional decrease in height-indexed left ventricular mass. Every 10% loss in body weight increased the odds of a 10% or greater drop in left ventricular mass by 50%. Finally, regression models suggested linear decreases in left ventricular mass regression with increasing weight loss, suggesting no threshold effect for weight loss on cardiac remodeling.

Conclusions

Weight loss is associated with significant beneficial effects on cardiac remodeling, even after adjustment for baseline obesity-related cardiometabolic risk. There is no threshold for the weight change needed before benefits on cardiac occur, suggesting that any degree of weight loss may be beneficial to the heart.

Funding

MESA was supported by contracts NO1-HC-95159 through NO1-HC-95169 from the National Heart, Lung, and Blood Institute. Dr. Abbasi is supported by a T32 fellowship. Dr. Shah is supported by an American

Heart Association Post-Doctoral Fellowship Award (11POST000002) and a training grant from the Heart Failure National Institutes of Health Clinical Research Network (U01-HL084877). Dr. Jeresch-Herold receives support through R01-HL-65580. All other authors have no financial disclosures relevant to the content of this manuscript.

Authors' details

¹Section of Non-invasive Cardiovascular Imaging (Department of Internal Medicine/Division of Cardiology and Department of Radiology), Brigham and Women's Hospital/Harvard Medical School, Boston, Massachusetts, USA. ²Department of Internal Medicine (Division of Cardiology), Massachusetts General Hospital, Boston, Massachusetts, USA. ³Department of Medicine (Cardiovascular Medicine Division) and Department of Radiology (Nuclear Medicine and Cardiothoracic Imaging Divisions), University of Michigan, Ann Arbor, Michigan, USA. ⁴Office of Biostatistics Research, National Heart, Lung, and Blood Institute, Bethesda, Maryland, USA. ⁵Department of Internal Medicine, Division of Cardiology, Johns Hopkins Bayview Medical Center, Baltimore, Maryland, USA. ⁶Department of Internal Medicine/Division of Endocrinology, Joslin Diabetes Center/Brigham and Women's Hospital, Boston, Massachusetts, USA. ⁷Radiology and Imaging Sciences, National Institutes of Health Clinical Center, National Institute of Biomedical Imaging and Bioengineering, Bethesda, Maryland, USA. ⁸Department of Radiology, Brigham and Women's Hospital, Boston, Massachusetts, USA.

Published: 16 January 2014

doi:10.1186/1532-429X-16-S1-O2

Cite this article as: Abbasi et al.: Weight loss decreases progressive left ventricular remodeling: The Multi-Ethnic Study of Atherosclerosis. *Journal of Cardiovascular Magnetic Resonance* 2014 **16**(Suppl 1):O2.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

