



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Generation Times in Wild Chimpanzees and Gorillas Suggest Earlier Divergence Times in Great Ape and Human Evolution

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Langergraber, Kevin E., Kay Prufer, Carolyn Rowney, Christophe Boesch, Catherine Crockford, Katie Fawcett, Eiji Inoue, et al. 2012. Generation Times in Wild Chimpanzees and Gorillas Suggest Earlier Divergence Times in Great Ape and Human Evolution. <i>Proceedings of the National Academy of Sciences</i> 109, no. 39: 15716–15721.
Published Version	doi:10.1073/pnas.1211740109
Accessed	February 16, 2015 5:53:33 PM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:12748565
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

**Generation times in wild chimpanzees and gorillas suggest earlier divergence times
in great ape and human evolution**

Kevin E. Langergraber^{a,b,1}, Kay Prüfer^{c,1}, Carolyn Rowney^b, Christophe Boesch^b, Catherine Crockford^d, Katie Fawcett^e, Eiji Inoue^f, Miho Inoue-Muruyama^g, John C. Mitani^h, Martin N. Mullerⁱ, Martha M. Robbins^b, Grit Schubert^{b,2}, Tara Stoinski^e, Bence Viola^j, David Watts^k, Roman M. Wittig^{b,d}, Richard W. Wrangham^l, Klaus Zuberbühler^d, Svante Pääbo^c, Linda Vigilant^{b,3}

^a Department of Anthropology, Boston University, Boston, MA 02215, USA

^b Primatology Department and

^c Genetics Department, Max Planck Institute for Evolutionary Anthropology, Leipzig, 04103, Germany

^d School of Psychology, University of St Andrews, St Andrews, KY16 9JP, UK

^e Dian Fossey Gorilla Fund International, 800 Cherokee Ave SE, Atlanta, GA 30315, USA

^f Graduate School of Science, Kyoto University, Kitashirakawa Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan

^g Wildlife Research Center, Kyoto University, 2-24 Tanaka-Sekiden-cho, Sakyo-ku, Kyoto 606-820, Japan

^h Department of Anthropology, University of Michigan, Ann Arbor, MI 48109, USA

ⁱ Department of Anthropology, University of New Mexico, Albuquerque, New Mexico 87131, USA

^j Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Leipzig, 04103, Germany

^k Department of Anthropology, Yale University, New Haven, CT 06511, USA

^l Department of Human Evolutionary Biology, Harvard University, Peabody Museum, 11 Divinity Avenue, Cambridge, MA 02138, USA

Keywords: speciation, hominin, molecular dating, primate

¹ K.E.L. and K.P. contributed equally to this work.

² Present address: Robert Koch Institute, Nordufer 20, Berlin 13353, Germany

³ to whom correspondence should be addressed. E-mail: vigilant@eva.mpg.de

Abstract

Fossils and molecular data are two independent sources of information that should in principle provide consistent inferences of when evolutionary lineages diverged. Here we employ a novel approach to genetic inference of species split times in recent human and ape evolution that is independent of the fossil record. We first use genetic parentage information on a large number of wild chimpanzees and mountain gorillas to directly infer their average generation times. We then compare these generation time estimates with those of humans and apply recent estimates of the human mutation rate per generation to derive estimates of split times of great apes and humans that are independent of fossil calibration. We date the human-chimpanzee split to at least 7 to 8 million years and the population split between Neandertals and modern humans to 400,000 to 800,000 years ago. This suggests that molecular divergence dates may not be in conflict with the attribution of 6 to 7 million-year-old fossils to the human lineage and 400,000-year-old fossils to the Neandertal lineage.

1 \body

2 **Introduction**

3

4 Over 40 years ago, Sarich and Wilson used immunological data to propose that humans
5 and African great apes diverged only about 5 million years ago, some three to four times
6 more recently than had been assumed based on the fossil record (1). Although
7 contentious at the time (e.g., (2)), this divergence has since been repeatedly estimated
8 from DNA sequence data at 4 - 6 million years ago (Ma) (3-8). However, this estimate is
9 incompatible with the attribution of fossils older than 6 Ma to the human lineage.
10 Although the assignment of fossils such as the approximately 6 Ma *Orrorin* (9) and the 6
11 - 7 Ma *Sahelanthropus* (10) to the human lineage remains controversial (11), it is also
12 possible that the divergence dates inferred from DNA sequence data are too recent.

13 The total amount of sequence differences observed today between two
14 evolutionary lineages can be expressed as the sum of two values: the sequence
15 differences that accumulated since gene flow ceased between the lineages (“split time”)
16 and the sequence differences that correspond to the diversity in the common ancestor of
17 both lineages. The extent of variation in the ancestral species may be estimated from the
18 variance of DNA sequence differences observed across different parts of the genome
19 between the species today, which will be larger the greater the level of variation in the
20 ancestral population. By subtracting this value from the total amount of sequence
21 differences, the sequence differences accumulated since the split can be estimated. The
22 rate at which DNA sequence differences accumulate in the genome (“mutation rate”) is
23 needed to then convert DNA sequence differences into split times.

24 In prior research, mutation rates have been calculated using species split times
25 estimated from the fossil record as calibration points. For calculating split times between
26 present-day humans and great apes, calibration points which assume that DNA sequence
27 differences between humans and orangutans have accumulated over 13 Ma (12) or 18 Ma
28 (5, 8) or between chimpanzees and humans over 7 Ma (13, 14) have been used. Recently,
29 researchers have commonly employed a mutation rate of 1×10^{-9} mutations per site per
30 year (e.g., (4, 6, 8, 15)) derived from the observed DNA sequence difference of around
31 1.3% between the human and chimpanzee genomes (8, 15, 16) and an assumed DNA
32 sequence divergence between these species at 7 Ma, as well as from an observed
33 sequence difference of 6.46% between the human and macaque genomes (17) and an
34 assumption of their DNA sequence divergence at 25 Ma. Although ubiquitous, this
35 approach has an inherent circularity and is subject to possible error because it relies on
36 the accuracy of the ages of fossils. While approaches to account for uncertainty in the
37 fossil record have been proposed (18), a means to avoid the use of fossil calibration
38 points would be useful.

39 An alternative approach to determine mutation rates is to compare genome
40 sequences from children and their parents (19-21). This has the advantage of not relying
41 on the fossil record. However, direct observation of mutation rates per site per generation
42 need to be converted to mutations rates per year in order to arrive at population split
43 times. For this we need the relevant generation times, which are the average maternal and
44 paternal age at reproduction in the lineages under consideration. Genetic studies of
45 humans have commonly used a generation time of 20 or 25 years (e.g., references in (5,
46 22)). However, genealogical data spanning the last two or three centuries from three

47 human populations suggest that the average generation time is 30-32 years (23-25). In
48 agreement with this, a comprehensive review considering estimated maternal age at first
49 and last childbirth and age differences between spouses in contemporary hunter-gatherers
50 as well as in food-producing countries with varying levels of industrialization inferred an
51 average human generation time of 29 years, with female- and male-specific values of 26
52 and 32 years, respectively (22). Thus, both direct genealogical and indirect demographic
53 studies conducted in a variety of societies, including those practicing a lifestyle thought
54 to be representative of that of the human lineage for much of its evolutionary history (i.e.,
55 hunter-gatherer), are fairly consistent in suggesting that the average present-day human
56 generation time is approximately 29 years and that it differs substantially between the
57 sexes.

58 Previous estimates of split times have used a wide variety of generation times for
59 great apes, including 25 years for chimpanzee, gorilla and orangutan (5), 20 years for
60 chimpanzee (13, 14) and orangutan, (6) or 15 years for chimpanzees (26), gorillas (27)
61 and chimpanzee, gorilla and orangutan (3). These estimates appear to lack any explicit
62 justification. A recent analysis used information from captive and wild populations
63 regarding female age of first reproduction, interbirth interval, age of last reproduction and
64 survivorship to estimate female generation times of 22 years for chimpanzees and 20 for
65 gorillas (28). These findings hint that some of the generation times commonly assumed in
66 studies of great apes are excessively short. Furthermore, it is possible that, as is the case
67 in humans, generation times of female great apes may not be representative of those of
68 males

69 Here we derive female and male generation times for present-day chimpanzees
70 and gorillas from genetic parentage data collected from large numbers of offspring born
71 into several wild social groups. We consider whether our data are consistent with the
72 suggestion of a positive correlation between body size and generation time in great apes
73 and humans, and explore the implications of our results for dating population split times
74 among these lineages.

75

76 **Results**

77
78 **Chimpanzee generation times.** Using parentage information for 226 offspring born in
79 eight wild chimpanzee communities, we find that the average age of parents is 24.6 years
80 (**Table 1**). Among communities, the generation times range from 22.5 to 28.9 years, but
81 no consistent difference is observed between western and eastern chimpanzees,
82 suggesting that the variation may arise due to demographic stochasticity rather than
83 consistent ecological or genetic differences between western and eastern chimpanzees.

84 Some of the chimpanzee communities are known to have experienced substantial
85 mortality in the recent past due to epidemic disease. To check if this may have altered
86 reproductive patterns, we compared the average generation intervals for groups known to
87 have experienced high infection-induced mortality (Tai North and South communities,
88 Mahale M community, Gombe Kasekela community) with those that have not (Budongo
89 Sonso group, Kibale Kanyawara and Ngogo communities). The average generation time
90 for the former groups was 24.9 while it was 24.3 for the latter. Thus, epidemic diseases
91 are not likely to have drastically affected generation times in these chimpanzee
92 communities.

93 The age of chimpanzee fathers ranged from 9.3 to 50.4 years, while age of
94 mothers ranged from 11.7 to 45.4 years (**SI Figure 1**). Thus, the potential reproductive
95 span of males (41.1 years) is some seven years, or 22%, longer than that of females (33.7
96 years). Nonetheless, because more than half (56.2%) of the offspring are produced by
97 fathers between the ages of 15 and 25, while most offspring (77%) have mothers between
98 the ages of 15 and 34, the average generation time for males and females is essentially
99 the same (24.1 and 25.2 years, respectively).

100
101 **Gorilla generation times.** Using information on the parentage of 105 mountain gorilla
102 offspring from two research sites, the average female and male generation times were
103 18.2 and 20.4 years, respectively, with an average of 19.3 years for both sexes (**Table 1**).
104 Thus, generation times in gorillas are substantially shorter than in chimpanzees.

105 The ages of gorilla fathers ranged from 10.8 to 30.9 years, while the ages of
106 gorilla mothers ranged from 7.3 to 38.0 years, suggesting that female gorillas reproduce
107 over substantially longer periods than do males. In fact, we found that more than 75% of
108 offspring were sired by males between the ages of 15 and 24, while the distribution of
109 gorilla maternal ages varied considerably more (**SI Figure 1**). Thus, in contrast to
110 chimpanzees, the potential reproductive lifespan of gorilla females is longer than for
111 gorilla males.

112
113 **Generation times and body mass.** Several life history characteristics, such as age of
114 weaning, female age at maturity, and female age at first breeding, exhibit a positive
115 relationship with body mass across primates (29). To evaluate whether generation time
116 also increases with body size in the great apes, we compared generation times and body
117 mass estimates. Supplementing our data with a recent estimate of orangutan female
118 generation time based on demographic information (28), we find that humans,
119 chimpanzees, and female orangutans display similar masses and generation times, while
120 male and female gorillas have more than twice as large body masses yet short generation
121 times, resulting in an overall negative association between mass and generation times in

122 these taxa (females, generation time = $-0.102\text{mass} + 33.5$, $R^2 = 0.88$; males, generation
123 time = $-0.059\text{mass} + 30.88$, $R^2 = 0.48$) (**SI Table 1**).

124
125 **Generation times and mutation rates.** DNA sequencing of human families has recently
126 yielded four direct estimates of mutation rates ranging from 0.97×10^{-8} to 1.36×10^{-8}
127 ⁸/site/generation (19-21). When considering the average present-day human generation
128 time of 29 years, this results in rates ranging from 0.33 to 0.47×10^{-9} /site/year.

129 Unfortunately, estimates of mutation rates per generation do not yet exist for apes.
130 However, if we assume that they are similar to those in humans, we can apply the rates of
131 0.97×10^{-8} to 1.36×10^{-8} /site/generation to the generation time of 19 years derived from
132 the gorilla, which yields mutation rates of 0.51 to 0.72×10^{-9} /site/year. Similarly,
133 application of the human mutation rate per generation to the chimpanzee with a
134 generation time of 25 years yields mutation rates of 0.39 to 0.54×10^{-9} /site/year. Because
135 the gorilla has the shortest and the human the longest generation time among the great
136 apes, this suggests that mutation rate for African apes and humans is between 0.33×10^{-9}
137 and 0.72×10^{-9} /site/year.

138
139 **Species split times.** We can use the observed generation times in apes and humans as well
140 as observed mutation rates in human families to recalibrate the previously published split
141 times among the human and ape evolutionary lineages. We assume that the common
142 ancestor at each branch point had a generation time and mutation rate within the range
143 described by the most extreme values of the present-day descendant species (see
144 Methods). **Table 2** shows that the resulting estimates are all substantially older than those
145 based on fossil calibrations of mutation rates. For example, we estimate the bonobo and
146 chimpanzee split time at 1.5 to 2.6 million years, while previous estimates put it at less
147 than 1 million years. We estimate the split time between the human and chimpanzee
148 lineages at between 7 and 13 million years while previous estimates range from 4 to 6
149 million years. We estimate the split between the gorilla lineage and the lineage leading to
150 humans, chimpanzees and bonobos at 8 to 19 million years, while previous estimates
151 range between 6 and 7 million years.

152 153 154 **Discussion**

155
156 By using direct observations of generation times in gorillas and chimpanzees and rates of
157 mutation per generation from direct observation of mutations in human families, we
158 estimate the species split times of humans and apes without relying on external fossil
159 calibration points. At 7 to 13 Ma our estimate of the chimpanzee-human split time is
160 earlier than those previously derived from molecular dating using fossil calibration points
161 but similar to the range of 6.5 – 10 Ma suggested by the fossil record (30).

162 While the earliest fossil universally accepted to belong to the lineage leading to
163 present-day humans rather than to chimpanzees, *Australopithecus anamensis*, is 4.2 Ma
164 (31) and thus reconcilable with a molecularly-inferred human-chimpanzee split time as
165 recent as 5 Ma, the attribution of late Miocene (5 to 7 Ma) fossils to the hominin lineage
166 has posed a problem. Our estimates make it possible to reconcile attribution of fossils
167 such as *Ardipithecus kaddaba* (5.2-5.8 Ma) (32), *Orrorin tugenensis* (6 Ma) (9) and

168 *Sahelanthropus tchadensis* (6-7 Ma) (10) to the hominin lineage with speciation times
169 inferred from genetic evidence (**Figure 1**). However, our estimates cannot address the
170 controversy of whether specimens such as these truly belong to the lineage leading to
171 present-day humans or to other, closely related lineages (11).

172 For the deeper time period of 7 to 13 Ma, the fossil record is even more limited
173 and difficult to interpret (31, 33). Fossils from between 8 and 11 Ma in Africa include
174 mainly *Gorilla*-sized forms, such as *Samburupithecus* (34), *Nakalipithecus* (35) and
175 *Chororapithecus*, the last of which is dated to 10 – 10.5 Ma and suggested to represent an
176 early member of the gorilla clade (36). Our estimate of 8 – 19 Ma for the split of the
177 gorilla lineage from the human-chimpanzee ancestor would be largely consistent with the
178 attribution of such forms to the gorilla lineage.

179 Even though not quantified here, our results also significantly push back the date
180 of the split between orangutans and African apes. Palaeontological data (e.g. (37)) have
181 been combined with genetic data (38) to suggest that this split occurred outside of Africa,
182 with a later “Back to Africa” migration of the common ancestor of African apes. The
183 purported “early great ape” *Pierolapithecus catalaunicus* from Spain, dating to about
184 12.5 - 13 Ma (39), and the presence of numerous derived African ape traits in Late
185 Middle Miocene fossils from Europe such as *Rudapithecus* and *Hispanopithecus* fit well
186 with this hypothesis. A split between African apes and orangutans that predates 15 Ma
187 would challenge this model, and would either put these fossils on the orangutan lineage,
188 or place them as unrelated to present-day great apes.

189 For more recent periods of hominin evolution the more recent dates provided here
190 for the human-chimpanzee split resolve an apparent contradiction between genetic and
191 paleontological data. Using a chimpanzee/human split of 5.6-8.3 Ma for calibration,
192 analyses of the Neandertal genome indicated a population split between present-day
193 humans and Neandertals at 270-440 Ka (40). This date appears to conflict with fossil
194 evidence tracing the emergence of Neandertal morphological characters over the course
195 of the Middle Pleistocene in Europe (41). The earliest evidence for Neandertal traits was
196 proposed to date to 600 \pm 66 ka at the Sima de Los Huesos (Atapuerca, Spain) (42),
197 thus predating the genetically-estimated population divergence times, but this date has
198 been disputed based both on the apparent conflict with the genetic data and on
199 stratigraphic grounds (43). However, even if the early dates for Sima are disregarded, it is
200 clear that fossils from Oxygen Isotope Stage 11 (around 400 ka), such as the
201 Swanscombe cranium, already show clear Neanderthal traits (44). Using the new human-
202 chimpanzee split estimate and assuming generation times between 25 and 29 years would
203 push back the human/Neandertal split to 423,000-781,000 years, resolving this apparent
204 conflict.

205 Recent attempts to model uncertainty in the fossil data used for molecular
206 calibration also suggest earlier split times in the evolutionary history of apes with
207 estimates of 6 – 10 Ma for the human-chimpanzee divergence and 7 – 12 Ma for the
208 divergence of the gorilla (18). Our estimates of divergence dates have the advantage that
209 they avoid fossil calibration points. However, it is possible that other aspects of our
210 analysis may lead to unreliable split time inferences. First, because of the limited
211 availability of data from the western gorilla species, we make the assumption that the
212 average generation interval of mountain gorillas is applicable to both present-day species
213 of the *Gorilla* genus. Although reliant primarily upon herbaceous vegetation, western

214 gorillas also eat fruit much of the year, while fruit is nearly absent from the mountain
215 gorilla habitat (45). More folivorous anthropoid primates are known to mature more
216 quickly than similarly-sized non-folivorous primates (46), and indeed limited data from
217 western gorillas suggest that females and males attain adulthood 2 and 3 years later,
218 respectively, than the more folivorous mountain gorilla (47). This implies that the
219 generation time in western gorillas may be on the order of 21 years, in contrast to the 19
220 years used here for gorilla generation time. However, because 19 is the shortest
221 generation time observed among present-day mountain gorillas, chimpanzees, and
222 humans, our use of this value is more conservative and simply contributes to a slightly
223 broader range for the inferred split time for the divergence of the gorilla lineage from that
224 leading to humans and chimpanzee, as well as to a broader range for the split time
225 between the two gorilla species. As with western gorillas, parentage data for calculation
226 of generation times in bonobos are lacking. However, neither extensive dietary
227 differences between bonobos and chimpanzees nor substantial differences in
228 developmental timing are apparent for these species and it is also relevant that we found
229 no consistent differences in generation times between chimpanzees from western and
230 eastern Africa. With regard to humans, highly similar estimates of generation time were
231 obtained from demographic analysis of a large sample of less- and more-developed
232 countries, a large sample of hunter-gatherer societies, and direct analysis of genealogies
233 (22). In sum, except for the gorillas where marked ecological differences may contribute
234 to a small degree of variation in generation time within the genus, the generation times
235 used here seem reliable estimates for present-day great apes and humans.

236 A further notable assumption of our work is that the generation times calculated
237 for present-day humans and great apes are valid proxies for their ancestors. It was
238 recently suggested that a slowdown in mutation rate concomitant with an increase in
239 body sizes and generation times has occurred in these lineages (8). However, there is an
240 extraordinarily diversity of ape body sizes in the fossil record since the Miocene (24 Ma
241 to 5 Ma) and it is difficult to know which ones may represent ancestors of present-day
242 apes and humans (32). Even if fossil evidence strongly suggested an increase in the size
243 of the ancestors of present-day apes and humans in the past, it is not clear that body mass
244 is a good correlate of life history parameters related to generation time (48). Although our
245 number of data points is necessarily limited, we found no correlation between mass and
246 generation time in present-day apes and humans, and the notably short generation time
247 for the relatively large mountain gorilla is consistent with the expectation that highly
248 folivorous (46) as well as more terrestrial (49) species are expected to reproduce earlier
249 than more frugivorous, arboreal primates. In accordance with the importance of diet and
250 habitat use in influencing life-history parameters it has been suggested that chimpanzees
251 and orangutans represent the most appropriate living models for the potential life history
252 variables of archaic hominins, and that the common ancestor of humans and chimpanzees
253 exhibited a slow life history similar to that of present-day chimpanzees (50). Skeletal and
254 dental analyses suggest that early hominins had growth patterns like those of present-day
255 great apes, while *Homo erectus* and Neandertals evolved slower development, but not to
256 the extent seen in present-day humans (51, 52). Given the information available at this
257 time, we suggest that the use of the ranges of the observed generation times in the
258 present-day species, including the extremes represented by gorillas (with their
259 comparatively fast life history and consequently short generation time) and humans (with

260 their comparatively slow life history and consequently long generation time), results in
261 conservatively broad estimates of hominid mutation rates and split times as shown (Table
262 2, Figure 1). Specifically, if we alternatively consider the human generation time of 29
263 years to be a recent phenomenon, and consider the chimpanzee generation time of 25
264 years to characterize the vast majority of evolution since the split between the gorilla and
265 the chimpanzee/human lineages, we would infer the date of this split at 10.9 to 17.2 Ma,
266 while the split between the lineages leading to chimpanzees and humans would be dated
267 at 6.8 to 11.6 Ma.

268 We also note that we explicitly assume that the mutation rates estimated by
269 sequencing members of present-day human families are also applicable to our closest
270 great ape relatives. This assumption, which is based on our close evolutionary
271 relationship and lack of evidence for differences in rates of evolution among the human
272 and African great ape lineages (7, 53), can be explicitly tested in the future by sequencing
273 of great ape family trios. As an additional point for future consideration, we note that the
274 original publications which provide the population split times that we recalibrate here use
275 various approaches for filtering the data analyzed, for example exclusion of repetitive
276 sequences or highly mutable sites. Refinements of our population split time estimates
277 may involve reexamination of the data, including consideration of different parts of the
278 genome, or different types of substitutions. For example, it will be interesting to compare
279 inferences from substitutions at CpG sites, which may accumulate in a time-dependent
280 fashion, with other classes of substitutions which may accumulate in a generation-
281 dependent fashion. However, studies which compared results for the human-chimpanzee
282 split obtained with and without inclusion of CpG sites found this to have little impact (3,
283 7).

284 Finally, we note that the estimation of generation times in chimpanzees and
285 gorillas derives from the long-term efforts of researchers who have invested years in
286 habituating the animals to human observation in order to collect information on their
287 natural behavior and life histories. This study illustrates the value of such approaches in
288 aiding interpretation of genomic data, and suggests that continued behavioral study of
289 wild apes, in addition to increased understanding of their behavior and cultures, is
290 necessary to complement genomic studies for a fuller understanding of the evolutionary
291 history of our closest living relatives as well as our own species.

Material and methods

Details regarding the analyses can be found in **SI Materials and Methods**. In brief, we compiled the ages of the genetically-confirmed mothers and fathers of offspring born into eight chimpanzee groups and six mountain gorilla groups habituated to human observation. We did not limit our sample to individuals whose ages are exactly known because this would lead to a downward bias in the estimation of the generation length, as older individuals are more likely to have been born before the start of long-term research on a particular group. Instead, we included in our study individuals whose ages were estimated using standard morphological, behavioral and life history criteria established from known-aged individuals and systematically incorporated estimation of ranges of minimum and maximum birthdates symmetrical about the assigned birthdate.

For the split time estimation, we first took the lowest and highest estimates of mutation rates in human families of 0.97×10^{-8} to 1.36×10^{-8} /site/generation and applied the estimated generation times of 19, 25, and 29 years for gorillas, chimpanzees and humans to arrive at low and high estimates of yearly mutation rates given each of these generation times. For example, the chimpanzee generation time of 25 years yields a rate of 0.39 to 0.54×10^{-9} mutations/site/year, while the human generation time of 29 years yields a rate of 0.33 to 0.46×10^{-9} mutations/site/year. For each split we then chose lower and upper bounds for the yearly mutation rates based upon the extreme values inferred for the taxa under consideration. For example, we assumed that the generation time of the common ancestor of chimpanzees and humans was between 25 and 29 years, the values for present-day chimpanzees and humans, respectively, and thus used the mutation rates of 0.33 and 0.54 mutations/site/year (Table 2). Similarly, the common ancestor of gorillas, chimpanzees and humans is assumed to have a generation time between 19 and 29 years and we thus used a correspondingly broader set of mutation rates. We adjusted previously published split times (Table 2) by multiplying with the factor $\mu_{\text{old}}/\mu_{\text{new}}$, where μ_{old} corresponds to the previously used mutation rate per year and μ_{new} to our upper and lower bounds based on the range of per generation mutation rates and generation intervals appropriate for the split under consideration.

No explicit mutation rate was assumed for the calculation of the split times of Neandertals and present-day humans in the original publication (41). However, the authors use a range of nuclear divergence times for orangutan-human to arrive at a human-chimpanzee divergence time of 5.6-8.3 million years. In order to recalibrate the Neandertal split time, we use the published nuclear divergence of ca. 1.3% between human and chimpanzee (8, 16) to convert these values to a mutation rate per year (corresponding to $1.1 - 0.7 \times 10^{-9}$).

Acknowledgements

We thank A. Abraham for laboratory assistance and R. Mundry for assistance with estimation of uncertainties in average parental ages. We thank the three reviewers and W.-H. Li for helpful comments on the manuscript. This project was funded by the Max Planck Society. K.E.L was supported by a fellowship from the Alexander von Humboldt Foundation. We also thank the many agencies and governments that support the field research (Supplemental Information).

Author contributions

Designed research: K.E.L., K.P., S.P., L.V.

Performed research: K.E.L., C.R., G.S.

Contributed new reagents/analytical tools: C.B., C.C., K.F., E.I., M.I.-M., J.C.M., M.N.M., M.M.R., T.S.S., D.W., R.M.W., R.W.W., K.Z.

Analyzed data: K.E.L., K.P., L.V.

Wrote the paper: K.E.L., K.P., B.V., S.P., L.V.

References

1. Sarich VM & Wilson AC (1967) Immunological time scale for hominid evolution. *Science* 158:1200-1203.
2. Uzzell T & Pilbeam D (1971) Phyletic divergence dates of hominoid primates: a comparison of fossil and molecular data. *Evolution* 25:615-635.
3. Burgess R & Yang Z (2008) Estimation of hominoid ancestral population sizes under bayesian coalescent models incorporating mutation rate variation and sequencing errors. *Molec Biol Evol* 25:1979-1994.
4. Dutheil JY, *et al.* (2009) Ancestral population genomics: the coalescent hidden Markov model approach. *Genetics* 183:259-274.
5. Hobolth A, Christensen OF, Mailund T, & Schierup MH (2007) Genomic relationships and speciation times of human, chimpanzee, and gorilla inferred from a coalescent hidden Markov model. *PLoS Genet* 3:e7.
6. Hobolth A, Dutheil JY, Hawks J, Schierup MH, & Mailund T (2011) Incomplete lineage sorting patterns among human, chimpanzee, and orangutan suggest recent orangutan speciation and widespread selection. *Genome Res* 21:349-356.
7. Patterson N, Richter DJ, Gnerre S, Lander ES, & Reich D (2006) Genetic evidence for complex speciation of humans and chimpanzees. *Nature* 441:1103-1108.
8. Scally A, *et al.* (2012) Insights into hominid evolution from the gorilla genome sequence. *Nature* 483:169-175.
9. Senut BP, M.; Gommery, D.; Mein, P.; Cheboi, K.; Coppens, Y. (2001) First hominid from the Miocene (Lukeino Formation, Kenya). *Earth and Planetary Sciences* 332:137-144.
10. Brunet M, *et al.* (2002) A new hominid from the Upper Miocene of Chad, Central Africa. *Nature* 418:145-151.
11. Wood B & Harrison T (2011) The evolutionary context of the first hominins. *Nature* 470:347-352.
12. Glazko GV & Nei M (2003) Estimation of divergence times for major lineages of primate species. *Molec Biol Evol* 20:424-434.
13. Becquet C & Przeworski M (2007) A new approach to estimate parameters of speciation models with application to apes. *Genome Res* 17:1505-1519.
14. Caswell JL, *et al.* (2008) Analysis of chimpanzee history based on genome sequence alignments. *PLoS Genet* 4:e1000057.
15. Prüfer K, *et al.* (2012) The bonobo genome compared with the chimpanzee and human genomes. *Nature* 486: 527-531.
16. Consortium TCSaA (2005) Initial sequence of the chimpanzee genome and comparison with the human genome. *Nature* 437:69-87.
17. Gibbs RA, *et al.* (2007) Evolutionary and biomedical insights from the rhesus macaque genome. *Science* 316:222-234.
18. Wilkinson RD, *et al.* (2011) Dating primate divergences through an integrated analysis of palaeontological and molecular data. *Syst Biol* 60: 16-31.
19. Roach JC, *et al.* (2010) Analysis of genetic inheritance in a family quartet by whole-genome sequencing. *Science* 328:636-639.

20. Awadalla P, *et al.* (2010) Direct measure of the de novo mutation rate in autism and schizophrenia cohorts. *Am J Hum Genet* 87:316-324.
21. Consortium GP (2010) A map of human genome variation from population-scale sequencing. *Nature* 467:1061-1073.
22. Fenner JN (2005) Cross-cultural estimation of the human generation interval for use in genetics-based population divergence studies. *Am J Phys Anthropol* 128:415-423.
23. Helgason A, Hrafnkelsson B, Gulcher JR, Ward R, & Stefansson K (2003) A populationwide coalescent analysis of Icelandic matrilineal and patrilineal genealogies: evidence for a faster evolutionary rate of mtDNA lineages than Y chromosomes. *Am J Hum Genet* 72:1370-1388.
24. Matsumura S & Forster P (2008) Generation time and effective population size in Polar Eskimos. *Proc Royal Society Biol* 275:1501-1508.
25. Tremblay M & Vezina H (2000) New estimates of intergenerational time intervals for the calculation of age and origins of mutations. *Am J Hum Genet* 66:651-658.
26. Won YJ & Hey J (2005) Divergence population genetics of chimpanzees. *Mol Biol Evol* 22:297-307.
27. Thalmann O, Fischer A, Lankester F, Paabo S, & Vigilant L (2007) The complex evolutionary history of gorillas: insights from genomic data. *Mol Biol Evol* 24:146-158.
28. Wich SA, *et al.* (2009) Orangutan life history variation. *Orangutans: Geographic Variation in Behavioral Ecology and Conservation*, eds Wich SA, Utami Atmoko SS, Mitra Setia T, & van Schaik CP (Oxford University Press, Oxford).
29. Harvey PH & Clutton-Brock TH (1985) Life history variation in primates. *Evolution* 39:559-581.
30. Benton MJ & Donoghue PCJ (2007) Paleontological evidence to date the tree of life. *Mol Biol Evol* 24:26-53.
31. Harrison T (2010) Apes among the tangled branches of human origins. *Science* 327:532-533.
32. Haile-Selassie Y, Suwa G, & White TD (2004) Late Miocene teeth from Middle Awash, Ethiopia, and early hominid dental evolution. *Science* 303:1503-1505.
33. Bernor RL (2007) New apes fill the gap. *Proceedings of the National Academy of Sciences of the United States of America* 104:19661-19662.
34. Ishida H & Pickford M (1997) A new Late Miocene hominoid from Kenya: *Samburupithecus kiptalami* gen. et sp. nov. *Comptes Rendus de l'Académie des Sciences - Series IIA - Earth and Planetary Science* 325:823-829.
35. Kunitatsu Y, *et al.* (2007) A new Late Miocene great ape from Kenya and its implications for the origins of African great apes and humans. *Proc Natl Acad Sci USA* 104:19220-19225.
36. Suwa G, Kono RT, Katoh S, Asfaw B, & Beyene Y (2007) A new species of great ape from the late Miocene epoch in Ethiopia. *Nature* 448:921-924.
37. Begun DR, Ward CW, & Rose MD (1997) Miocene Hominoid Evolution and Adaptations. *Function, Phylogeny and Fossils*, eds Begun DR, Ward CW, & Rose MD (Plenum Press, New York), pp 389-415.
38. Stewart CB & Disotell TR (1998) Primate evolution - in and out of Africa. *Current Biol* 8:R582-588.

39. Moya-Sola S, Kohler M, Alba DM, Casanovas-Vilar I, & Galindo J (2004) *Pierolapithecus catalaunicus*, a new Middle Miocene great ape from Spain. *Science* 306:1339-1344.
40. Green RE, *et al.* (2010) A draft sequence of the Neandertal genome. *Science* 328:710-722.
41. Hublin JJ (2009) Out of Africa: modern human origins special feature: the origin of Neandertals. *Proc Natl Acad Sci USA* 106:16022-16027.
42. Bischoff JL, *et al.* (2007) High-resolution U-series dates from the Sima de los Huesos yields 600 + ∞ —66 kyrs: Implications for the evolution of the Neanderthal lineage. *J Archaeol Sci* :763-770.
43. Endicott P, Ho SY, & Stringer C (2010) Using genetic evidence to evaluate four palaeoanthropological hypotheses for the timing of Neanderthal and modern human origins. *J Human Evol* 59:87-95.
44. Stringer CB & Hublin J (1999) New age estimates for the Swanscombe hominid, and their significance for human evolution. *J Human Evol* 37:873-877.
45. Rogers ME, *et al.* (2004) Western gorilla diet: a synthesis from six sites. *Am J Primatol* 64:173-192.
46. Leigh SR (1994) Ontogenetic correlates of diet in anthropoid primates. *Am J Phys Anthropol* 94:499-522.
47. Breuer T, Hockemba MB, Olejniczak C, Parnell RJ, & Stokes EJ (2009) Physical maturation, life-history classes and age estimates of free-ranging western gorillas—insights from Mbeli Bai, Republic of Congo. *Am J Primatol* 71:106-119.
48. Kelley J & Schwartz GT (2010) Dental development and life history in living African and Asian apes. *Proc Natl Acad Sci USA* 107:1035-1040.
49. Van Schaik C & Deaner R (2003) Life history and cognitive evolution in primates. *Animal Social complexity: Intelligence, culture, and Individualized Societies*, eds de Waal F & Tyack P (Harvard University Press, Cambridge, MA), pp 5-25.
50. Robson SL & Wood B (2008) Hominin life history: reconstruction and evolution. *J Anatom* 212:394-425.
51. Dean MC & Lucas VS (2009) Dental and skeletal growth in early fossil hominins. *Annals Human Biol* 36:545-561.
52. Smith TM, *et al.* (2010) Dental evidence for ontogenetic differences between modern humans and Neanderthals. *Proc Natl Acad Sci USA* 107:20923-20928.
53. Chen F-C & Li W-H (2001) Genomic divergences between human and other hominoids and the effective population size of the common ancestor of human and chimpanzee. *Am Journal Hum Genet* 68:444-456.

Figure legends

Figure 1: A diagram illustrating the branching pattern and timing of the splits between humans, chimpanzees, bonobos, western gorillas and eastern gorillas. The paler shading indicates the range of split times inferred in this study. Cartoon skulls indicate approximate age of the indicated fossil remains, but do not imply that these fossils were necessarily on those ancestral lineages, nor that entire crania actually exist for these forms.

Table 1. Generation intervals for each chimpanzee and gorilla study community

Taxa	Study site	No. Offspring	Mean Generation Interval in years					
			Female	CI	Male	CI	Both sexes	CI
Western chimpanzees	Tai-North	28	23.03	22.19-23.80	23.05	22.31-23.81	23.04	22.48-23.58
	Tai-Middle	4	31.71	28.34-35.15	26.06	23.90-28.32	28.89	26.80-31.03
	Tai-South	28	28.76	27.54-29.98	25.36	24.46-26.30	27.06	26.29-27.84
Eastern chimpanzees	Gombe-Kasekela	31	24.62	24.24-25.00	21.84	21.75-21.93	23.23	23.03-23.43
	Mahale-M	14	25.03	23.95-26.08	19.87	19.62-20.13	22.45	21.90-23.00
	Kibale-Ngogo	72	24.5	23.80-25.22	23.57	23.05-24.13	24.04	23.60-24.48
	Kibale-Kanyawara	15	23.34	22.43-24.25	28.42	27.15-29.75	25.88	25.04-26.68
	Budongo-Sonso	34	26.08	25.03-27.08	26.66	25.93-27.34	26.37	25.72-26.95
All chimpanzees		226	25.18	24.86-25.54	24.08	23.83-24.34	24.63	24.42-24.85
Mountain gorillas	Karisoke	97	18.18	17.97-18.37	20.27	20.23-20.30	19.22	19.12-19.32
	Bwindi	8	18.26	16.87-19.64	21.67	20.37-22.93	19.97	18.96-20.88
All gorillas		105	18.19	18.00-18.39	20.37	20.27-20.47	19.28	19.17-19.39
Humans*	Hunter-gatherers	157 societies	25.6		31.5		28.6	
	Countries	360 societies	27.3		30.8		29.1	

*Fenner, 2005 (22)

Table 2. Original and recalibrated population split times from several recent studies.

Speciation event	Original yearly mutation rate	Original split estimate (Ma)	Generation times		New yearly mutation rate		New split estimate (Ma)		Publication
			lower	upper	lower	upper	lower	upper	
HCG	1.0x10e-9	5.95	19	29	0.33	0.72	8.31	11.79	Scally et al. 2012 (8)
HCG	1.0x10e-9	6.69	19	29	0.33	0.72	9.35	20.00	Dutheil et al. 2009 (4)
HC	1.0x10e-9	3.69	25	29	0.33	0.54	6.78	11.03	Scally et al. 2012 (8)
HC	1.0x10e-9	4.22	25	29	0.33	0.54	7.76	12.62	Hobolth et al. 2011 (6)
HC	1.0x10e-9	4.5	25	29	0.33	0.54	8.27	13.45	Prüfer et al. 2012 (15)
HC	1.0x10e-9	4.38	25	29	0.33	0.54	8.05	13.09	Dutheil et al. 2009 (4)
BC	1.0x10e-9	0.99	25	25	0.39	0.54	1.82	2.55	Prüfer et al. 2012 (15)
BC	1.0x10e-9	0.79-0.92	25	25	0.39	0.54	1.45-1.69	2.04-2.37	Becquet and Przeworski 2007 (13)
wG-eG	0.96x10e-9	0.9-1.6	19	19	0.51	0.72	1.20-2.13	1.69-3.01	Thalmann et al. 2007 (27)
wG-eG	1.33x10e-9	0.92	19	19	0.51	0.72	1.29	1.80	Becquet and Przeworski 2007 (13)

HCG, human chimpanzee gorilla split; HC, human chimpanzee split; BC, bonobo chimpanzee split; wG-eG, western gorilla-eastern gorilla split.

