



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Genetic Circuitry of Survival Motor Neuron, the Gene Underlying Spinal Muscular Atrophy

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Sen, Anindya, Douglas N. Dimlich, K. G. Guruharsha, Mark W. Kankel, Kazuya Hori, Takakazu Yokokura, Sophie Brachat, et al. 2013. Genetic Circuitry of Survival Motor Neuron, the Gene Underlying Spinal Muscular Atrophy. Proceedings of the National Academy of Sciences 110, no. 26: E2371–E2380.
Published Version	doi:10.1073/pnas.1301738110
Accessed	February 16, 2015 7:38:14 PM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:12872186
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

The genetic circuitry of Survival Motor Neuron, the gene underlying Spinal Muscular Atrophy

Anindya Sen^{1*}, Douglas N. Dimlich^{1*}, K. G. Guruharsha^{1*}, Mark W. Kankel^{1*}, Kazuya Hori¹, Takakazu Yokokura^{1,2}, Sophie Brachat^{3,4}, Delwood Richardson³, Joseph Loureiro³, Rajeev Sivasankaran³, Daniel Curtis³, Lance S. Davidow⁵, Lee L. Rubin⁵, Anne C. Hart⁶, David Van Vactor¹, and Spyros Artavanis-Tsakonas¹. * Equal contribution

1.Department of Cell Biology, Harvard Medical School, Boston, MA 02115, USA. 2.Current affiliation: Okinawa Science and Technology Graduate University, 1919-1 Tancha, Onna-son, Okinawa 904-0495, Japan 3.Developmental and Molecular Pathways, Novartis Institutes for Biomedical Research, 250 Massachusetts Avenue, Cambridge, Massachusetts 02139, USA. 4.Current affiliation: Musculoskeletal Diseases, Novartis Institutes for Biomedical Research, Novartis Campus, CH-4002 Basel, Switzerland 5.Department of Stem Cell and Regenerative Biology, Harvard Medical School, Boston, MA 02115, USA. 6.Department of Neuroscience, Brown University, 185 Meeting Street Box GL-N, Providence, RI 02912, USA.

Submitted to Proceedings of the National Academy of Sciences of the United States of America

The clinical severity of the neurodegenerative disorder Spinal Muscular Atrophy (SMA) is dependent on the levels of functional Survival Motor Neuron (SMN) protein. Consequently, current strategies for developing treatments for SMA generally focus on augmenting SMN levels. To identify additional potential therapeutic avenues and achieve a greater understanding of SMN, we applied in vivo, in vitro, and in silico approaches to identify genetic and biochemical interactors of the Drosophila SMN homolog. We identified more than three hundred candidate genes that alter an Smn-dependent phenotype in vivo. Integrating the results from our genetic screens, large-scale protein interaction studies and bioinformatics analysis, we define a unique interactome for SMN which provides a knowledge base for a better understanding of SMA.

Genetic screen | Interactome | Proteomics | Spinal Muscular Atrophy | Survival Motor Neuron

INTRODUCTION

Spinal Muscular Atrophy (SMA), the leading genetic cause of infant mortality, results from the partial loss of Survival Motor Neuron (SMN) gene activity (1). Numerous studies indicate that SMN functions as a central component of a complex which is responsible for the assembly of spliceosomal small nuclear ribonucleoproteins (snRNPs) [reviewed in (2)]. SMN is also reported to play additional roles, including mRNA trafficking in the axon (3). In humans, SMN is encoded by two nearly identical genes, *SMN1* and *SMN2*, which are located on chromosome 5 (4). *SMN2* differs from *SMN1* in that only 10% of *SMN2* transcripts produce functional SMN due to a single nucleotide polymorphism that results in inefficient splicing of exon 7 and translation of a truncated, unstable SMN protein (1, 5, 6). The clinical severity of SMA correlates with *SMN2* copy number, which varies between individuals (7). As the small amount of functional SMN2 protein produced by each copy is capable of partially compensating for the loss of the *SMN1* gene function, higher copy numbers of *SMN2* typically result in milder forms of SMA. Therefore, genetic modifiers capable of increasing the abundance and/or specific activity of SMN hold promise as therapeutic targets.

The *Drosophila* genome harbors a single, highly conserved ortholog of SMN1/2, the *Survival motor neuron (Smn)* gene. SMN is essential for cell viability in vertebrates and *Drosophila* (8, 9). In *Drosophila*, zygotic loss of *Smn* function results in recessive larval lethality (not embryonic as might be expected) due to the rescue of early development by maternal contribution of *Smn*. The larval phenotype includes neuromuscular junction (NMJ) abnormalities that are reminiscent of those associated with the human disease, rendering this invertebrate organism an excellent system to model SMN biology (10-12). We previously described a genetic screen for modifiers of the lethal phenotype resulting from a complete loss of function *Smn* allele (13). This screen,

though it probed half of the *Drosophila* genome, identified only a relatively small number of genes that affected the NMJ phenotype associated with *Smn* loss of function (13). In particular, it did not identify genes involved in snRNP biogenesis, the molecular functionality that is most clearly associated with SMN.

As the human disease state results from partial loss of SMN function, we reasoned that a screening paradigm using a hypomorphic *Smn* background, (as opposed to a background that completely eliminates SMN function) would more closely resemble the genetic condition in SMA. Such a screen would therefore enhance our ability to detect novel elements of the *Smn* genetic network, and, consequently, add significantly to our efforts to both dissect the *Smn* genetic circuitry as well as identify potential clinically relevant targets with novel mode of action.

This complementary screen proved to be more sensitive than our previous screen and led to the identification of over 300 genetic interactors. Taking advantage of the recently established *Drosophila* Protein Interaction Map (DPiM) (14), we related the newly identified genetic interactors to the SMN protein interactome, producing an integrated *Drosophila* SMN biological network. Finally, the *Drosophila* SMN network was evaluated for its relevance to human biology by mapping *Drosophila* SMN network genes to their human homologs, and analyzing the human network using computational biology tools. The projection of the *Drosophila* SMN network derived from this study onto the human network derived from prior knowledge provides a rational basis for novel SMN functional hypotheses and network

Significance

Spinal Muscular Atrophy (SMA), the leading genetic cause of infant mortality, is a devastating neurodegenerative disease caused by reduced levels of Survival Motor Neuron (SMN) gene activity. Despite well-characterized aspects of the involvement of SMN in snRNP biogenesis, the gene circuitry affecting SMN activity remains obscure. Here, we use *Drosophila* as a model system to integrate results from large-scale genetic and proteomic studies, and bioinformatics analyses to define a unique SMN interactome to provide a basis for a better understanding of SMA. Such efforts not only help dissect the *Smn* biology but may also point to potential clinically relevant targets.

Reserved for Publication Footnotes

137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204




Fig. 1. Genetic modifiers of *Smn* using pupal lethality to screen the Exelixis collection of transposon insertions and their functional roles(A) *tubulinGAL4* (*tub-GAL4*) directed expression of an inducible *Smn*-RNAi construct (*UAS-Smn-RNAi*^{FL26B}) leads to a fully penetrant pupal lethality where approximately 40% of the pupae reach a pigmented developmental stage (Control). The remaining 60% die at an earlier unpigmented developmental stage. Introduction of an *Smn* deficiency into this background causes the entire population of pupae to die at the unpigmented stage (*Smn* deficiency), while ectopic *Smn* expression leads to survival to adulthood of the vast majority of pupae (*Smn* rescue). Introduction of previously isolated enhancers (d02492 and d09801) and suppressors (f05549 and c05057) of *Smn* (13) lead to quantitative changes in the fractions of pigmented vs. unpigmented pupae. (B) The screening strategy to identify genetic modifiers of the *Smn* pupal lethality phenotype using the Exelixis collection (illustrated for insertions on the 3rd chromosome). The lethal phase for all *Smn* *Tb*+ TE (16) pupae in individual test crosses are scored and compared to those observed in control crosses (more survival = enhancers, more lethality = suppressors). (C) *Drosophila* functional categories over-represented in the genetic modifier list. GO biological functions with the highest significance relate to known *Smn* functions such as alternative splicing or SMA affected processes (neuronal and muscular). Enrichment significance is expressed as the $-\log_{10}$ (p-values).

205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272

intervention points that carry potential for so far unexplored clinical applications.

RESULTS

A genetic screen for modifiers of *Smn*-dependent lethality

We examined several *Smn*-RNAi strains to identify a hypomorphic *Smn* allele that could be used to model SMA in *Drosophila* more faithfully than alleles that completely abolish *Smn* function. We identified a transgenic strain, *UAS-Smn-RNAi*^{FL26B} (FL26B) that displays a less severe phenotype than the allele used in our previous screen (13). Specifically, expression of FL26B under the control of *tubulinGAL4* (*tubGAL4*:FL26B) results in late pupal lethality whereby approximately 50% of the pupae reach a more mature (pigmented) developmental stage prior to death (Figure 1A) than their less mature, unpigmented siblings.

We determined that this phenotype, measured by the ratio of pigmented to unpigmented pupae, is sensitive to *Smn* dosage, as reducing or increasing *Smn* copy number in the *tubGAL4*:FL26B genetic background resulted in enhancement or suppression, respectively (Figure 1A). In addition, the ability of wild type *Smn* (expressed by a *UAS-Smn-GFP* transgene) to rescue the lethality indicates that this phenotype does not result from off target RNAi effects. These results were corroborated using an independent *Smn* RNAi strain. Finally, we demonstrated that previously identified *Smn* modifiers altered the *Smn* RNAi phenotype in the ex-

pected fashion (Figure 1A). Together, these results demonstrate that the *tubGAL4*:FL26B phenotype is useful to detect changes in *Smn* functional activity and is thus suitable assay on which to base a modifier screen that will define and dissect the *Smn* genetic network.

Using this novel assay, we screened the Exelixis collection of genome-wide insertional mutations [https://drosophila.med.harvard.edu/ and (15, 16)] for dominant modification of the lethality associated with the *tubGAL4*:FL26B strain (see Figure 1B for scheme) and identified nearly 1600 candidate strains. To eliminate false positives, all interacting insertions were retested. Only those that were not lethal in *trans* with *tubGAL4* (13) and for which results could be clearly repeated were finally designated as modifiers. From this analysis, we identified 303 modifying strains (129 enhancers and 174 suppressors), which represents nearly 2% of the collection and a greater than tenfold increase in hit recovery in comparison to our previous screening method. As the genomic location of some insertions in the Exelixis collection may be near multiple coding regions, unambiguous gene assignments are not always possible. Given this consideration, we determined that these 303 insertions potentially affected 340 *Drosophila* genes. In most cases, single genes were affected by single insertions, though 14 genes scored in the screen were represented by 2 or more alleles. No gene could be assigned for 36 of the 303 insertions. Careful human to *Drosophila* homology mapping using a combination of

273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340

341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408





Fig. 2. The extended *Drosophila* genetic sub-network The sub-network of proteins connected to Smn and its genetic modifiers in the *Drosophila* Protein Interaction Map (DPIIM). A total of 62 Smn genetic modifiers (diamonds with red border) are directly connected to 361 other proteins (circles), also known as first-degree neighbors through 3,800 interactions. The thickness of the gray lines connecting the proteins is proportional to the interaction score in the DPIIM. Proteins belonging to individual clusters with GO term enrichment are shown with different colors. Proteins colored gray are part of clusters that are not enriched for any specific GO terms. Smn protein (indicated by an arrow head) itself is only connected to Nmdmc and shown as an interacting pair at the bottom.

several prediction algorithms show that out of the 340 *Drosophila* genes, at least 229 have human orthologs. Since a fraction of

409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476



- activation — green
- expression — cyan
- inhibition — red
- localization — purple
- modification — orange
- phosphorylation — pink
- protein-DNA interactions — grey
- protein-protein interactions — black
- proteolysis — red
- reaction — yellow
- regulation of binding — blue
- transcription — green
- translocation — light blue

Fig. 3. Human Snn genetic modifiers network Ingenuity Pathway Analysis indicates that about one-third of the human orthologs of *Drosophila* Snn genetic modifiers (103 genes, green circles) are connected in a network involving 282 interactions with other modifiers and SMN1/SMN2 (yellow). Different types of interactions are indicated with distinct colored lines. A small number of modifiers (19 genes) have only interactions with other modifiers but not SMN1/SMN2 (4 pairs are two human orthologs of the same *Drosophila* gene). Nearly two-thirds of the modifiers (177 genes, not shown) have no interactions that would connect them to SMN1/SMN2 or other modifiers

477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544

these genes are represented by multiple paralogs, we identified a total of 322 human genes corresponding to the 229 *Drosophila* modifiers (see Materials and Methods, **Dataset S1**).

To assess the biological space covered by the novel *Snn* modifiers they were evaluated using the **Database for Annotation, Visualization and Integrated Discovery (DAVID)** (17, 18) (<http://david.abcc.ncifcrf.gov/>), which identifies biological processes statistically overrepresented in the set of genetic modifiers. Analysis of the *Drosophila* SMN network with DAVID identified known SMN molecular activity (alternative splicing) and SMN-dependent processes (“neuron differentiation”, “axon guidance”, “axonogenesis”, “muscle organ development” and “dendrite morphogenesis/development”) (**Figure 1C**) (13, 19). The predominant processes enriched in the modifier set reflect broad effects on morphogenesis and development reinforcing the notion that *Snn* depletion has pleiotropic consequences.

Integration of *Drosophila* genetic and proteomic interactors

To determine whether the genetic modifiers are interconnected through physical interactions, we placed them in the context of the recently generated *Drosophila* Protein Interaction Map (DPiM) (14). We first retrieved the set of proteins that co-purify with *Drosophila* Snn in DPiM, and asked whether any of the modifiers belong to this Snn sub-network. From all *Drosophila* proteins tested, 8 form protein complexes with Snn and interestingly one, *NAD-dependent methylenetetrahydrofolate dehydrogenase (Nmdmc)*, is also a genetic modifier. Expansion of the Snn subnetwork to include proteins that form complexes with each of the eight members in turn identified 35 additional proteins. Some of these proteomic interactors have been previously associated with Snn function (e.g. Gemin2, Gemin3 and several snRNPs), and nearly half (20/43) are known to be involved

in mRNA processing, a functionality closely linked to the documented biochemical role of SMN (**Figure S1, Dataset S2**).

To better understand the biological functions identified by the Snn modifiers, we extracted the *Drosophila* protein complexes, which include the Snn genetic modifiers in DPiM. We found that 62 of the proteins corresponding to genetic modifiers passed the stringent statistical criteria necessary for inclusion in the DPiM interactome (**Materials and Methods**) which includes only the top 5% of the total interactions scored in the coAP/MS analysis. These 62 proteins were associated with 50 separate Markov clusters, a statistical definition of significantly associated proteins (20), each of which may define a functional protein complex (see **Figure 2** for the sub network of complexes identified by this analysis). Of these 50, we focused our attention on the 24 that are enriched for specific biological functions based on Gene Ontology (GO) terms – a system that provides a controlled vocabulary of terms for describing gene cellular and molecular functions (21). The majority of these clusters harbor a single modifier, but 9 contained two or more (**Table S3**).

Inspecting the GO biological function terms of these 24 clusters (annotated in **Figure 2**), we find that many contain annotated functions previously linked to *Snn* activity including RNA metabolism (“RNA splicing”, “mRNA binding”) [reviewed in (3)], translation control (“eIF3 complex”) (22-25), endocytosis (“Snap/SNARE complex”) (26, 27), and protein transport (“flotillin complex”) (28-30). Importantly, several genetic modifiers fall within protein complexes whose functions have not been previously associated with *Snn* function. These include complexes with phosphatase and kinase activities as well as those involved in intracellular signaling (“Toll signaling pathway” and “Hedgehog signaling”) (**Figure 3B**). Two independent loss-of-function alleles of the *Ect4/dsarm* gene suppress *Snn* dependent

545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612

613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680




Fig. 4. Extended human Smn genetic sub-network Adding SMN1/2 first-degree neighbors to the network shown in Figure 3 generated an extended sub-network. In this interactome, 151 human orthologs of *Drosophila* Smn genetic modifiers (green circles) directly or indirectly connected to SMN1/SMN2. A total of 48 modifiers are connected to SMN1/SMN2 through 71 additional intermediate proteins (pink circles) from literature, seven among them (blue circles) were also identified in replicate SMN bait purifications in *Drosophila*. Different types of interactions are indicated with distinct colored lines

lethality. The recovery of *Ect4/dsarm* may provide additional evidence linking the Toll signaling pathway to *Smn* activity as it

681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748




Fig. 5. Network of 36 high-priority genetic interactors of SmnThe network shows 36 human orthologs of *Drosophila* Smn genetic modifiers (green circles) connected to SMN1/SMN2 in human. These 36 modifiers are present in *Drosophila* DPIIM as well as human IPA based network and were selected for functional validation in NMJs. The intermediate proteins (pink circles) shown provide the shortest path to connect the modifiers to SMN1/SMN2. Different types of interactions are indicated with distinct colored lines.

749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816




Fig. 6. Genetic modifiers of Smn regulate NMJ morphology. (A) An NMJ derived from muscle 6/7 of a *tubulinGAL4::UAS-Smn-RNAi^{FL26B}* (*tubGAL4::FL26B*) 3rd instar larva. (B, C) A reduction in NMJ size is observed upon introduction of enhancers c02456 (B) and d02738 (C) into the *tubGAL4::FL26B* background. (D, E, F) Introduction of a suppressors c05501, c06705 into this screening background leads to an increase in NMJ size, whereas suppressor d05711 (F) does not result in significant modification of the NMJ (G) Quantitation of bouton numbers/muscle in individuals of indicated genotypes, which include enhancers (red) and suppressors (green), normalized per muscle surface area (MSA) and expressed as percentage change as compared to Tub Gal4:Smn RNAi alone. The ANOVA multiple comparison test was used for statistical analysis of the bouton numbers/muscle. Significance P < 0.05. Scale Bar = 50 mm. n = 20. All preparations were stained with anti-HRP (red) and anti-Dlg (green). The muscle nucleus was labeled using DAPI.

encodes a Toll/interleukin-1 receptor homology (TIR) domain. Intriguingly, loss-of-function *Ect4/dsarm* mutations also suppress Wallerian degeneration phenotypes observed in *Drosophila* and

mouse models (31). Together, these data suggest that the Wallerian degeneration pathway may also affect Smn pathobiology, an effect that may be mediated through Toll signaling. Hence this

approach both confirmed and expanded the functional categories and pathways associated with SMN.

To further explore the relationships of the 62 proteins and their functional context within DPiM, we carried out a first-degree neighbor analysis to identify other proteins directly connected in the network that may represent potential biochemical interactors. This retrieved 361 additional proteins that are linked to the 62 *Smn* modifier interactors (Figure 2, Dataset S3). These 361 proteins include 128 that are directly linked to at least two of the 62 modifiers (Table S3). A GO term (i.e. functionalities) analysis of these proteins reveals additional connections to the spliceosome, RNA binding and Snap/SNARE functions. Thus, considering genetic modifiers in the context of the DPiM provides us with a novel perspective of the diverse molecular functions that can modulate SMN activity *in vivo*.

Overlaying the genetic modifiers on the human interactome

To study the *Smn* genetic circuitry in the human context, we generated a human view of the genetic *Smn* interactome taking advantage of the manually curated source of human molecular interactions from IPA (Ingenuity® Systems, www.ingenuity.com). This database integrates human gene relationships derived from a variety of experimental approaches, including proteomic studies. Using the human *Smn* proteins and the 322 human genes corresponding to the genetic modifiers identified in *Drosophila* (see above and Materials and Methods), we used the IPA knowledgebase to derive a human SMN interaction network. Unlike DPiM, IPA is not limited to physical interactions thus allowing consideration of other functional interactions including, for example, expression, localization, modification, and regulation. Such an approach allowed us to evaluate potential indirect relationships between the modifiers and SMN, and uncover molecular functions beyond its canonical role in the SMN complex.

Based on the generated network, we found that orthologs of five modifiers HNRNPR, SNRPD1, SYNCRIP, TRA2B and ZNF259 are directly related to SMN1/2 (Figure 3). HNRNPR, SNRPD1 and SYNCRIP proteins physically interact with SMN1 and 2 and have a role in RNA splicing (32). *Trab2* (also known as SFRS10 or Htra2-beta) was shown to regulate *Smn2* protein levels by being a potent splicing enhancer (33). Finally, ZNF259 also known as ZPR1 was shown to be necessary for the localization of *Smn1* to nuclear bodies (34) and more recently emerged as a key modifier of SMA pathology in patients (35). These findings support the relevance of the identified *Drosophila* modifiers in understanding the human pathways underlying SMA pathology.

Furthermore, 98 modifiers are indirectly related via these five interactors to human SMN. Together these 103 proteins representing one-third of the identified modifiers are interconnected in a human IPA database. In addition, we find another group of 19 proteins that make pair-wise functional interactions with other SMN genetic modifiers, but do not connect to the human *Smn* interactome. The remaining 177 proteins that are not connected in the human interactome (and the 19 that have pair-wise connections, 4 pairs of which are between two orthologs of the same *Drosophila* gene) potentially represent functions that have not been linked to SMN biology in human studies so far.

Expansion of the human SMN interactome beyond the 103 modifiers, by incorporating first-degree neighbor proteins of SMN in the database, connects an additional 48 modifiers (Figure 4). This expanded human SMN network contains intermediates that are known to associate physically with SMN (GEMINs, HNRNPs and LSM and SNRP family members) (3, 14) and signaling pathway elements known to affect SMN activity (FGF2, GSK3B, MAP3K5) (19, 36-39).

Validation of genetic modifiers at the larval NMJ

We chose to prioritize the modifiers for further functional characterization by using membership within both the *Smn* modifier network in DPiM (Figure S1) and the expanded IPA Human

SMN network (Figure 4) as the primary criterion. A total of 36 genes are shared between these interactomes (Figure 5, Dataset S4). The list includes 4 previously analyzed modifiers (*Actn*, *Moes*, *Fim*, *cut up*) (13, 27), 13 enhancers, (*Sod*, *Hsp68*, *Hsf*, *step*, *CG17838*, *ns1*, *shrb*, *VhaSFD*, *Rel*, *Hexo2*, *osa*, *CG13902*, *cathD*) and 19 suppressors (*CG30194*, *Nedd4*, *Pka-R2*, *Rho1*, *Tango7*, *Argk*, *14-3-3-epsilon*, *Zpr1*, *CG9769*, *cenG1A*, *flw*, *comt*, *CG9062*, *l(3)72Ab*, *Karybeta3*, *HmgZ*, *Rbsn-5*, *sel*, *Paip2*).

Our previous analyses (13) indicated a strong correlation between the strength of the lethal *Smn* phenotype with the severity of NMJ abnormalities. Therefore, examination of the effects of *Drosophila* modifiers on the *Smn* NMJ phenotype was used to validate their role in *Drosophila* and prioritize the corresponding orthologs for further investigation in vertebrate model systems. We used NMJ assays (13, 19) to sample the ability of a subset of these modifiers to alter the *tubGAL4::FL26B* NMJ phenotype. Examination of third instar larvae carrying a combination of *tubGAL4::FL26B* and each of 20 modifier strains revealed that 11 out of the 20 (55%) strains reveal a statistically significant change in the number of synaptic boutons and are modifiers of the *Smn* NMJ phenotype (Figure 6).

Effect of genetic modifiers on *Smn* protein levels and localization

Given that the severity of the disease phenotypes, in both patients and *Drosophila* models, correlates with SMN protein levels, we examined whether the prioritized genes affected SMN levels in *Drosophila* S2R+ cells (40), the same cell line used to generate DPiM. We used an image-based analysis (37) to quantify SMN protein levels in S2R+ cells expressing inducible FLAG-HA tagged constructs corresponding to 21 *Smn* modifying genes available from the Universal Proteomics Resources (41). Untransfected cells within the same wells were used as controls. Surprisingly, we found that none of these ectopically expressed modifier genes significantly altered total *Smn* protein expression (Figure S2A). Since *Smn* is localized in both the cytoplasm and the nucleus, we also used this assay to evaluate whether any of these modifiers altered its distribution between these two compartments. We found seven modifiers significantly increased the nuclear *Smn* levels (Figure S2B and Dataset S5), consistent with the notion that some modifiers from the screen, which affect *Smn* lethality and NMJ phenotype, may directly affect *Smn* distribution between the nucleus and cytoplasm. It is worth noting that, a recent study (42) showed that mutant superoxide dismutase-1 (SOD1), known to cause familial ALS, alters the sub-cellular localization of the SMN protein and disrupts its recruitment to Cajal bodies thereby preventing the formation of nuclear 'gems'. *Sod* was identified in our screen as an enhancer and was also shown to affect NMJ phenotype (Figure 6). A subset of modifiers does not alter either *Smn* levels or its localization. How these modifiers affect the functional *Smn* remains to be explored. Given these results, however, it is important to note that small changes in SMN function may have an important biological impact given that the severity of clinical manifestation in SMA patients correlates with small changes in SMN expression (1).

DISCUSSION

Different animal models for SMA-associated neuromuscular defects contributed significantly to a better understanding of the Spinal Muscular Atrophy etiology and genetics over the last few years. However, despite the well-characterized role for SMN in snRNP biogenesis, the links between its molecular function and the defects observed in SMA patients remain unclear. One of the key features of SMA is that the severity of the disease is dependent on SMN dosage, prompting the development of therapeutic strategies designed to increase SMN protein levels in patients. Still, it is essential to identify alternative approaches to modulate SMN activity. For this purpose, genetically tractable invertebrate

953 systems may help to identify so far undiscovered elements of
954 the SMN genetic circuitry. In particular, these organisms provide
955 more flexible avenues to investigate the poorly understood role
956 of SMN at the NMJ.

957 We have used *Drosophila* as our experimental system and
958 previously described a genetic screen which uncovered a small
959 number of *Smn* modifiers (13) of a strong loss of function mutant
960 phenotype. In this screen, we identified functional links between
961 *Smn* and the FGF pathway (13, 19), a relationship corroborated
962 and extended by recent evidence in a severe mouse model of
963 SMA, which demonstrated widespread alterations of the FGF-
964 system in both muscle and spinal cord (38).

965 The relatively small number of modifiers recovered suggested
966 a more sensitive genetic screen could provide extended informa-
967 tion about the *Smn* genetic network. Our assessment of the lethal
968 phase exhibited by a mild loss of function *Smn* RNAi allele, which
969 more closely resembles the SMA hypomorphic condition, pro-
970 vided us with a more sensitive and quantifiable assay for genetic
971 interaction. In comparison to our previous results, the RNAi-
972 based screen described here provided us with a broader spectrum
973 of modifiers including those related to the canonical role of *Smn*
974 in snRNP biogenesis as well as additional elements of FGF and
975 BMP signaling (13). Our careful mining of the screening mod-
976 ifier list based on functional term enrichment, and interactome
977 analysis both in *Drosophila* and human, suggest that loss of *Smn*
978 function may impact a range of developmental and maintenance-
979 related programs of the whole neuromuscular system, including
980 synaptic vesicle recycling, ion channels and signaling pathways
981 that regulate intrinsic cellular functions. Finally, this analysis also
982 uncovered biological processes not previously associated with
983 *Smn*.

984 Among the newly recovered genes, many are associated with
985 RNA metabolism; however, the majority is not involved with
986 canonical SMN activity of snRNP biogenesis and includes fac-
987 tors involved in transcription, post-transcriptional modifications,
988 RNA transport and translation regulation. Intriguingly, *CG17838*
989 is the *Drosophila* homolog of two closely related vertebrate RNA-
990 binding proteins, hnRNP-R and SYNCRIP/hnRNP-Q, both of
991 which bind to SMN in a yeast two-hybrid assay (32) and localize
992 to mRNA containing granules that are transported in cultured
993 neurons (28, 32, 43). Since both SMN and hnRNP-R affect lo-
994 calization of mRNA in axons (44, 45), this could have profound
995 consequences on local translation in neurons (45).

996 Given the complexity of motor neuronal sub-cellular domains
997 and their distance from the neuromuscular synapses, local reg-
998 ulation of the translation of synaptic proteins is likely to be
999 important in synaptic plasticity and neurological diseases. In fact,
1000 many mRNA binding proteins (RBPs) that function as key regu-
1001 lators of local RNA translation are associated with neurological
1002 diseases, including FMRP in Fragile X Syndrome, ATXN-2 in
1003 Spinocerebellar Ataxia, and TDP-43, FUS (fused in sarcoma),
1004 ANG and ATXN-2 in Amyotrophic Lateral Sclerosis. Consistent
1005 with a possible role for *Smn* in affecting local translation, we
1006 recovered *pumilio* and *eIF-4E*, which are thought to be a part
1007 of the local translational apparatus in neuromuscular synapses
1008 (46). Furthermore, we recovered another translation regulator,
1009 *eIF-4A*, which negatively regulates BMP signaling components.
1010 Components of BMP signaling pathway have been shown to play
1011 a role in retrograde signaling in the NMJ (47, 48). Our results
1012 support the relationship between *Smn* and local translation and
1013 also provide an additional link to the retrograde signaling present
1014 in the neuromuscular system. Interestingly, perturbation of RNA
1015 translational control may result in defects in endocytosis (49, 50),
1016 a process that has been suggested play a key role in neurode-
1017 generative diseases, including Alzheimer's (51) and Huntington's
1018 diseases (52). Consistent with this notion, aberrant synaptic ves-
1019 cle release at the NMJs in severe SMA mice may be evidence of

1020 impaired synaptic vesicle dynamics and/or abnormal active zone
1021 architecture (53). Further supporting a link between endocyto-
1022 sis and *Smn* (27), we identified *Synaptotagmin1*, *Synaptotagmin-*
1023 *alpha*, *Syntaxin4* and *comatose*, the *Drosophila* homolog of N-
1024 ethylmaleimide sensitive factor (NSF), which are core compo-
1025 nents of synaptic vesicle recycling. We also recovered genes that
1026 are directly required for synaptic transmitter release, such as
1027 *methuselah*, or indirectly, such as *bruchpilot*, which plays a role
1028 in constructing the active zone.

1029 Though many of the recovered genes broadly impact the neu-
1030 romuscular system, a subset includes the *Drosophila* homologs for
1031 several disease related genes, including *Nrx-1* (schizophrenia and
1032 Autism Spectrum Disorders) (54, 55), *Dystrophin* (Duchenne's
1033 Muscular Dystrophy) (56, 57), *Superoxide dismutase* (42, 58),
1034 *RhoA*, (Amyotrophic Lateral Sclerosis) (59) and *Ect4/dSarm*
1035 (Wallerian degeneration) (31). Our recovery of these genes sug-
1036 gests the genetic network identified by our screen may overlap,
1037 perhaps significantly, the genetic networks impacted by other
1038 human neurological disorders. If true, the use of *Drosophila* to
1039 explore other neurological disease networks via genetic screens,
1040 combined with the integration of additional genome-wide ap-
1041 proaches, could identify common therapeutic targets which could
1042 potentially be tested in other disease models.

1043 Since such genetic modifier screens are very sensitive and
1044 are able to recover a large number of modifiers that span a
1045 broad range of molecular functions, prioritization of candidates
1046 for further validation is essential. Here, bioinformatics mining
1047 allowed us to assemble a list of 36 *Drosophila* genes with hu-
1048 man homologs for continued investigation. The majority of these
1049 tested genes showed a functional role in the structure and/or
1050 development of the NMJ in *Drosophila*, and some can alter the
1051 distribution of *Smn* in S2R+ cells, making them good candidates
1052 to pursue in vertebrate models of SMA. In addition, candidates
1053 may be drawn from a pool of modifiers that include members of
1054 signaling pathways such as GPCR, Kinases and Proteases, which
1055 are considered to be plausible small molecule targets, or secreted
1056 or membrane proteins, which may be targeted by antibodies. Our
1057 results thus provide an extensive list of novel genes and pathways
1058 that have now been functionally linked to an *Smn*-dependent
1059 phenotype and therefore represent potentially novel therapeutic
1060 targets.

1061 MATERIALS AND METHODS

1062 *Drosophila* stocks and culture

1063 All *Drosophila* stocks were maintained on standard *Drosophila* medium
1064 at 25°C. The generation of the *Smn* alleles and constructs used in this
1065 study (*Smn*^{X7}, *UAS-Smn-RNAi*^{FL26B}, *UAS-Smn-GFP*) were originally described
1066 in (13). The tubulinGAL4 and TM6B, *Tb Hu tubulinGAL80* chromosomes
1067 used to generate the screening stock were obtained from the Bloomington
1068 *Drosophila* Stock Center (Bloomington, IN). The Exelixis Collection is housed
1069 in the Artavanis-Tsakonas laboratory in the Department of Cell Biology at
1070 Harvard Medical School (Boston, MA).

1071 Genetic Modifier Screen

1072 Individual strains from the Exelixis Collection were tested for the ability
1073 to genetically modify the *tubGAL4*-induced *UAS-Smn-RNAi*^{FL26B} pupal lethal
1074 phenotype by mating 3-5 males of the strain to 3 females of the *w*; *UAS-*
1075 *Smn-RNAi*^{FL26B}; *tubGAL4/TM6B*, *Hu Tb tubGAL80* screening stock. After two
1076 days, adults were transferred to a fresh vial to create a duplicate cross and
1077 to maintain optimal culture density. Accordingly, adults were discarded from
1078 the duplicate after an additional two days had passed. 15 days after being
1079 initiated, crosses were scored by counting the number of pigmented and
1080 unpigmented *Tb*+ pupae along with any *Hu*+ adult escapers. The ratio of
1081 unpigmented pupae to the total of pigmented pupae + escapers adults
1082 was compared to that derived from control crosses using males from the
1083 isogenic strain in which the Exelixis Collection was generated. Two control
1084 crosses were performed for each set of approximately 100 strains that were
1085 tested. Crosses that failed to produce 40 experimental animals were repeated
1086 as above. A change in the ratio of unpigmented to pigmented individuals
1087 of 20% corresponded to an approximate 1.5 standard deviation from the
1088 mean. Enhancers were defined as those mutations causing a reduction in
1089 this percentage ($\leq 30\%$), while those that increased this percentage ($\geq 70\%$)
1090 were classified as suppressors.

1091 Gene Assignments for the Exelixis Collection of Transposon Insertions

1089 Data for *Drosophila* genes and Exelixis transposon insertion sites were
1090 obtained from FlyBase version 5.39, which was current as of August 2011. Of
1091 the 15,952 Exelixis stocks screened, 14,621 stocks were mapped in FlyBase
1092 to 15,326 transposons with specific insertion sites within the *Drosophila*
1093 genome. To determine the coordinates of insertion sites of transposons
1094 present in the remaining 331 strains, sequences from the region flanking the
1095 insertion sites (15) were searched against the *Drosophila* genome using the
1096 blastn program of the Basic Local Alignment Search Tool [BLAST, (60)]. The
1097 insertion sites of the transposons were then used to create gene assignments
1098 according to the following criteria: a transposon was considered to map to
1099 a particular gene if its insertion site is located in the transcription unit of the
gene itself or within either 1 kb upstream of the transcription start site or
100 bp downstream of the transcription termination site.

Mapping *Drosophila* genes to human orthologs

1100 FlyBase version v5.39 identifies 15,233 *Drosophila* genes, which were
1101 iteratively mapped to human orthologs using predictions made by several
1102 prediction algorithms. Multiple predictions were combined into a single
1103 prediction by ordering the algorithms based upon lowest false positive and
1104 highest false negative rates [see (61)], and choosing the first prediction. The
1105 methods used (in order) were inParanoid version 7 (62), orthoMCL version 5
1106 (63), Homologene build 65 (64) and, orthoMCL version 2 (65). The inParanoid
1107 predictions were selected using a probability score of 0.4. As a result, of the
1108 15,233 *Drosophila* genes considered, 6,821 could be mapped to 6,703 human
1109 gene ids. This dataset was used to assign the human orthologs of *Drosophila*
Snn modifiers as shown in Dataset S1.

Function and network analysis

1110 The functional enrichment Gene Ontology terms of the *Drosophila*
1111 genetic modifiers was assessed using EASE statistics available through the
1112 Database for Annotation, Visualization, and Integrated Discovery (DAVID)
1113 Bioinformatics Resources using the Exelixis collection as a reference set (17,
1114 18). Human and *Drosophila*, protein-protein and genetic interactions were
1115 visualized and analyzed (first neighbors) using Cytoscape (66). Cytoscape
1116 BinGo plugin was used to evaluate the Functional categories of the retrieved
1117 clusters in the *Drosophila* sub-networks. The human network was generated
1118 through the use of IPA (Ingenuity Systems, www.ingenuity.com) and further
1119 visualized and mined in Cytoscape.

Neuromuscular junction analyses

1120 Third instar larvae were dissected in cold 1X phosphate buffered saline
1121 (PBS) and fixed at room temperature (RT) for 20 min. in 4% Paraformalde-
1122 hyde (PFA). The samples were washed in 0.1% Triton-X100 in PBS (PTX) and
1123 incubated overnight at 4°C with primary antibody. The primary antibody was
1124 washed off with PTX at RT. The samples were incubated at RT with secondary
1125 antibody for 90 min. This was followed by PTX wash, and the tissues were
1126 mounted in Vectashield Mounting Media with DAPI (Vector Laboratories).

- 1127 1. Lefebvre S, et al. (1997) Correlation between severity and SMN protein level in spinal
1128 muscular atrophy. *Nat Genet* 16(3):265-269.
- 1129 2. Workman E, Kolb SJ, & Battle DJ (2012) Spliceosomal small nuclear ribonucleoprotein
1130 biogenesis defects and motor neuron selectivity in spinal muscular atrophy. *Brain Res* 1462:93-
1131 99.
- 1132 3. Burghes AH & Beattie CE (2009) Spinal muscular atrophy: why do low levels of survival
1133 motor neuron protein make motor neurons sick? *Nat Rev Neurosci* 10(8):597-609.
- 1134 4. Lefebvre S, et al. (1995) Identification and characterization of a spinal muscular atrophy-
1135 determining gene. *Cell* 80(1):155-165.
- 1136 5. Lorson CL, Hahnen E, Androphy EJ, & Wirth B (1999) A single nucleotide in the SMN gene
1137 regulates splicing and is responsible for spinal muscular atrophy. *Proc Natl Acad Sci U S A*
1138 96(11):6307-6311.
- 1139 6. Monani UR, et al. (1999) A single nucleotide difference that alters splicing patterns distin-
1140 guishes the SMA gene SMN1 from the copy gene SMN2. *Hum Mol Genet* 8(7):1177-1183.
- 1141 7. Feldkotter M, Schwarzer V, Wirth R, Wienker TF, & Wirth B (2002) Quantitative analyses of
1142 SMN1 and SMN2 based on real-time lightCycler PCR: fast and highly reliable carrier testing
1143 and prediction of severity of spinal muscular atrophy. *Am J Hum Genet* 70(2):358-368.
- 1144 8. Schrank B, et al. (1997) Inactivation of the survival motor neuron gene, a candidate gene for
1145 human spinal muscular atrophy, leads to massive cell death in early mouse embryos. *Proc*
1146 *Natl Acad Sci U S A* 94(18):9920-9925.
- 1147 9. Chan YB, et al. (2003) Neuromuscular defects in a *Drosophila* survival motor neuron gene
1148 mutant. *Human molecular genetics* 12(12):1367-1376.
- 1149 10. Chan YB, et al. (2003) Neuromuscular defects in a *Drosophila* survival motor neuron gene
1150 mutant. *Hum Mol Genet* 12(12):1367-1376.
- 1151 11. Miguel-Aliaga I, Chan YB, Davies KE, & van den Heuvel M (2000) Disruption of SMN
1152 function by ectopic expression of the human SMN gene in *Drosophila*. *FEBS Lett* 486(2):99-
1153 102.
- 1154 12. Rajendra TK, et al. (2007) A *Drosophila melanogaster* model of spinal muscular atrophy
1155 reveals a function for SMN in striated muscle. *J Cell Biol* 176(6):831-841.
- 1156 13. Chang HC, et al. (2008) Modeling spinal muscular atrophy in *Drosophila*. *PLoS ONE*
3(9):e3209.
14. Guruharsha KG, et al. (2011) A Protein Complex Network of *Drosophila melanogaster*. *Cell*
147(3):690-703.
15. Thibault ST, et al. (2004) A complementary transposon tool kit for *Drosophila melanogaster*
using P and piggyBac. *Nat Genet* 36(3):283-287.
16. Artavanis-Tsakonas S (2004) Accessing the Exelixis collection. *Nat Genet* 36(3):207.
17. Huang da W, Sherman BT, & Lempicki RA (2009) Systematic and integrative analysis of large
gene lists using DAVID bioinformatics resources. *Nat Protoc* 4(1):44-57.

Bouton numbers were counted using a Nikon TE2000 microscope, based
on the Discs large and anti-HRP staining in the A3 segment muscle 4 as
indicated. The muscle area for every animal was measured, and no significant
difference was observed among different genotypes. At least 20-25 animals
of each genotype were dissected for the bouton analysis. The ANOVA
multiple comparison test was used for statistical analysis of the bouton
number/muscle.

Microscopy

1162 All images were collected with a Nikon C1si spectral point scanning
1163 confocal connected to a Nikon TE2000 inverted microscope equipped with
1164 DIC, phase, and epi-fluorescence optics, 40x Plan Fluor NA 1.4 objective lens
1165 and the Perfect Focus System for continuous maintenance of focus. 100mW
1166 mercury arc lamp illumination for viewing fluorescence by eye, and confocal
1167 scanning using Melles Griot solid state diode lasers: 405nm, 488nm (10mW),
1168 and 561nm (10mW). The image acquisition software used was Nikon EZ-C1.
1169 All samples were mounted and imaged in Vectashield mounting medium with
1170 DAPI (Vector Laboratories) at room temperature. Adobe Photoshop CS5 was
1171 used to pseudocolor images.

Analysis of Snn levels in S2R+ cells

1172 *Drosophila* S2R+ cells (40), a derivative of Schneider S2 cells, were
1173 cultured in Schneiders *Drosophila* medium (Gibco) with 10% fetal bovine
1174 serum, 100 U/ml of penicillin, and 100 µg/ml of streptomycin at 25°C. FLAG-
1175 HA constructs (Guruharsha et al., 2011) were transfected using TransIT-2020
1176 (Mirus). One day after transfection, plasmid expression was induced with
1177 0.35 mM CuSO₄ overnight. Harvested cells were plated on concanavalin A
1178 (0.5 mg/ml, Sigma) -coated plate and fixed at room temperature for 20
1179 min in 4% paraformaldehyde (Electron microscopy sciences). The cells were
1180 washed in PBS-DT (0.3% sodium deoxycholate, 0.3% Triton X-100 in PBS) and
1181 incubated overnight at 4°C with rabbit anti-Snn (1:2000) (Sen et al., 2011)
1182 and mouse anti-Flag (1:1000, Sigma). After washes in PBS-DT, the cells were
1183 incubated with mouse Alexa 488- and rabbit Alexa 568-conjugated secondary
1184 antibodies (1:500, Molecular Probes), followed by washing in PBS-T (0.1%
1185 Triton X-100 in PBS). The samples were mounted in VECTASHIELD mounting
1186 medium with DAPI (Vector Laboratories).

ACKNOWLEDGEMENTS

1185 This work was initially supported by a grant from the SMA Foundation
1186 to S. Artavanis-Tsakonas, A.C. Hart, and D. van Vactor. Additional sup-
1187 port from Spinal Muscular Atrophy Program Project (P01N5066888) to S.
1188 Artavanis-Tsakonas, L. Rubin, A.C. Hart, and D. van Vactor. Work on DPiM
1189 project was supported by a grant from the National Institutes of Health
1190 (NIH 5RO1HG003616) to S. Artavanis-Tsakonas. A. Sen was supported by a
1191 postdoctoral fellowship from the Families of Spinal Muscular Atrophy. The
1192 authors thank Nina Makhortova for help with cell imaging and the Nikon
1193 Imaging Center at Harvard Medical School for help with light microscopy.

- 1194 18. Huang da W, Sherman BT, & Lempicki RA (2009) Bioinformatics enrichment tools: paths
1195 toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 37(1):1-
1196 13.
- 1197 19. Sen A, et al. (2011) Modeling spinal muscular atrophy in *Drosophila* links Snn to FGF
1198 signaling. *J Cell Biol* 192(3):481-495.
- 1199 20. Enright AJ, Van Dongen S, & Ouzounis CA (2002) An efficient algorithm for large-scale
1200 detection of protein families. *Nucleic Acids Res* 30(7):1575-1584.
- 1201 21. Ashburner M, et al. (2000) Gene ontology: tool for the unification of biology. The Gene
1202 Ontology Consortium. *Nat Genet* 25(1):25-29.
- 1203 22. Sharma A, et al. (2005) A role for complexes of survival of motor neurons (SMN) protein
1204 with gemins and profilin in neurite-like cytoplasmic extensions of cultured nerve cells.
1205 *Experimental cell research* 309(1):185-197.
- 1206 23. Leung KM, et al. (2006) Asymmetrical beta-actin mRNA translation in growth cones medi-
1207 ates attractive turning to netrin-1. *Nat Neurosci* 9(10):1247-1256.
- 1208 24. Al-Ramahi I, et al. (2007) dAtaxin-2 mediates expanded Ataxin-1-induced neurodegenera-
1209 tion in a *Drosophila* model of SCA1. *PLoS Genet* 3(12):e234.
- 1210 25. Piazzon N, et al. (2008) In vitro and in cellulo evidences for association of the survival of
1211 motor neuron complex with the fragile X mental retardation protein. *The Journal of biological*
1212 *chemistry* 283(9):5598-5610.
- 1213 26. Oprea GE, et al. (2008) Plastin 3 is a protective modifier of autosomal recessive spinal
1214 muscular atrophy. *Science* 320(5875):524-527.
- 1215 27. Dimitriadis M, et al. (2010) Conserved genes act as modifiers of invertebrate SMN loss of
1216 function defects. *PLoS Genet* 6(10):e1001172.
- 1217 28. Rossoll W, et al. (2003) Snn, the spinal muscular atrophy-determining gene product, mod-
1218 ulates axon growth and localization of beta-actin mRNA in growth cones of motoneurons.
1219 *The Journal of cell biology* 163(4):801-812.
- 1220 29. Zhang HL, et al. (2003) Active transport of the survival motor neuron protein and the role
1221 of exon-7 in cytoplasmic localization. *J Neurosci* 23(16):6627-6637.
- 1222 30. Bowerman M, Beauvais A, Anderson CL, & Kothary R (2010) Rho-kinase inactivation pro-
1223 longes survival of an intermediate SMA mouse model. *Human molecular genetics* 19(8):1468-
1224 1478.
31. Osterloh JM, et al. (2012) dSarm/Sarm1 is required for activation of an injury-induced axon
death pathway. *Science* 337(6093):481-484.
32. Rossoll W, et al. (2002) Specific interaction of Snn, the spinal muscular atrophy determining
gene product, with hnRNP-R and gry-rbp/hnRNP-Q: a role for Snn in RNA processing in
motor axons? *Human molecular genetics* 11(1):93-105.
33. Hofmann Y, Lorson CL, Stamm S, Androphy EJ, & Wirth B (2000) Htra2-beta 1 stimulates
an exonic splicing enhancer and can restore full-length SMN expression to survival motor

1225	neuron 2 (SMN2). <i>Proc Natl Acad Sci U S A</i> 97(17):9618-9623.	
1226	34. Gangwani L, Mikrut M, Theroux S, Sharma M, & Davis RJ (2001) Spinal muscular atrophy disrupts the interaction of ZPR1 with the SMN protein. <i>Nature cell biology</i> 3(4):376-383.	1294
1227		1295
1228	35. Ahmad S, Wang Y, Shaik GM, Burghes AH, & Gangwani L (2012) The zinc finger protein ZPR1 is a potential modifier of spinal muscular atrophy. <i>Human molecular genetics</i> 21(12):2745-2758.	1296
1229		1297
1230	36. Claus P, Bruns AF, & Grothe C (2004) Fibroblast growth factor-2(23) binds directly to the survival of motoneuron protein and is associated with small nuclear RNAs. <i>Biochem J</i> 384(Pt 3):559-565.	1298
1231		1299
1232	37. Makhortova NR, et al. (2011) A screen for regulators of survival of motor neuron protein levels. <i>Nat Chem Biol</i> 7(8):544-552.	1300
1233		1301
1234	38. Hensel N, et al. (2012) Analysis of the fibroblast growth factor system reveals alterations in a mouse model of spinal muscular atrophy. <i>PLoS ONE</i> 7(2):e31202.	1302
1235		1303
1236	39. Kwon JE, Kim EK, & Choi EJ (2011) Stabilization of the survival motor neuron protein by ASK1. <i>FEBS Lett</i> 585(9):1287-1292.	1304
1237		1305
1238	40. Yanagawa S, Lee JS, & Ishimoto A (1998) Identification and characterization of a novel line of Drosophila Schneider S2 cells that respond to wingless signaling. <i>J Biol Chem</i> 273(48):32353-32359.	1306
1239		1307
1240	41. Yu C, et al. (2011) Development of expression-ready constructs for generation of proteomic libraries. <i>Methods Mol Biol</i> 723:257-272.	1308
1241		1309
1242	42. Kariya S, et al. (2012) Mutant superoxide dismutase 1 (SOD1), a cause of amyotrophic lateral sclerosis, disrupts the recruitment of SMN, the spinal muscular atrophy protein to nuclear Cajal bodies. <i>Human molecular genetics</i> 21(15):3421-3434.	1310
1243		1311
1244	43. Gu W, Pan F, Zhang H, Bassell GJ, & Singer RH (2002) A predominantly nuclear protein affecting cytoplasmic localization of beta-actin mRNA in fibroblasts and neurons. <i>The Journal of cell biology</i> 156(1):41-51.	1312
1245		1313
1246	44. Glinka M, et al. (2010) The heterogeneous nuclear ribonucleoprotein-R is necessary for axonal beta-actin mRNA translocation in spinal motor neurons. <i>Human molecular genetics</i> 19(10):1951-1966.	1314
1247		1315
1248	45. Fallini C, et al. (2011) The survival of motor neuron (SMN) protein interacts with the mRNA-binding protein HuD and regulates localization of poly(A) mRNA in primary motor neuron axons. <i>J Neurosci</i> 31(10):3914-3925.	1316
1249		1317
1250	46. Menon KP, et al. (2004) The translational repressor Pumilio regulates presynaptic morphology and controls postsynaptic accumulation of translation factor eIF-4E. <i>Neuron</i> 44(4):663-676.	1318
1251		1319
1252	47. Aberle H, et al. (2002) wishful thinking encodes a BMP type II receptor that regulates synaptic growth in Drosophila. <i>Neuron</i> 33(4):545-558.	1320
1253		1321
1254	48. McCabe BD, et al. (2003) The BMP homolog Gbb provides a retrograde signal that regulates synaptic growth at the Drosophila neuromuscular junction. <i>Neuron</i> 39(2):241-254.	1322
1255		1323
1256	49. Gibbins DJ, Ciaudo C, Erhardt M, & Voinnet O (2009) Multivesicular bodies associate with components of miRNA effector complexes and modulate miRNA activity. <i>Nature cell biology</i> 11(9):1143-1149.	1324
1257		1325
1258		1326
1259		1327
1260		1328
1261		1329
1262		1330
1263		1331
1264		1332
1265		1333
1266		1334
1267		1335
1268		1336
1269		1337
1270		1338
1271		1339
1272		1340
1273		1341
1274		1342
1275		1343
1276		1344
1277		1345
1278		1346
1279		1347
1280		1348
1281		1349
1282		1350
1283		1351
1284		1352
1285		1353
1286		1354
1287		1355
1288		1356
1289		1357
1290		1358
1291		1359
1292		1360