



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Global Burden of HIV among Men Who Engage in Transactional Sex: A Systematic Review and Meta-Analysis

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Oldenburg, Catherine E., Amaya G. Perez-Brumer, Sari L. Reisner, Jason Mattie, Till Bärnighausen, Kenneth H. Mayer, and Matthew J. Mimiaga. 2014. "Global Burden of HIV among Men Who Engage in Transactional Sex: A Systematic Review and Meta-Analysis." PLoS ONE 9 (7): e103549. doi:10.1371/journal.pone.0103549. http://dx.doi.org/10.1371/journal.pone.0103549 .
Published Version	doi:10.1371/journal.pone.0103549
Accessed	February 16, 2015 8:12:00 PM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:12785814
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)



Global Burden of HIV among Men Who Engage in Transactional Sex: A Systematic Review and Meta-Analysis

Catherine E. Oldenburg^{1,2*}, Amaya G. Perez-Brumer³, Sari L. Reisner^{1,2}, Jason Mattie², Till Bärnighausen^{4,5}, Kenneth H. Mayer^{2,6}, Matthew J. Mimiaga^{1,2,7}

1 Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts, United States of America, **2** The Fenway Institute, Fenway Community Health, Boston, Massachusetts, United States of America, **3** Department of Sociomedical Sciences, Columbia Mailman School of Public Health, New York, New York, United States of America, **4** Department of Global Health and Population, Harvard School of Public Health, Boston, Massachusetts, United States of America, **5** Africa Centre for Health and Population Science, Mtubatuba, South Africa, **6** Department of Medicine, Beth Israel Deaconess, Boston, Massachusetts, United States of America, **7** Department of Psychiatry, Massachusetts General Hospital, Boston, Massachusetts, United States of America

Abstract

Background: Men who engage in transactional sex, the exchange of sex for money, goods, or other items of value, are thought to be at increased risk of HIV, but there have been no systematic attempts to characterize HIV burden in this population. We undertook a systematic review and meta-analysis to quantify the burden in this population compared with that of men in the general population to better inform future HIV prevention efforts.

Methods: We searched seven electronic databases, national surveillance reports, and conference abstracts for studies of men who engage in transactional sex published between 2004–2013. Random effects meta-analysis was used to determine pooled HIV prevalence and prevalence ratios (PR) for the difference in HIV prevalence among men who engage in transactional sex as compared to general population men.

Findings: Of 66 studies included representing 31,924 men who had engaged in transactional sex in 28 countries, pooled biological assay-confirmed HIV prevalence was 10.5% (95% CI = 9.4 to 11.5%). The highest pooled HIV prevalence was in Sub-Saharan Africa (31.5%, 95% CI = 21.6 to 41.5%), followed by Latin America (19.3%, 95% CI = 15.5 to 23.1%), North America (16.6%, 95% CI = 3.7 to 29.5%), and Europe (12.2%, 95% CI = 6.0 to 17.2%). Men who engaged in transactional sex had an elevated burden of HIV compared to the general male population (PR = 20.7, 95% CI = 16.8 to 25.5).

Conclusions: The global burden of HIV is disproportionately high among men who engage in transactional sex compared with the general male population. There is an urgent need to include this population in systematic surveillance as well as to scale-up access to quality HIV prevention programs.

Citation: Oldenburg CE, Perez-Brumer AG, Reisner SL, Mattie J, Bärnighausen T, et al. (2014) Global Burden of HIV among Men Who Engage in Transactional Sex: A Systematic Review and Meta-Analysis. PLoS ONE 9(7): e103549. doi:10.1371/journal.pone.0103549

Editor: Garrett Prestage, The University of New South Wales, Australia

Received: April 9, 2014; **Accepted:** June 29, 2014; **Published:** July 28, 2014

Copyright: © 2014 Oldenburg et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. UNAIDS data: http://www.unaids.org/documents/20101123_GlobalReport_Annexes1_em.pdf US Census Bureau Data: <http://www.census.gov/population/international/data/idb/informationGateway.php> Data necessary for all calculations extracted from individual studies is presented in Table S1.

Funding: CEO is supported by a National Institute of Allergy and Infectious Disease T32 NRSA grant (T32AI007535; PI: Seage). APB is supported by a Eunice Kennedy Shriver National Institute of Child Health & Human Development T32 NRSA grant (T32 HD049339; PI: Nathanson). MJM, JM and KHM were supported in part by the National Institute On Drug Abuse of the National Institutes of Health under Award Numbers R21DA035113 and R21DA033720 (PI for both awards: Mimiaga). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* Email: ceo242@mail.harvard.edu

Introduction

Transactional sex, or the exchange of sex for money, goods, drugs, or other items of value (including protection, housing, or food), is thought to be associated with an increased vulnerability to HIV. Men who engage in transactional sex are considered a subpopulation of the men who have sex with men (MSM) community, but current research indicates that they may have differential HIV risk compared to MSM who do not engage in transactional sex, although there is conflicting evidence as to

whether engagement in transactional sex is associated with increased risk-taking behavior. Some studies have shown that MSM who engage in transactional sex may have a higher prevalence of HIV than those who do not report transactional sex. [1–3] Drivers of both engagement in transactional sex and HIV risk among this population are complex and vary by geographic location. For example, motivations for engaging in transactional sex may be different in regions where homosexuality and/or sex work is criminalized. Structural factors, such as low educational

attainment and few opportunities for gainful employment, may motivate transactional sex. [4] However, evidence from regions such as Australia, that have relatively less MSM stigma, suggests that transactional sex may also be a form of expression of sexuality. [4] There are also likely large regional differences in risk-taking behavior and transactional sex that are reflective of local culture and environments.

Psychosocial factors that may increase HIV risk in this population include an increased burden of depression and mental health distress [5,6], sexual violence [6], a history of childhood sexual abuse [6], and substance use disorders. [1,6] In particular, substance use may encourage men to engage in transactional sex and may be a motivator for continuing to rely on income from transactional sex. Men who engage in transactional sex may have differential condom use patterns with commercial and non-commercial partners, which can increase risk for HIV. [3,7–10] Finally, clients of men who engage in transactional sex may offer greater monetary incentive for unprotected sex, or may react violently if men insist on using a condom. [11].

Men who engage in transactional sex span the sexual orientation spectrum, and a large proportion may identify as bisexual or heterosexual, with wide-ranging estimates from different settings of 30 to 75%. [8,12] Some men who engage in transactional sex do so with women in addition to, or instead of, other men. Those who engage in transactional sex primarily with women may have different risk as compared to men who engage in transactional sex primarily with men. HIV prevention programming that aims to recruit homosexual/gay-identified men only may not reach non-gay identified or “closeted” men who sell sex. In addition, men who engage in transactional sex primarily with men may be married to women or have female primary and/or casual sexual partners [3,13–18], and may act as an HIV transmission “bridge” to the general population.

To characterize the worldwide burden of HIV among men who engage in transactional sex, we undertook a systematic review of the literature and meta-analysis to assess the distribution of HIV prevalence across the globe and to assess the prevalence of HIV as compared to the general male population. An increased understanding of the burden of HIV among men who engage in transactional sex, including quantifying the burden of HIV in this population globally compared with that of other men, is crucial to understanding the evolving nature of the HIV epidemic and informing corresponding policy and HIV prevention efforts.

Methods

Search Strategy

This meta-analysis was conducted and reported according to Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Guidelines (Checklist S1). We searched seven electronic databases for studies published between January 1, 2004 and July 31, 2013 (to identify articles published in the preceding 10-year period) including PubMed, EMBASE, PsycINFO, Sociological Abstracts, POPLINE, CINAHL, and Web of Science. Search terms related to transactional sex included “commercial sex”, “sex work*”, “male sex worker*”, “prostitution”, “exchange sex”, and “transactional sex”. In addition, we performed a search including the terms “HIV” and “men who have sex with men”. We also searched abstracts from the International AIDS Society (IAS), the American Public Health Association (APHA), the Conference on Retroviruses and Opportunistic Infections (CROI), and the International Society for Sexually Transmitted Disease Research (ISSTD). HIV surveillance reports including demographic and health surveys (DHS) and integrated biological and

behavioral surveillance (IBBS) reports were also searched. Reference lists of all included articles were reviewed for additional articles.

Inclusion and Exclusion Criteria

Studies were included if they contained primary, quantitative data on HIV prevalence among men (individuals assigned a male sex at birth and presently identified as a male/man) who reported exchanging any sex act for anything of value, including money, goods, or drugs. Studies were included regardless of whether HIV status was determined by laboratory methods or via self-report. Studies published in English, Spanish, French, or Portuguese, or if enough study information was published in an English-language abstract, were included. For studies that reported results from overlapping cohorts, the study with the most complete data was included in the review and meta-analysis. Studies reporting the larger sample size, or in a peer-reviewed journal versus an abstract or surveillance report were considered the “more complete” study. In cases in which no delineation was made between reporting HIV prevalence among male and transgender male-to-female sex workers, the study was included if the majority ($\geq 80\%$) of participants in the study were not transgender male-to-female. Three studies were excluded because they did not differentiate men from transgender women in HIV estimates [19–21].

Data Extraction

Data were extracted independently on a standardized data collection form by two separate reviewers (CEO, APB, and/or JM) with $>90\%$ agreement. Adjudication for inconsistencies was done through discussion and if necessary a third reviewer (SLR) served as the tiebreaker. Extracted data included total number of men who reported transactional sex in the study, total number of men who were tested for HIV or self-reported their HIV serostatus, and total number testing or reporting HIV infection. Additional data extracted included sampling methodology, study design, definition of transactional sex (categorized as male sex worker, exchanged sex for money at least once in the previous 12 months, and ever exchanged sex for money), country, and region, and if the study reported individual-level factors that increase HIV risk including drug use (injection or non-injection), engagement in unprotected anal sex (UAS) with commercial and/or noncommercial partners, clinical depression, experience of childhood sexual abuse (CSA), and sexually transmitted infection (STI) prevalence.

Data Analysis

Meta-analysis using a DerSimonian-Laird random effects model [22] was used to generate an overall pooled point estimate and 95% confidence interval for HIV prevalence separately by biological assay and by self-report for all eligible studies, and then restricted to studies which reported ≥ 50 participants. After the initial assessment of HIV prevalence by self-report and of any sample size, all analyses included only studies that assessed HIV prevalence by biological assay and reported ≥ 50 participants. Pooled point estimates of biological assay-confirmed HIV prevalence were calculated by country, region, and definition of transactional sex (“male sex workers”, “exchanged sex in the last 12 months”, and “ever exchanged sex”). A random effects model was used to allow for heterogeneity between studies. [22] Random effects meta-regression was used to assess differences in HIV prevalence by definition of transactional sex. Publication bias was assessed with Egger’s test [23] and Begg’s test. [24] Methodological quality of the studies was determined using a modified GRADE scoring assessing sampling methodology, study design,




Figure 1. Flow diagram of included studies.
doi:10.1371/journal.pone.0103549.g001

HIV measure, if the study population was well-defined, and generalizability of the study (Table S1). [25,26].

Prevalence ratios comparing the prevalence of biological assay-confirmed HIV among men reporting engagement in transactional sex to the prevalence of HIV among men aged 15 and over in the general population were calculated to assess the excess burden of HIV associated with transactional sex among men. HIV prevalence estimates for men of reproductive age were calculated using UNAIDS 2009 data for the total estimated number of HIV cases as the numerator and the United States Census Bureau International Division data for the total number of men aged 15 and over in the global population in 2009 as the denominator. [27] The midpoint of the uncertainty bounds for number of HIV-

infected males aged 15 and over was used as the estimate of absolute number of HIV-infected males in China because no estimates for the absolute number were available in the UNAIDS 2009 report. We chose the 2009 report since the studies represented in this analysis were published between 2004–2013, and thus this provided a midpoint. Prevalence ratios were calculated with a random-effects model. A standard correction of 0.5 was added to any zero cells. Heterogeneity was assessed with an I^2 and τ^2 statistics, by region of study origin and overall. All analyses were conducted in Stata 12.0 (StataCorp, College Station, TX).

Table 1. Pooled HIV prevalence by HIV measurement and sample size.

	N (k)	Pooled HIV Prevalence (95% CI)
Overall	34,531 (88)	11.9% (10.9 to 12.9%)
Biological assay	32,007 (71)	10.7% (9.7 to 11.8%)
Self-report, untested or unknown status excluded from calculation	2,524 (17)	20.6% (14.5 to 26.8%)
Self-report, untested or unknown status assumed to be negative	3,109 (17)	13.3% (9.6 to 17.0%)
Biological assay, ≥ 50 participants in study	31,924 (66)	10.5% (9.4 to 11.5%)

N = number of subjects; k = number of studies; 95% CI = 95% Confidence Interval.
doi:10.1371/journal.pone.0103549.t001




Figure 2. Pooled HIV prevalence by country among studies reporting biologically-confirmed HIV prevalence with a sample size of ≥ 50 (N = 66).

doi:10.1371/journal.pone.0103549.g002

Results

Of 20,193 titles and abstracts, 547 conference abstracts, and 165 surveillance reports, 446 titles and abstracts were selected for further review, and a total of 88 articles, abstracts, or surveillance reports representing 34,531 individuals in 30 countries were included in the review. Of the 88 studies, 14 were in East Asia (all of which were from China)[12,13,28–39], 16 in South Asia [3,16,17,40–52], 15 in Southeast Asia [7,9,15,53–64], 11 in Latin America [2,14,18,65–72], 9 in Sub-Saharan Africa [73–81], 9 in Europe [8,10,82–88], and 14 in North America [1,5,89–99]. Of these, 66 [1–3,7–9,12,14–17,28,29,31–39,41–51,53,55,57,59–61,63–67,70–72,74–77,79–88,93–95,100] included a biological HIV assay and had a sample size of ≥ 50 , and thus were included in the analyses, representing 31,924 men who engaged in transactional sex with other men in 28 countries (Figure 1).

Table 1 shows the pooled HIV prevalence among men who engaged in transactional sex by HIV measurement and sample size. The HIV prevalence among men engaging in transactional sex across all 88 studies was 11.9% (95% CI = 10.9 to 12.9%). Across 71 studies reporting biological assay-confirmed HIV serostatus, including small samples, HIV prevalence was 10.7% (95% CI 9.7 to 11.8%). Across 66 studies with biological assay and restricted to studies with ≥ 50 participants, the pooled HIV prevalence was 10.5% (95% CI 9.4% to 11.5%).

The pooled biological assay-confirmed HIV prevalence among men who engaged in transactional sex by region, country, and definition of transactional sex is presented in Table 2. By region, the highest pooled HIV prevalence was in Sub-Saharan Africa (31.5%, 95% CI 21.6 to 41.5%), followed by Latin America (19.3%, 95% CI 15.5 to 23.1%), North America (16.6%, 95% CI 3.7 to 29.5%), and Southeast Asia (12.9%, 95% CI 8.8 to 17.0%). The lowest prevalence region was South Asia (2.7%, 95% CI 1.7 to 3.6%). Figure 2 graphically depicts HIV prevalence by country. Overall heterogeneity of the pooled estimates was high ($\tau^2 = 2800.08$). Begg's test ($P = 0.002$) and Egger's test ($P < 0.001$) indicated the possibility of publication bias, which could be due to small studies or high heterogeneity. With the exception of South Asia, these tests suggested no publication bias in analyses stratified by geographic region.

Table 3 presents prevalence ratios for biological assay-confirmed HIV prevalence among men engaging in transactional sex compared to the general population of males age 15 or older. The pooled prevalence ratio (PR) for HIV infection for men who engage in transactional sex compared to the general population of men aged 15 and older was 20.7 (95% CI 16.8 to 25.5). The regions with the highest PRs included East Asia (PR 51.3, 95% CI 38.5 to 68.2), Latin America (PR 35.0, 95% CI 27.5 to 44.7), and Europe (PR 32.4, 95% CI 25.9 to 40.6). The overall heterogeneity of the prevalence ratio estimates was high ($I^2 = 96.9\%$, $\tau^2 = 2076.67$). Begg's test ($P = 0.18$) and Egger's test ($P = 0.89$) suggested no evidence of publication bias.

The distribution of studies reporting individual risk factors related to HIV vulnerability by region is shown in Table 4. Reporting of individual risk factors varied widely by type of risk factor. The majority of studies reported UAS (77.3% among men who engage in transactional sex); less than half of studies reported substance use, injection drug use, and STI history or current infection (43.2%, 43.2%, and 48.9% respectively). Few studies reported depression (8.0%) or childhood sexual abuse (12.5%). The only risk factor whose reporting differed significantly by region among men who engaged in transactional sex was injection drug use ($P < 0.001$), with most studies in North America including injection drug use history as a risk behavior (78.6%), and few in East Asia (7.1%) and Latin America (9.1%).

A single study reported HIV incidence, among men who engage in transactional sex in Thailand, and found an incidence of 7.4 per 100 person-years (95% CI 4.7–11.1). [64].

Discussion

The burden of HIV infection is disproportionately high among men who engage in transactional sex globally, with these men having more than 20 times the prevalence of HIV infection as compared to the general male population. These data underscore the urgent global need to scale up access to quality HIV prevention programs specific to this population. Previous systematic reviews have demonstrated a disproportionate burden of HIV among female sex workers (FSW), transgender women, and MSM as compared to the general population. [101–104] The findings of

Table 2. Pooled HIV prevalence among men who engage in transactional sex by region, country, and definition of transactional sex.

	Overall (95% CI) ¹		Male sex workers		Exchanged sex in last 12 months		Ever exchanged sex		p ²
	N (k)	HIV Prevalence (95%CI)	N (k)	HIV Prevalence (95%CI)	N (k)	HIV Prevalence (95%CI)	N (k)	HIV Prevalence (95%CI)	
East Asia (China)	7,221 (12)	4.1% (2.8 to 5.4%)	2,200 (7)	4.7% (2.5 to 6.8%)	5,001 (5)	3.7% (1.7 to 5.7%)	-	-	0.64
South Asia	14,453 (14)	2.7% (1.7 to 3.6%)	11,433 (11)	1.0% (0.5 to 1.5%)	809 (1)	2.0% (1.0 to 2.9%)	2,211 (2)	11.8% (6.0 to 17.6%)	0.0001
Bangladesh	985 (1)	0.0% (0.0 to 0.0%)	985 (1)	0.0% (0.0 to 0.0%)	-	-	-	-	-
India	2,211 (2)	11.8% (6.0 to 17.6%)	100 (2)	24.0% (15.6 to 32.4%)	-	-	2,211 (2)	11.8% (6.0 to 17.6%)	-
Nepal	218 (2)	3.5% (1.0 to 5.9%)	218 (2)	3.5% (1.0 to 5.9%)	-	-	-	-	-
Pakistan	11,039 (9)	1.2% (0.6 to 1.7%)	10,230 (8)	1.0% (0.5 to 1.6%)	809 (1)	2.0% (1.0 to 2.9%)	-	-	-
Southeast Asia	4,477 (12)	12.9% (8.8 to 17.0%)	3,292 (8)	12.6% (8.1 to 17.1%)	778 (2)	13.2% (4.3 to 22.0%)	407 (2)	13.9% (0.0 to 41.1%)	0.98
Indonesia	250 (1)	3.6% (1.3 to 5.9%)	250 (1)	3.6% (1.3 to 5.9%)	-	-	-	-	-
Laos	119 (1)	8.4% (3.4 to 13.4%)	-	-	119 (1)	8.4% (3.4 to 13.4%)	-	-	-
Thailand	3,201 (7)	17.5% (13.7 to 21.2%)	2,211 (5)	15.6% (11.6 to 19.6%)	659 (1)	17.5% (14.6 to 20.3%)	484 (2)	23.2% (13.9 to 32.5%)	-
Vietnam	907 (3)	6.8% (0.0 to 13.8%)	831 (2)	9.9% (2.8 to 17.0%)	-	-	76 (1)	0.0% (0.0 to 0.0%)	-
Latin America	1,704 (9)	19.3% (15.5 to 23.1%)	1,416 (6)	20.4% (15.5 to 25.2%)	-	-	288 (3)	16.7% (18.7 to 15.1%)	0.46
Argentina	170 (2)	11.2% (6.4 to 15.9%)	114 (1)	11.4% (5.6 to 17.2%)	-	-	56 (1)	10.7% (2.6 to 18.8%)	-
Ecuador	76 (1)	19.7% (10.8 to 28.7%)	-	-	-	-	76 (1)	19.7% (10.8 to 28.7%)	-
El Salvador	156 (1)	19.2% (13.0 to 25.4%)	-	-	-	-	156 (1)	19.2% (13.0 to 25.4%)	-
Mexico	507 (2)	17.2% (12.0 to 22.5%)	507 (2)	17.2% (12.0 to 22.5%)	-	-	-	-	-
Peru	478 (2)	27.3% (20.2 to 34.4%)	478 (2)	27.3% (20.2 to 34.4%)	-	-	-	-	-
Uruguay	317 (1)	21.8% (17.2 to 26.3%)	317 (1)	21.8% (17.2 to 26.3%)	-	-	-	-	-
Sub-Saharan Africa	1,608 (7)	31.5% (21.6 to 41.5%)	1,386 (5)	36.3% (24.9 to 47.6%)	93 (1)	26.9% (17.9 to 35.9%)	129 (1)	12.4% (6.7 to 18.1%)	0.34
Cote d'Ivoire	96 (1)	50.0% (40.0 to 60.0%)	96 (1)	50.0% (40.0 to 60.0%)	-	-	-	-	-
Kenya	1,290 (4)	33.2% (21.0 to 45.3%)	1,290 (4)	33.2% (21.0 to 45.3%)	-	-	-	-	-
Senegal	93 (1)	26.9 (17.9 to 35.9%)	-	-	93 (1)	26.9 (17.9 to 35.9%)	-	-	-
Uganda	129 (1)	12.4% (6.7 to 18.1%)	-	-	-	-	129 (1)	12.4% (6.7 to 18.1%)	-
Europe	1,854 (8)	10.2% (5.2 to 15.2%)	988 (6)	12.2% (9.1 to 15.3%)	636 (1)	9.3% (7.0 to 11.5%)	230 (1)	0.9% (0.0 to 2.1%)	0.05
Belgium	120 (1)	10.8% (5.3 to 16.4%)	120 (1)	10.8% (5.3 to 16.4%)	-	-	-	-	-
Czech Republic	230 (1)	0.9% (0.0 to 2.1%)	-	-	-	-	230 (1)	0.9% (0.0 to 2.1%)	-
Israel	53 (1)	5.7% (0.0 to 11.9%)	53 (1)	5.7% (0.0 to 11.9%)	-	-	-	-	-
Netherlands	99 (1)	11.1% (4.9 to 17.3%)	99 (1)	11.1% (4.9 to 17.3%)	-	-	-	-	-
Russia	50 (1)	18.0% (7.4 to 28.6%)	50 (1)	18.0% (7.4 to 28.6%)	-	-	-	-	-
Spain	666 (2)	14.2% (9.6 to 18.8%)	666	14.2% (9.6 to 18.8%)	-	-	-	-	-
United Kingdom	636 (1)	9.3% (7.0 to 11.5%)	-	-	636 (1)	9.3% (7.0 to 11.5%)	-	-	-
North America	607 (4)	16.6% (3.7 to 29.5%)	-	-	108 (1)	26.9% (18.5 to 35.2%)	499 (3)	13.4% (0.0 to 27.3%)	0.43
United States	349 (2)	19.3% (8.0 to 30.6%)	-	-	-	-	349 (2)	19.5% (13.7 to 25.3%)	-
Canada	258 (2)	14.1% (0.0 to 38.4%)	-	-	108 (1)	26.9% (18.5 to 35.2%)	150 (1)	2.0% (0 to 4.2%)	-

Table 2. Cont.

	Overall (95% CI) ¹		Male sex workers		Exchanged sex in last 12 months		Ever exchanged sex		P ²
	N (k)	HIV Prevalence (95%CI)	N (k)	HIV Prevalence (95%CI)	N (k)	HIV Prevalence (95%CI)	N (k)	HIV Prevalence (95%CI)	
OVERALL	31,924 (66)	10.5% (9.4 to 11.5%)	20,715 (43)	10.4% (9.2 to 11.5%)	7,445 (11)	8.4% (5.7 to 11.1%)	3,917 (12)	12.6% (7.4 to 17.8%)	0.78

¹Analyses restricted to individual studies reporting ≥ 50 participants and biological assay-confirmed HIV prevalence;

²Type-3 P-value from random effects meta-regression adjusted for country-level HIV prevalence among males aged 15+ (presented for regions).

N = number of subjects; k = number of studies; 95% CI = 95% Confidence Interval.
doi:10.1371/journal.pone.0103549.t002

the current study indicate that men who engage in transactional sex bear a similarly higher burden of HIV. Drivers of engagement in transactional sex are multifaceted, and include drug use, economic motivation, stigma, harassment and discrimination from multiple sources and levels of influence, likely impact the health and well-being (e.g., can lead to increased depression and lower self-acceptance) of men who engage in transactional sex and may limit access to and uptake of HIV prevention and treatment services. [105] Moreover, resulting distress may reduce their ability to uptake and incorporate HIV prevention counseling information and behavioral skills, leading to continued sexual risk. Ensuring that this vulnerable population is included in strategic plans and resource allocation to address both structural and individual levels of influence is essential to adequately curb the spread of HIV globally.

These results demonstrate a consistently disproportionate burden of HIV globally among men who engage in transactional sex as compared to men in the general population. Regions in which the HIV epidemic is largely driven by sexual transmission and concentrated among MSM had the largest disparities in the burden of HIV. In Latin America, North America, and Europe, regions in which the HIV epidemic is concentrated among MSM [106], both absolute and relative measures of HIV burden among men engaging in transactional sex were high. The absolute HIV prevalence in Latin America was nearly 20%, and was second only to Sub-Saharan Africa. The regions with the third and fourth-highest HIV prevalence were North America and Europe, respectively. Despite these regions being culturally, socially, economically, and politically diverse, these data show a consistently higher burden of HIV among men who engage in transactional sex. While North America and Europe were the only regions that represented higher-income countries in this study, their inequitable burden was second only to that of China and Latin America, underscoring the association of the social marginalization of sex work with conditions that potentiated HIV spread. Importantly, men who engage in transactional sex should not be overlooked in national HIV prevention strategies in countries of a wide range of economic development.

Sub-Saharan Africa had the highest absolute HIV prevalence among men who engage in transactional sex, with an estimated pooled HIV prevalence of nearly one-third, but this region also had the lowest pooled prevalence ratio of all regions included in this meta-analysis. This is reflective of the generalized background HIV epidemic of many countries in that region. Male-male sexual behavior is highly stigmatized and often criminalized in many Sub-Saharan African countries, and in countries such as Uganda there are extreme penalties for engaging in male-male sex. [107,108] These penalties limit access to healthcare and HIV prevention services among MSM, including men who engage in transactional sex with other men, thereby increasing vulnerability to HIV, and late entry into care leading to increased spread because of uncontrolled viremia (i.e. higher community viral load). [107,108] These men may also engage in transactional sex with women, and may represent a substantially different population than those who engage in transactional sex with men.

China was the only country in East Asia represented in this study, and had the largest disparity in HIV prevalence among men who engage in transactional sex as compared to the general population, with more than 50 times the risk of HIV infection. However, China had a lower pooled HIV prevalence than any region except South Asia. China has recently experienced evidence of a growing HIV epidemic among MSM, and previous studies have demonstrated elevated HIV prevalence among men who sell sex to men as compared to MSM who do not sell sex.

Table 3. Prevalence ratios comparing pooled HIV prevalence among men who engage in transactional sex to general population men aged 15 and older.

	N (k)	HIV Prevalence ¹ (95%CI)	Country-Wide HIV Prevalence, Males 15+	I ²	Prevalence Ratio ² (95% CI)
East Asia (China)	7,221 (12)	4.1% (2.8 to 5.4%)	0.10%	73.4%	51.3 (38.5 to 68.2)
South Asia	14,453 (14)	2.7% (1.7 to 3.6%)		95.0%	12.3 (7.1 to 21.2)
Bangladesh	985 (1)	0.0% (0.0 to 0.0%)	0.0093%		5.5 (0.3 to 87.3)
India	2,211 (3)	11.8% (6.0 to 17.6%)	0.34%		34.4 (20.6 to 57.6)
Nepal	218 (2)	3.5% (1.0 to 5.9%)	0.46%		8.2 (4.2 to 16.2)
Pakistan	11,039 (9)	1.2% (0.6 to 1.7%)	0.11%		11.0 (6.1 to 19.8)
Southeast Asia	4,477 (12)	12.9% (8.8 to 17.0%)		83.9%	15.3 (12.6 to 18.5)
Indonesia	250 (1)	3.6% (1.3 to 5.9%)	0.24%		14.7 (7.7 to 27.9)
Laos	119 (1)	8.4% (3.4 to 13.4%)	0.25%		33.5 (18.5 to 60.7)
Thailand	3,201 (7)	17.5% (13.7 to 21.2%)	1.21%		14.4 (11.8 to 17.7)
Vietnam	907 (3)	6.8% (0.0 to 13.8%)	0.59%		12.7 (5.3 to 30.5)
Latin America	1,704 (9)	19.3% (15.5 to 23.1%)		83.1%	35.0 (27.5 to 44.7)
Argentina	170 (2)	11.2% (6.4 to 15.9%)	0.53%		21.2 (13.9 to 32.4)
Ecuador	76 (1)	19.7% (10.8 to 28.7%)	0.51%		39.0 (24.8 to 61.4)
El Salvador	156 (1)	19.2% (13.0 to 25.4%)	1.10%		17.5 (12.7 to 24.1)
Mexico	507 (2)	17.2% (12.0 to 22.5%)	0.42%		41.0 (30.2 to 55.8)
Peru	478 (2)	27.3% (20.2 to 34.4%)	0.56%		49.2 (37.9 to 63.9)
Uruguay	317 (1)	21.8% (17.2 to 26.3%)	0.55%		39.7 (32.2 to 49.0)
Sub-Saharan Africa	1,608 (7)	31.5% (21.6 to 41.5%)		97.7%	8.9 (5.4 to 14.5)
Cote d'Ivoire	96 (1)	50.0% (40.0 to 60.0%)	2.56%		19.5 (16.0 to 23.8)
Kenya	1,290 (4)	33.2% (21.0 to 45.3%)	4.60%		6.9 (4.7 to 10.0)
Senegal	93 (1)	26.9 (17.9 to 35.9%)	0.69%		38.7 (27.7 to 54.1)
Uganda	129 (1)	14.2% (7.7 to 20.6%)	5.81%		2.4 (1.5 to 3.8)
Europe	1,854 (8)	12.2% (6.0 to 17.2%)		52.5%	32.4 (25.9 to 40.6)
Belgium	120 (1)	10.8% (5.3 to 16.4%)	0.23%		46.4 (27.8 to 77.6)
Czech Republic	230 (1)	0.9% (0.0 to 2.1%)	0.033%		26.5 (6.7 to 105.4)
Israel	53 (1)	5.7% (0.0 to 11.9%)	0.20%		27.7 (9.2 to 83.2)
Netherlands	99 (1)	11.1% (4.9 to 17.3%)	0.24%		46.3 (26.5 to 80.8)
Russia	50 (1)	18.0% (7.4 to 28.6%)	0.89%		20.3 (11.2 to 36.6)
Spain	666 (2)	14.2% (9.6 to 18.8%)	0.52%		27.6 (20.0 to 38.2)
United Kingdom	636 (1)	9.3% (7.0 to 11.5%)	0.23%		39.6 (31.1 to 50.5)
North America	607 (4)	16.6% (3.7 to 29.5%)		93.9%	24.6 (11.3 to 53.7)
United States	349 (2)	19.3% (8.0 to 30.6%)	0.77%		24.6 (13.3 to 45.3)
Canada	258 (2)	14.1% (0.0 to 38.4%)	0.38%		19.8 (0.67 to 581.5)
OVERALL	31,924 (66)	10.5% (9.4 to 11.5%)		96.9%	20.7 (16.8 to 25.5)

¹Among studies confirming HIV infection with a biological assay and among studies reporting ≥ 50 participants;

²All prevalence ratios significant ($P < 0.05$) except for Bangladesh and Canada.

N = number of subjects; k = number of studies; I² = variation in pooled prevalence ratio due to heterogeneity; 95% CI = 95% Confidence Interval.

doi:10.1371/journal.pone.0103549.t003

[109] This result demonstrates a need to enhance the HIV prevention response among this population in China.

In an effort to report relevant data on those who do and do not identify as a sex worker, we used a broad definition of transactional sex in this meta-analysis, and included both male sex workers (MSWs) who self-identify as sex workers and engage in transactional sex regularly, as well as men who engage in transactional sex occasionally and may not identify as such. Men who engage in transactional sex more often or with more partners may have a greater HIV risk as compared to those who exchange sex less frequently. [2] Studies reported HIV prevalence by a

variety of definitions and temporal periods in which participants had engaged in transactional sex. An inherent difficulty in assessing differences using these definitions is the degree to which there is overlap among them. Individual studies may have lumped "male sex workers" in the "ever exchange sex" category, which could not be distinguished in this analysis and may have made the results appear not to differ by strata. Reliance on self-report may affect estimates, since sex work is highly stigmatized, so some individuals in these studies may have engaged in it, but were unwilling to disclose, or others might only have admitted a lesser level of activity. Finally, the source of HIV infection in this

Table 4. Proportion of studies reporting individual factors that affect vulnerability to HIV by region, overall and specifically among men who engage in transactional sex.

	Total Number of Studies	Any Drug Use	Injection Drug Use	Depression	Childhood Sexual Abuse	Unprotected Anal Sex	STI ¹ History/Prevalence
East Asia (China)	14						
Overall ²	8 (57.1%)	4 (28.6%)	2 (14.3%)	2 (14.3%)	14 (100%)	12 (85.7%)	
Transactional Sex ³	4 (28.6%)	1 (7.1%)	2 (14.3%)	2 (14.3%)	12 (85.7%)	7 (50.0%)	
South Asia	16						
Overall ²	6 (37.5%)	11 (68.8%)	0 (0%)	1 (6.3%)	16 (100%)	12 (75.0%)	
Transactional Sex ³	6 (37.5%)	10 (62.5%)	0 (0%)	1 (6.3%)	15 (93.8%)	10 (62.5%)	
Southeast Asia	15						
Overall ²	10 (66.7%)	6 (40.0%)	1 (6.7%)	0 (0%)	14 (93.3%)	10 (66.7%)	
Transactional Sex ³	8 (53.3%)	6 (40.0%)	1 (6.7%)	0 (0%)	13 (86.7%)	7 (46.7%)	
Latin America	11						
Overall ²	7 (63.6%)	2 (18.2%)	1 (9.1%)	1 (9.1%)	9 (81.8%)	8 (72.7%)	
Transactional Sex ³	2 (18.2%)	1 (9.1%)	1 (9.1%)	1 (9.1%)	7 (63.6%)	4 (36.4%)	
Sub-Saharan Africa	9						
Overall ²	6 (66.7%)	5 (55.6%)	1 (11.1%)	1 (11.1%)	9 (100%)	6 (66.7%)	
Transactional Sex ³	4 (44.4%)	3 (33.3%)	0 (0%)	1 (11.1%)	7 (77.8%)	4 (44.4%)	
Europe	9						
Overall ²	5 (55.6%)	6 (66.7%)	0 (0%)	2 (22.2%)	5 (55.6%)	6 (66.7%)	
Transactional Sex ³	5 (55.6%)	6 (66.7%)	0 (0%)	2 (22.2%)	5 (55.6%)	6 (66.7%)	
North America	14						
Overall ²	10 (71.4%)	12 (85.7%)	3 (21.4%)	4 (28.6%)	10 (71.4%)	6 (42.9%)	
Transactional Sex ³	9 (64.3%)	11 (78.6%)	3 (21.4%)	4 (28.6%)	9 (64.3%)	5 (35.7%)	
Total	88						
Overall ²	52 (59.1%)	46 (52.3%)	8 (9.1%)	11 (12.5%)	77 (87.5%)	60 (68.2%)	
Transactional Sex ³	38 (43.2%)	38 (43.2%)	7 (8.0%)	11 (12.5%)	68 (77.3%)	43 (48.9%)	
P⁴-value	-						
Overall ²	0.61	0.005	0.45	0.29	0.003	0.39	
Transactional Sex ³	0.25	<0.001	0.33	0.29	0.18	0.69	

¹Including syphilis, gonorrhoea, chlamydia, human papilloma virus (HPV), and/or herpes simplex virus (HSV);

²Risk factor reported among any population in the study (i.e., men who have sex with men);

³Risk factor reported specifically among men who engage in transactional sex;

⁴Fisher's exact test comparing frequency of reporting factors by geographic region of study, overall and among men who report transactional sex.
doi:10.1371/journal.pone.0103549.t004

population is not necessarily due to commercial partnerships, which could mitigate the importance of definition of transactional sex. This review was unable to analyze source of HIV infection. However, in total, the results of this review point to the need for standardized definitions of sex work and transactional sex, including identification of reliable measures of engaging in transactional sex and/or buying sex.

Consideration of individual-level risk factors that affect biological and behavioral vulnerability to HIV is critically needed to understand how HIV prevention interventions may best serve the needs of this population. Studies reporting individual risk factors such as unprotected anal sex (UAS) reported very different definitions of UAS (i.e., with the last partner, any in the last 3 months, with commercial versus non-commercial partners), and as such could not be meta-analyzed. While most studies reported UAS among men who engaged in transactional sex, fewer than half of studies reported STIs or substance use; very few studies reported psychosocial factors such as depression and childhood sexual abuse, which can have sequelae that lead to increased risk for HIV. [6,110] Future studies should consider these syndemic (i.e., co-occurring and potentially interacting) factors within cultural contexts, and future prevention interventions should consider specifically how these syndemic conditions increase vulnerability to HIV and how prevention interventions can best mitigate risk. Interestingly, there was a statistically significant geographic difference in reporting injection drug use. Studies in North America, Europe, and South Asia in particular were significantly more likely to report injection drug use, whereas studies in Latin America much less frequently reported injection drug use. Given that engagement in transactional sex may be driven by drug use, and injection drug use is a strong risk factor for HIV, characterizing the degree of this behavior will be important to better understand the epidemic. In addition, in general studies less frequently reported risk factors for HIV infection among men who engaged in transactional sex specifically as compared to reporting among MSM overall. Some of this was likely driven by the fact that many of these studies were primarily focused on MSM, and reported transactional sex secondarily. Despite this, to fully understand the dynamics of the HIV epidemic in this population, a better characterization of factors that influence HIV transmission among men who engage in transactional sex will be necessary.

There are some limitations to consider in this analysis. The pooled HIV prevalence estimates likely are biased by inherent difficulties in accessing and sampling men who engage in transactional sex. The degree of stigma, harassment and discrimination that this population faces in many contexts, which includes physical threats and extreme punishments in some cases, may limit research and surveillance of this key risk group in many global settings. There were few studies from regions in which male-male sexual behavior is illegal with very strong punishments. Sampling strategies varied widely, and could have influenced the results. In addition, it is likely that modality of meeting commercial partners has changed over time, with an increasing reliance on the Internet and mobile phones. Some evidence suggests that men who meet commercial partners violence in the Internet have differential HIV risk as compared to those who meet partners in physical venues (i.e., street-based or other physical venues). [5] We were unable to analyze HIV prevalence trends by partner-meeting modality, since typically HIV estimates were not reported by venue. However, attention should be given HIV prevention needs for men who meet commercial partners via the Internet, as they may differ substantially than those who meet partners in physical venues.

Pooled HIV prevalence could be underestimated if lower-risk individuals are more accessible and therefore more likely to be included in research and surveillance. These results may also have been affected by publication bias, which could result in over- or underestimates of pooled HIV prevalence as well as biased prevalence ratios. This study included only estimates that have been published in the peer-reviewed scientific literature or country-level surveillance data. Ultimately, the quality and accuracy of the pooled estimates is limited by the quality and accuracy of the data from which they arose. Restricting analyses only to studies reporting biologically-confirmed HIV prevalence with sample sizes of greater than 50 likely improves the quality and generalizability of these data. We used general population HIV prevalence among men as the comparison group as opposed to HIV prevalence among MSM, due to the lack of systematic surveillance of HIV among MSM which precluded comparable comparison. Furthermore, men who engage in transactional sex may be included in general population estimates, which could bias the results of corresponding prevalence ratios. Despite these limitations, this study presents the first comprehensive global overview of the current status of literature relating to the HIV epidemic among men who engage in transactional sex with other men, providing important evidence for the urgent need to scale up HIV prevention interventions among this group.

This study demonstrates a consistent elevation in HIV burden among men who engage in transactional sex globally in comparison with men in the general population. An understanding of populations of individuals who engage in transactional sex would be improved by routinely asking about transactional sex history in surveillance studies in diverse settings, which would allow for better monitoring of and response to the epidemic. The results of this study provide a macroscopic view of the state of the literature on this population, and demonstrate a need for additional studies worldwide, and in particular demonstrates gaps in knowledge in regions where MSM behavior is more highly stigmatized. This study also underscores the need for development of structural interventions, including economic interventions that may reduce reliance on income from transactional sex, as well as individual-level interventions such as increased access to substance use treatment. While drivers of the HIV epidemic differ geographically within countries and regions, the disproportionate burden of HIV shouldered by this population clearly demonstrates a need for a multisectoral response to increase efforts in HIV surveillance, research, and prevention, as well as increased allocation of resources for such a response, in this highly vulnerable yet understudied group.

Ethics Statement

An ethics statement was not required for this work.

Supporting Information

Table S1 Modified GRADE table describing quality of included studies.

(DOCX)

Checklist S1 PRISMA Checklist.

(DOCX)

Protocol S1 Study protocol.

(PDF)

Author Contributions

Conceived and designed the experiments: CEO AGPB SLR TB KHM MJM. Performed the experiments: CEO AGPB SLR JM. Analyzed the

data: CEO AGPB SLR JM TB KHM MJM. Contributed reagents/materials/analysis tools: TB KHM MJM. Wrote the paper: CEO AGPB SLR JM TB KHM MJM.

References

- Bacon O, Lum P, Hahn J, Evans J, Davidson P, et al. (2006) Commercial Sex Work and Risk of HIV Infection Among Young Drug-Injecting Men Who Have Sex With Men in San Francisco. *Sexually Transmitted Diseases* 33: 228–234. doi:10.1097/01.olq.00000204914.91923.ad.
- Lama JR, Lucchetti A, Suarez L, Laguna-Torres A, Guanira JV, et al. (2006) Association of Herpes Simplex Virus Type 2 Infection and Syphilis with Human Immunodeficiency Virus Infection among Men Who Have Sex with Men in Peru. *JID* 194: 1459–1466.
- Brahmam GN, Kodavalla V, Rajkumar H, Rachakulla HK, Kallam S, et al. (2008) Sexual practices, HIV and sexually transmitted infections among self-identified men who have sex with men in four high HIV prevalence states of India. *AIDS* 22: S45–S57.
- Prestage G, Jin F, Bavinton B, Hurley M (2014) Sex Workers and Their Clients Among Australian Gay and Bisexual Men. *AIDS Behav*. doi:10.1007/s10461-014-0755-6.
- Mimiaga MJ, Reisner SL, Tinsley JP, Mayer KH, Safren SA (2008) Street Workers and Internet Escorts: Contextual and Psychosocial Factors Surrounding HIV Risk Behavior among Men Who Engage in Sex Work with Other Men. *Journal of Urban Health: Bulletin of the New York Academy of Medicine* 86: 54–66. doi:10.1007/s11524-008-9316-5.
- Biello KB, Colby D, Closson E, Mimiaga MJ (2013) The Syndemic Condition of Psychosocial Problems and HIV Risk Among Male Sex Workers in Ho Chi Minh City, Vietnam. *AIDS Behav*. doi:10.1007/s10461-013-0632-8.
- Guadamuz T, Kunawararak P, Beyrer C, Pumpaisanchai J, Wei C, et al. (2010) HIV prevalence, sexual and behavioral correlates among Shan, Hill tribe, and Thai male sex workers in Northern Thailand. *AIDS Care* 22: 597–605. doi:10.1080/09540120903280935.
- Sethi G (2006) HIV, sexually transmitted infections, and risk behaviours in male sex workers in London over a 10 year period. *Sexually Transmitted Infections* 82: 359–363. doi:10.1136/sti.2005.019257.
- Pisani E (2004) HIV, syphilis infection, and sexual practices among transgenders, male sex workers, and other men who have sex with men in Jakarta, Indonesia. *Sexually Transmitted Infections* 80: 536–540. doi:10.1136/sti.2003.007500.
- Ballester R, Salmerón P, Gil MD, Gómez S (2011) Sexual Risk Behaviors for HIV Infection in Spanish Male Sex Workers: Differences According to Educational Level, Country of Origin and Sexual Orientation. *AIDS Behav* 16: 960–968. doi:10.1007/s10461-011-9964-4.
- Harcourt C (2005) The many faces of sex work. *Sexually Transmitted Infections* 81: 201–206. doi:10.1136/sti.2004.012468.
- Cai WD, Zhao J, Zhao JK, Raymond HF, Feng YJ, et al. (2010) HIV prevalence and related risk factors among male sex workers in Shenzhen, China: results from a time-location sampling survey. *Sexually Transmitted Infections* 86: 15–20. doi:10.1136/sti.2009.037440.
- Wong FY, Huang ZJ, He N, Smith BD, Ding Y, et al. (2008) HIV risks among gay- and non-gay-identified migrant money boys in Shanghai, China. *AIDS Care* 20: 170–180. doi:10.1080/09540120701534707.
- Gayet C, Magis-Rodríguez C, Fernandez AC, Donna S, Ramirez-Aranda J, et al. (n.d.) Men who sell sex in urban Mexico: HIV prevalence, sexual practices, and condom use - results from a biological and behavioral surveillance.
- Toledo CA, Varangrat A, Wimolsate W, Chemnasiri T, Phanuphak P, et al. (2010) Examining HIV infection among male sex workers in Bangkok, Thailand: A comparison of participants recruited at entertainment and street venues. *AIDS Education and Prevention* 22: 299–311.
- Bokhari A, Nizamani NM, Jackson DJ, Rehan NE, Rahman M, et al. (2007) HIV risk in Karachi and Lahore, Pakistan: an emerging epidemic in injecting and commercial sex networks. *International Journal of STD & AIDS* 18: 486–492. doi:10.1258/095646207781147201.
- Shaw SY, Emmanuel F, Adrien A, Holte-Mckenzie M, Archibald CP, et al. (2011) The descriptive epidemiology of male sex workers in Pakistan: a biological and behavioural examination. *Sexually Transmitted Infections* 87: 73–80. doi:10.1136/sti.2009.041335.
- Prado Cortez FC, Boer DP, Baltieri DA (2011) A Psychosocial Study of Male-to-Female Transgendered and Male Hustler Sex Workers in São Paulo, Brazil. *Arch Sex Behav* 40: 1223–1231. doi:10.1007/s10508-011-9776-7.
- Tun W, de Mello M, Pinho A, Chinaglia M, Diaz J (2008) Sexual risk behaviours and HIV seroprevalence among male sex workers who have sex with men and non-sex workers in Campinas, Brazil. *Sexually Transmitted Infections* 84: 455–457. doi:10.1136/sti.2008.031336.
- Chariyalertsak S, Kosachunhanan N, Saokhieo P, Songsupa R, Wongthance A, et al. (2011) HIV Incidence, Risk Factors, and Motivation for Biomedical Intervention among Gay, Bisexual Men, and Transgender Persons in Northern Thailand. *PLoS ONE* 6: e24295. doi:10.1371/journal.pone.0024295.t004.
- Reza-Paul S, Beattie T, Pasha A, Venugopal MS, Ramesh BM, et al. (2008) High HIV prevalence among male sex workers in Mysore, India – need for integrating care and support with prevention.
- DerSimonian R, Laird N (1986) Meta-Analysis in Clinical Trials*. *Controlled Clinical Trials* 7: 177–188.
- Egger M, Smith GD, Schneider M, Minder C (1997) Bias in meta-analysis detected by a simple, graphical test. *BMJ* 315: 629.
- Begg CB (1994) Operating Characteristics of a Rank Correlation Test for Publication Bias. *Biometrics* 50: 1088–1101.
- Oldenburg CE, Bärnighausen T, Harling G, Mimiaga MJ, Mayer KH (2013) Adherence to Post-Exposure Prophylaxis for Non-forcible Sexual Exposure to HIV: A Systematic Review and Meta-Analysis. *AIDS Behav*. doi:10.1007/s10461-013-0567-0.
- Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, et al. (2011) GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *Journal of Clinical Epidemiology* 64: 383–394. doi:10.1016/j.jclinepi.2010.04.026.
- Global Report: UNAIDS Report on the Global AIDS Epidemic: 2010 (2010) Global Report: UNAIDS Report on the Global AIDS Epidemic: 2010. UN Joint Programme on HIV/AIDS.
- Cheng WB, Zhong F, Wen F, Gao K, Liu JW, et al. (2010) [Investigation of HIV and syphilis infection and AIDS-related behaviors among money boys, in Guangzhou, China]. *Zhonghua Yu Fang Yi Xue Za Zhi*. 5 pp.
- Ngo TD, Laeyendecker O, Li C, Tai H, Cui M, et al. (2008) Herpes simplex virus type 2 infection among commercial sex workers in Kunming, Yunnan Province, China. *International Journal of STD & AIDS* 19: 694–697. doi:10.1258/ijsa.2008.008072.
- He N, Wong FY, Huang ZJ, Ding Y, Fu C, et al. (2007) HIV risks among two types of male migrants in Shanghai, China: money boys vs. general male migrants. *AIDS* 21: S73–S79.
- Feng L, Ding X, Lu R, Liu J, Sy A, et al. (2009) High HIV Prevalence Detected in 2006 and 2007 Among Men Who Have Sex With Men in China's Largest Municipality: An Alarming Epidemic in Chongqing, China. *J Acquir Immune Defic Syndr* 52: 79–85.
- Wu Z, Xu J, Liu E, Mao Y, Xiao Y, et al. (2013) HIV and Syphilis Prevalence Among Men Who Have Sex With Men: A Cross-Sectional Survey of 61 Cities in China. *Clinical Infectious Diseases* 57: 298–309. doi:10.1093/cid/cit210.
- Tao X, Gai R, Zhang N, Zheng W, Zhang X, et al. (2010) HIV infection and mental health of “money boys”: A pilot study in Shandong Province, China. *Southeast Asian J Trop Med Public Health* 41: 358–368.
- Ruan S, Yang H, Zhu Y, Ma Y, Li J, et al. (2008) HIV Prevalence and Correlates of Unprotected Anal Intercourse Among Men Who Have Sex with Men, Jinan, China. *AIDS Behav* 12: 469–475. doi:10.1007/s10461-008-9361-9.
- Liu S, Zhao J, Rou K, Chen L, Cai W, et al. (2011) A Survey of Condom Use Behaviors and HIV/STI Prevalence Among Venue-Based Money Boys in Shenzhen, China. *AIDS Behav* 16: 835–846. doi:10.1007/s10461-011-9978-y.
- Zhang L, Ding X, Lu R, Feng L, Li X, et al. (2012) Predictors of HIV and Syphilis among Men Who Have Sex with Men in a Chinese Metropolitan City: Comparison of Risks among Students and Non-Students. *PLoS ONE* 7: e37211. doi:10.1371/journal.pone.0037211.t002.
- Zhao J, Cai W-D, Gan Y-X, Zhang Y, Yang Z-R, et al. (2012) A Comparison of HIV Infection and Related Risk Factors Between Money Boys and Noncommercial Men Who Have Sex With Men in Shenzhen, China. *Sexually Transmitted Diseases* 39: 942–948. doi:10.1097/OLQ.0b013e31826f356f.
- Yu BN, Wang XA, Yu F, Wang J, Blanchard JF (2011) An exploratory survey of male sex workers and HIV risk in an urban area of Southwest China.
- Xiao Y, Sun J, Li C, Lu F, Allen KL, et al. (2010) Prevalence and Correlates of HIV and Syphilis Infections Among Men Who Have Sex With Men in Seven Provinces in China With Historically Low HIV Prevalence. *J Acquir Immune Defic Syndr* 53: S66–S73.
- Gupta A, Mehta S, Godbole SV, Sahay S, Walshe L, et al. (2006) Same-Sex Behavior and High Rates of HIV Among Men Attending Sexually Transmitted Infection Clinics in Pune, India (1993–2002). *J Acquir Immune Defic Syndr* 43: 483–490.
- Altaf A, Emmanuel F, Archibald CP, Baloch CR, Uzma Q (2006) Behavioral characteristics of male and eunuch (hijra) sex workers in Karachi, Pakistan.
- Altaf A (2008) Explosive expansion of HIV and associated risk factors among male and hijra sex workers in Sindh, Pakistan 1–190.
- Chan PA, Khan OA (2007) Risk factors for HIV infection in Males who have Sex with Males (MSM) in Bangladesh. *BMC Public Health* 7: 153. doi:10.1186/1471-2458-7-153.
- Hawkes S, Collumbien M, Platt L, Lalji N, Rizvi N, et al. (2009) HIV and other sexually transmitted infections among men, transgenders and women selling sex in two cities in Pakistan: a cross-sectional prevalence survey. *Sexually Transmitted Infections* 85: ii8–ii16. doi:10.1136/sti.2008.033910.

45. Hernandez AL, Lindan CP, Mathur M, Ekstrand M, Madhivanan P, et al. (2006) Sexual Behavior Among Men Who Have Sex with Women, Men, and Hijras in Mumbai, India—Multiple Sexual Risks. *AIDS Behav* 10: 5–16. doi:10.1007/s10461-006-9129-z.
46. Integrated Bio-behavioral Survey (IBBS) among Men who have Sex with Men in the Kathmandu Valley—2007 (2008) Integrated Bio-behavioral Survey (IBBS) among Men who have Sex with Men in the Kathmandu Valley—2007: 1–101.
47. Integrated Bio-behavioral Survey (IBBS) among MSM Population in Kathmandu Valley (2005) Integrated Bio-behavioral Survey (IBBS) among MSM Population in Kathmandu Valley. Center for Research on Environment Health and Population Activities: 1–97.
48. HIV Second Generation Surveillance in Pakistan: National Report Round I (2005) HIV Second Generation Surveillance in Pakistan: National Report Round I. National AIDS Control Program: 1–89.
49. HIV Second Generation Surveillance in Pakistan: National Report Round III (2008) HIV Second Generation Surveillance in Pakistan: National Report Round III. National AIDS Control Program: 1–87.
50. HIV Second Generation Surveillance in Pakistan - National report Round IV 2011 (2011) HIV Second Generation Surveillance in Pakistan - National report Round IV 2011. National AIDS Control Program: 1–120.
51. Saleem NH, Adrien A, Razaque A (2008) Risky sexual behavior, knowledge of sexually transmitted infections and treatment utilization among a vulnerable population in Rawalpindi, Pakistan. *Southeast Asian J Trop Med Public Health* 39: 642–648.
52. Shinde S, Setia MS, Row-Kavi A, Anand V, Jerajani H (2009) Male sex workers: Are we ignoring a risk group in Mumbai, India? *Indian J Dermatol Venereol Lepros* 75: 41–46.
53. Chemnasiri T, Netwong T, Visarutratana S, Varangrat A, Li A, et al. (2010) Inconsistent condom use among young men who have sex with men, male sex workers, and transgenders in Thailand. *AIDS Education and Prevention* 22: 100–109.
54. Clatts MC, Gian LM, Goldsamt LA, Yi H (2007) Male sex work and HIV risk among young heroin users in Hanoi, Vietnam. *Sex Health* 4: 261–267.
55. Colby D, Minh TT, Toan TT (2008) Down on the farm: homosexual behaviour, HIV risk and HIV prevalence in rural communities in Khanh Hoa province, Vietnam. *Sexually Transmitted Infections* 84: 439–443. doi:10.1136/sti.2008.031294.
56. Colby D, Trang N, Lan H, Thien DD (2010) HIV risk and access to prevention services among male sex workers in Ho Chi Minh City, Vietnam.
57. Colby D, Trang NNN, Lan HTX, Nguyen T, Thien DD, et al. (2012) Prevalence of sexually transmitted diseases, HIV, and hepatitis among male sex workers in Ho Chi Minh City, Vietnam. *International Journal of Infectious Diseases* 16: e332. doi:10.1016/j.ijid.2012.05.388.
58. Hoang TV, Tuan NA, Mills SJ, Tung ND, Thang BD, et al. (2006) Results from the HIV/STI integrated biological and behavioral surveillance (IBBS) in Vietnam: 2005–2006.
59. Kladsawad K (2008) Results of the 3rd round of surveillance of HIV infection and risk behavior in populations of men who have sex with men in Thailand.
60. Li A, Varangrat A, Wimonasate W, Chemnasiri T, Sinthuwattanawibool C, et al. (2008) Sexual Behavior and Risk Factors for HIV Infection Among Homosexual and Bisexual Men in Thailand. *AIDS Behav* 13: 318–327. doi:10.1007/s10461-008-9448-3.
61. HIV Prevalence Among Populations of Men Who Have Sex with Men – Thailand, 2003 and 2005 (2005) HIV Prevalence Among Populations of Men Who Have Sex with Men – Thailand, 2003 and 2005. *MMWR Recomm Rep* 55: 844–848.
62. Nguyen TA, Nguyen HT, Le GT, Detels R (2007) Prevalence and Risk Factors Associated with HIV Infection Among Men Having Sex with Men in Ho Chi Minh City, Vietnam. *AIDS Behav* 12: 476–482. doi:10.1007/s10461-007-9267-y.
63. Sheridan S, Phimpachanh C, Chanlivong N, Manivong S, Khamsvolsong S, et al. (2009) HIV prevalence and risk behaviour among men who have sex with men in Vientiane Capital, Lao People's Democratic Republic, 2007. *AIDS* 23: 409–414. doi:10.1097/QAD.0b013e32831ef510.
64. van Griensven F, Thienkrua W, McNicholl J, Wimonasate W, Chaikummao S, et al. (2013) Evidence of an explosive epidemic of HIV infection in a cohort of men who have sex with men in Thailand. *AIDS* 27: 825–832. doi:10.1097/QAD.0b013e32835c546e.
65. Creswell J, Guardado ME, Lec J, Nieto AI, Kim AA, et al. (2012) HIV and STI control in El Salvador: results from an integrated behavioural survey among men who have sex with men. *Sexually Transmitted Infections* 88: 633–638. doi:10.1136/sextrans-2012-050521.
66. Jacobson JO, Sánchez-Gómez A, Montoya O, Soria E, Tarupi W, et al. (2013) A Continuing HIV Epidemic and Differential Patterns of HIV-STI Risk among MSM in Quito, Ecuador: An Urgent Need to Scale Up HIV Testing and Prevention. *AIDS Behav*. doi:10.1007/s10461-013-0478-0.
67. Magis C, Campos P, Soler C, Morales S, Gayet C, et al. (n.d.) HIV prevalence and factors associated with the possession of condoms among male sex workers in two cities: Guadalajara and Mexico City, Mexico.
68. Montano SM, Sanchez JL, Laguna-Torres A, Cuchi P, Avila MM, et al. (2005) Prevalences, Genotypes, and Risk Factors for HIV Transmission in South America: 1–8.
69. Schuelter-Trevisol F, Custodio G, Barreto da Silva AC, de Oliveira MB, Wolfart A, et al. (2013) HIV, hepatitis B and C, and syphilis prevalence and coinfection among sex workers in Southern Brazil. *Revista da Sociedade Brasileira de Medicina Tropical* 46: 493–497. doi:10.1590/0037-8682-1364-2013.
70. Ramos FarÃ as dos MAS, Garcia MN, Reynaga E, Romero M, Vaulet MLG, et al. (2011) First report on sexually transmitted infections among trans (male to female transvestites, transsexuals, or transgender) and male sex workers in Argentina: high HIV, HPV, HBV, and syphilis prevalence. *International Journal of Infectious Diseases* 15: e635–e640. doi:10.1016/j.ijid.2011.05.007.
71. Konda KA, Clark JL, Segura E, Salvatierra J, Galea J, et al. (2008) Male sex workers among men who have sex with men in Lima, Peru.
72. Segura M, Estani SS, Marone R, Bautista CT, Pando MA, et al. (2007) Buenos Aires Cohort of Men Who Have Sex with Men: Prevalence, Incidence, Risk Factors, and Molecular Genotyping of HIV Type 1. *AIDS Research and Human Retroviruses* 23: 1322–1329. doi:10.1089/aid.2007.0063.
73. Baral S, Burrell E, Scheibe A, Ben Brown, Beyrer C, et al. (2011) HIV Risk and Associations of HIV Infection among men who have sex with men in Peri-Urban Cape Town, South Africa. *BMC Public Health* 11: 766. doi:10.1186/1471-2458-11-766.
74. Gakii G, Kimani J, Gelmon L, Izulla P, Swop Clinic SS (2010) Male sex workers who have sex with men (MSM-SW): marginalized and high risk population with poor access to STI/HIV health services in Nairobi, Kenya AIDS 2010 - XVIII International AIDS Conference.
75. Hladik W, Barker J, Ssenkusu JM, Opio A, Tappero JW, et al. (2012) HIV Infection among Men Who Have Sex with Men in Kampala, Uganda—A Respondent Driven Sampling Survey. *PLoS ONE* 7: e38143. doi:10.1371/journal.pone.0038143.t003.
76. McKinnon L, Gakii G, Juno J, Izulla P, Munyao J, et al. (2013) High HIV prevalence, incidence and diverse sexual practices in a cohort of male sex workers in Nairobi, Kenya.
77. Muraguri N, Tun W, Okal J, Broz D, Raymond HF, et al. (2012) Burden of HIV and sexual behavior among men who have sex with men and male sex workers in Nairobi, Kenya.
78. Pitipan EV, Kalichman SC, Eaton LA, Watt MH, Sikkema KJ, et al. (2013) Men (and Women) as “Sellers” of Sex in Alcohol-Serving Venues in Cape Town, South Africa. *Prev Sci*. doi:10.1007/s11121-013-0381-y.
79. Vuylsteke B, Semde G, Sika L, Crucitti T, Etiéngue Traore V, et al. (2012) High prevalence of HIV and sexually transmitted infections among male sex workers in Abidjan, Cote d'Ivoire: need for services tailored to their needs. *Sexually Transmitted Infections* 88: 288–293. doi:10.1136/sextrans-2011-050276.
80. Wade AS, Kane CT, Diallo PAN, Diop AK, Gueye K, et al. (2005) HIV infection and sexually transmitted infections among men who have sex with men in Senegal. *AIDS* 19: 2133–2140.
81. van der Elst EM, Okuku HS, Nakanya P, Muhaari A, Davies A, et al. (2009) Is Audio Computer-Assisted Self-Interview (ACASI) Useful in Risk Behaviour Assessment of Female and Male Sex Workers, Mombasa, Kenya? *PLoS ONE* 4: e5340. doi:10.1371/journal.pone.0005340.t004.
82. Baral S, Kizub D, Masenior NF, Peryskina A, Stachowiak J, et al. (2010) Male sex workers in Moscow, Russia: a pilot study of demographics, substance use patterns, and prevalence of HIV-1 and sexually transmitted infections. *AIDS Care* 22: 112–118. doi:10.1080/09540120903012551.
83. Belza MJ (2005) Risk of HIV infection among male sex workers in Spain. *Sexually Transmitted Infections* 81: 85–88. doi:10.1136/sti.2003.008649.
84. Bruckova M, Bautista CT, Graham RR, Maly M, Vandasova J, et al. (2006) HIV infection among commercial sex workers and injecting drug users in the Czech Republic. *Am J Trop Med Hyg* 75: 1017–1020.
85. Cuyppers WJ, Niekamp AM, Keesmekers R, Spauwen L, Hollman D, et al. (2011) P1-S2.11 High prevalence of HIV, other sexually transmitted infections and risk profile in male commercial sex workers who have sex with men in the Netherlands. *Sexually Transmitted Infections* 87: A127–A127. doi:10.1136/sextrans-2011-050108.68.
86. Leuridan E, Wouters K, Stalpaert M, Van Damme P (2005) Male sex workers in Antwerp, Belgium: a descriptive study. *International Journal of STD & AIDS* 16: 744–748. doi:10.1258/095646205774763072.
87. Mor Z (2012) Knowledge, attitudes, sexual practices and STI/HIV prevalence in male sex workers and other men who have sex in Tel Aviv, Israel: a cross-sectional study. *Sex Transm Infect* 88: 574–580. doi:10.1136/sextrans-2011.
88. Del Romero Guerrero J, Mengel M, Rodriguez-Martin C, Ballesteros Martin J, Clavo Escribano P, et al. (2011) HIV prevalence in male and transsexual sex workers in Madrid, Spain.
89. Bimbi DS, Parsons JT (2005) Barebacking Among Internet Based Male Sex Workers. *Journal of Gay & Lesbian Psychotherapy* 9: 85–105. doi:10.1300/J236v09n03_06.
90. Burnette ML, Lucas E, Ilgen M, Frayne SM, Mayo J, et al. (2008) Prevalence and Health Correlates of Prostitution Among Patients Entering Treatment for Substance Use Disorders. *Arch Gen Psychiatry* 65: 337–344.
91. Cohan D (2006) Sex worker health: San Francisco style. *Sexually Transmitted Infections* 82: 418–422. doi:10.1136/sti.2006.020628.
92. Fujimoto K, Williams ML, Ross MW (2013) Venue-Based Affiliation Networks and HIV Risk-Taking Behavior Among Male Sex Workers. *Sexually Transmitted Diseases* 40: 453–458. doi:10.1097/OLQ.0b013e31829186e5.

93. Haley N (2004) HIV risk profile of male street youth involved in survival sex. *Sexually Transmitted Infections* 80: 526–530. doi:10.1136/sti.2004.010728.
94. Kral AH (2005) HIV Prevalence and Risk Behaviors Among Men Who Have Sex with Men and Inject Drugs in San Francisco. *Journal of Urban Health: Bulletin of the New York Academy of Medicine* 82: i43–i50. doi:10.1093/jurban/jti023.
95. Kuypers LM (2004) Factors associated with sex trade involvement among male participants in a prospective study of injection drug users. *Sexually Transmitted Infections* 80: 531–535. doi:10.1136/sti.2004.011106.
96. Smith MD, Seal DW (2008) Sexual Behavior, Mental Health, Substance Use, and HIV Risk Among Agency-Based Male Escorts in a Small U.S. City. *International Journal of Sexual Health* 19: 27–39. doi:10.1300/J514v19n04_04.
97. Timpson SC, Ross MW, Williams ML, Atkinson J (2007) Characteristics, Drug Use, and Sex Partners of a Sample of Male Sex Workers. *Am J Drug Alcohol Abuse* 33: 63–69. doi:10.1080/00952990601082670.
98. Williams ML, Bowen AM, Timpson SC, Ross MW, Atkinson JS (2006) HIV prevention and street-based male sex workers: an evaluation of brief interventions. *AIDS Education and Prevention* 18: 204–215.
99. Washington TA, Meyer-Adams N (2010) HIV Prevention Needs of Sex-Trading Injection Drug-Using Black Men Who Have Sex With Both Men and Women. *American Journal of Men's Health* 4: 104–110. doi:10.1177/1557988308330249.
100. Hoang TV, Tuan NA, Vi LNL, Mills SJ, Ha TTT, et al. (2009) Results from the HIV/STI integrated biological and behavioral surveillance (IBBS) in Vietnam - Round II 2009.
101. Baral S, Beyrer C, Muessig K, Potat T, Wirtz AL, et al. (2012) Burden of HIV among female sex workers in low-income and middle-income countries: a systematic review and meta-analysis. *The Lancet Infectious Diseases* 12: 538–549. doi:10.1016/S1473-3099(12)70066-X.
102. Baral S, Sifakis F, Cleghorn F, Beyrer C (2007) Elevated risk for HIV infection among men who have sex with men in low- and middle-income countries 2000–2006: A systematic review. *PLoS Medicine* 4: e339. doi:10.1371/journal.pmed.
103. Operario D, Soma T, Underhill K (2008) Sex Work and HIV Status Among Transgender Women: Systematic Review and Meta-Analysis. *J Acquir Immune Defic Syndr* 48: 97–103.
104. Baral S, Potat T, Stromdahl S, Wirtz AL, Guadamuz T, et al. (2013) Worldwide burden of HIV in transgender women: a systematic review and meta-analysis. *The Lancet Infectious Diseases* 13: 214–222. doi:10.1016/S1473-3099(12)70315-8.
105. Baral S, Logie CH, Grosso A, Wirtz AL, Beyrer C (2013) Modified social ecological model: a tool to guide the assessment of the risks and risk contexts of HIV epidemics. *BMC Public Health* 13: 1–1. doi:10.1186/1471-2458-13-482.
106. Beyrer C, Baral SD, van Griensven F, Goodreau SM, Chariyalertsak S, et al. (2012) HIV in men who have sex with men 1: Global epidemiology of HIV infection in men who have sex with men. *The Lancet* 380: 367–377. doi:10.1016/S0140-6736(12)60821-6.
107. Semugoma P, Beyrer C, Baral S (2012) Assessing the effects of anti-homosexuality legislation in Uganda on HIV prevention, treatment, and care services. *SAHARA-J: Journal of Social Aspects of HIV/AIDS* 9: 173–176. doi:10.1080/17290376.2012.744177.
108. Fay H, Baral SD, Trapence G, Motimedi F, Umar E, et al. (2010) Stigma, Health Care Access, and HIV Knowledge Among Men Who Have Sex With Men in Malawi, Namibia, and Botswana. *AIDS Behav* 15: 1088–1097. doi:10.1007/s10461-010-9861-2.
109. Chow EPF, Iu KI, Fu X, Wilson DP, Zhang L (2012) HIV and Sexually Transmissible Infections among Money Boys in China: A Data Synthesis and Meta-Analysis. *PLoS ONE* 7: e48025. doi:10.1371/journal.pone.0048025.s004.
110. Mimiaga MJ, Biello KB, Sivasubramanian M, Mayer KH, Anand VR, et al. (2013) Psychosocial risk factors for HIV sexual risk among Indian men who have sex with men. *AIDS Care* 25: 1109–1113. doi:10.1080/09540121.2012.749340.