



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Translating neuronal activity at the synapse: presynaptic calcium sensors in short-term plasticity

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters.

Citation	de Jong, Arthur P. H., and Diasynou Fioravante. 2014. "Translating neuronal activity at the synapse: presynaptic calcium sensors in short-term plasticity." <i>Frontiers in Cellular Neuroscience</i> 8 (1): 356. doi:10.3389/fncel.2014.00356. http://dx.doi.org/10.3389/fncel.2014.00356 .
Published Version	doi:10.3389/fncel.2014.00356
Accessed	February 17, 2015 6:50:40 AM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:13454838
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)



Translating neuronal activity at the synapse: presynaptic calcium sensors in short-term plasticity

Arthur P. H. de Jong^{1*} and Diasynou Fioravante^{2*}

¹ Department of Neurobiology, Harvard Medical School, Boston, MA, USA

² Department of Neurobiology, Physiology and Behavior, Center for Neuroscience, University of California Davis, Davis, CA, USA

Edited by:

Philippe Isope, Centre National pour la Recherche Scientifique, France

Reviewed by:

Marco Canepari, Institut National de la Santé et de la Recherche Médicale, France

Erwin Neher, Max Planck Institute for Biophysical Chemistry, Germany

*Correspondence:

Arthur P. H. de Jong, Department of Neurobiology, Harvard Medical School, 220 Longwood Ave, Boston, MA 02115, USA

e-mail: arthur_de_jong@hms.harvard.edu;

Diasynou Fioravante, Department of Neurobiology, Physiology and Behavior, Center for Neuroscience, University of California Davis, 1544 Newton Court, Davis, CA 95618, USA

e-mail: dfioravante@ucdavis.edu

The complex manner in which patterns of presynaptic neural activity are translated into short-term plasticity (STP) suggests the existence of multiple presynaptic calcium (Ca^{2+}) sensors, which regulate the amplitude and time-course of STP and are the focus of this review. We describe two canonical Ca^{2+} -binding protein domains (C2 domains and EF-hands) and define criteria that need to be met for a protein to qualify as a Ca^{2+} sensor mediating STP. With these criteria in mind, we discuss various forms of STP and identify established and putative Ca^{2+} sensors. We find that despite the multitude of proposed sensors, only three are well established in STP: Munc13, protein kinase C (PKC) and synaptotagmin-7. For putative sensors, we pinpoint open questions and potential pitfalls. Finally, we discuss how the molecular properties and modes of action of Ca^{2+} sensors can explain their differential involvement in STP and shape net synaptic output.

Keywords: C2 domain, protein kinase C, Munc13, synaptotagmin, calmodulin, post-tetanic potentiation, residual calcium, short-term plasticity

INTRODUCTION

Synaptic transmission is initiated by action potential-evoked influx of calcium (Ca^{2+}) into the presynaptic terminal, which triggers fusion of vesicles by binding to a specialized Ca^{2+} sensor. Bursts of action potentials lead to the buildup of residual Ca^{2+} ($[\text{Ca}^{2+}]_{\text{residual}}$) in the terminal, which outlives neuronal activity, and induce multiple forms of short-term presynaptic plasticity (STP), including facilitation, depression, augmentation and post-tetanic potentiation (PTP) (reviewed in Fioravante and Regehr, 2011). STP plays a crucial role in synaptic computations and shapes the properties of microcircuits (reviewed in Abbott and Regehr, 2004; Regehr, 2012).


The dynamics of some forms of STP are dictated by the kinetics of $[\text{Ca}^{2+}]_{\text{residual}}$ (Delaney et al., 1989; Kamiya and Zucker, 1994) and can be explained by changes in vesicular release probability (Katz and Miledi, 1968; Zucker and Stockbridge, 1983) or by depletion of the readily releasable pool (RRP) of vesicles (Bailey and Chen, 1988; Liu and Tsien, 1995; von Gersdorff and Matthews, 1997). However, at several synapses the magnitude of facilitation is higher than can be explained by $[\text{Ca}^{2+}]_{\text{residual}}$ alone, and both facilitation and PTP decay slower than the $[\text{Ca}^{2+}]_{\text{residual}}$ signal (Regehr et al., 1994; Atluri and Regehr, 1996; Brager et al., 2003; Felmy et al., 2003; Fioravante et al., 2011; **Figure 1**). Furthermore, many types of STP rely on the regulation of steps upstream of vesicle fusion (Dittman and Regehr, 1998;

Wang and Kaczmarek, 1998), including RRP refilling and Ca^{2+} influx through voltage-gated Ca^{2+} channels (VGCCs; Stevens and Wesseling, 1998; Xu and Wu, 2005; Mochida et al., 2008; Müller et al., 2008; Leal et al., 2012). These events are strongly Ca^{2+} -dependent, and thus Ca^{2+} sensors must be activated to induce and sustain STP. The Ca^{2+} sensors that mediate STP are the topic of this mini-review. First, we will discuss the molecular structure and function of two Ca^{2+} -binding domains employed by Ca^{2+} sensors: C2 domains and EF-hands. Subsequently, we will define the criteria for establishing Ca^{2+} sensors for STP, and, guided by these criteria, discuss a body of recent literature on well accepted and putative sensors that regulate STP.

Ca^{2+} -BINDING MOTIFS

C2 DOMAINS

The best described Ca^{2+} sensors in the context of synapses are C2 domains, which are found in many signal transduction and membrane trafficking proteins (Rizo and Südhof, 1998). C2 domains consist of ~ 130 amino acids that form a compact β -sandwich of two 4-stranded β -sheets. Three loops connecting the β -sheets at the top of the domain contain 4–5 highly conserved aspartates that coordinate the binding of 2 to 3 Ca^{2+} ions (Shao et al., 1996; Ubach et al., 1998; Fernandez et al., 2001). The Ca^{2+} -binding properties of C2 domains have been described in detail in synaptotagmin (syt), which acts as the Ca^{2+} sensor for



synchronous vesicle fusion at most synapses (Pang and Südhof, 2010). Mutations that interfere with Ca²⁺ binding on syt-1 alter the Ca²⁺-sensitivity of vesicle fusion (Nishiki and Augustine, 2004; Shin et al., 2009; Kochubey and Schneggenburger, 2011; Kochubey et al., 2011). Analogous mutation analyses of other C2 domains showed similar effects on Ca²⁺ binding (Shin et al., 2010; Fioravante et al., 2014; Liu et al., 2014). Some C2 domains naturally lack these aspartate residues and cannot bind Ca²⁺ (e.g., Pappa et al., 1998). Ca²⁺ binding increases the affinity of C2 domains for phospholipids (Brose et al., 1992; Fernandez et al., 2001), thus recruiting the domain to the plasma membrane. In addition, it may trigger a conformational change that increases association with effector proteins (for instance syt-1

binding to SNAREs (Bai et al., 2004)) or exposes a domain within the protein (e.g., the MUN domain of Munc13 (Shin et al., 2010)). Many C2 domains display a steep increase in Ca²⁺ affinity in the presence of phosphatidylinositol 4,5-bisphosphate (PIP2; van den Bogaart et al., 2012), which helps localize the domain to the PIP2-enriched active zone (Rohrbough and Broadie, 2005).

EF-HANDS

The EF-hand is the most common Ca²⁺-binding motif, with diverse cellular functions including cytoplasmic Ca²⁺ buffering and signal transduction (Skelton et al., 1994; Schaub and Heizmann, 2008; Schwaller, 2009). The motif consists of two

α -helices connected by a linker of 12 amino acids (Lewit-Bentley and Réty, 2000). Six residues within this linker coordinate binding to a single Ca^{2+} ion, and their mutation abolishes Ca^{2+} binding (Maune et al., 1992). Examples of EF-hand-containing proteins with proposed Ca^{2+} -sensing roles in STP include calmodulin (CaM), neuronal calcium sensor 1 (NCS-1) and visin-like proteins (VILIPs).

CaM is the prototypical EF-hand protein that interacts with numerous effector proteins in a Ca^{2+} -dependent manner (Xia and Storm, 2005). Important presynaptic effectors are CaM-dependent kinase II (CaMKII), myosin light chain kinase (MLCK), adenylyl cyclase, the protein phosphatase calcineurin, Munc13, VGCCs and Ca^{2+} -activated potassium channels, all of which regulate presynaptic function (de Jong and Verhage, 2009; Adelman et al., 2012). Because the Ca^{2+} affinity of CaM is differentially regulated by its binding partners, different CaM-protein complexes vary in their Ca^{2+} sensitivity (Olwin and Storm, 1985; Xia and Storm, 2005) and could therefore be differentially engaged during various forms of STP. Direct assessment of the role of CaM as a Ca^{2+} sensor for STP has proven difficult because manipulations of CaM levels alter expression of >200 genes (Pang et al., 2010) and rescue experiments in neuronal preparations with Ca^{2+} -binding mutants of CaM have not been conducted thus far.

DEFINITION OF A Ca^{2+} SENSOR FOR STP

With a plethora of C2- and EF-hand-containing proteins in the presynaptic terminal, there are numerous candidate Ca^{2+} sensors for STP. We propose that in order to qualify as a sensor for STP, a protein must fulfill the following three criteria:

1. *Ca^{2+} must bind directly to the protein.* An obvious requirement for a Ca^{2+} sensor is that it must bind Ca^{2+} . Some EF-hands and C2 domains lack the Ca^{2+} -coordinating residues and cannot bind Ca^{2+} . Therefore, Ca^{2+} binding must be experimentally established for each protein.
2. *Protein must be part of, or directly modulate, vesicle availability or the vesicle release machinery.* Changes in availability and/or fusogenicity of synaptic vesicles and in presynaptic Ca^{2+} influx shape STP (Dutta Roy et al., 2014). A Ca^{2+} sensor for STP must therefore directly affect vesicle availability (recruitment, docking, priming) and/or the vesicle fusion machinery, including VGCCs and SM proteins (for a discussion of release machinery, see Südhof, 2013). This definition includes enzymes like kinases, which directly regulate the properties of these components. For the purpose of this review, we do not consider Ca^{2+} buffers (e.g., parvalbumin) and pumps, which indirectly affect STP by changing the spatiotemporal distribution of free Ca^{2+} through binding or extrusion (Müller et al., 2007; Scullin and Partridge, 2010), or components of the endocytotic machinery, which can affect vesicle or release site availability after prolonged episodes of exocytosis (Wilkinson and Lin, 2004; Hosoi et al., 2009).
3. *Mutations that interfere with Ca^{2+} binding affect STP.* Even if a protein satisfies criteria 1 and 2, it is not a Ca^{2+} sensor for STP unless Ca^{2+} binding is required for the protein's function in STP. For instance, whether Ca^{2+} binding to Doc2 is required for spontaneous release is debated and the role of Doc2 as a

Ca^{2+} sensor for spontaneous release remains unclear (Groffen et al., 2010; Pang et al., 2011). Therefore, it is necessary to show that mutation of the Ca^{2+} binding site abolishes function (for example using a knockout/rescue or knockin approach) in order to conclude that a protein is a Ca^{2+} sensor mediating STP. It could even be argued that a requirement for Ca^{2+} binding *during plasticity* must be demonstrated in order to establish a protein as a Ca^{2+} sensor, but the technology for this type of experiments is currently lacking.

Ca^{2+} SENSORS IN STP

FACILITATION

At synapses with low initial release probability, brief bursts of activity can induce transient facilitation of release, which relies on increased release probability due to elevated $[\text{Ca}^{2+}]_{\text{residual}}$ (Katz and Miledi, 1968; Kamiya and Zucker, 1994; Regehr et al., 1994). However, this mechanism alone cannot fully explain the magnitude of facilitation at all synapses (Atluri and Regehr, 1996; Felmy et al., 2003), and additional Ca^{2+} -dependent processes have been suggested (Zucker and Regehr, 2002), including the existence of a yet unidentified presynaptic Ca^{2+} sensor distinct from syt-1 (Bain and Quastel, 1992; Saraswati et al., 2007). Enhancement of Ca^{2+} currents is an attractive mechanism to mediate facilitation, and the capability of Ca^{2+} /CaM to modulate overexpressed VGCCs during strong depolarization has been studied extensively (Catterall et al., 2013). Ca^{2+} /CaM binds to a regulatory domain of $\text{Ca}_v2.1$, the VGCC that mediates the P/Q type Ca^{2+} current driving synaptic transmission in most synapses. In heterologous cell lines, this interaction leads to enhancement of Ca^{2+} currents, which depends on Ca^{2+} binding to CaM (Lee et al., 1999; DeMaria et al., 2001). Several EF-hand-containing proteins including VILIPs, CaBPs and NCS-1 (collectively named neuronal Ca^{2+} sensors, or nCaS) also modulate Ca^{2+} influx through VGCCs (Few et al., 2005; Lautermilch et al., 2005; Burgoyne, 2007; Dason et al., 2012; Catterall et al., 2013) and may affect facilitation in a manner that depends on the nCaS binding domain of VGCCs (Tsujimoto et al., 2002; Sippy et al., 2003; Mochida et al., 2008; Leal et al., 2012). For none of these protein functions, however, has a Ca^{2+} binding requirement been established, and some of them may actually be independent of Ca^{2+} (Few et al., 2005). In addition, due to the lack of suitable genetic models, most experiments rely on overexpression of exogenous proteins (Mochida et al., 2003). Whether nCaS are specifically involved in the regulation of STP, or the altered STP is a consequence of altered basal synaptic properties, remains controversial (Dason et al., 2012).

DEPRESSION AND RECOVERY FROM DEPRESSION

Prolonged high-frequency stimulation leads to transient decrease in presynaptic strength, which can be due to depletion of the RRP (Elmqvist and Quastel, 1965; Liu and Tsien, 1995; Schneggenburger et al., 2002) and activity-dependent decrease in Ca^{2+} influx (Forsythe et al., 1998; Xu and Wu, 2005) (for a complete review of known mechanisms of depression, see Regehr, 2012). CaM, CaBP1 and NCS-1 have been proposed as putative Ca^{2+} sensors to mediate the latter effect (Xu and Wu, 2005; Catterall and Few, 2008; Mochida et al., 2008). Depression can be slowed by Ca^{2+} -dependent replenishment of the RRP

(Stevens and Wesseling, 1998; Wang and Kaczmarek, 1998). The vesicle priming factor Munc13 acts as a Ca^{2+} sensor to determine the rate of depression, via its C2B and CaM-binding domains. Ca^{2+} binding to the C2B domain of Munc13 activates its MUN domain that promotes assembly of the machinery responsible for vesicle fusion, thereby increasing refilling of the RRP (Shin et al., 2010; Ma et al., 2011). Indeed, Munc13 knockout neurons expressing a variant of the protein with mutated Ca^{2+} -coordinating aspartates display increased synaptic depression without affecting initial release probability (Shin et al., 2010). In addition, Munc13 binds Ca^{2+} /CaM, and this interaction also accelerates RRP refilling (Junge et al., 2004; Lipstein et al., 2012, 2013). In line with this observation, CaM inhibitors slow the RRP refilling rate (Sakaba and Neher, 2001; Hosoi et al., 2007). Although a Ca^{2+} -binding CaM mutant has not been studied in this context, the CaM/Munc13 interaction is strongly Ca^{2+} -dependent (Junge et al., 2004; Dimova et al., 2006; Lipstein et al., 2012), thus making the Ca^{2+} /CaM-Munc13 complex a likely Ca^{2+} -sensor for STP.

Synaptotagmin-7 has also been identified as a sensor that regulates depression, operating via its two Ca^{2+} -binding C2 domains (Liu et al., 2014). At the zebrafish neuromuscular junction, syt-7 regulates desynchronized release (Wen et al., 2010), but its function in mammalian neurons has been debated (Maximov et al., 2008; Bacaj et al., 2013; Liu et al., 2014). A recent study showed that in syt-7 knockout mice, initial release probability is unaffected but the rate of vesicle replenishment during and after bursts of activity is significantly reduced (Liu et al., 2014). This phenotype is rescued by wild-type syt-7 but not by syt-7 carrying mutations of the Ca^{2+} binding sites, demonstrating that syt-7 is a Ca^{2+} sensor that mediates RRP refilling. Syt-7 also probably interacts with Ca^{2+} /CaM (Liu et al., 2014), but the functional significance of this complex remains to be identified.

In contrast to the proteins discussed above that promote recovery from depression, rabphilin is thought to slow down recovery from depression (Deák et al., 2006). Rabphilin is a synaptic vesicle protein with two Ca^{2+} -sensing C2 domains (Yamaguchi et al., 1993; Ubach et al., 1999; Coudevylle et al., 2008), but whether Ca^{2+} binding is required for its role in STP has not been determined.

AUGMENTATION AND PTP

Augmentation and PTP are two closely related forms of STP that require prolonged high-frequency stimulation (Magleby, 1973; Magleby and Zengel, 1976a; Stevens and Wesseling, 1999; Habets and Borst, 2005; Korogod et al., 2005). For augmentation, varying stimulus duration increases the peak amplitude of the enhancement without significantly affecting the time course of decay (Magleby, 1979). The mechanisms underlying augmentation are not well understood and changes in both release probability and Ca^{2+} -dependent replenishment of the RRP have been proposed (Magleby and Zengel, 1976b; Stevens and Wesseling, 1999; Rosenmund et al., 2002; Kalkstein and Magleby, 2004). Munc13 and syt-7 have been suggested as Ca^{2+} sensors for augmentation (Shin et al., 2010; Lipstein et al., 2013; Liu et al., 2014),

but since both sensors affect depression as well, dissociation of their roles in synaptic depression vs. augmentation has not been possible. Various phospholipase C (PLC) isoforms could also act as Ca^{2+} sensors because they require binding of a Ca^{2+} ion for activation of their catalytic domain (Grobler and Hurley, 1998; Rebecchi and Pentylala, 2000). Pharmacological studies suggest that PLC activation is required for augmentation (Rosenmund et al., 2002) but not PTP (Genc et al., 2014). PLC hydrolyses PIP2 to diacylglycerol, which could lead to potentiation of synaptic transmission via Munc13 and protein kinase C (PKC; de Jong and Verhage, 2009).

PTP typically lasts longer than augmentation and shows a progressive increase in the time course of decay with increased duration and frequency of stimulation (Magleby, 1979; Korogod et al., 2005). Pharmacological (e.g., Alle et al., 2001; Brager et al., 2002; Beierlein et al., 2007; Korogod et al., 2007) and genetic (Fioravante et al., 2011, 2012, 2014; Chu et al., 2014) studies at several synapses have firmly established the requirement for PKC in PTP. Three PKC isoforms (α , β and γ) possess a C2 domain and bind Ca^{2+} with low micromolar affinity (Torrecillas et al., 2004; Newton, 2010; **Figure 1**). PKCs enhance release through phosphorylation of effectors, including components of the vesicular release machinery such as Munc18 (Wierda et al., 2007; de Jong and Verhage, 2009; Genc et al., 2014). Mutations of the Ca^{2+} -coordinating aspartates in the C2 domain of PKC β abolish its ability to support PTP, without affecting basal synaptic function (Fioravante et al., 2014).

PKC β is probably not the only Ca^{2+} sensor for PTP. At the immature calyx of Held, PTP depends on PKC γ (Chu et al., 2014). Moreover, at the parallel fiber-Purkinje cell synapse in the cerebellum, PKC α can readily support PTP in the absence of PKC β and γ (Fioravante et al., 2012). It remains to be tested whether Ca^{2+} binding to PKC α and γ is necessary for PTP and whether all PKC isoforms act through Munc18 phosphorylation. Finally, pharmacological studies suggest that Ca^{2+} /CaM, acting via MLCK, makes a small contribution to PTP at immature, but not functionally mature, synapses (Lee et al., 2008; Fioravante et al., 2011).

Tetanic stimulation enhances not only evoked responses (i.e., PTP) but also spontaneous events in a Ca^{2+} -dependent manner. The frequency (Zengel and Magleby, 1981; Zucker and Lara-Estrella, 1983; Eliot et al., 1994; Habets and Borst, 2005), and at some synapses also the amplitude (He et al., 2009), of spontaneous events increase after tetanization. Because of similarities in the time course of these effects with PTP, a common mechanism has been speculated (Zengel and Magleby, 1981). However, the effects of $[\text{Ca}^{2+}]_{\text{residual}}$ on spontaneous transmission were recently shown to be independent of PKC (Xue and Wu, 2010; Fioravante et al., 2011; but see Brager et al., 2003) and the increase in amplitude requires syt-2 (He et al., 2009). The Ca^{2+} sensors remain unknown.

DIFFERENTIAL ENGAGEMENT OF Ca^{2+} SENSORS AND IMPLICATIONS FOR STP

Different patterns of neuronal activity result in variable Ca^{2+} signals stretching over an order of magnitude (**Figure 1**). Diverse

sensors are therefore needed to translate the Ca^{2+} signals into distinct forms of STP. Factors such as Ca^{2+} affinity, specific (sub-)cellular expression and mechanisms of action contribute to the specialization of sensors for different forms of STP. For example, NCS-1 has high affinity for Ca^{2+} and localizes at the plasma membrane (O'Callaghan et al., 2002; Burgoyne, 2007) where it could rapidly respond to local Ca^{2+} signals. PKC β , on the other hand, has lower Ca^{2+} affinity, is cytoplasmic at rest (Newton, 2010) and likely has to phosphorylate more than one substrates to induce plasticity; therefore, sustained, global Ca^{2+} increases are likely required for its activation, in agreement with the prolonged stimulation requirement for PTP (Habets and Borst, 2005; Korogod et al., 2005). Even for the same sensor, Ca^{2+} affinity can vary as a result of effector binding, phospholipid binding, and post-translational modifications (Xia and Storm, 2005; Li et al., 2011; van den Bogaart et al., 2012). Finally, specific expression patterns of Ca^{2+} sensors could help explain why identical activation regimes do not always lead to the same STP across synapses or during development (Rosenmund et al., 2002; Chu et al., 2014).

Most synapses exhibit multiple forms of STP and the net synaptic output reflects the interaction between these different forms (de Jong and Verhage, 2009). It is therefore likely that different Ca^{2+} sensors interact, and might even compete (Chu et al., 2014), during STP. The dynamics of these interactions should be considered when building computational models of STP. Traditionally, such models combine use-dependent depletion and Ca^{2+} -dependent facilitation to explain synaptic output (Tsodyks et al., 1998; Fuhrmann et al., 2002; Pfister et al., 2010). Introduction of additional components such as vesicle replenishment, which are engaged under conditions that activate the corresponding Ca^{2+} sensors, more accurately reflects our understanding of the underlying biology and allows better prediction of synaptic and network behavior (Hennig, 2013).

ACKNOWLEDGMENTS

We would like to thank Drs. P.S. Kaeser, M. Verhage, M. Thanawala, W. Regehr and E. Antzoulatos for critically reading the manuscript. This work was funded by the Netherlands Organization for Scientific Research (NWO, 825.12.028) to Arthur P. H. de Jong and UC Davis College of Biological Sciences Dean's start-up award to Diasynou Fioravante.

REFERENCES

- Abbott, L. F., and Regehr, W. G. (2004). Synaptic computation. *Nature* 431, 796–803. doi: 10.1038/nature03010
- Adelman, J. P., Maylie, J., and Sah, P. (2012). Small-Conductance Ca^{2+} -Activated K^+ channels: form and function. *Annu. Rev. Physiol.* 74, 245–269. doi: 10.1146/annurev-physiol-020911-153336
- Alle, H., Jonas, P., and Geiger, J. R. (2001). PTP and LTP at a hippocampal mossy fiber-interneuron synapse. *Proc. Natl. Acad. Sci. U S A* 98, 14708–14713. doi: 10.1073/pnas.251610898
- Aravind, P., Chandra, K., Reddy, P. P., Jeromin, A., Chary, K. V. R., and Sharma, Y. (2008). Regulatory and structural EF-hand motifs of neuronal calcium sensor-1: Mg^{2+} modulates Ca^{2+} binding, Ca^{2+} -induced conformational changes and equilibrium unfolding transitions. *J. Mol. Biol.* 376, 1100–1115. doi: 10.1016/j.jmb.2007.12.033
- Atluri, P. P., and Regehr, W. G. (1996). Determinants of the time course of facilitation at the granule cell to Purkinje cell synapse. *J. Neurosci.* 16, 5661–5671.
- Bacaj, T., Wu, D., Yang, X., Morishita, W., Zhou, P., Xu, W., et al. (2013). Synaptotagmin-1 and synaptotagmin-7 trigger synchronous and asynchronous phases of neurotransmitter release. *Neuron* 80, 947–959. doi: 10.1016/j.neuron.2013.10.026
- Bai, J., Wang, C.-T., Richards, D. A., Jackson, M. B., and Chapman, E. R. (2004). Fusion Pore dynamics are regulated by Synaptotagmin-SNARE interactions. *Neuron* 41, 929–942. doi: 10.1016/s0896-6273(04)00117-5
- Bailey, C. H., and Chen, M. (1988). Morphological basis of short-term habituation in *Aplysia*. *J. Neurosci.* 8, 2452–2459.
- Bain, A. I., and Quastel, D. M. (1992). Multiplicative and additive Ca^{2+} -dependent components of facilitation at mouse endplates. *J. Physiol.* 455, 383–405.
- Beierlein, M., Fioravante, D., and Regehr, W. G. (2007). Differential expression of posttetanic potentiation and retrograde signaling mediate target-dependent short-term synaptic plasticity. *Neuron* 54, 949–959. doi: 10.1016/j.neuron.2007.06.002
- Brager, D. H., Cai, X., and Thompson, S. M. (2003). Activity-dependent activation of presynaptic protein kinase C mediates post-tetanic potentiation. *Nat. Neurosci.* 6, 551–552. doi: 10.1038/nn1067
- Brager, D. H., Capogna, M., and Thompson, S. M. (2002). Short-term synaptic plasticity, simulation of nerve terminal dynamics, and the effects of protein kinase C activation in rat hippocampus. *J. Physiol.* 541, 545–559. doi: 10.1113/jphysiol.2001.015842
- Brose, N., Petrenko, A. G., Südhof, T. C., and Jahn, R. (1992). Synaptotagmin: a calcium sensor on the synaptic vesicle surface. *Science* 256, 1021–1025. doi: 10.1126/science.1589771
- Burgoyne, R. D. (2007). Neuronal calcium sensor proteins: generating diversity in neuronal Ca^{2+} signalling. *Nat. Rev. Neurosci.* 8, 182–193. doi: 10.1038/nrn2093
- Catterall, W. A., and Few, A. P. (2008). Calcium channel regulation and presynaptic plasticity. *Neuron* 59, 882–901. doi: 10.1016/j.neuron.2008.09.005
- Catterall, W. A., Leal, K., and Nanou, E. (2013). Calcium channels and short-term synaptic plasticity. *J. Biol. Chem.* 288, 10742–10749. doi: 10.1074/jbc.R112.411645
- Chu, Y., Fioravante, D., Leitges, M., and Regehr, W. G. (2014). Calcium-dependent PKC isoforms have specialized roles in short-term synaptic plasticity. *Neuron* 82, 859–871. doi: 10.1016/j.neuron.2014.04.003
- Coudeville, N., Montaville, P., Leonov, A., Zweckstetter, M., and Becker, S. (2008). Structural determinants for Ca^{2+} and phosphatidylinositol 4,5-bisphosphate binding by the C2A domain of rabphilin-3A. *J. Biol. Chem.* 283, 35918–35928. doi: 10.1074/jbc.M804094200
- Dason, J. S., Romero-Pozuelo, J., Atwood, H. L., and Ferrús, A. (2012). Multiple roles for Frequentin/NCS-1 in synaptic function and development. *Mol. Neurobiol.* 45, 388–402. doi: 10.1007/s12035-012-8250-4
- Deák, F., Shin, O.-H., Tang, J., Hanson, P., Ubach, J., Jahn, R., et al. (2006). Rabphilin regulates SNARE-dependent re-priming of synaptic vesicles for fusion. *EMBO J.* 25, 2856–2866. doi: 10.1038/sj.emboj.7601165
- de Jong, A. P. H., and Verhage, M. (2009). Presynaptic signal transduction pathways that modulate synaptic transmission. *Curr. Opin. Neurobiol.* 19, 245–253. doi: 10.1016/j.conb.2009.06.005
- Delaney, K. R., Zucker, R. S., and Tank, D. W. (1989). Calcium in motor nerve terminals associated with posttetanic potentiation. *J. Neurosci.* 9, 3558–3567.
- DeMaria, C. D., Soong, T. W., Alseikhan, B. A., Alvania, R. S., and Yue, D. T. (2001). Calmodulin bifurcates the local Ca^{2+} signal that modulates P/Q-type Ca^{2+} channels. *Nature* 411, 484–489. doi: 10.1038/35078091
- Dimova, K., Kawabe, H., Betz, A., Brose, N., and Jahn, O. (2006). Characterization of the Munc13-calmodulin interaction by photoaffinity labeling. *Biochim. Biophys. Acta* 1763, 1256–1265. doi: 10.1016/j.bbamer.2006.09.017
- Dittman, J. S., and Regehr, W. G. (1998). Calcium dependence and recovery kinetics of presynaptic depression at the climbing fiber to Purkinje cell synapse. *J. Neurosci.* 18, 6147–6162.
- Dutta Roy, R., Stefan, M. I., and Rosenmund, C. (2014). Biophysical properties of presynaptic short-term plasticity in hippocampal neurons: insights from electrophysiology, imaging and mechanistic models. *Front. Cell. Neurosci.* 8:141. doi: 10.3389/fncel.2014.00141
- Eliot, L. S., Kandel, E. R., and Hawkins, R. D. (1994). Modulation of spontaneous transmitter release during depression and posttetanic potentiation of *Aplysia* sensory-motor neuron synapses isolated in culture. *J. Neurosci.* 14, 3280–3292.
- Elmqvist, D., and Quastel, D. M. (1965). A quantitative study of end-plate potentials in isolated human muscle. *J. Physiol.* 178, 505–529.

- Felmy, F., Neher, E., and Schneggenburger, R. (2003). Probing the intracellular calcium sensitivity of transmitter release during synaptic facilitation. *Neuron* 37, 801–811. doi: 10.1016/s0896-6273(03)00085-0
- Fernandez, I., Araç, D., Ubach, J., Gerber, S. H., Shin, O., Gao, Y., et al. (2001). Three-dimensional structure of the synaptotagmin 1 C2B-domain: synaptotagmin 1 as a phospholipid binding machine. *Neuron* 32, 1057–1069. doi: 10.1016/s0896-6273(01)00548-7
- Few, A. P., Lautermilch, N. J., Westenbroek, R. E., Scheuer, T., and Catterall, W. A. (2005). Differential regulation of CaV2.1 channels by calcium-binding protein 1 and visinin-like protein-2 requires N-terminal myristoylation. *J. Neurosci.* 25, 7071–7080. doi: 10.1523/jneurosci.0452-05.2005
- Fioravante, D., Chu, Y., de Jong, A. P., Leitges, M., Kaeser, P. S., and Regehr, W. G. (2014). Protein kinase C is a calcium sensor for presynaptic short-term plasticity. *Elife* 3:e03011. doi: 10.7554/elifelife.03011
- Fioravante, D., Chu, Y., Myoga, M. H., Leitges, M., and Regehr, W. G. (2011). Calcium-dependent isoforms of protein kinase C mediate posttetanic potentiation at the calyx of held. *Neuron* 70, 1005–1019. doi: 10.1016/j.neuron.2011.04.019
- Fioravante, D., Myoga, M. H., Leitges, M., and Regehr, W. G. (2012). Adaptive regulation maintains posttetanic potentiation at cerebellar granule cell synapses in the absence of calcium-dependent PKC. *J. Neurosci.* 32, 13004–13009. doi: 10.1523/jneurosci.0683-12.2012
- Fioravante, D., and Regehr, W. G. (2011). Short-term forms of presynaptic plasticity. *Curr. Opin. Neurobiol.* 21, 269–274. doi: 10.1016/j.conb.2011.02.003
- Forsythe, I. D., Tsujimoto, T., Barnes-Davies, M., Cuttle, M. F., and Takahashi, T. (1998). Inactivation of presynaptic calcium current contributes to synaptic depression at a fast central synapse. *Neuron* 20, 797–807. doi: 10.1016/s0896-6273(00)81017-x
- Fuhrmann, G., Segev, I., Markram, H., and Tsodyks, M. (2002). Coding of temporal information by activity-dependent synapses. *J. Neurophysiol.* 87, 140–148.
- Genc, O., Kochubey, O., Toonen, R. F., Verhage, M., and Schneggenburger, R. (2014). Munc18–1 is a dynamically regulated PKC target during short-term enhancement of transmitter release. *Elife* 3:e01715. doi: 10.7554/elifelife.01715
- Grobler, J. A., and Hurley, J. H. (1998). Catalysis by phospholipase C delta1 requires that Ca²⁺ bind to the catalytic domain, but not the C2 domain. *Biochemistry* 37, 5020–5028. doi: 10.1021/bi972952w
- Groffen, A. J., Martens, S., Diez Arazola, R., Cornelisse, L. N., Lozovaya, N., de Jong, A. P. H., et al. (2010). Doc2b is a high-affinity Ca²⁺ sensor for spontaneous neurotransmitter release. *Science* 327, 1614–1618. doi: 10.1126/science.1183765
- Habets, R. L. P., and Borst, J. G. G. (2005). Post-tetanic potentiation in the rat calyx of Held synapse. *J. Physiol.* 564, 173–187. doi: 10.1113/jphysiol.2004.079160
- He, L., Xue, L., Xu, J., McNeil, B. D., Bai, L., Melicoff, E., et al. (2009). Compound vesicle fusion increases quantal size and potentiates synaptic transmission. *Nature* 459, 93–97. doi: 10.1038/nature07860
- Hennig, M. H. (2013). Theoretical models of synaptic short term plasticity. *Front. Comput. Neurosci.* 7:45. doi: 10.3389/fncom.2013.00154
- Hosoi, N., Holt, M., and Sakaba, T. (2009). Calcium dependence of exo- and endocytotic coupling at a glutamatergic synapse. *Neuron* 63, 216–229. doi: 10.1016/j.neuron.2009.06.010
- Hosoi, N., Sakaba, T., and Neher, E. (2007). Quantitative analysis of calcium-dependent vesicle recruitment and its functional role at the calyx of Held synapse. *J. Neurosci.* 27, 14286–14298. doi: 10.1523/jneurosci.4122-07.2007
- Junge, H. J., Rhee, J.-S., Jahn, O., Varoqueaux, F., Spiess, J., Waxham, M. N., et al. (2004). Calmodulin and Munc13 form a Ca²⁺ sensor/effector complex that controls short-term synaptic plasticity. *Cell* 118, 389–401. doi: 10.1016/j.cell.2004.06.029
- Kalkstein, J. M., and Magleby, K. L. (2004). Augmentation increases vesicular release probability in the presence of masking depression at the Frog neuromuscular junction. *J. Neurosci.* 24, 11391–11403. doi: 10.1523/jneurosci.2756-04.2004
- Kamiya, H., and Zucker, R. S. (1994). Residual Ca²⁺ and short-term synaptic plasticity. *Nature* 371, 603–606. doi: 10.1038/371603a0
- Katz, B., and Miledi, R. (1968). The role of calcium in neuromuscular facilitation. *J. Physiol.* 195, 481–492.
- Kochubey, O., Lou, X., and Schneggenburger, R. (2011). Regulation of transmitter release by Ca²⁺ and synaptotagmin: insights from a large CNS synapse. *Trends Neurosci.* 34, 237–246. doi: 10.1016/j.tins.2011.02.006
- Kochubey, O., and Schneggenburger, R. (2011). Synaptotagmin increases the dynamic range of synapses by driving Ca²⁺-evoked release and by clamping a near-linear remaining Ca²⁺ sensor. *Neuron* 69, 736–748. doi: 10.1016/j.neuron.2011.01.013
- Korogod, N., Lou, X., and Schneggenburger, R. (2005). Presynaptic Ca²⁺ requirements and developmental regulation of posttetanic potentiation at the calyx of held. *J. Neurosci.* 25, 5127–5137. doi: 10.1523/jneurosci.1295-05.2005
- Korogod, N., Lou, X., and Schneggenburger, R. (2007). Posttetanic potentiation critically depends on an enhanced Ca²⁺ sensitivity of vesicle fusion mediated by presynaptic PKC. *Proc. Natl. Acad. Sci. U S A* 104, 15923–15928. doi: 10.1073/pnas.0704603104
- Lautermilch, N. J., Few, A. P., Scheuer, T., and Catterall, W. A. (2005). Modulation of CaV2.1 channels by the neuronal calcium-binding protein visinin-like protein-2. *J. Neurosci.* 25, 7062–7070. doi: 10.1523/jneurosci.0447-05.2005
- Leal, K., Mochida, S., Scheuer, T., and Catterall, W. A. (2012). Fine-tuning synaptic plasticity by modulation of Ca(V)2.1 channels with Ca²⁺ sensor proteins. *Proc. Natl. Acad. Sci. U S A* 109, 17069–17074. doi: 10.1073/pnas.1215172109
- Lee, J. S., Kim, M.-H., Ho, W.-K., and Lee, S.-H. (2008). Presynaptic release probability and readily releasable pool size are regulated by two independent mechanisms during posttetanic potentiation at the calyx of held synapse. *J. Neurosci.* 28, 7945–7953. doi: 10.1523/JNEUROSCI.2165-08.2008
- Lee, A., Wong, S., Gallagher, D., Li, B., Storm, D., Scheuer, T., et al. (1999). Ca²⁺/calmodulin binds to and modulates P/Q-type calcium channels. *Nature* 399, 155–159. doi: 10.1038/20194
- Lewit-Bentley, A., and Réty, S. (2000). EF-hand calcium-binding proteins. *Curr. Opin. Struct. Biol.* 10, 637–643. doi: 10.1016/S0959-440X(00)00142-1
- Li, C., Pan, W., Braunewell, K. H., and Ames, J. B. (2011). Structural analysis of Mg²⁺ and Ca²⁺ binding, myristoylation and dimerization of the neuronal calcium sensor and visinin-like protein 1 (VILIP-1). *J. Biol. Chem.* 286, 6354–6366. doi: 10.1074/jbc.M110.173724
- Lipstein, N., Sakaba, T., Cooper, B. H., Lin, K.-H., Strenzke, N., Ashery, U., et al. (2013). Dynamic control of synaptic vesicle replenishment and short-term plasticity by Ca(2+)-calmodulin-Munc13-1 signaling. *Neuron* 79, 82–96. doi: 10.1016/j.neuron.2013.05.011
- Lipstein, N., Schaks, S., Dimova, K., Kalkhof, S., Ihling, C., Kölbl, K., et al. (2012). Nonconserved Ca(2+)/calmodulin binding sites in Munc13s differentially control synaptic short-term plasticity. *Mol. Cell. Biol.* 32, 4628–4641. doi: 10.1128/MCB.00933-12
- Liu, H., Bai, H., Hui, E., Yang, L., Evans, C. S., Wang, Z., et al. (2014). Synaptotagmin 7 functions as a Ca²⁺-sensor for synaptic vesicle replenishment. *Elife* 3:e01524. doi: 10.7554/elifelife.01524
- Liu, G., and Tsien, R. W. (1995). Properties of synaptic transmission at single hippocampal synaptic boutons. *Nature* 375, 404–408. doi: 10.1038/375404a0
- Ma, C., Li, W., Xu, Y., and Rizo, J. (2011). Munc13 mediates the transition from the closed syntaxin-Munc18 complex to the SNARE complex. *Nat. Struct. Mol. Biol.* 18, 542–549. doi: 10.1038/nsmb.2047
- Magleby, K. L. (1973). The effect of repetitive stimulation on facilitation of transmitter release at the frog neuromuscular junction. *J. Physiol.* 234, 327–352.
- Magleby, K. L. (1979). Facilitation, augmentation and potentiation of transmitter release. *Prog. Brain Res.* 49, 175–182. doi: 10.1016/s0079-6123(08)64631-2
- Magleby, K. L., and Zengel, J. E. (1976a). Augmentation: a process that acts to increase transmitter release at the frog neuromuscular junction. *J. Physiol.* 257, 449–470.
- Magleby, K. L., and Zengel, J. E. (1976b). Long term changes in augmentation, potentiation and depression of transmitter release as a function of repeated synaptic activity at the frog neuromuscular junction. *J. Physiol.* 257, 471–494.
- Maune, J. F., Klee, C. B., and Beckingham, K. (1992). Ca²⁺ binding and conformational change in two series of point mutations to the individual Ca(2+)-binding sites of calmodulin. *J. Biol. Chem.* 267, 5286–5295.
- Maximov, A., Lao, Y., Li, H., Chen, X., Rizo, J., Sorensen, J. B., et al. (2008). Genetic analysis of synaptotagmin-7 function in synaptic vesicle exocytosis. *Proc. Natl. Acad. Sci. U S A* 105, 3986–3991. doi: 10.1073/pnas.0712372105
- Mochida, S., Few, A. P., Scheuer, T., and Catterall, W. A. (2008). Regulation of Presynaptic CaV2.1 Channels by Ca²⁺ sensor proteins mediates short-term synaptic plasticity. *Neuron* 57, 210–216. doi: 10.1016/j.neuron.2007.11.036
- Mochida, S., Westenbroek, R. E., Yokoyama, C. T., Itoh, K., and Catterall, W. A. (2003). Subtype-selective reconstitution of synaptic transmission in sympathetic

- ganglion neurons by expression of exogenous calcium channels. *Proc. Natl. Acad. Sci. U S A* 100, 2813–2818. doi: 10.1073/pnas.262787299
- Müller, M., Felmy, F., and Schneggenburger, R. (2008). A limited contribution of Ca^{2+} current facilitation to paired-pulse facilitation of transmitter release at the rat calyx of held. *J. Physiol.* 586, 5503–5520. doi: 10.1113/jphysiol.2008.155838
- Müller, M., Felmy, F., Schwaller, B., and Schneggenburger, R. (2007). Parvalbumin is a mobile presynaptic Ca^{2+} buffer in the calyx of held that accelerates the decay of Ca^{2+} and short-term facilitation. *J. Neurosci.* 27, 2261–2271. doi: 10.1523/jneurosci.5582-06.2007
- Newton, A. C. (2010). Protein kinase C: poised to signal. *Am. J. Physiol. Endocrinol. Metab.* 298, E395–E402. doi: 10.1152/ajpendo.00477.2009
- Nishiki, T., and Augustine, G. J. (2004). Dual roles of the C2B domain of synaptotagmin I in synchronizing Ca^{2+} -dependent neurotransmitter release. *J. Neurosci.* 24, 8542–8550. doi: 10.1523/jneurosci.2545-04.2004
- O'Callaghan, D. W., Ivings, L., Weiss, J. L., Ashby, M. C., Tepikin, A. V., and Burgoyne, R. D. (2002). Differential use of Myristoyl groups on neuronal calcium sensor proteins as a determinant of spatio-temporal aspects of Ca^{2+} signal transduction. *J. Biol. Chem.* 277, 14227–14237. doi: 10.1074/jbc.M111750200
- Olwin, B. B., and Storm, D. R. (1985). Calcium binding to complexes of calmodulin and calmodulin binding proteins. *Biochemistry* 24, 8081–8086. doi: 10.1021/bi00348a037
- Pang, Z. P., Bacaj, T., Yang, X., Zhou, P., Xu, W., and Südhof, T. C. (2011). Doc2 supports spontaneous synaptic transmission by a Ca^{2+} -independent mechanism. *Neuron* 70, 244–251. doi: 10.1016/j.neuron.2011.03.011
- Pang, Z. P., and Südhof, T. C. (2010). Cell biology of Ca^{2+} -triggered exocytosis. *Curr. Opin. Cell Biol.* 22, 496–505. doi: 10.1016/jceb.2010.05.001
- Pang, Z. P., Xu, W., Cao, P., and Südhof, T. C. (2010). Calmodulin suppresses Synaptotagmin-2 transcription in cortical neurons. *J. Biol. Chem.* 285, 33930–33939. doi: 10.1074/jbc.M110.150151
- Pappa, H., Murray-Rust, J., Dekker, L. V., Parker, P. J., and McDonald, N. Q. (1998). Crystal structure of the C2 domain from protein kinase C-delta. *Structure* 6, 885–894. doi: 10.1016/s0969-2126(98)00090-2
- Pfister, J.-P., Dayan, P., and Lengyel, M. (2010). Synapses with short-term plasticity are optimal estimators of presynaptic membrane potentials. *Nat. Neurosci.* 13, 1271–1275. doi: 10.1038/nn.2640
- Rebecchi, M. J., and Pentylala, S. N. (2000). Structure, function and control of phosphoinositide-specific Phospholipase C. *Physiol. Rev.* 80, 1291–1335.
- Regehr, W. G. (2012). Short-term presynaptic plasticity. *Cold Spring Harb. Perspect. Biol.* 4:a005702. doi: 10.1101/cshperspect.a005702
- Regehr, W. G., Delaney, K. R., and Tank, D. W. (1994). The role of presynaptic calcium in short-term enhancement at the hippocampal mossy fiber synapse. *J. Neurosci.* 14, 523–537.
- Rizo, J., and Südhof, T. C. (1998). C2-domains, structure and function of a universal Ca^{2+} -binding domain. *J. Biol. Chem.* 273, 15879–15882. doi: 10.1074/jbc.273.26.15879
- Rohrbough, J., and Broadie, K. (2005). Lipid regulation of the synaptic vesicle cycle. *Nat. Rev. Neurosci.* 6, 139–150. doi: 10.1038/nrn1608
- Rosenmund, C., Sigler, A., Augustin, I., Reim, K., Brose, N., and Rhee, J.-S. (2002). Differential control of vesicle priming and short-term plasticity by Munc13 isoforms. *Neuron* 33, 411–424. doi: 10.1016/s0896-6273(02)00568-8
- Sakaba, T., and Neher, E. (2001). Calmodulin mediates rapid recruitment of fast-releasing synaptic vesicles at a calyx-type synapse. *Neuron* 32, 1119–1131. doi: 10.1016/s0896-6273(01)00543-8
- Saraswati, S., Adolfsen, B., and Littleton, J. T. (2007). Characterization of the role of the Synaptotagmin family as calcium sensors in facilitation and asynchronous neurotransmitter release. *Proc. Natl. Acad. Sci. U S A* 104, 14122–14127. doi: 10.1073/pnas.0706711104
- Schaub, M. C., and Heizmann, C. W. (2008). Calcium, troponin, calmodulin, S100 proteins: from myocardial basics to new therapeutic strategies. *Biochem. Biophys. Res. Commun.* 369, 247–264. doi: 10.1016/j.bbrc.2007.10.082
- Schneggenburger, R., Sakaba, T., and Neher, E. (2002). Vesicle pools and short-term synaptic depression: lessons from a large synapse. *Trends Neurosci.* 25, 206–212. doi: 10.1016/s0166-2236(02)02139-2
- Schwaller, B. (2009). The continuing disappearance of “pure” Ca^{2+} buffers. *Cell. Mol. Life Sci.* 66, 275–300. doi: 10.1007/s00018-008-8564-6
- Scullin, C. S., and Partridge, L. D. (2010). Contributions of SERCA pump and ryanodine-sensitive stores to presynaptic residual Ca^{2+} . *Cell Calcium* 47, 326–338. doi: 10.1016/j.ceca.2010.01.004
- Shao, X., Davletov, B. A., Sutton, R. B., Südhof, T. C., and Rizo, J. (1996). Bipartite Ca^{2+} -binding motif in C2 domains of synaptotagmin and protein kinase C. *Science* 273, 248–251. doi: 10.1126/science.273.5272.248
- Shin, O.-H., Lu, J., Rhee, J.-S., Tomchick, D. R., Pang, Z. P., Wojcik, S. M., et al. (2010). Munc13 C2B domain is an activity-dependent Ca^{2+} regulator of synaptic exocytosis. *Nat. Struct. Mol. Biol.* 17, 280–288. doi: 10.1038/nsmb.1758
- Shin, O.-H., Xu, J., Rizo, J., and Südhof, T. C. (2009). Differential but convergent functions of Ca^{2+} binding to synaptotagmin-1 C2 domains mediate neurotransmitter release. *Proc. Natl. Acad. Sci. U S A* 106, 16469–16474. doi: 10.1073/pnas.0908798106
- Sippy, T., Cruz-Martín, A., Jeromin, A., and Schweizer, F. E. (2003). Acute changes in short-term plasticity at synapses with elevated levels of neuronal calcium sensor-1. *Nat. Neurosci.* 6, 1031–1038. doi: 10.1038/nn1117
- Skelton, N. J., Kördel, J., Akke, M., Forsén, S., and Chazin, W. J. (1994). Signal transduction versus buffering activity in Ca^{2+} -binding proteins. *Nat. Struct. Biol.* 1, 239–245. doi: 10.1038/nsb0494-239
- Stevens, C. F., and Wesseling, J. F. (1998). Activity-dependent modulation of the rate at which synaptic vesicles become available to undergo exocytosis. *Neuron* 21, 415–424. doi: 10.1016/s0896-6273(00)80550-4
- Stevens, C. F., and Wesseling, J. F. (1999). Augmentation is a potentiation of the Exocytotic process. *Neuron* 22, 139–146. doi: 10.1016/s0896-6273(00)80685-6
- Südhof, T. C. (2013). Neurotransmitter release: the last millisecond in the life of a synaptic vesicle. *Neuron* 80, 675–690. doi: 10.1016/j.neuron.2013.10.022
- Sugita, S., Shin, O.-H., Han, W., Lao, Y., and Südhof, T. C. (2002). Synaptotagmins form a hierarchy of exocytotic Ca^{2+} sensors with distinct Ca^{2+} affinities. *EMBO J.* 21, 270–280. doi: 10.1093/emboj/21.3.270
- Torreillas, A., Laynez, J., Menéndez, M., Corbalán-García, S., and Gómez-Fernández, J. C. (2004). Calorimetric study of the interaction of the C2 domains of classical protein kinase C isoenzymes with Ca^{2+} and phospholipids. *Biochemistry* 43, 11727–11739. doi: 10.1021/bi0489659
- Tsodyks, M., Pawelzik, K., and Markram, H. (1998). Neural networks with dynamic synapses. *Neural Comput.* 10, 821–835. doi: 10.1162/089976698300017502
- Tsujimoto, T., Jeromin, A., Saitoh, N., Roder, J. C., and Takahashi, T. (2002). Neuronal calcium sensor 1 and activity-dependent facilitation of P/Q-type calcium currents at presynaptic nerve terminals. *Science* 295, 2276–2279. doi: 10.1126/science.1068278
- Ubach, J., García, J., Nittler, M. P., Südhof, T. C., and Rizo, J. (1999). Structure of the Janus-faced C2B domain of rabphilin. *Nat. Cell Biol.* 1, 106–112. doi: 10.1038/10076
- Ubach, J., Zhang, X., Shao, X., Südhof, T. C., and Rizo, J. (1998). Ca^{2+} binding to synaptotagmin: how many Ca^{2+} ions bind to the tip of a C2-domain? *EMBO J.* 17, 3921–3930. doi: 10.1093/emboj/17.14.3921
- van den Bogaart, G., Meyenberg, K., Diederichsen, U., and Jahn, R. (2012). Phosphatidylinositol 4,5-bisphosphate increases Ca^{2+} affinity of synaptotagmin-1 by 40-fold. *J. Biol. Chem.* 287, 16447–16453. doi: 10.1074/jbc.M112.343418
- von Gersdorff, H., and Matthews, G. (1997). Depletion and Replenishment of vesicle pools at a ribbon-type synaptic terminal. *J. Neurosci.* 17, 1919–1927.
- Wang, L. Y., and Kaczmarek, L. K. (1998). High-frequency firing helps replenish the readily releasable pool of synaptic vesicles. *Nature* 394, 384–388. doi: 10.1038/28645
- Wen, H., Linhoff, M. W., McGinley, M. J., Li, G.-L., Corson, G. M., Mandel, G., et al. (2010). Distinct roles for two synaptotagmin isoforms in synchronous and asynchronous transmitter release at zebrafish neuromuscular junction. *Proc. Natl. Acad. Sci. U S A* 107, 13906–13911. doi: 10.1073/pnas.1008598107
- Wierda, K. D. B., Toonen, R. F. G., de Wit, H., Brussaard, A. B., and Verhage, M. (2007). Interdependence of PKC-dependent and PKC-independent pathways for presynaptic plasticity. *Neuron* 54, 275–290. doi: 10.1016/j.neuron.2007.04.001
- Wilkinson, R. S., and Lin, M. Y. (2004). Endocytosis and synaptic plasticity: might the tail wag the dog? *Trends Neurosci.* 27, 171–174. doi: 10.1016/j.tins.2004.01.011
- Xia, Z., and Storm, D. R. (2005). The role of calmodulin as a signal integrator for synaptic plasticity. *Nat. Rev. Neurosci.* 6, 267–276. doi: 10.1038/nrn1647

- Xu, J., and Wu, L.-G. (2005). The decrease in the presynaptic calcium current is a major cause of short-term depression at a calyx-type synapse. *Neuron* 46, 633–645. doi: 10.1016/j.neuron.2005.03.024
- Xue, L., and Wu, L.-G. (2010). Post-tetanic potentiation is caused by two signalling mechanisms affecting quantal size and quantal content. *J. Physiol.* 588, 4987–4994. doi: 10.1113/jphysiol.2010.196964
- Yamaguchi, T., Shirataki, H., Kishida, S., Miyazaki, M., Nishikawa, J., Wada, K., et al. (1993). Two functionally different domains of rabphilin-3A, Rab3A p25/smg p25A-binding and phospholipid- and Ca(2+)-binding domains. *J. Biol. Chem.* 268, 27164–27170.
- Zengel, J. E., and Magleby, K. L. (1981). Changes in miniature endplate potential frequency during repetitive nerve stimulation in the presence of Ca^{2+} , Ba^{2+} and Sr^{2+} at the frog neuromuscular junction. *J. Gen. Physiol.* 77, 503–529. doi: 10.1085/jgp.77.5.503
- Zucker, R. S., and Lara-Estrella, L. O. (1983). Post-tetanic decay of evoked and spontaneous transmitter release and a residual-calcium model of synaptic facilitation at crayfish neuromuscular junctions. *J. Gen. Physiol.* 81, 355–372. doi: 10.1085/jgp.81.3.355
- Zucker, R. S., and Regehr, W. G. (2002). Short-term synaptic plasticity. *Annu. Rev. Physiol.* 64, 355–405. doi: 10.1146/annurev.physiol.64.092501.114547
- Zucker, R. S., and Stockbridge, N. (1983). Presynaptic calcium diffusion and the time courses of transmitter release and synaptic facilitation at the squid giant synapse. *J. Neurosci.* 3, 1263–1269.
- Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.
- Received: 18 September 2014; accepted: 09 October 2014; published online: 29 October 2014.
- Citation: de Jong APH and Fioravante D (2014) Translating neuronal activity at the synapse: presynaptic calcium sensors in short-term plasticity. *Front. Cell. Neurosci.* 8:356. doi: 10.3389/fncel.2014.00356
- This article was submitted to the journal *Frontiers in Cellular Neuroscience*.
- Copyright © 2014 de Jong and Fioravante. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.