



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Pathology of rituximab-induced Kaposi sarcoma flare: role of B-cell depletion

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Pantanowitz, L, K Früh, D Aboulafia, S Marconi, and BJ Dezube. 2009. "Pathology of rituximab-induced Kaposi sarcoma flare: role of B-cell depletion." <i>Infectious Agents and Cancer</i> 4 (Suppl 2): P33. doi:10.1186/1750-9378-4-S2-P33. http://dx.doi.org/10.1186/1750-9378-4-S2-P33 .
Published Version	doi:10.1186/1750-9378-4-S2-P33
Accessed	February 17, 2015 8:33:36 AM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:13581177
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

Poster presentation

Open Access

Pathology of rituximab-induced Kaposi sarcoma flare: role of B-cell depletion

L Pantanowitz*¹, K Früh², D Aboulafia³, S Marconi¹ and BJ Dezube⁴

Address: ¹Department of Pathology, Baystate Medical Center, Tufts University School of Medicine, Springfield, Massachusetts, USA, ²Vaccine and Gene Therapy Institute, Oregon Health and Science University, Portland, Oregon, USA, ³Virginia Mason Medical Center, Seattle, Washington, USA and ⁴Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, USA

* Corresponding author

from 11th International Conference on Malignancies in AIDS and Other Acquired Immunodeficiencies (ICMAOI): Basic, Epidemiologic, and Clinical Research
Bethesda, MD, USA. 6–7 October 2008

Published: 17 June 2009

Infectious Agents and Cancer 2009, **4**(Suppl 2):P33 doi:10.1186/1750-9378-4-S2-P33

This abstract is available from: <http://www.infectagentscancer.com/content/4/S2/P33>

© 2009 Pantanowitz et al; licensee BioMed Central Ltd.

Background

Kaposi sarcoma (KS) flare (exacerbation) may occur following therapy with corticosteroids, as part of the immune reconstitution inflammatory syndrome seen with highly active antiretroviral therapy (HAART), and after rituximab therapy. The exact mechanism responsible for iatrogenic KS flare is unclear. Therefore, the aim of this study was to investigate the pathologic features in cases of AIDS-associated KS flare.

Methods

Two cases of AIDS-associated cutaneous KS flare were identified following rituximab therapy. Tissue biopsies obtained from these KS flare lesions were compared to similar controls (AIDS-KS samples of similar stage from patients who had not received rituximab) by means of immunohistochemistry using vascular makers (CD34, CD31), monoclonal antibodies to human Herpesvirus-8 (HHV-8) gene products (LNA-1, K5), as well as B-lymphocyte (CD20) and T-lymphocyte (CD3, CD4, CD8) markers.

Results

One patient was a 36-year-old male, HIV-positive for 2 years (CD4 T-lymphocyte count 363 cells/mm³; HIV viral load >100,000 copies/ml), and the other a 44-year-old male, HIV-positive for 5 years (CD4 T-lymphocyte count 443 cells/mm³; HIV viral load <75 copies/ml), who had both received rituximab therapy for multicentric Castle-

man disease. The former manifested with cutaneous KS after 7 days, and the later patient with skin and lymphadenopathic KS flares within 3 months of receiving rituximab. In the control cases both CD3 and CD20 cells were present, with a preponderance of T-lymphocytes identified. In the KS flare specimens, there were only T-cells present with a notable absence of B-lymphocytes. In all control and flare KS cases T-cell subsets showed CD8 > CD4. CD20+ B-cell depletion in the KS flare case of the 36-year-old patient occurred concomitantly with activation of the HHV-8 immediate early gene protein K5. KS flares in both patients were successfully treated with liposomal doxorubicin and valganciclovir.

Conclusion

Rituximab-induced KS flare appears to be related to a modification in the host immune system, most notably with complete absence of B lymphocytes in the KS lesion. B-cell depletion with rituximab in patients at risk for KS development may activate HHV-8, resulting in unwanted KS flare. Effective management of iatrogenic KS flare therefore depends upon the control of HHV-8 viremia in conjunction with specific chemotherapy for KS.