



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Hearty miR-363 controls HAND1 in cardiac cell specification

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Gupta, Manoj K., and Tata Nageswara Rao. 2014. "Hearty miR-363 controls HAND1 in cardiac cell specification." <i>Stem Cell Research & Therapy</i> 5 (4): 89. doi:10.1186/scrt478. http://dx.doi.org/10.1186/scrt478 .
Published Version	doi:10.1186/scrt478
Accessed	February 17, 2015 12:46:32 PM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:13890805
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

COMMENTARY

Hearty miR-363 controls *HAND1* in cardiac cell specification

Manoj K Gupta^{1*} and Tata Nageswara Rao^{1,2}

See related research by Wagh *et al.*, <http://stemcellres.com/content/5/3/75>

Abstract

MicroRNAs regulate target gene expression post-transcriptionally in a myriad of cell types and play critical roles in diverse physiological and pathological processes, including cardiomyocyte development, differentiation, and regeneration. The recent publication in *Stem Cell Research and Therapy* by Wagh and colleagues reports a novel regulatory role for miR-363 in cardiomyocyte specification. By employing microRNA expression profiling and functional knockdown studies on human embryonic stem cell-derived cardiomyocytes, the authors identified miR-363 as an upstream negative regulator of left ventricular specification transcription factor *HAND1*.

The efficient derivation of specialized and functional cardiomyocytes (CMs) from pluripotent stem cells is a primary goal for stem cell-based cardiac regenerative therapies. Such a prospect is currently hampered, however, in part by an incompletely defined complex of molecular regulators of cardiac cell development and differentiation.

This commentary discusses the findings from Wagh and colleagues published in this issue of *Stem Cell Research and Therapy* demonstrating a critical role for miR-363 in post-transcriptional regulation of CM differentiation via the hand and neural crest derivative expressed *HAND1* transcription factor [1]. Teleost fishes and amphibians show robust regeneration abilities of heart tissues, whereas the adult mammalian heart poorly renews itself, showing extremely limited regenerative capacity. CM proliferative capacity rapidly ceases shortly after birth in mammals [2]. There is little evidence for

cardiac cell renewal in humans [3], indicating the limited proliferation capacity of adult CMs. During myocardium infarction or heart failure, the extensive loss of CMs leads to progressive cardiac dysfunction and chronic heart failure, a leading cause of morbidity and mortality in the developed world. Identifying molecular regulators and critical mediators of cardiac cell type development, proliferation, and differentiation is of great clinical importance, and unraveling such molecular horizons could lead to the development of therapeutic strategies for successful regeneration of the human adult heart.

MicroRNAs (miRNAs) are small noncoding RNAs (~22 nucleotides in length) that regulate gene expression post-transcriptionally by imperfect binding to the 3' untranslated region of target mRNAs in a wide variety of cell types. Given the notion that a single miRNA may have multiple cellular targets and given the existence of vast numbers of miRNAs (~1,500 in humans), we can expect to witness the discovery of novel miRNA-dependent regulation in the modulation of versatile biological functions [4]. Recently, miRNAs have been recognized as important players in cardiac development, pathology, and regeneration. Disruption of the miRNA pathway in CMs leads to heart failure and cardiomyopathy [5]. Some miRNAs have been demonstrated to suppress CM proliferation, including miR-1, miR-133, and the miR-15 family, while other miRNAs (miR-199a and miR-590) have been shown to promote *in vivo* CM proliferation in rat and mouse models [6,7]. miRNAs could thus be the best targets for understanding cardiac specialization during differentiation of human embryonic stem cells (hESCs). Further exploring the role of miRNAs in cardiac cells during development and disease may therefore hold great promise for cardiac therapy applications.

hESCs are a unique source for deriving cardiac cells in a specialized manner and also offer a unique opportunity to study the mechanisms and derivation of different

* Correspondence: manoj.gupta@joslin.harvard.edu

¹Section of Islet Cell and Regenerative Biology, Joslin Diabetes Center, Harvard Medical School, Boston, MA 01125, USA

Full list of author information is available at the end of the article

cardiac cell types *in vitro* [8,9]. The recent publication by Wagh and colleagues sought to identify the novel miRNAs regulating cardiac-specific transcription factors that determine left versus right ventricular determination [1]. To this end, the authors performed miRNA expression profiling in undifferentiated hESCs and CMs at day 8 and day 14 after differentiation. This screening identified 18 differentially expressed candidate miRNAs, from which they focused on miRNAs that could potentially target the 3' untranslated regions (as predicted by *in silico* analysis) of the CM subtype specification transcription factors (*HAND1*, *HAND2*, *TBX3*, *GJA1*, *NPPA*, *RYR2*, *SLN*), and found that a subset of miRNAs (miR-363, miR-367, miR-181a and miR-181c) was expressed at a higher level in hESC-derived non-CMs compared with CMs. These novel findings were confirmed by real-time polymerase chain reaction analysis: higher expression of *HAND* genes correlated with lower expression of miR-363, miR-367, miR-181a and miR-181c. Using an elegant dual-reporter assay, the authors further showed that miR-363 specifically inhibits *HAND1* gene expression through 3' untranslated region binding. Perhaps more heartening, overexpression of antisense oligonucleotide directed against miR-363 (anti-miR-363) during hESC differentiation promoted more *HAND1*-expressing CMs, confirming the role of miR-363 in dorso-ventral patterning and interventricular septum formation in the embryonic heart [10,11] (Figure 1).

In summary, Wagh and colleagues demonstrated for the first time that miR-363 acts as an upstream negative regulator of *HAND1* transcription factor and extends the role of miRNAs in cardiac cell biology (Figure 1). This study has important implications for the human clinical perspective. For instance, since it is especially of

great interest to derive specific cardiac subtypes from embryonic stem cell/induced pluripotent stem cell cultures and ventricular heart cells, which are the most affected cells during cardiac injury, the reversible pharmacological suppression of miR-363 may provide a novel strategy for generating functional left ventricular CMs for cardiac regenerative therapy.

Nevertheless, Wagh and colleagues' study opens new areas of research and presents several unanswered questions for future investigations. The current study is impressive, but does the miR-363-mediated repression of *HAND1* occur *in vivo*? To what extent does this regulation affect the cardiac cell specification during development? Is it safe to use anti-miR-363 for cardiac regenerative therapies? What are the additional miRNA networks that cooperate with miR-363 to form a high-order regulatory complex in determining cardiac cell specification? Most importantly, how can we derive the desired functional cardiac subtype without any genetic manipulation? These are just a few of the myriad questions that are emerging from this exciting study. Answering these questions may identify more druggable targets and might help push cardiac research much closer to human cardiac regenerative therapy.

Abbreviations

CM: Cardiomyocyte; *HAND1*: Hand and neural crest derivative expressed 1; hESCs: Human embryonic stem cells; miRNA/miR: microRNA.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Section of Islet Cell and Regenerative Biology, Joslin Diabetes Center, Harvard Medical School, Boston, MA 01125, USA. ²Department of Stem Cell and Regenerative Biology, Harvard University, Harvard Stem Cell Institute, Cambridge, MA 01238, USA.




Figure 1 miR-363 negatively regulates the left ventricular cardiomyocyte specification transcription factor *HAND1*. BMP, bone morphogenetic protein; *HAND*, hand and neural crest derivative expressed; miR, microRNA; NKX2.5, NK2 transcription factor related, locus 5.

Published: 29 Jul 2014

References

1. Wagh V, Pomorski A, Wilschut KJ, Sebastian P, Bernstein HS: **MicroRNA-363 negatively regulates the left ventricular determining transcription factor HAND1 in human embryonic stem cell-derived cardiomyocytes.** *Stem Cell Res Ther* 2014, **5**:75.
2. Ahuja P, Sdek P, MacLellan WR: **Cardiac myocyte cell cycle control in development, disease, and regeneration.** *Physiol Rev* 2007, **87**:521–544.
3. Bergmann O, Bhardwaj RD, Bernard S, Zdunek S, Barnabe-Heider F, Walsh S, Zupicich J, Alkass K, Buchholz BA, Druid H, Jovinge S, Frisén J: **Evidence for cardiomyocyte renewal in humans.** *Science* 2009, **324**:98–102.
4. Bartel DP: **MicroRNAs: target recognition and regulatory functions.** *Cell* 2009, **136**:215–233.
5. van Rooij E, Sutherland LB, Liu N, Williams AH, McAnally J, Gerard RD, Richardson JA, Olson EN: **A signature pattern of stress-responsive microRNAs that can evoke cardiac hypertrophy and heart failure.** *Proc Natl Acad Sci U S A* 2006, **103**:18255–18260.
6. Porrello ER, Johnson BA, Aurora AB, Simpson E, Nam YJ, Matkovich SJ, Dorn GW 2nd, van Rooij E, Olson EN: **MiR-15 family regulates postnatal mitotic arrest of cardiomyocytes.** *Circ Res* 2011, **109**:670–679.
7. Zhao Y, Ransom JF, Li A, Vedantham V, von Drehle M, Muth AN, Tsuchihashi T, McManus MT, Schwartz RJ, Srivastava D: **Dysregulation of cardiogenesis, cardiac conduction, and cell cycle in mice lacking miRNA-1-2.** *Cell* 2007, **129**:303–317.
8. Zhang J, Wilson GF, Soerens AG, Koonce CH, Yu J, Palecek SP, Thomson JA, Kamp TJ: **Functional cardiomyocytes derived from human induced pluripotent stem cells.** *Circ Res* 2009, **104**:e30–e41.
9. Gupta MK, Illich DJ, Gaarz A, Matzkies M, Nguemo F, Pfannkuche K, Liang H, Classen S, Reppel M, Schultze JL, Hescheler J, Sarić T: **Global transcriptional profiles of beating clusters derived from human induced pluripotent stem cells and embryonic stem cells are highly similar.** *BMC Dev Biol* 2010, **10**:98.
10. McFadden DG, Barbosa AC, Richardson JA, Schneider MD, Srivastava D, Olson EN: **The Hand1 and Hand2 transcription factors regulate expansion of the embryonic cardiac ventricles in a gene dosage-dependent manner.** *Development* 2005, **132**:189–201.
11. Togi K, Kawamoto T, Yamauchi R, Yoshida Y, Kita T, Tanaka M: **Role of Hand1/eHAND in the dorso-ventral patterning and interventricular septum formation in the embryonic heart.** *Mol Cell Biol* 2004, **24**:4627–4635.

10.1186/scrt478

Cite this article as: Gupta and Rao: Hearty miR-363 controls *HAND1* in cardiac cell specification. *Stem Cell Research & Therapy* 2014, **5**:89