



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Carbon monoxide coupling and functionalisation at a simple uranium coordination complex

Citation for published version:

Arnold, PL, Turner, ZR, Bellabarba, RM & Tooze, RP 2011, 'Carbon monoxide coupling and functionalisation at a simple uranium coordination complex' *Chemical Science*, vol. 2, no. 1, pp. 77-79. DOI: 10.1039/c0sc00452a

Digital Object Identifier (DOI):

[10.1039/c0sc00452a](https://doi.org/10.1039/c0sc00452a)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Chemical Science

Publisher Rights Statement:

Copyright © 2010 by the Royal Society of Chemistry. All rights reserved.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Post-print of a peer-reviewed article published by the Royal Society of Chemistry.

Published article available at: <http://dx.doi.org/10.1039/C0SC00452A>

Cite as:

Arnold, P. L., Turner, Z. R., Bellabarba, R. M., & Tooze, R. P. (2011). Carbon monoxide coupling and functionalisation at a simple uranium coordination complex. *Chemical Science*, 2(1), 77-79.

Manuscript received: 20/08/2010; Accepted: 08/10/2010; Article published: 28/10/2010

Carbon monoxide coupling and functionalisation at a simple uranium coordination complex**

Polly L. Arnold,^{1,*} Zoë R. Turner,¹ Ronan M. Bellabarba² and Robert P. Tooze²

^[1]EaStCHEM, School of Chemistry, Joseph Black Building, University of Edinburgh, West Mains Road, Edinburgh, EH9 3JJ, UK.

^[2]Sasol Technology UK, Purdie Building, North Haugh, St Andrews, UK.


^[*]Corresponding author; e-mail: polly.arnold@ed.ac.uk

^[**]We thank Dr Stephen M. Mansell for collection of some of the analytical data, and Sasol Technology UK (studentship for Z. R. T.), the UK EPSRC (fellowship for P. L. A.) and the University of Edinburgh for funding.

Supporting information:

^[†] Electronic supplementary information (ESI) available: Experimental and crystallographic details (CCDC reference numbers 775504 and 775505) and spectroscopic data. For ESI and crystallographic data in CIF or other electronic format see <http://dx.doi.org/10.1039/C0SC00452A>

Graphical abstract:



Abstract

A simple coordination complex of uranium(III), a uranium tris(amide), can selectively couple gaseous CO to the linear enediolate $[\text{OCCO}]^{2-}$ dianion, at room temperature and pressure, regardless of the reagent stoichiometry. This product exhibits further reactivity upon warming in the form of the addition of a C–H bond of a methyl group across the $\text{C}\equiv\text{C}$ triple bond, this second carbon–carbon bond forming reaction generating a functionalised enediolate dianion.

Introduction

Carbon monoxide is the key C_1 component in the Fischer–Tropsch process and the chemical industry worldwide. It is generated from biomass or coal in the form of synthesis gas, or syngas, $(\text{CO} + n\text{H}_2)$ and used to make millions of tonnes of solvents, fuels, and bulk chemicals every year. The bond in CO is the strongest in the periodic table, and whilst many metal complexes are capable of catalysing the insertion of CO into other organic substrates, only a handful of complexes have demonstrated the C–C coupling of two metal-coordinated CO ligands. Molten potassium was reported to reductively couple CO to form salts of the oxocarbon anions from homologues to $[\text{C}_2\text{O}_2]^{2-}$ up to $[\text{C}_5\text{O}_5]^{2-}$ and $[\text{C}_6\text{O}_6]^{2-}$ in the early nineteenth century.^{1,2} The reduction of $\text{Ta}(\text{CO})_2(\text{dmpe})_2\text{Cl}$ by magnesium in the presence of Cp_2ZrCl_2 affords coupled and silylated $\text{Ta}(\text{Me}_3\text{SiOC}\equiv\text{COSiMe}_3)(\text{dmpe})_2\text{Cl}$ ($\text{dmpe} = 1,2\text{-bis}\{\text{dimethylphosphino}\}\text{ethane}$)³ and a study on the insertion of one and two equivalents of CO into the weak Rh–Rh bond in the rhodium octaethylporphyrin (oep) dimer $[\text{Rh}(\text{oep})]_2$ used 12 atm pressures of CO to force an equilibrium reaction over to a double insertion product characterised by ^1H NMR spectroscopy as $(\text{oep})\text{Rh}(\text{CO})(\text{CO})\text{Rh}(\text{oep})$.⁴ A heterotrimetallic tantalum hydride was shown to reductively couple six CO molecules at ambient pressure and temperature; protonolysis with $[\text{Me}_3\text{NH}][\text{BPh}_4]$ yielded $[\text{Ta}(2,6\text{-(CH}_2\text{-3-}^t\text{Bu-5-Me-2-OPh)}(\text{thf})_4(\text{C}_6\text{O}_6)]$.⁵ Reactions that can form carbon–carbon bonds are rare in general, and the direct dimerisation of CO to afford OCCO is energetically uphill by $+73 \text{ kcal mol}^{-1}$, and particularly rare.¹ Though the insertion of CO into metal alkyl, amido and hydride bonds is now well documented for Group 4 and f-block metal complexes,^{6–14} there are only two complexes which bind and reductively couple carbon monoxide to date; the sterically congested, trivalent, organometallic uranium complexes $[\text{U}(\eta\text{-C}_8\text{H}_6\{\text{Si}^i\text{Pr}_3\text{-1,4}\}_2)(\eta^5\text{-Cp}^R)]$ ($\text{Cp}^R = \text{C}_5(\text{CH}_3)_5$ or $\text{C}_5(\text{CH}_3)_4\text{H}$).^{15–17} The former demonstrated unprecedented reductive trimerisation of CO. The same U^{III} complex is also capable of reductively dimerising CO if exactly one equivalent of CO gas is used, although no interconversion between the $(\text{CO})_n$ homologues, or subsequent reactivity, has been demonstrated to date. Reductive homologation by the strongly reducing La^{III} and Sm^{II} complexes, $[\text{La}(\eta\text{-Cp}^*)_2]_2(\text{N}_2)$ and $[\text{Sm}(\eta\text{-Cp}^*)_2(\text{thf})_2]$, affords a ketene carboxylate dianion from three CO molecules at 90 psi pressures in 19 and 20% isolated yield, respectively.^{18,19} The incorporation of a pendant borane group to provide a hydride source has recently been used to address the second challenge in CO coupling – the formation of the reduced C–C and C–H bonds, the carbonyl $[(\text{Ph}_2\text{P}(\text{CH}_2)_2\text{B}(\text{C}_8\text{H}_{14}))_2\text{Re}(\text{CO})_4][\text{BF}_4]$ is borane activated to form $\text{Re-CH}_2\text{-O-BR}_3$ which couples


with a second carbonyl equivalent.²⁰ As yet, no catalytic C–C coupling or oligomerisation reactions exist for carbon monoxide at low pressures or temperatures.²¹

The tris(amido) complex of uranium UN''_3 , where N'' is the hydrocarbon-soluble, sterically demanding, bis(alkylsilyl)amido monoanion $N(\text{SiMe}_3)_2^-$, is widely used as a reagent in coordination chemistry, and despite its pyramidal structure, it has generally been regarded not to possess any interesting small molecule activation chemistry of its own. The 1979 report of its synthesis reported the authors' surprise at the absence of reactivity with CO or similar Lewis bases such as tetrahydrofuran (THF), phosphines or isonitriles.²² Since then, the 1 : 1 adduct with the strong nucleophilic donor N-heterocyclic carbene tetramethylimidazol-2-ylidene (NHC),²³ and with a further equivalent of the potassium amide KN'' have been reported;²⁴ the products $(\text{NHC})UN''_3$ and $K[UN''_4]$, respectively, demonstrating that there is sufficient space at the metal centre for further reaction chemistry.


Herein, we report how hydrocarbon solutions of the simple trivalent uranium amide UN''_3 react at ambient temperatures and pressures with carbon monoxide to afford a reductively coupled ynediolate $[\text{OCCO}]^{2-}$ unit, and how, for the first time, further reactivity is demonstrated for the linear ynediolate fragment. The simplicity, selectivity and redox-reactivity of this system suggest interesting catalytic carbon monoxide functionalisation chemistry may soon be accessible with related uranium(III) salts.

Results and discussion

Exposure of a dark purple hydrocarbon solution of the uranium amide complex UN''_3 ($N'' = N(\text{SiMe}_3)_2$) to carbon monoxide, CO, at ambient temperature and pressure results in a gradual decolourisation of the solution to amber. Golden-coloured crystals of the U^{IV} complex $N''_3U\text{--}OC\equiv CO\text{--}UN''_3$ **1** were formed in 82% yield, as the product of a reductive coupling of CO between two U centres, Scheme 1.‡ We note that the original report of the inertness of UN''_3 towards CO did not include details of solvent,²² the formation of **1** works equally well in hexanes or toluene, but not in the donor solvents tetrahydrofuran (THF) or pyridine. The ^1H NMR spectrum of a benzene solution of **1** contains a single paramagnetically shifted resonance for the silylmethyl H nuclei at -8.6 ppm (parts per million). The reaction can be carried out readily on a gram scale, and if repeated with ^{13}C -carbon monoxide, a resonance at 171 ppm is observed in the ^{13}C NMR spectrum for the ynediolate carbon nuclei in $N''_3U\text{--}O^{13}\text{C}\equiv^{13}\text{CO}\text{--}UN''_3$ **1a**. The carbon resonance was not visible in the absence of ^{13}C labelling. A single-crystal X-ray diffraction experiment confirms the one-electron oxidation of the uranium to U^{IV} , which we assume drives the reaction, and the linearity of the UOCCOU unit in **1** ($\text{U}\text{--}\text{O}\text{--}\text{C} = 178^\circ$); Fig. 1 shows the core of the molecular structure of **1**. The formerly pyramidal geometry at the uranium centre has flattened out with the U cation dropping from 0.46 Å above the plane described by the three N atoms (in UN''_3)²⁵ to 0.16 Å (in **1**) and the average U–N bond distance has shortened from 2.320 Å (in UN''_3) to 2.253 Å (in **1**), in accordance with the oxidation of the metal to U^{IV} .




(a)



(b)

Figure 1. Displacement ellipsoid drawing of the molecular structure of (a) **1** and (b) **2** (50% probability) with all silyl methyl groups omitted for clarity. Selected distances (Å) and angles (°): **1**: U1–O1 2.102(2), O1–C19 1.301(4), C19–C19ⁱ 1.183(7), U1–N1 2.251(3), O1–U1–N1 95.12(10), N1–U1–N2 122.01(10); **2**: U2–O2 2.093(7), O2–C38 1.402(11), C38–C39 1.310(13), U2–N5 2.242(7), U2–O2–C38 171.6(6), U3–O3–C39 170.9(6), O2–U2–N5 98.9(3), C56–C38–C39 125.8(9), N1–U1–N2 116.3(3).



Scheme 1. Coupling and functionalisation of carbon monoxide by the trivalent uranium amide to form a uranium-coordinated ynediolate **1**, and then an ene-diolate **2**.

The first example of CO coordination to uranium was observed in 1986; tris(cyclopentadienyl) U^{III} was shown to bind CO reversibly,²⁶ with substituted cyclopentadienyl and organometallic analogues later demonstrated to show stronger binding.^{27–29} The first non-organometallic uranium carbonyl complex was formed by reductive activation of a single CO molecule by two $\text{U}^{\text{III}}\text{L}$ complexes, in which L is a bulky tris(aryloxy) ligand, forming $[\text{LU}^{3.5}]_2(\mu\text{-}\eta^1, \eta^1\text{-CO})$.³⁰ No further reactivity of these interesting complexes has yet been reported. In the sole previous example of the reductive coupling of carbon monoxide to the coordinated ynediolate no further reaction was observed; a result corroborated by computational analysis.¹⁶ In contrast, heating a sample of **1** in benzene results in the formation of asymmetric **2**, Scheme 1, in which one of the amido Si–CH₃ groups has reacted formally to add C–H across the alkyne group intramolecularly, affording a seven-membered metallacyclic ring containing the newly formed alkene. Identified by ¹H NMR spectroscopy (containing six paramagnetically shifted resonances), and a single-crystal X-ray diffraction study, Fig. 1b, the gold-coloured complex still contains two U^{IV} cations, with close to linear U–O–C moieties, a planar O–C=C–O group, and an average $\text{U}^{\text{IV}}\text{–N}$ distance of 2.266 Å.

Both the reactions to form **1** and **2** with ¹³C-labelled carbon monoxide were monitored by *in situ* NMR spectroscopies, but no evidence of any intermediates was observed in either case. Further monitoring was hampered by the fact that both products tend to crystallise out of the solution as they form, and the range of solvents in which the formation of **1** can be monitored is limited to very non-polar, non-coordinating solvents.

Further chemistry of both **1** and **2** has also been investigated, to identify the potential for intermolecular functionalisation or substitution of the coupled oxycarbonanion. No reaction was observed between either **1** or **2** and dihydrogen at 1 atm pressure in benzene or THF solvent. The exposure of solutions of UN''_3 and the

chlorosilane ClSiMe_3 to an atmosphere of CO result in the formation of coupled **1** before any other appreciable reaction occurs, and warming the mixture still leads to the formation of **2** in preference to any other reactivity, presumably due to the obvious steric crowding in the complex. No reaction between **1** and other silanes, boranes and haloboranes (see ESI†) was observed, and heating of these reaction mixtures again afforded only **2**.

Conclusions

Hydrocarbon solutions of the simple trivalent uranium amide UN''_3 react at ambient temperatures and pressures with carbon monoxide to afford a reductively coupled ynediolate $[\text{OCCO}]^{2-}$ unit. The silylamido ligands are too sterically encumbering to allow any intermolecular reaction to occur in this system with reagents tried so far, but the electronic structure of this tris(amido) uranium system is clearly able to support further reactivity of the ynediolate fragment, exemplified here by the second carbon–carbon bond and carbon–hydrogen bond forming reactions occurring. To our knowledge, this is the first time that a metal complex has been able to reductively couple exclusively two molecules of gaseous CO at atmospheric pressure, and the first time that this reductively coupled $(\text{CO})_2$ has been demonstrated to form bonds to sp^3 carbon and hydrogen atoms.

An increasing range of reductive activation reactions of small, traditionally inert molecules such as dinitrogen and carbon dioxide are being demonstrated for trivalent uranium complexes,^{31–35} and the ready interconversion between the U^{III} and U^{IV} oxidation states involved here suggests that catalytic systems based on this coupling and functionalisation are viable. It is notable that the reaction occurs with such a simple coordination compound – an amide that is made from simple commercially available ligands (the precursor amide salt currently costs under €100 per mol). Work is in progress to develop mild routes to remove the coupled diolate with other reagents, and complete such a cycle.

Notes and references

[[†]] To a freeze-pump-thaw degassed solution of UN³ (1.00 g, 1.39 mmol) in toluene (20 mL) was added CO (1 atm) at room temperature. After 1 h, golden coloured microcrystalline solid started to precipitate and the reaction mixture was then kept at room temperature for 3 days. The solid was isolated by filtration, washed with hexane (3 × 10 mL) and then dried *in vacuo* to afford N³UOCCOUN³**1** as a golden-coloured crystalline solid. Yield 0.853 g (82%). Anal. (%): found (calc. for C₃₈H₁₀₈N₆O₂Si₁₂U₂): C, 30.37 (30.54); H, 7.35(7.28); N, 5.55 (5.62). ¹H NMR (C₆D₆, 599.8 MHz, 298 K): δ 8.55 (108 H, s, N{Si(CH₃)₃}₂) (fwhm = 1297 Hz) ppm. No ynediolate stretch was visible in the FTIR spectrum. Heating a slurry of N³UOCCOUN³**1** (0.107 g, 0.0716 mmol) into toluene (10 mL) to 80 °C for 24 h resulted in the conversion to N³UOCC(H)OU{N(SiMe₂CH₂)(SiMe₃)}N²**2**, also a golden-coloured solid, in 0.0713 g (67%) yield. Single crystals suitable for X-ray structure determinations were grown from room temperature benzene and toluene solutions of **1** and **2**, respectively.

[1] B. Wayland and X. Fu, *Science*, 2006, **311**, 790.

[2] W. Büchner, *Helv. Chim. Acta.*, 1963, **46**, 2111.

[3] P. A. Bianconi, I. D. Williams, M. P. Engeler and S. J. Lippard, *J. Am. Chem. Soc.*, 1986, **108**, 311.

[4] V. L. Coffin, W. Brennen and B. B. Wayland, *J. Am. Chem. Soc.*, 1988, **110**, 6063.

[5] T. Watanabe, Y. Ishida, T. Matsuo and H. Kawaguchi, *J. Am. Chem. Soc.*, 2009, **131**, 3474.

[6] P. T. Wolczanski and J. E. Bercaw, *Acc. Chem. Res.*, 1980, **13**, 121.

[7] T. J. Marks, *Science*, 1982, **217**, 989.

[8] W. J. Evans, J. W. Grate and R. J. Doedens, *J. Am. Chem. Soc.*, 1985, **107**, 1671.

[9] K. G. Moloy, P. J. Fagan, J. M. Manriquez and T. J. Marks, *J. Am. Chem. Soc.*, 1986, **108**, 56.

[10] K. G. Moloy and T. J. Marks, *J. Am. Chem. Soc.*, 1984, **106**, 7051.

[11] K. Tatsumi, A. Nakamura, P. Hofmann, R. Hoffmann, K. G. Moloy and T. J. Marks, *J. Am. Chem. Soc.*, 1986, **108**, 4467.

[12] W. J. Evans, S. A. Kozimor, G. W. Nyce and J. W. Ziller, *J. Am. Chem. Soc.*, 2003, **125**, 13831.

[13] E. L. Werkema, L. Maron, O. Eisenstein and R. A. Andersen, *J. Am. Chem. Soc.*, 2007, **129**, 6662.

- [14] O. Bénaud, J. C. Berthet, P. Thuéry and M. Ephritikhine, *Inorg. Chem.*, 2010, DOI: 10.1021/ic1012044.
- [15] O. T. Summerscales, F. G. N. Cloke, P. B. Hitchcock, J. C. Green and N. Hazari, *J. Am. Chem. Soc.*, 2006, **128**, 9602.
- [16] O. T. Summerscales, F. G. N. Cloke, P. B. Hitchcock, J. C. Green and N. Hazari, *Science*, 2006, **311**, 829.
- [17] A. S. Frey, F. G. N. Cloke, P. B. Hitchcock, I. J. Day, J. C. Green and G. Aitken, *J. Am. Chem. Soc.*, 2008, **130**, 13816.
- [18] W. J. Evans, J. W. Grate, L. A. Hughes, H. Zhang and J. L. Atwood, *J. Am. Chem. Soc.*, 1985, **107**, 3728.
- [19] W. J. Evans, D. S. Lee, J. W. Ziller and N. Kaltsoyannis, *J. Am. Chem. Soc.*, 2006, **128**, 14176.
- [20] A. J. M. Miller, J. A. Labinger and J. E. Bercaw, *J. Am. Chem. Soc.*, 2008, **130**, 11874.
- [21] E. M. Carnahan, J. D. Protasiewicz and S. J. Lippard, *Acc. Chem. Res.*, 1993, **26**, 90.
- [22] R. A. Andersen, *Inorg. Chem.*, 1979, **18**, 1507.
- [23] H. Nakai, X. L. Hu, L. N. Zakharov, A. L. Rheingold and K. Meyer, *Inorg. Chem.*, 2004, **43**, 855.
- [24] W. J. Evans, D. S. Lee, D. B. Rego, J. M. Perotti, S. A. Kozimor, E. K. Moore and J. W. Ziller, *J. Am. Chem. Soc.*, 2004, **126**, 14574.
- [25] J. L. Stewart and R. A. Andersen, *Polyhedron*, 1998, **17**, 953.
- [26] J. G. Brennan, R. A. Andersen and J. L. Robbins, *J. Am. Chem. Soc.*, 1986, **108**, 335.
- [27] M. D. Conejo, J. S. Parry, E. Carmona, M. Schultz, J. G. Brennann, S. M. Beshouri, R. A. Andersen, R. D. Rogers, S. Coles and M. Hursthouse, *Chem. Eur. J.*, 1999, **5**, 3000.
- [28] J. Parry, E. Carmona, S. Coles and M. Hursthouse, *J. Am. Chem. Soc.*, 1995, **117**, 2649.
- [29] W. J. Evans, S. A. Kozimor, G. W. Nyce and J. W. Ziller, *J. Am. Chem. Soc.*, 2003, **125**, 13831.
- [30] I. Castro-Rodriguez and K. Meyer, *J. Am. Chem. Soc.*, 2005, **127**, 11242.
- [31] A. R. Fox, S. C. Bart, K. Meyer and C. C. Cummins, *Nature*, 2008, **455**, 341.
- [32] O. T. Summerscales and F. G. N. Cloke, *Struct. Bonding*, 2008, **127**, 87.
- [33] I. Korobkov, S. Gorelsky and S. Gambarotta, *J. Am. Chem. Soc.*, 2009, **131**, 10406.

[34] M. Ephritikhine, *Dalton Trans.*, 2006, 2501.

[35] I. Castro-Rodriguez, H. Nakai, L. N. Zakharov, A. L. Rheingold and K. Meyer, *Science*, 2004, **305**, 1757.