



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Enhanced ectodysplasin-A receptor (EDAR) signaling alters multiple fiber characteristics to produce the East Asian hair form

Citation for published version:

Mou, C, Thomason, HA, Willan, PM, Clowes, C, Harris, WE, Drew, CF, Dixon, J, Dixon, MJ & Headon, DJ 2008, 'Enhanced ectodysplasin-A receptor (EDAR) signaling alters multiple fiber characteristics to produce the East Asian hair form' Human Mutation, vol. 29, no. 12, pp. 1405-11. DOI: 10.1002/humu.20795

Digital Object Identifier (DOI):

[10.1002/humu.20795](https://doi.org/10.1002/humu.20795)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Early version, also known as pre-print

Published In:

Human Mutation

General rights


Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy


The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Supplementary Figure S1. EDAR370A displays more potent signaling than EDAR370V under autoregulatory conditions. EDAR370V and EDAR370A were expressed from an NF- κ B responsive promoter. 100 ng/ml EDA was added to medium 12 hours post-transfection and NF- κ B responsive luciferase reporter activity was measured 3 or 16 hours after EDA administration. Error bars indicate s.e.m.




Supplementary Figure S2. Dose effect of the *Edar*^{Tg951} phenotype and its suppression by mutation of *Edaradd*. **A:** Gross appearance of wild type; *Edar*^{Tg951} heterozygous (labelled *Edar*^{Tg951} het) and *Edar*^{Tg951} homozygous (labelled *Edar*^{Tg951}) mice. Elevated expression of *Edar* in the heterozygous transgenic produces a coarse hair coat, though not to the extent seen in the homozygote. **B:** Gross appearance of: wild type; *Edar*^{Tg951}; *Edaradd*^{cr/cr} and *Edar*^{Tg951},*Edaradd*^{cr/cr} animals. The *Edaradd*^{cr/cr} and *Edar*^{Tg951},*Edaradd*^{cr/cr} mice have the same gross phenotype. **C:** PCR genotyping of the mice in **B**, detecting the *Edar* YAC transgene, *Edaradd* and *Actin* positive control.



Supplementary Figure S3. All hair follicle types are present in *Edar*^{Tg951} skin and cycle normally, though hair follicle density is reduced.

A and B: Longitudinal sections of newborn mouse dorsal skin showing the presence of primary (1°), secondary (2°) and tertiary (3°) follicle primordia, formed in three distinct waves during development. Scale bar, 100 μm. **C and D:** Transverse sections of P28 dorsal skin. Three follicle sizes are indicated. Scale bar 100 μm. **E:** Quantification of the density of each hair follicle type in P28 wild type and *Edar*^{Tg951} skin. The density of tertiary follicles, which produce zigzag hairs, is reduced. In wild type skin tertiary follicles comprise 80% of the total, while this follicle type constitutes 72% of all follicles in transgenic skin. In humans, scalp hair density in Chinese and Korean subjects has been reported to be significantly lower than that of Europeans, but reports disagree regarding the relative hair density of African subjects (Lee, et al., 2002; Loussouarn, et al., 2005; Tsai, et al., 2002). The formation of follicles in distinct waves in mouse, in addition to the effect of postnatal skin growth on hair density, make it difficult to compare mechanisms of reduced hair density in mouse to human skin. **F:** Quantitative RT-PCR determination of *Shh* and *Edar* expression in dissected hair follicle bulbs at P28, normalized to *βActin* levels. Error bars show s.e.m. **G and H:** *Shh* expression in P28 anagen hair bulbs detected by in situ hybridisation. The dashed red line denotes the central axis of the follicle. **G:** Most wild type follicles have asymmetric *Shh* expression that is restricted to the matrix on one side of the dermal papilla. **H:** Hair bulbs of *Edar*^{Tg951} mice display a broader, symmetric *Shh* expression domain. Scale bar, 50 μm. **I-L:** Hair cycling characteristics of wild type and transgenic animals. **I and J:** Hair follicles enter catagen appropriately at P15 and **K and L:** are initiating the first anagen at P21. Scale bar, 200 μm.



References

- Lee HJ, Ha SJ, Lee JH, Kim JW, Kim HO, Whiting DA. 2002. Hair counts from scalp biopsy specimens in Asians. *J Am Acad Dermatol* 46(2):218-21.
- Loussouarn G, El Rawadi C, Genain G. 2005. Diversity of hair growth profiles. *Int J Dermatol* 44 Suppl 1:6-9.
- Tsai RY, Lee SH, Chan HL. 2002. The distribution of follicular units in the Chinese scalp: implications for reconstruction of natural-appearing hairlines in Orientals. *Dermatol Surg* 28(6):500-3.