



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Metrics with Galilean Conformal Isometry

Citation for published version:

Bagchi, A & Kundu, A 2010, 'Metrics with Galilean Conformal Isometry', *Physical Review D*, vol. 83, no. 6, 066018, pp. -. <https://doi.org/10.1103/PhysRevD.83.066018>

Digital Object Identifier (DOI):

[10.1103/PhysRevD.83.066018](https://doi.org/10.1103/PhysRevD.83.066018)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Published In:

Physical Review D

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Metrics with Galilean Conformal Isometry

Arjun Bagchi ¹, Arnab Kundu ²

¹ *School of Mathematics,
University of Edinburgh
Edinburgh EH9 3JZ, UK.*

² *Theory Group
Department of Physics
University of Texas at Austin
Austin, TX 78712, USA.*

arjun.bagchi@ed.ac.uk, arnab@physics.utexas.edu

ABSTRACT: The Galilean Conformal Algebra (GCA) arises in taking the non-relativistic limit of the symmetries of a relativistic Conformal Field Theory in any dimensions. It is known to be infinite-dimensional in all spacetime dimensions. In particular, the 2d GCA emerges out of a scaling limit of linear combinations of two copies of the Virasoro algebra. In this paper, we find metrics in dimensions greater than two which realize the finite 2d GCA (the global part of the infinite algebra) as their isometry by systematically looking at a construction in terms of cosets of this finite algebra. We list all possible sub-algebras consistent with some physical considerations motivated by earlier work in this direction and construct all possible higher dimensional non-degenerate metrics. We briefly study the properties of the metrics obtained. In the standard one higher dimensional “holographic” setting, we find that the only non-degenerate metric is Minkowskian. In four and five dimensions, we find families of non-trivial metrics with a rather exotic signature. A curious feature of these metrics is that all but one of them are Ricci-scalar flat.

Contents

1. Introduction	1
2. A Review of the GCA	3
2.1 GCA in arbitrary dimensions	3
2.2 GCA in 2d	3
3. Construction of Metrics on Coset Spaces	5
4. Homogeneous spaces with 2d Galilean Conformal isometry	6
4.1 3-dimensional Minkowski space	7
4.2 4-dimensional metrics	8
4.3 5-dimensional metrics	10
5. Summary and conclusions	13
A. The list of sub-algebras	15

1. Introduction

Non-relativistic conformal theories have received a lot of recent attention in connection with the AdS/CFT conjecture, more generally the gauge-gravity duality. The most popular of the versions of this non-relativistic gauge-gravity duality has been the one studied in the context of the Schrodinger algebra. The Schrodinger algebra is the largest symmetry algebra of the free Schrodinger equation [1, 2] and has been observed in cold atom systems at unitarity [3]. Gravity duals of a certain class of field theories possessing Schrodinger symmetry have been proposed in [4, 5] and now there is an extensive literature in this line of research, some of which can be found in the excellent review [6]. Another popular venue of research in this field has been in relation to spacetime with Lifshitz symmetry proposed in [7], which unlike the Schrodinger case, does not exhibit invariance under Galilean boosts and hence does not contain the Galilean group as a part of the symmetry algebra.

In [8], a different direction to non-relativistic AdS/CFT was proposed by focusing on a systematic limiting procedure of the relativistic symmetry group. The relativistic conformal algebra on the boundary was parametrically contracted to what was called the Galilean Conformal Algebra (GCA). One of the remarkable observations

here was that the GCA could be given an infinite dimensional lift for any spacetime dimensions. It was also observed that the GCA was important to the study of non-relativistic hydrodynamics. Specifically, the finite dimensional GCA is the symmetry algebra of the Euler equations, which is valid in cases where the fluid viscosity can be neglected. There have been further studies of the various aspects of the GCA in [9] – [12].

The gravity dual of the GCA was proposed initially to be a novel Newton-Cartan like $AdS_2 \times R^d$ in [8]. The systematic limit when performed on the parent AdS metric leads to a degeneration and hence the proposal was that when one looked for a standard one dimension higher holographic construction, there would be no non-degenerate space-time metric and the theory described in terms of connections would be a geometrized version of Newtonian gravity. We should, at this point, remind the reader that in the case of the Schrodinger algebra, the gravity dual was found in a two-dimensional higher space-time. The question of finding a metric with the Galilean Conformal isometry in higher dimensions remained. Recently, in [13], a connection between asymptotically flat spaces and the GCA has been established. The 2d infinite dimensional GCA was shown to be isomorphic to the Bondi-Metzner-Sachs (BMS) algebra [14] in 3 dimensions which is the group of asymptotic isometries of flat three dimensional space at null infinity [15]. The two different points of view are seemingly at loggerheads and one of the issues that we address in this paper is this apparent confusion.

The basic philosophy behind constructing the gravity duals of non-relativistic field theories is to realize the corresponding symmetry group as the isometry group of a spacetime metric. We attempt to find all possible higher (greater than two) dimensional metrics possessing the Galilean Conformal isometry by a process of coset construction. In the context of non-relativistic Gauge-gravity duality, the authors of [16] have shown that under some “physical” conditions the metrics obtained by this method uniquely reproduces the holographic constructions with Schrödinger and Lifshitz isometries. This procedure has also been followed in [17], in relation to the aging algebra, an algebra of relevance to some non-equilibrium statistical mechanical systems without time translation symmetry. We conduct a case by case exhaustive study of all possible metrics that can arise out of this coset construction for the 2d GCA, using the finite part of the algebra. We look to implement the two “physical” conditions as outlined in [16] and then make our search more extensive by relaxing one of them. We find that when we are looking at metrics with one extra direction, the “physical” conditions do not lead to any non-degenerate spacetime metric in 3-d, adding strength to the claim that the correct structure to look for is indeed a Newton-Cartan like $AdS_2 \times R$. Interestingly, when one of the two “physical” conditions are relaxed, we obtain a flat 3d metric in keeping with the connection discussed in [13]. We find other non-degenerate metrics for higher dimensional spaces. Curiously, most of these metrics turn out to be Ricci-scalar flat, although (except for the Minkowskian

one) they source non-trivial Ricci tensors.

The outline of the paper is as follows: we first review, in Sec. 2, the Galilean Conformal Algebra with special emphasis on the 2d GCA which shall be the focus of the paper. In Sec. 3, we outline the procedure of constructing metrics on homogeneous coset spaces that we would use. Sec. 4 contains the main results of the paper. We sub-divide the section according to the dimension of the space-time metric that we construct and make several comments. The main results are also summarized in a table in this section. We end with some concluding remarks. An appendix contains a list of all possible sub-algebras for the 2d GCA.

2. A Review of the GCA

2.1 GCA in arbitrary dimensions

The maximal set of conformal isometries of Galilean spacetime generates the infinite dimensional Galilean Conformal Algebra [8]. The notion of Galilean spacetime is a little subtle since the spacetime metric degenerates into a spatial part and a temporal piece. Nevertheless there is a definite limiting sense (of the relativistic spacetime) in which one can define the conformal isometries (see [18]) of the nonrelativistic geometry. Algebraically, the set of vector fields generating these symmetries are given by

$$\begin{aligned} L^{(n)} &= -(n+1)t^n x_i \partial_i - t^{n+1} \partial_t, \\ M_i^{(n)} &= t^{n+1} \partial_i, \\ J_a^{(n)} \equiv J_{ij}^{(n)} &= -t^n (x_i \partial_j - x_j \partial_i), \end{aligned} \quad (2.1)$$

for integer values of n . Here $i = 1 \dots (d-1)$ range over the spatial directions. These vector fields obey the algebra

$$\begin{aligned} [L^{(m)}, L^{(n)}] &= (m-n)L^{(m+n)}, & [L^{(m)}, J_a^{(n)}] &= -nJ_a^{(m+n)}, \\ [J_a^{(n)}, J_b^{(m)}] &= f_{abc}J_c^{(n+m)}, & [L^{(m)}, M_i^{(n)}] &= (m-n)M_i^{(m+n)}. \end{aligned} \quad (2.2)$$

There is a finite dimensional subalgebra of the GCA (also sometimes referred to as the GCA) which consists of taking $n = 0, \pm 1$ for the $L^{(n)}, M_i^{(n)}$ together with $J_a^{(0)}$. This algebra is obtained by considering the nonrelativistic contraction of the usual (finite dimensional) global conformal algebra $SO(d, 2)$ (in $d > 2$ spacetime dimensions) (see for example [8, 19]).

2.2 GCA in 2d

In two spacetime dimensions, as is well known, the situation is special. The relativistic conformal algebra is infinite dimensional and consists of two copies of the Virasoro algebra. One expects this to be related to the infinite dimensional GCA

algebra [20]. In two dimensions the non-trivial generators in (2.2) are the L_n and the M_n :

$$L^{(n)} = -(n+1)t^n x \partial_x - t^{n+1} \partial_t, \quad M^{(n)} = t^{n+1} \partial_x, \quad (2.3)$$

which obey

$$[L^{(m)}, L^{(n)}] = (m-n)L^{(m+n)}, \quad [M^{(m)}, M^{(n)}] = 0, \quad [L^{(m)}, M^{(n)}] = (m-n)M^{(m+n)}. \quad (2.4)$$

These generators in (2.3) arise precisely from a nonrelativistic contraction of the two copies of the Virasoro algebra. To see this, let us remember that the non-relativistic contraction consists of taking the scaling

$$t \rightarrow t, \quad x \rightarrow \epsilon x, \quad (2.5)$$

with $\epsilon \rightarrow 0$. This is equivalent to taking the velocities $v \sim \epsilon$ to zero (in units where $c = 1$). Consider the vector fields which generate (two copies of) the centre-less Virasoro Algebra in two dimensions:

$$\mathcal{L}^{(n)} = -z^{n+1} \partial_z, \quad \bar{\mathcal{L}}^{(n)} = -\bar{z}^{n+1} \partial_{\bar{z}}. \quad (2.6)$$

In terms of space and time coordinates, $z = t + x$, $\bar{z} = t - x$. Expressing $\mathcal{L}_n, \bar{\mathcal{L}}_n$ in terms of t, x and taking the above scaling (2.5) reveals that in the limit the combinations

$$\mathcal{L}^{(n)} + \bar{\mathcal{L}}^{(n)} = -t^{n+1} \partial_t - (n+1)t^n x \partial_x + \mathcal{O}(\epsilon^2); \quad \mathcal{L}^{(n)} - \bar{\mathcal{L}}^{(n)} = -\frac{1}{\epsilon} t^{n+1} \partial_x + \mathcal{O}(\epsilon). \quad (2.7)$$

Therefore we see that as $\epsilon \rightarrow 0$ [20]

$$\mathcal{L}^{(n)} + \bar{\mathcal{L}}^{(n)} \longrightarrow L^{(n)}, \quad \epsilon(\mathcal{L}^{(n)} - \bar{\mathcal{L}}^{(n)}) \longrightarrow -M^{(n)}. \quad (2.8)$$

Let us now rewrite the (1+1)-dimensional (finite) algebra generated by $\{L^{(\pm 1)}, L^{(0)}\}$ and $\{M^{(\pm 1)}, M^{(0)}\}$. The non-trivial commutators resulting from (2.4) are given by

$$\begin{aligned} [D, H] &= H, & [D, K_0] &= -K_0, & [D, K_1] &= -K_1, & [D, P] &= P, \\ [K_0, H] &= 2D, & [B, H] &= P, & [K_1, H] &= 2B, \\ [K_0, B] &= K_1, & [K_0, P] &= 2B, \end{aligned} \quad (2.9)$$

where we have made the following identifications

$$\begin{aligned} L^{(-1)} &\equiv H, & L^{(0)} &\equiv D, & L^{(+1)} &\equiv K_0, \\ M^{(-1)} &\equiv P, & M^{(0)} &\equiv B, & M^{(+1)} &\equiv K_1. \end{aligned} \quad (2.10)$$

Here H is the time translation generator, D is the dilatation operator, P is the spatial translation generator, B is the Galilean boost and K_0, K_1 are the two components of the special conformal generator. These identifications naturally arise when one

considers the contraction of the relativistic conformal algebra [8]. In the rest of the paper we will entirely focus on the algebra written in (2.9) and not be concerned about the infinite dimensional extension. We would look to realize this finite algebra as the isometries of spacetime metrics in dimensions greater than two. It is natural to expect that only the finite GCA would play the role of the true isometries and the other higher modes may correspond to asymptotic isometries of the metrics that we would obtain¹. This is something that we would not address in the current paper.

3. Construction of Metrics on Coset Spaces

Here we briefly review the construction of metrics on coset spaces that we will use in the rest of the paper. We closely follow the notation and conventions of [16]. We would like to consider a coset $\mathcal{M} = \mathcal{G}/\mathcal{H}$, where \mathcal{G} is the Galilean Conformal group and \mathcal{H} is a subgroup of \mathcal{G} . The corresponding Lie algebras are denoted by \mathfrak{g} and \mathfrak{h} respectively and for each $g \in \mathfrak{g}$ there is a corresponding element denoted by $[g] \in \mathfrak{g}/\mathfrak{h}$. As vector spaces, we can always decompose

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m} . \quad (3.1)$$

The coset \mathcal{M} is called a reductive coset if there exists a choice of $\mathfrak{m} \in \mathcal{M}$ such that $[\mathfrak{h}, \mathfrak{m}] \subset \mathfrak{m}$. We will see that for the GCA generically we do not have such reductive cosets.

Our goal here will be to construct a \mathcal{G} -invariant metric on the homogeneous space \mathcal{M} . Given a Lie group the Cartan-Killing form is given by

$$\Omega_{ab} \equiv \frac{1}{I_{\text{adj}}} f_{ac}{}^d f_{bd}{}^c , \quad (3.2)$$

where $f_{ab}{}^c$ are the structure constants and I_{adj} is the Dynkin index. For a semi-simple Lie group the Cartan-Killing form is non-degenerate and therefore induces a non-degenerate \mathcal{G} -invariant metric on \mathcal{M} . However, the GCA is not a semi-simple algebra and thus the corresponding Killing form is degenerate. We would like to point out here that there is the possibility of constructing a non-degenerate two-form over the whole group manifold *via* a procedure called “double extension”. We would have more to say about this later.

Following [21], there exists a one-to-one correspondence between \mathcal{G} -invariant metric on $\mathcal{M} = \mathcal{G}/\mathcal{H}$ and $\text{Ad}(\mathcal{H})$ -invariant non-degenerate symmetric bilinear forms Ω on $\mathfrak{g}/\mathfrak{h}$. When \mathcal{H} is connected, this invariance takes the following form

$$\Omega_{[m][n]} f_{[k]p}{}^{[m]} + \Omega_{[k][m]} f_{[n]p}{}^{[m]} = 0 , \quad (3.3)$$

¹For a brief review on asymptotic isometries see *e.g.* [27].

where $[m], [n], \dots$ are indices corresponding to \mathfrak{m} and p indicates the index corresponding to \mathfrak{h} . Given the structure constants for a particular choice of \mathfrak{h} and \mathfrak{m} , we can solve for the bilinear Ω from this equation.

However, the existence or the uniqueness of a solution for Ω is not guaranteed and we will observe later that for the GCA only a few choices for the sub-algebra \mathfrak{h} we have a non-degenerate Ω . Moreover, a typical solution of (3.3) does not fix Ω completely, rather gives a symmetric bilinear in terms of a bunch of arbitrary real numbers. This therefore will result in redundancies in the description of the \mathcal{G} -invariant (family of) metrics that we will eventually obtain.

Now let us choose an explicit coordinate basis as in [16]. First we fix a linear space decomposition (3.1) and denote that t_m, t_n, \dots are the basis of \mathfrak{h} and t_p, t_q, \dots are the basis of \mathfrak{m} . Then an element $[g] \in \mathcal{G}/\mathcal{H}$ can be represented by

$$[g] = [\exp(x_m t_m) \exp(x_n t_n) \dots] \quad \text{modulo } \mathcal{H} . \quad (3.4)$$

The Maurer-Cartan one-form given by $J_g = g^{-1} dg$ can then be computed according to the linear space decomposition in (3.1)

$$J_g = e_m t_m + e_p t_p , \quad (3.5)$$

where e_m and e_p are the vielbein. The metric on the coset is then constructed by contracting the symmetric bilinear Ω with the vielbein

$$G = \Omega^{pq} e_p e_q . \quad (3.6)$$

4. Homogeneous spaces with 2d Galilean Conformal isometry

In this section we will discuss and present the non-trivial homogeneous spaces (and the corresponding choice of the sub-algebra) that we obtain *via* the coset construction. For the interested reader, we have presented a complete list of all possible sub-algebras of the 2d GCA in appendix A.

Let us mention our guiding principles for the choices of sub-algebra here. In [16], the authors uniquely determined the metrics for the Schrodinger and the Lifshitz algebras by imposing the following ‘‘physical’’ conditions:

1. \mathfrak{h} does not contain the translation generator P .
2. \mathfrak{h} contains the boost generator B .

As argued in [16], condition (1) is natural in the sense that P would induce infinitesimal translations in the resulting geometry and should not be included in the stabilizer of a point in \mathcal{G}/\mathcal{H} . We shall strictly follow condition (1) in all our examples. Condition (2) is derived from the higher dimensional analogue of Lorentz

invariance. For a d dimensional algebra, the authors of [16] proposed to keep J_{ij}, B_i in \mathfrak{h} to preserve Lorentz invariance in d dimensions. We, however, do not believe in the sanctity of this condition in our analysis and would proceed to relax it in our exhaustive study.

4.1 3-dimensional Minkowski space

We begin by considering the case when $\dim \mathcal{M} = 3$. In this case, the only choice that gives a non-degenerate symmetric bilinear Ω (and therefore a non-degenerate metric) is the coset $\mathcal{M} = \mathcal{G}/\{H, D, K_0\}$. Note that in this case the sub-algebra does not contain the boost generator B and thus it falls under the category where we relax one of the ‘‘physical’’ conditions outlined above (and in [16]).

The structure constants are given by

$$f_{[i]H}^{[j]} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 2 \\ 1 & 0 & 0 \end{pmatrix}, \quad f_{[i]D}^{[j]} = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad f_{[i]K_0}^{[j]} = \begin{pmatrix} 0 & 0 & -2 \\ 0 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}, \quad (4.1)$$

which gives

$$\Omega = \begin{pmatrix} 0 & -2\omega_{33} & 0 \\ -2\omega_{33} & 0 & 0 \\ 0 & 0 & \omega_{33} \end{pmatrix}, \quad \omega_{ij} \in \mathbb{R}. \quad (4.2)$$

Now the coset element is parametrized as

$$[g] = [e^{x_P P} e^{x_1 K_1} e^{x_B B}], \quad (4.3)$$

which gives the following vielbein

$$e_P = dx_P, \quad e_{K_1} = dx_1, \quad e_B = dx_B. \quad (4.4)$$

So, the resulting metric reads:

$$ds_3^2 = \Omega^{pq} e_p e_q = \omega_{33} (-4dx_P dx_1 + dx_B). \quad (4.5)$$

This is the flat 3-d Minkowski space². As we remarked earlier based on the observation recently made in [13], this is a consequence of the isomorphism between the finite Galilean Conformal group in $(1+1)$ -dimensions and the Poincaré group in $(2+1)$ -dimensions. The isomorphism actually extends beyond the finite GCA and encompasses the full infinite extension of the GCA on one side and the infinite-dimensional BMS group in 3 dimensions which is the asymptotic symmetric group of flat 3d space-times at null infinity [13].

²Clearly we can set $\omega_{33} = 1$ without any loss of generality.

We observe that the strict imposition of both the “physical conditions” above does not lead to any non-degenerate spacetime metric. As remarked in the introduction, the original proposal for the dual gravitational description of a system with the GCA was given in terms of a Newton-Cartan like AdS [8]. In the case of the three dimensional bulk dual, the structure of the spacetime would be a fibre bundled $AdS_2 \times R$. The space-time metric degenerates and the dynamical quantities are the Christoffel symbols which “talk” to the separate metrics of the base AdS_2 and the fibres. The imposition of Lorentz symmetry in two dimensions (condition (2)) in our present construction rules out a non-degenerate spacetime metric and this is in keeping with the claim that the correct structure to look for is a Newton-Cartan like $AdS_2 \times R$.

Let us comment on a couple of things here about the flat metric that we have obtained. Firstly, we know that if an n -dimensional manifold admits $\frac{1}{2}n(n+1)$ Killing vectors, it must be a manifold of constant curvature. We were looking for spacetimes in 3 dimensions admitting the 6 dimensional GCA as an isometry. So, we would have ended up with spacetimes of constant curvature, our only choices are: flat, de-Sitter, or anti de-Sitter in three dimensions. That we get a flat spacetime is thus not a surprise.

Another point to note is that this seems to be the metric that is picked out by the method of contractions on that gave rise to the GCA from the relativistic conformal algebra from the point of view of AdS/CFT [8], both on the boundary and in the bulk. To see this, let us remind ourselves that the AdS_3 metric is obtained by the following coset construction (see *e.g.* [22]):

$$AdS_3 = \frac{SL(2, R) \times SL(2, R)}{SL(2, R)_{\text{diag}}} . \quad (4.6)$$

The above construction of the Minkowskian metric of the GCA is precisely the contraction of (4.6)³. The finite GCA is obtained by contracting the global $SL(2, R) \times SL(2, R)$ of the Virasoro algebra and $SL(2, R)_{\text{diag}}$, the diagonal $SL(2, R)$ subgroup of the relativistic theory, is parent of the $\{H, D, K_0\}$ subalgebra of the GCA.

4.2 4-dimensional metrics

Next we consider the case when $\dim \mathcal{M} = 4$. The first non-trivial case is the coset $\mathcal{M} = \mathcal{G}/\{B, D\}$, which obeys both the “physical” conditions outlined in [16]. In this case the structure constants are given by

$$f_{[i]B}^{[j]} = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & +1 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad f_{[i]D}^{[j]} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & +1 & 0 \\ 0 & 0 & 0 & +1 \end{pmatrix}, \quad (4.7)$$

³We would like to thank Rajesh Gopakumar for pointing this out to us.

which yields

$$\Omega = \begin{pmatrix} 0 & 0 & \omega_{13} & \omega_{14} \\ 0 & 0 & \omega_{14} & 0 \\ \omega_{13} & \omega_{14} & 0 & 0 \\ \omega_{14} & 0 & 0 & 0 \end{pmatrix}, \quad \omega_{ij} \in \mathbb{R}. \quad (4.8)$$

This is non-degenerate as long as $\omega_{14} \neq 0$.

Since we get a non-degenerate bilinear, let us compute the vielbein in this case. We parametrize the coset element as

$$[g] = [e^{x_H H} e^{x_P P} e^{x_0 K_0} e^{x_1 K_1}], \quad (4.9)$$

which gives the following vielbein

$$\begin{aligned} e_H &= dx_H, & e_P &= dx_P, \\ e_{K_0} &= x_0^2 dx_H + dx_0, \\ e_{K_1} &= 2x_0 x_1 dx_H + x_0^2 dx_P + dx_1. \end{aligned} \quad (4.10)$$

For the sake of visualization, let us write down the full metric. We define $x_H = t$, $x_P = x$, $x_0 = y$, $x_1 = z$, $w_{31} = a$, $w_{41} = b$. The metric, then, can be written as:

$$ds_{4(1)}^2 = (2ay^2 + 4byz)dt^2 + 4by^2 dt dx + 2adtdy + 2bdtdz + 2bdxdy. \quad (4.11)$$

Note that here we have two arbitrary real numbers a, b which parametrize a family of metrics. This family of metrics has vanishing Ricci-scalar.

The other non-trivial result comes from taking the coset $\mathcal{M} = \mathcal{G}/\{B, \alpha_1 D + \alpha_2 K_1\}$. The structure constants are given by

$$f_{[i]B}^{[j]} = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -\alpha_1/\alpha_2 & 0 \end{pmatrix}, \quad f_{[i]\alpha_1 D + \alpha_2 K_1}^{[j]} = \alpha_1 \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & +1 & 0 \\ 0 & 0 & 0 & +1 \end{pmatrix}, \quad (4.12)$$

which yields the following

$$\Omega = \begin{pmatrix} 0 & 0 & \omega_{31} & \omega_{41} \\ 0 & 0 & 0 & -\frac{\alpha_1}{\alpha_2} \omega_{31} \\ \omega_{31} & 0 & 0 & 0 \\ \omega_{41} & -\frac{\alpha_1}{\alpha_2} \omega_{31} & 0 & 0 \end{pmatrix}, \quad \omega_{ij} \in \mathbb{R}. \quad (4.13)$$

The above Ω is non-degenerate for $\alpha_1 \neq 0$. The vielbein are obtained to be:

$$\begin{aligned} e_H &= e^{-x_D} dx_H, & e_P &= e^{-x_D} dx_P, \\ e_D &= dx_D - 2x_0 e^{-x_D} dx_H - \frac{\alpha_1}{\alpha_2} dx_P, \\ e_{K_0} &= e^{-x_D} x_0^2 dx_H - x_0 dx_D + dx_0. \end{aligned} \quad (4.14)$$

It can be checked that without any loss of generality we can set⁴ $\omega_{31} = 1 = \alpha_2$. Hence we get a family of metrics parametrized by two real numbers ω_{41} and α_1 . Again, for clarity, it is useful to write the metric down explicitly. We make the following redefinitions: $x_H = t, x_P = x, x_0 = y, e^{x^D} = r, \omega_{41} = a, \alpha_2 = \alpha$.

$$ds_{4(2)}^2 = \frac{2}{r^2} \{ (1-ay)drdt + (ay^2 - 2y)dt^2 - \alpha(r+y^2)dtdx + ardt dy + \alpha y dx dr - \alpha r dx dy \} . \quad (4.15)$$

It is trivial to check that this metric also has vanishing Ricci-scalar. This is the only non-reductive example that we encounter in the coset construction of the 2-dimensional Galilean Conformal symmetry.

Let us offer some comments regarding the signature of these 4-dimensional metrics. It can be observed that the two distinct families of metrics we obtained take the following generic form

$$ds^2 = 2\Omega_{13}e_1e_3 + 2\Omega_{14}e_1e_4 + 2\Omega_{2,(3/4)}e_2e_{(3/4)} , \quad (4.16)$$

where Ω_{ij} are the corresponding matrix entries in (4.8) or (4.13) and e_i 's are the vielbein given in (4.10) or (4.14). If we introduce a local orthonormal frame $\{E_1, E_2, E_3, E_4\}$, where E_i 's are appropriate linear combinations of e_i 's, the particular form of the metric in (4.16) is strongly suggestive that the signature of the metric should be $(2, 2)$.⁵ It is worth noting at this point that in [11], a geometric realization of the ‘‘exotic’’ Galilean Conformal Isometry in $(2 + 1)$ -dimensions (called ‘‘exotic’’ because of the existence of a central charge in the commutator of the boost generators on the plane which is special to these dimensions) was found in terms of an AdS_7 -metric with $(3, 4)$ signature.

4.3 5-dimensional metrics

Finally we present the 5-dimensional metrics obtained *via* the coset construction. The first non-trivial case is the coset $\mathcal{M} = \mathcal{G}/\{B\}$. This gives the following symmetric bilinear

$$\Omega = \begin{pmatrix} \omega_{11} & 0 & \omega_{13} & \omega_{14} & \omega_{15} \\ 0 & 0 & 0 & \omega_{15} & 0 \\ \omega_{13} & 0 & \omega_{33} & \omega_{34} & 0 \\ \omega_{14} & \omega_{15} & \omega_{34} & \omega_{44} & 0 \\ \omega_{15} & 0 & 0 & 0 & 0 \end{pmatrix} , \quad \omega_{ij} \in \mathbb{R} , \quad (4.17)$$

⁴This is achieved by computing the Ricci tensor and observing that only the ratios ω_{41}/ω_{31} and α_2/α_1 appear.

⁵It is easy to check that one cannot write $ds^2 = -E_1^2 + E_2^2 + E_3^2 + E_4^2$; however one can write $ds^2 = -E_1^2 - E_2^2 + E_3^2 + E_4^2$.

which is non-degenerate if $\omega_{15} \neq 0$ and $\omega_{33} \neq 0$ and without any loss of generality we can set $\omega_{33} = 1 = \omega_{13} = \omega_{14} = \omega_{15} = \omega_{34}$. In this case we get the following vielbein

$$\begin{aligned} e_H &= e^{-x_D} dx_H, & e_P &= e^{-x_D} dx_P, & e_D &= -2x_0 e^{-x_D} dx_H + dx_D, \\ e_{K_0} &= x_0^2 e^{-x_D} dx_H - x_0 dx_D + dx_0, \\ e_{K_1} &= 2x_0 x_1 e^{-x_D} dx_H + x_0^2 e^{-x_D} dx_P - x_1 dx_D + dx_1. \end{aligned} \quad (4.18)$$

The resulting two-parameter family of metrics is Ricci-scalar flat. Clearly, this construction obeys both the ‘‘physical’’ conditions.

The only other non-trivial example in 5-dimensions is the coset $\mathcal{M} = \mathcal{G}/\{D\}$, which does not obey the ‘‘physical’’ condition (2). In this case we get

$$\Omega = \begin{pmatrix} 0 & 0 & \omega_{31} & \omega_{41} & 0 \\ 0 & 0 & \omega_{32} & \omega_{42} & 0 \\ \omega_{31} & \omega_{32} & 0 & 0 & 0 \\ \omega_{41} & \omega_{42} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \omega_{55} \end{pmatrix}, \quad \omega_{ij} \in \mathbb{R}. \quad (4.19)$$

This is also non-degenerate provided $\omega_{55} \neq 0$ and $\omega_{32}\omega_{41} \neq \omega_{31}\omega_{42}$. The vielbein are given by

$$\begin{aligned} e_H &= dx_H, & e_P &= dx_P, & e_{K_0} &= x_0^2 dx_H + dx_0, \\ e_{K_1} &= (2x_0 x_1 + x_0^2 x_B) dx_H + x_0^2 dx_P + x_B dx_0 + dx_1, \\ e_B &= -2x_0 dx_P + dx_B. \end{aligned} \quad (4.20)$$

This actually gives a four-parameter family of metrics. This family generically has coordinate dependent Ricci scalar which diverges as $R \sim x_1^2$ for $x_1 \rightarrow \infty$. If $\omega_{32} = 0$, we still get a non-degenerate metric but the Ricci-scalar vanishes identically. On the other hand, if $\omega_{32} \neq 0$, then the Ricci-scalar can vanish at a particular point in x_1 .

As in the examples with 4-dimensional metrics, it can be also argued that the existence of a local orthonormal frame and the precise structure of these 5-dimensional metrics strongly suggests the signature be (2, 3).

Note that the 2-dimensional GCA has 6 generators; hence a homogeneous space of 5-dimensions is constructed by choosing a sub-algebra which consists of only one generator. This is a rather trivial choice which nonetheless yields a family of non-trivial metrics.

Finally, we summarize some of our results in the following table:

Here R denotes the curvature scalar defined by $R = g^{\mu\nu} R_{\mu\nu}$; $R_{\mu\nu}^2 \equiv R^{\mu\nu} R_{\mu\nu}$; $R_{\mu\nu\rho\sigma}^2 \equiv R^{\mu\nu\rho\sigma} R_{\mu\nu\rho\sigma}$ and finally the curvature of the Weyl tensor is defined as $C_{\mu\nu\rho\sigma}^2 = C^{\mu\nu\rho\sigma} C_{\mu\nu\rho\sigma}$. The metrics that we obtain in this construction (except the Minkowski one) do yield fairly non-trivial Ricci tensor. Thus it is not clear to us what matter fields will source such backgrounds. It is therefore not obvious that

Table 1: The summary

Choice of subalgebra	dim \mathcal{M}	Properties
$\langle B \rangle$	5	$R = R_{\mu\nu}^2 = R_{\mu\nu\rho\sigma}^2 = 0 = C_{\mu\nu\rho\sigma}^2$
$\langle D \rangle$	5	$R \neq 0, R_{\mu\nu}^2 \neq 0, R_{\mu\nu\rho\sigma}^2 \neq 0$ and $C_{\mu\nu\rho\sigma}^2 \neq 0$ singularity appears as $x_0, x_1, x_B \rightarrow \infty$
$\langle B, D \rangle$	4	$R = R_{\mu\nu}^2 = R_{\mu\nu\rho\sigma}^2 = 0 = C_{\mu\nu\rho\sigma}^2$
$\langle B, \alpha_1 D + \alpha_2 K_1 \rangle,$ $\alpha_{1,2} \neq 0$	4	$R = R_{\mu\nu}^2 = R_{\mu\nu\rho\sigma}^2 = 0 = C_{\mu\nu\rho\sigma}^2;$ a (non-trivial) non-reductive coset
$\langle B, D, K_0 \rangle$	3	Minkowski

such matter fields will preserve the Galilean Conformal isometry. Thus although our metrics do possess the desired isometry, the full background (the metric along with the matter fields sourcing it) may not.

Before we leave this section altogether, a few comments are in order: First, as we remarked earlier in this construction we get a family of metrics parametrized by arbitrary real numbers. The redundancy in this description does not fix the sign of these parameters and hence does not fix the signature of the metric. However, by assuming the existence of a local orthonormal frame we seem to be able to fix the signature of these metrics and they turn out to be rather non-standard.

Second, note that once we know a metric with the Galilean Conformal isometry in a given dimension, it is straightforward to construct a higher dimensional metric with the same isometry by fibering the lower dimensional metric over a base manifold

$$ds^2 = f_1(\zeta)d\zeta^2 + f_2(\zeta)ds_{\text{GCA}}^2, \quad (4.21)$$

where $f_1(\zeta)$ and $f_2(\zeta)$ are two arbitrary functions and ds_{GCA}^2 is the metric with the Galilean Conformal isometry. This isometry acts non-trivially on the metric ds_{GCA}^2 but has no natural action on the base manifold. However, a spacetime thus constructed is not a homogeneous space since the Galilean Conformal isometry group does not act transitively on the whole manifold. Therefore the homogeneous spaces we obtained in 4 and 5-dimensions are not related in any obvious manner to the 3-dimensional Minkowski space and are thus truly non-trivial⁶.

Finally let us return to a point which was made in the initial sections. The 2d GCA has a degenerate Cartan-Killing form given by

$$\Omega \sim \begin{pmatrix} 0 & 0 & -2 \\ 0 & 1 & 0 \\ -2 & 0 & 0 \\ & & & 0 \end{pmatrix}, \quad (4.22)$$

⁶We would like to thank J. Simon and J. Figueroa-O'Farill for discussions related to this issue.

where the upper left 3×3 non-degenerate block comes from the $SL(2, R)$ sub-algebra spanned by $\{L^{(\pm)}, L^{(0)}\}$. The rest of the matrix entries are all zeroes.

However, the 2d GCA actually allows for a non-degenerate 2-form over the whole group manifold. The situation is similar to the well-known Nappi-Witten algebra [23] (the centrally extended 2d Euclidean algebra) or the Abelian extension of d -dimensional Euclidean algebra considered in *e.g.* [24]. The general construction of an invariant non-degenerate metric for non semi-simple Lie algebra goes by the name of “double extension” introduced in [25]⁷. Below we briefly review this.

Let \mathfrak{h} be any Lie algebra and \mathfrak{h}^* be its dual. Let the basis for \mathfrak{h} and \mathfrak{h}^* be respectively denoted by $\{X_a\}$ and $\{X^a\}$ obeying the relation: $\langle X_a, X^b \rangle = \delta_a^b$. Using the fact that \mathfrak{h} acts on \mathfrak{h}^* *via* the coadjoint representation, one can define the following Lie algebra structure on the vector space $\mathfrak{h} \oplus \mathfrak{h}^*$

$$\begin{aligned} [X_a, X_b] &= f_{ab}{}^c X_c, \\ [X^a, X^b] &= 0, \\ [X_a, X^b] &= -f_{ac}{}^b X^c, \end{aligned} \tag{4.23}$$

where $f_{ab}{}^c$ are the structure constants for the Lie algebra \mathfrak{h} . This defines a semidirect product of \mathfrak{h} and \mathfrak{h}^* . It is now possible to define an invariant metric on this semidirect product algebra.

From the definition of the finite 2d GCA in (2.4) and the Lie algebra structure defined in (4.23), it is obvious that $X_a \equiv L^{(m)}$ and $X^a \equiv M^{(m)}$, where $m = 0, \pm$. Thus the GCA is isomorphic to the semidirect product of $SL(2, R)$ with its coadjoint representation. We can define a two parameter family of invariant inner products in the following manner:

$$\langle X_a, X_b \rangle = \alpha \Omega_{ab}, \quad \langle X_a, X^b \rangle = \beta \delta_a^b, \quad \langle X^a, X^b \rangle = 0, \tag{4.24}$$

where α and β are non-zero real numbers and Ω_{ab} is the non-degenerate Cartan-Killing form for $SL(2, R)$. This construction works for the semidirect product of any simple Lie algebra G with its coadjoint representation. It is called the double extension of the trivial metric Lie algebra by G [25].

5. Summary and conclusions

In this paper we have systematically constructed metrics in dimensions greater than two which realize the two-dimensional Galilean Conformal Algebra as their isometry. We classified all the relevant sub-algebras of the 2D GCA and in order to construct these metrics looked at a formulation in terms of cosets. Though many

⁷We would like to thank J. Figueroa-O’Farill for explaining this issue to us and bringing this reference to our attention.

choices of these cosets turned out to produce degenerate metrics, we were able to get some non-trivial higher dimensional metrics. In three dimensions, we obtained a flat Minkowskian metric which we observed to be the contracted limit of the metric on AdS_3 . In higher dimensions, *viz.* four and five, we found several families of metrics, all except one of which turned out to be Ricci-scalar flat.

It is curious that most of the metrics we have obtained are Ricci-scalar flat. It would be worthwhile trying to understand if there is any deeper reason behind this, or if it is a mere co-incidence. One would also like to understand if there is any fundamental difference between these Ricci-scalar flat metrics and the family which is not, given that they were obtained in by similar methods.

Despite the fact that these metrics (except the Minkowski one) seem to have a “wrong” signature, a further analysis may turn out to be useful in understanding their structure. It will be very interesting to determine the matter fields which source such backgrounds. However since these metrics are neither Lorentzian nor Euclidean, it may be difficult to interpret such “matter fields” physically.

In the spirit of the gauge/gravity duality, one could look to try and reproduce the correlation functions of the 2d GCA [10, 20] from a gravity analysis. This might actually be a challenging task as there is little chance that modes would separate into normalizable and non-normalizable ones as in the usual AdS case. But if one is able to perform such computations, then one could claim that these metrics are actually holographically dual to the non-relativistic field theories with the GCA as their symmetry algebra.

Another speculation made earlier was that the metrics obtained by this method might realize the infinite dimensional GCA as asymptotic symmetries. It has been observed in [15] that the infinite BMS algebra in three dimensions, which is isomorphic to the 2d GCA [13], arises as the asymptotic symmetries of flat space at null infinity. So, this speculation indeed holds for our construction in three dimensions. The expectation is that the other metrics which have the finite 2d GCA as their isometries would also realize the infinite GCA in a manner similar to the BMS case. In [27], following the general scheme of calculating asymptotic symmetries outlined in [28], the authors constructed the asymptotic symmetry algebra for metrics with Schrodinger symmetry and found that the infinite extension of the Schrodinger algebra indeed emerges as the asymptotic symmetries of those metrics. The obstruction for applying the general formalism of [28] to the GCA was the absence of a spacetime metric. Now that in this work we have derived a number of metrics with the finite GCA as the isometry algebra, it should in principle be possible to carry out a similar analysis to [27] and check whether our speculation is indeed correct.

A natural direction of extending this analysis is to construct the metrics for the higher dimensional GCAs by this method of cosets. But the problem of classifying relevant subalgebras quickly becomes intractable and the full analysis too unwieldy to attempt by a case-by-case study. This would involve a mathematical machinery

more elaborate and powerful than what we have used in the two dimensional analysis. Another natural extension is to consider the Super-GCA and construction of super-cosets. A natural place to begin would be again two dimensions [29]. The size of the finite algebra would provide a challenge which in this case may be overcome by imposing strict “physical conditions”.

To conclude, let us remark on a point we have only fleetingly looked at in this paper. The existence of a non-degenerate 2-form on the full finite GCA is an avenue of potential fruitful research. Given that there is no field theory known for the GCA, it would be nice to use the construction of Nappi-Witten [23] and its generalizations [25] to construct a WZW model with the GCA as its symmetry. It would also be useful to understand if the infinite extension of the GCA plays any interesting role in this context.

Acknowledgments

It is a pleasure to acknowledge discussions with Paul de Medeiros, Anindya Dey, Jacques Distler, Rajesh Gopakumar, Debashis Ghoshal, Joan Simon, Yuji Tachikawa, Masahito Yamazaki and especially Jose Figueroa-O’Farrill. AB would like to thank the Institute for Advanced Study, Princeton for their generous hospitality during the initial part of this work. AK would like to thank the Institute for Advanced Study for the stimulating environment during the PITP school on “Aspects of Supersymmetry” when a part of this work was carried out. AK is supported by a Simons Postdoctoral Fellowship awarded by the Simons Foundation and by the National Science Foundation under Grant Numbers PHY-0969020 and PHY-0455649.

A. The list of sub-algebras

Here we enlist the possible choices of the sub-algebras for the finite part of GCA in (2.9). We begin by imposing the “physical” conditions imposed in [16] and then relaxing it. Just to remind the reader, the “physical” conditions are:

1. \mathfrak{h} does not contain the translation generator P .
2. \mathfrak{h} contains the boost generator B .

However we do not impose any constraint on the dimensionality of $\mathcal{M} = \mathcal{G}/\mathcal{H}$.

Let us therefore enlist the possible choices in the descending order in $\dim\mathcal{M}$:

(i) $\dim\mathcal{H} = 1$, $\dim\mathcal{M} = 5$:

$$\mathfrak{h} = \langle B \rangle, \quad \mathfrak{m} = \langle H, P, D, K_0, K_1 \rangle. \quad (\text{A.1})$$

More generally, however we have

$$\begin{aligned} \mathfrak{h} &= \langle \alpha_1 B + \alpha_2 H + \alpha_3 D + \alpha_4 K_0 + \alpha_5 K_1 \rangle, \quad \alpha_1 \neq 0, \\ \mathfrak{m} &= \langle H, P, D, K_0, K_1 \rangle. \end{aligned} \quad (\text{A.2})$$

(ii) $\dim\mathcal{H} = 2, \dim\mathcal{M} = 4$:

$$\mathfrak{h}_{(1)} = \langle B, K_1 \rangle, \quad \mathfrak{m}_{(1)} = \langle H, P, D, K_0 \rangle, \quad (\text{A.3})$$

$$\mathfrak{h}_{(2)} = \langle B, D \rangle, \quad \mathfrak{m}_{(2)} = \langle H, P, K_0, K_1 \rangle. \quad (\text{A.4})$$

More generally we can have

$$\mathfrak{h}_{(3)} = \langle B, \alpha_1 D + \alpha_2 K_1 \rangle, \quad \mathfrak{m}_{(3)} = \langle H, P, D, K_0 \rangle, \quad \alpha_2 \neq 0, \quad (\text{A.5})$$

$$\mathfrak{h}_{(4)} = \langle B, \alpha_1 D + \alpha_2 K_1 \rangle, \quad \mathfrak{m}_{(4)} = \langle H, P, K_0, K_1 \rangle, \quad \alpha_1 \neq 0, \quad (\text{A.6})$$

(iii) $\dim\mathcal{H} = 3, \dim\mathcal{M} = 3$:

$$\mathfrak{h}_{(1)} = \langle B, K_1, \alpha_1 D + \alpha_2 K_0 \rangle, \quad \mathfrak{m}_{(1)} = \langle H, P, K_0 \rangle, \quad \alpha_1 \neq 0, \quad (\text{A.7})$$

$$\mathfrak{h}_{(2)} = \langle B, K_1, K_0 \rangle, \quad \mathfrak{m}_{(2)} = \langle H, P, D \rangle. \quad (\text{A.8})$$

More generally we have

$$\mathfrak{h}_{(3)} = \langle B, K_1, \alpha_1 D + \alpha_2 K_0 + \alpha_3 K_1 \rangle, \quad \mathfrak{m}_{(3)} = \langle H, P, K_0 \rangle, \quad \alpha_{1,3} \neq 0, \quad (\text{A.9})$$

$$\mathfrak{h}_{(4)} = \langle B, K_0, \alpha_1 D + \alpha_2 K_0 + \alpha_3 K_1 \rangle, \quad \mathfrak{m}_{(4)} = \langle H, P, D \rangle, \quad \alpha_{2,3} \neq 0. \quad (\text{A.10})$$

(iv) $\dim\mathcal{H} = 4, \dim\mathcal{M} = 2$:

$$\mathfrak{h} = \langle B, K_1, D, K_0 \rangle, \quad \mathfrak{m} = \langle H, P \rangle. \quad (\text{A.11})$$

Let us now enlist the possibilities relaxing the condition (2), *i.e.* we consider \mathfrak{h} not containing B . The choices are:

(v) $\dim\mathcal{H} = 1, \dim\mathcal{M} = 5$:

$$\mathfrak{h}_{(1)} = \langle H \rangle, \quad \mathfrak{m}_{(1)} = \langle P, D, K_0, K_1, B \rangle, \quad (\text{A.12})$$

$$\mathfrak{h}_{(2)} = \langle D \rangle, \quad \mathfrak{m}_{(2)} = \langle H, P, K_0, K_1, B \rangle, \quad (\text{A.13})$$

$$\mathfrak{h}_{(3)} = \langle K_0 \rangle, \quad \mathfrak{m}_{(3)} = \langle H, P, D, K_1, B \rangle, \quad (\text{A.14})$$

$$\mathfrak{h}_{(4)} = \langle K_1 \rangle, \quad \mathfrak{m}_{(4)} = \langle H, P, D, K_0, B \rangle. \quad (\text{A.15})$$

(vi) $\dim\mathcal{H} = 2, \dim\mathcal{M} = 4$:

$$\mathfrak{h}_{(1)} = \langle H, D \rangle, \quad \mathfrak{m}_{(1)} = \langle P, K_0, K_1, B \rangle, \quad (\text{A.16})$$

$$\mathfrak{h}_{(2)} = \langle K_0, K_1 \rangle, \quad \mathfrak{m}_{(2)} = \langle H, P, D, B \rangle, \quad (\text{A.17})$$

$$\mathfrak{h}_{(3)} = \langle K_0, D \rangle, \quad \mathfrak{m}_{(3)} = \langle H, P, K_1, B \rangle, \quad (\text{A.18})$$

$$\mathfrak{h}_{(4)} = \langle K_1, D \rangle, \quad \mathfrak{m}_{(4)} = \langle H, P, K_0, B \rangle. \quad (\text{A.19})$$

More generally we can have

$$\begin{aligned} \mathfrak{h}_{(5)} = \langle D, \alpha_1 K_0 + \alpha_2 K_1 \rangle, \quad \mathfrak{m}_{(5)} = \langle H, P, K_1, B \rangle, \quad \alpha_1 \neq 0, \\ \mathfrak{m}_{(5)} = \langle H, P, K_0, B \rangle, \quad \alpha_2 \neq 0, \end{aligned} \quad (\text{A.20})$$

$$\mathfrak{h}_{(6)} = \langle K_0, D + \alpha_1 K_1 \rangle, \quad \mathfrak{m}_{(6)} = \langle H, P, K_1, B \rangle, \quad (\text{A.21})$$

$$\mathfrak{h}_{(7)} = \langle K_1, D + \alpha_1 K_0 \rangle, \quad \mathfrak{m}_{(7)} = \langle H, P, K_0, B \rangle. \quad (\text{A.22})$$

(vii) $\dim \mathcal{H} = 3, \dim \mathcal{M} = 3$:

$$\mathfrak{h}_{(1)} = \langle H, D, K_0 \rangle, \quad \mathfrak{m}_{(1)} = \langle P, K_1, B \rangle, \quad (\text{A.23})$$

$$\mathfrak{h}_{(2)} = \langle K_0, K_1, D \rangle, \quad \mathfrak{m}_{(2)} = \langle H, P, B \rangle. \quad (\text{A.24})$$

More generally we can also have

$$\mathfrak{h}_{(3)} = \langle K_0, D + \alpha_1 K_1, \beta_1 K_0 + \beta_2 K_1 \rangle, \quad \mathfrak{m} = \langle H, P, B \rangle. \quad (\text{A.25})$$

For the sake of completeness, below we list the possible sub-algebras relaxing both the conditions (1) and (2):

(i) $\dim \mathcal{H} = 1, \dim \mathcal{M} = 5$:

$$\mathfrak{h} = \langle P \rangle. \quad (\text{A.26})$$

(ii) $\dim \mathcal{H} = 2, \dim \mathcal{M} = 4$:

$$\mathfrak{h} = \langle P, B \rangle, \quad \langle P, K_1 \rangle, \quad \langle P, H \rangle, \quad \langle P, D \rangle. \quad (\text{A.27})$$

(iii) $\dim \mathcal{H} = 3, \dim \mathcal{M} = 3$:

$$\mathfrak{h} = \langle P, B, K_1 \rangle, \quad \langle P, B, D \rangle, \quad \langle P, B, H \rangle, \quad \langle P, K_1, D \rangle. \quad (\text{A.28})$$

References

- [1] C. R. Hagen, “Scale and conformal transformations in galilean-covariant field theory,” *Phys. Rev. D* **5**, 377 (1972).
- [2] U. Niederer, “The maximal kinematical invariance group of the free Schrodinger equation,” *Helv. Phys. Acta* **45**, 802 (1972).
- [3] Y. Nishida, D. T. Son, “Nonrelativistic conformal field theories,” *Phys. Rev.* **D76**, 086004 (2007). [arXiv:0706.3746 [hep-th]].
- [4] D. T. Son, “Toward an AdS/cold atoms correspondence: a geometric realization of the Schrodinger symmetry,” *Phys. Rev. D* **78**, 046003 (2008) [arXiv:0804.3972 [hep-th]].
- [5] K. Balasubramanian and J. McGreevy, “Gravity duals for non-relativistic CFTs,” *Phys. Rev. Lett.* **101**, 061601 (2008) [arXiv:0804.4053 [hep-th]].
- [6] S. A. Hartnoll, “Lectures on holographic methods for condensed matter physics,” *Class. Quant. Grav.* **26**, 224002 (2009) [arXiv:0903.3246 [hep-th]].

- [7] S. Kachru, X. Liu and M. Mulligan, “Gravity Duals of Lifshitz-like Fixed Points,” *Phys. Rev. D* **78**, 106005 (2008) [arXiv:0808.1725 [hep-th]].
- [8] A. Bagchi and R. Gopakumar, “Galilean Conformal Algebras and AdS/CFT,” *JHEP* **0907**, 037 (2009) [arXiv:0902.1385 [hep-th]].
- [9] M. Alishahiha, A. Davody and A. Vahedi, “On AdS/CFT of Galilean Conformal Field Theories,” arXiv:0903.3953 [hep-th].
- [10] A. Bagchi and I. Mandal, “On Representations and Correlation Functions of Galilean Conformal Algebras,” *Phys. Lett. B* **675**, 393 (2009) [arXiv:0903.4524 [hep-th]].
- [11] D. Martelli and Y. Tachikawa, “Comments on Galilean conformal field theories and their geometric realization,” arXiv:0903.5184 [hep-th].
- [12] A. Bagchi and I. Mandal, “Supersymmetric Extension of Galilean Conformal Algebras,” *Phys. Rev. D* **80**, 086011 (2009) [arXiv:0905.0580 [hep-th]];
- J. A. de Azcarraga and J. Lukierski, “Galilean Superconformal Symmetries,” *Phys. Lett. B* **678**, 411 (2009) [arXiv:0905.0141 [math-ph]];
- M. Sakaguchi, “Super Galilean conformal algebra in AdS/CFT,” arXiv:0905.0188 [hep-th];
- A. Mukhopadhyay, “A Covariant Form of the Navier-Stokes Equation for the Galilean Conformal Algebra,” arXiv:0908.0797 [hep-th];
- A. Hosseiny and S. Rouhani, “Affine Extension of Galilean Conformal Algebra in 2+1 Dimensions,” arXiv:0909.1203 [hep-th];
- K. Hotta, T. Kubota and T. Nishinaka, “Galilean Conformal Algebra in Two Dimensions and Cosmological Topologically Massive Gravity,” arXiv:1003.1203 [hep-th].
- [13] A. Bagchi, “The BMS/GCA correspondence,” arXiv:1006.3354 [hep-th];
- A. Bagchi, “Correspondence between asymptotically flat spacetimes and non-relativistic CFTs,” *Phys. Rev. Lett.* **105**, 171601 (2010).
- [14] H. Bondi, M. G. van der Burg, and A. W. Metzner, “Gravitational waves in general relativity. 7. Waves from axisymmetric isolated systems,” *Proc. Roy. Soc. Lond. A* **269** (1962) 21.
- R. K. Sachs, “Gravitational waves in general relativity. 8. Waves in asymptotically flat space-times,” *Proc. Roy. Soc. Lond. A* **270** (1962) 103.
- R. K. Sachs, “Asymptotic symmetries in gravitational theory,” *Phys. Rev.* **128** (1962) 2851.
- [15] G. Barnich and G. Compere, “Classical central extension for asymptotic symmetries at null infinity in three spacetime dimensions,” *Class. Quant. Grav.* **24**, F15 (2007) [Erratum-ibid. **24**, 3139 (2007)] [arXiv:gr-qc/0610130].

- [16] S. Schafer-Nameki, M. Yamazaki and K. Yoshida, “Coset Construction for Duals of Non-relativistic CFTs,” JHEP **0905**, 038 (2009) [arXiv:0903.4245 [hep-th]].
- [17] J. I. Jottar, R. G. Leigh, D. Minic and L. A. P. Zayas, “Aging and Holography,” arXiv:1004.3752 [hep-th].
- [18] C. Duval and P. A. Horvathy, “Non-relativistic conformal symmetries and Newton-Cartan structures,” J. Phys. A **42**, 465206 (2009) [arXiv:0904.0531 [math-ph]].
- [19] J. Lukierski, P. C. Stichel and W. J. Zakrzewski, “Exotic Galilean conformal symmetry and its dynamical realisations,” Phys. Lett. A **357**, 1 (2006) [arXiv:hep-th/0511259];
J. Gomis, J. Gomis and K. Kamimura, “Non-relativistic superstrings: A new soluble sector of AdS(5) x S**5,” JHEP **0512**, 024 (2005) [arXiv:hep-th/0507036].
- [20] A. Bagchi, R. Gopakumar, I. Mandal and A. Miwa, “GCA in 2d,” JHEP **1008**, 004 (2010). [arXiv:0912.1090 [hep-th]].
- [21] S. Kobayashi and K. Nomizu, “Foundations of Differential Geometry, Vol 1,” Wiley Classics Library. John Wiley & Sons Inc., New York, 1966
- [22] P. Kraus, “Lectures on black holes and the AdS(3)/CFT(2) correspondence,” Lect. Notes Phys. **755**, 193 (2008) [arXiv:hep-th/0609074].
- [23] C. R. Nappi, E. Witten, “A WZW model based on a nonsemisimple group,” Phys. Rev. Lett. **71**, 3751-3753 (1993). [hep-th/9310112].
- [24] K. Sfetsos, “Gauging a nonsemisimple WZW model,” Phys. Lett. B **324**, 335 (1994) [arXiv:hep-th/9311010].
- [25] J. M. Figueroa-O’Farrill, S. Stanciu, “Nonsemisimple Sugawara constructions,” Phys. Lett. **B327**, 40-46 (1994). [hep-th/9402035].
- [26] M. Henkel, R. Schott, S. Stoimenov, and J. Unterberger, “The Poincare algebra in the context of ageing systems: Lie structure, representations, Appell systems and coherent states,” arXiv:math-ph/0601028.
- [27] G. Compere, S. de Buyl, S. Detournay *et al.*, “Asymptotic symmetries of Schrodinger spacetimes,” JHEP **0910**, 032 (2009). [arXiv:0908.1402 [hep-th]].
- [28] G. Barnich, F. Brandt, “Covariant theory of asymptotic symmetries, conservation laws and central charges,” Nucl. Phys. **B633**, 3-82 (2002). [hep-th/0111246].
- [29] I. Mandal, “Supersymmetric Extension of GCA in 2d,” [arXiv:1003.0209 [hep-th]].